

ON GEOMETRIC GRAPH RAMSEY NUMBERS

GYULA KÁROLYI¹ Department of Algebra and Number Theory, Eötvös University, Pázmány P. sétány 1/C, Budapest, H-1117 Hungary
VERA ROSTA Alfréd Rényi Institute of Mathematics, Reáltanoda utca 13-15, Budapest, H-1053 Hungary

ABSTRACT. For any two-colouring of the segments determined by $3n - 3$ points in general position in the plane, either the first colour class contains a triangle, or there is a noncrossing cycle of length n in the second colour class, and this result is tight. We also give a series of more general estimates on off-diagonal geometric graph Ramsey numbers in the same spirit. Finally we investigate the existence of large noncrossing monochromatic matchings in multicoloured geometric graphs.

1. INTRODUCTION

For any finite sequence G_1, G_2, \dots, G_t of simple graphs, $R(G_1, G_2, \dots, G_t)$ denotes the smallest integer r with the property that whenever the *edges* of a complete graph on at least r vertices are partitioned into t colour classes, there is an integer $1 \leq i \leq t$ such that the i th colour class contains a subgraph isomorphic to G_i . Such a subgraph will be referred to as a monochromatic subgraph in the i th colour.

¹Research partially supported by Hungarian Scientific Research Grant OTKA T043631.

In the special case, when each $G_i = K_{k_i}$ is a complete graph on k_i vertices, we will simply write $R(k_1, k_2, \dots, k_t)$ for $R(G_1, G_2, \dots, G_t)$. In general, if G_i has k_i vertices, then the existence of $R(G_1, G_2, \dots, G_t)$ follows directly from that of $R(k_1, k_2, \dots, k_t)$, the latter was first observed and applied to formal logic by Ramsey [13]. For more on Ramsey theory in general, we refer to the monograph [6] and the collection of survey articles [12, 14].

A *geometric graph* is a graph drawn in the plane so that every vertex corresponds to a point, and every edge is a closed straight-line segment connecting two vertices but not passing through a third. The $\binom{n}{2}$ segments determined by n points in the plane, no three of which are collinear, form a *complete* geometric graph with n vertices. A geometric graph is *convex* if its vertices correspond to those of a convex polygon. Further, we say that a subgraph of a geometric graph is *non-crossing*, if no two of its edges have an interior point in common.

For a sequence of graphs G_1, G_2, \dots, G_t , the *geometric* Ramsey number that we denote by $R_g(G_1, G_2, \dots, G_t)$ is defined as the smallest integer r with the property that whenever the edges of a complete geometric graph on at least r vertices are partitioned into t colour classes, the i th colour class contains a non-crossing copy of G_i , for some $1 \leq i \leq t$. The number $R_c(G_1, G_2, \dots, G_t)$ denotes the corresponding number if we restrict our attention to *convex* geometric graphs only. These concepts have been introduced by Károlyi, Pach and Tóth in [9] and further explored in [10] and [11].

These numbers exist if and only if each graph G_i is *outerplanar*, that is, can be obtained as a subgraph of a triangulated cycle (convex n -gon triangulated by non-crossing diagonals). The necessity of the condition is obvious, whereas the ‘if part’ is implied by the following theorem, based on a result of Gritzmann et al. [7].

Theorem 1. *Let, for each $1 \leq i \leq t$, G_i denote an outerplanar graph on k_i vertices. Then*

$$R(G_1, \dots, G_t) \leq R_c(G_1, \dots, G_t) \leq R_g(G_1, \dots, G_t) \leq R(k_1, \dots, k_t).$$

Proof. Only the last inequality needs verification. For that end, consider any complete geometric graph G on $R(k_1, k_2, \dots, k_t)$ vertices whose edges are coloured using t different colours. Then there is an integer $1 \leq i \leq t$ such that G contains a

monochromatic complete geometric graph H_i on k_i vertices in the i th colour. The fact that H_i contains a noncrossing subgraph isomorphic to G_i follows directly from the following remarkable property of outerplanar graphs:

Theorem 2 ([7]). *Let P be an arbitrary set of n points in the plane in general position. For any outerplanar graph H on n vertices, there is a straight-line embedding f of H into the plane such that the vertex set of $f(H)$ is P and no two edges of $f(H)$ cross each other.*

Throughout this paper we use the following notation. We denote by C_k a cycle of k vertices, D_k a cycle of k vertices triangulated from a vertex, P_k a path of k vertices, and S_k a star of k vertices. In addition, $M_{2k} = kP_2$ will stand for any perfect matching on $2k$ vertices. Results on geometric Ramsey numbers for paths and cycles were found by Károlyi et al. [10] and were extended in [11]. For example, if $k \geq 3$, then

$$2k - 3 = R_c(P_k, P_k) \leq R_g(P_k, P_k) = O(k^{3/2}).$$

Moreover, if k and ℓ are integers larger than 2, then

$$\begin{aligned} (k-1)(\ell-1) + 1 &= R_c(C_k, P_\ell) = R_g(C_k, P_\ell) \leq R_g(C_k, C_\ell) \\ &\leq (k-1)(\ell-2) + (k-2)(\ell-1) + 2. \end{aligned}$$

For points in convex position, a similar result was obtained independently by Harborth and Lefmann [8]. We will consider a few ramifications in the following section, which in turn complete the solution of a problem studied in [2]. In Section 3 we prove that in the case $k = 3$ the upper bound in the above estimate is sharp:

Theorem 3. *For any integer $\ell \geq 3$,*

$$R_c(C_3, C_\ell) = R_g(C_3, C_\ell) = R_g(D_3, D_\ell) = 3\ell - 3.$$

Note that the only further example when the exact value of either $R_c(C_k, C_\ell)$ or $R_g(C_k, C_\ell)$ is known is in the case $k = \ell = 4$, namely $R_c(C_4, C_4) = R_g(C_4, C_4) = 14$, see [2, 10]. It would be very interesting to see if the upper estimate in the above inequality is tight in general.

Finally we study the existence of large monochromatic matchings in multicoloured complete graphs. It is proved in [4] that

$$R(M_{2k_1}, M_{2k_2}, \dots, M_{2k_t}) = \sum_{i=1}^t k_i + \max_{1 \leq i \leq t} k_i - t + 1.$$

This result, in the case $t = 2$ has been extended to geometric graphs as follows:

Theorem 4 ([9]).

$$R_c(M_{2k}, M_{2\ell}) = R_g(M_{2k}, M_{2\ell}) = R(M_{2k}, M_{2\ell}) = k + \ell + \max\{k, \ell\} - 1.$$

It implies the following general upper bound for

$$R_g^{(t)}(M_{2k}) = R_g(\underbrace{M_{2k}, \dots, M_{2k}}_{t \text{ times}}).$$

Theorem 5.

$$R_g^{(t)}(M_{2k}) \leq \begin{cases} \frac{3t}{2}k - \frac{3t}{2} + 2 & \text{for } t \text{ even,} \\ \frac{3t+1}{2}k - \frac{3t+1}{2} + 2 & \text{for } t \text{ odd.} \end{cases}$$

Proof. The result is immediate for $t = 1$, and is true also for $t = 2$ in view of Theorem 4. Assume thus first that $t = 2s \geq 4$ is even. Let G be any t -coloured complete geometric graph on $N \geq \frac{3t}{2}k - \frac{3t}{2} + 2$ vertices. Recolouring the edges of G such that the edges coloured using any of the first s colours are now coloured with a first new colour and the remaining edges with another new colour, we find that one of the new colour classes contains $s(k-1) + 1$ pairwise disjoint edges, due to the fact that

$$N \geq 3sk - 3s + 2 = 3(s(k-1) + 1) - 1 = R_g(M_{2(s(k-1)+1)}, M_{2(s(k-1)+1)}).$$

It follows that one of the s original colour classes that combine to this new colour class contains k pairwise disjoint edges. Similarly, for $t = 2s + 1 \geq 3$, we have

$$R_g^{(t)}(M_{2k}) \leq R_g(M_{2(s(k-1)+1)}, M_{2((s+1)(k-1)+1)}) = \frac{3t+1}{2}k - \frac{3t+1}{2} + 2.$$

□

In particular, $R_g(M_{2k}, M_{2k}, M_{2k}) \leq 5k-3$ and $R_g(M_{2k}, M_{2k}, M_{2k}, M_{2k}) \leq 6k-4$.

In Section 4 we will prove that the latter bound is sharp:

Theorem 6. *For every positive integer k we have*

$$R_c(M_{2k}, M_{2k}, M_{2k}, M_{2k}) = R_g(M_{2k}, M_{2k}, M_{2k}, M_{2k}) = 6k - 4.$$

Combining this result with the obvious inequality

$$(1) \quad R_c^{(t+1)}(M_{2k}) \geq R_c^{(t)}(M_{2k}) + (k - 1)$$

we obtain the following general lower bound:

Theorem 7. *If $t \geq 4$ and k are positive integers, then $R_c^{(t)}(M_{2k}) \geq (t + 2)k - t$.*

We give the following essential improvement upon this result.

Theorem 8. *For arbitrary integers $t \geq 2$ and $k \geq 6t - 10$ we have*

$$R_c^{(t)}(M_{2k}) > \frac{6}{5}tk.$$

In [1], Araujo et al. studied the chromatic number of some geometric Kneser graphs. For example, given n points in convex position in the plane, let G_n be the graph whose vertices are the $\binom{n}{2}$ line segments determined by the points, two such vertices connected by an edge in G_n if and only if the corresponding line segments are disjoint.

Theorem 9 ([1]).

$$2 \left\lfloor \frac{n+1}{3} \right\rfloor - 1 \leq \chi(G_n) \leq \min \left\{ n - 2, n - \frac{\log n}{2} \right\}.$$

In this result, the lower bound is derived as a consequence of Theorem 4. Note that any improvement upon the upper bound in Theorem 5, even when R_g is replaced by R_c , would have an impact on the lower bound in the above theorem. However, Theorem 8 shows that no improvement beyond $\frac{5}{6}n$ could be obtained this way.

2. GENERAL ESTIMATES

In this section we apply the methods of [10, 11] to obtain a few general estimates concerning asymmetric geometric graph Ramsey numbers.

Theorem 10. *If $k, \ell \geq 2$ and H is an outerplanar graph with ℓ vertices, then*

- (i) $R_g(S_k, H) \leq (k-1)(\ell-1) + 1$ and
- (ii) $R_g(P_k, H) \leq (k-1)(\ell-1) + 1$.

Proof. The proof of the first inequality depends on a result of Chvátal [3] claiming that $R(T_k, K_\ell) = (k-1)(\ell-1) + 1$ for every fixed tree T_k on k vertices. Coupled with Theorem 2 it implies that any complete geometric graph with at least $(k-1)(\ell-1) + 1$ vertices whose edges are coloured with red and blue contains either a blue noncrossing copy of H or a red copy of S_k whose edges cannot cross anyway.

To prove the second statement, consider a complete geometric graph on at least $(k-1)(\ell-1) + 1$ vertices whose edges are coloured with red and blue. Let $p_1, p_2, \dots, p_{(k-1)(\ell-1)+1}$ be vertices of this graph, listed in increasing order of their x -coordinates, which we may assume to be all distinct. A path $p_{i_1} p_{i_2} \dots p_{i_j}$ is said to be *monotone* if $i_1 < i_2 < \dots < i_j$. Introduce a partial ordering on these vertices as follows. Let $p_i < p_j$ if and only if $i < j$, and there is a monotone red path connecting p_i to p_j . By Dilworth's theorem [5], one can find either k elements that form a totally ordered subset $Q \subseteq \{p_1, p_2, \dots, p_{(k-1)(\ell-1)+1}\}$, or ℓ elements that are pairwise incomparable. In the first case there is a monotone red path visiting every vertex of Q , it contains a red subpath on k vertices whose edges do not cross. Note that any two incomparable elements are connected by a blue edge. Thus one finds a complete blue subgraph on ℓ vertices in the second case. According to Theorem 2 it contains a noncrossing copy of H . \square

Theorem 11. *Let G be a connected outerplanar graph with $k \geq 2$ vertices and H any outerplanar graph that contains a cycle of length ℓ . Then*

$$(k-1)(\ell-1) + 1 \leq R_c(G, H).$$

Proof. Take $(k-1)(\ell-1)$ points on a circle and partition them into $\ell-1$ groups, each containing $k-1$ consecutive points. Colour with blue all edges between points in different groups, and colour with red all edges between points belonging to the same group. It follows that the two-coloured geometric graph thus obtained

does not contain a noncrossing blue copy of H . Indeed, any noncrossing blue cycle contains at most one point from each group, hence it can not have more than $\ell - 1$ points. On the other hand, all vertices of a red connected subgraph must be from the same group, so there is no such graph with more than $k - 1$ points. \square

Putting the lower and upper bounds together we obtain the following result.

Corollary 12. *Let $k \geq 2$ be an integer and H any outerplanar graph on ℓ vertices that contains a cycle of length ℓ . Then*

$$R_c(S_k, H) = R_c(P_k, H) = R_g(S_k, H) = R_g(P_k, H) = (k - 1)(\ell - 1) + 1.$$

In fact, the following general statement may be true.

Conjecture 13. *Let T_k denote a tree of $k \geq 2$ vertices and H any outerplanar graph on ℓ vertices that contains a cycle of length ℓ . Then*

$$R_c(T_k, H) = R_g(T_k, H) = (k - 1)(\ell - 1) + 1.$$

The lower bound follows directly from Theorem 11.

Finally, let G and H denote arbitrary outerplanar graphs on k and ℓ vertices, respectively. According to Theorem 1,

$$R_g(G, H) \leq R(k, \ell) \leq \binom{k + \ell - 2}{\ell - 1}.$$

We do not know, if this upper bound can be essentially reduced in general. It is the case, however, if each graph is part of a cycle triangulated from one vertex.

Theorem 14. *For arbitrary integers $k, \ell \geq 3$ we have*

$$R_g(D_k, D_\ell) \leq (k - 2)(\ell - 1) + (k - 1)(\ell - 2) + 2.$$

Proof. Let P denote the vertex set of a complete geometric graph of at least $(k - 2)(\ell - 1) + (k - 1)(\ell - 2) + 2$ vertices, whose edges are coloured with red and blue. Let p be a vertex of the convex hull of P . Consider the edges incident to p , either at least $(k - 2)(\ell - 1) + 1$ of them are red, or at least $(k - 1)(\ell - 2) + 1$ of them are blue. Suppose, without any loss of generality, that the first possibility is the case. Let $p_1, p_2, \dots, p_{(k-2)(\ell-1)+1}$ be vertices of P , listed in clockwise order of visibility from p , such that each edge pp_i is red. We say that a path $p_{i_1}p_{i_2} \dots p_{i_j}$ is monotone if $i_1 < i_2 < \dots < i_j$. Define a partial ordering on the vertices

$p_1, p_2, \dots, p_{(k-2)(\ell-1)+1}$ as follows. Let $p_i < p_j$ if and only if $i < j$, and there is a monotone red path connecting p_i to p_j . Applying Dilworth's theorem again, there are either $k-1$ elements that form a linearly ordered subset, or ℓ elements that are pairwise incomparable. In the first case there is a monotone red path $q_1 q_2 \dots q_{k-1}$, and we can complete it to a noncrossing red cycle $p q_1 q_2 \dots q_{k-1} p$ of length k , triangulated from the point p . In the second case there is a complete blue subgraph on ℓ vertices, and it follows from Theorem 2 that it contains a blue noncrossing copy of D_ℓ . \square

For any convex drawing D of a given graph, let $r_c(D)$ denote the smallest integer r such that every two-colouring of the diagonals and sides of a convex r -gon contains a monochromatic copy of D . In [2], Bialostocki and Harborth determined $r_c(D)$ for all convex drawings D of graphs with 4 vertices, except one. What they proved about that exceptional case can be rephrased as $14 \leq R_c(D_4, D_4) \leq 16$. From Theorem 14 it follows that

$$R_c(D_4, D_4) = R_g(D_4, D_4) = 14,$$

thus completing the investigation initiated in [2].

3. NONCROSSING CYCLES VERSUS TRIANGLES

Proof of Theorem 3. It follows from Theorem 14 that $R_c(C_3, C_\ell) \leq R_g(C_3, C_\ell) \leq R_g(D_3, D_\ell) \leq 3\ell - 3$. In order to prove that $R_c(C_3, C_\ell) \geq 3\ell - 3$, we construct a two-coloured geometric graph with $3\ell - 4$ vertices in convex position which does not have a monochromatic red triangle, nor a blue noncrossing cycle of ℓ vertices. According to the parity of ℓ , we distinguish between two cases.

Assume first that ℓ is odd. Put $N = 3\ell - 4$ and let $x_1, x_2, \dots, x_{3\ell-4}$ be the vertices of a convex N -gon P , listed in clockwise order. For an integer n , let n^* denote the integer between 0 and $N - 1$ obtained reducing n modulo N . Colour the edge $x_i x_j$ red if $\min\{(j - i)^*, (i - j)^*\} = 2s - 1$ for some integer s that satisfies $1 \leq 2s - 1 \leq \ell - 2$, and colour it blue otherwise. This way we obtain a well defined two-colouring of the complete convex geometric graph G whose vertices are $x_1, x_2, \dots, x_{3\ell-4}$. If x_i and x_j are consecutive vertices in the clockwise order of some convex polygon Q whose vertices are among the vertices of P , then we say that the (combinatorial) *length* of the side $x_i x_j$ is $(j - i)^*$.

We say that a side is *short* if its length does not exceed $\ell - 2$, it is called *long* otherwise. Note that all short red edges are of odd length and all short blue edges are of even length.

Suppose that x_i, x_j, x_k are the vertices of a triangle in the clockwise order. Since $3(\ell - 2) < 3\ell - 4$, it cannot be that all edges are short. If $x_k x_i$ is a long red edge, then $(k - i)^*$ is an odd integer not exceeding $\ell - 2$. Then both edges $x_i x_j$ and $x_j x_k$ must be short and one of them has an even length, which implies it cannot be red. Consequently, G does not contain a red triangle.

Next let $x_{i_1} x_{i_2}, \dots, x_{i_\ell}$ be the vertices of a noncrossing cycle of length ℓ . If it has a long edge, then the sum of the lengths of the remaining edges is at most $(3\ell - 4) - (\ell - 1) = 2\ell - 3$, thus one of the remaining edges is of length 1, hence cannot be blue. On the other hand, if all edges are short, then since the sum of their lengths, $3\ell - 4$ is an odd number, one of them must have an odd length and thus cannot be blue. All in all, G contains no non-crossing blue C_ℓ either.

The case when ℓ is even is slightly more complicated. First we construct a two-colouring of a complete convex geometric graph G' on $N = 3\ell - 3$ vertices, obtained by inserting an auxiliary vertex x_0 between $x_{3\ell-4}$ and x_1 . Using the terminology of the previous case, we colour all short edges of odd length red. This time we will call *medium* edges those whose length is $\ell - 1$, and the following medium edges we also color red: $x_{2i+1} x_{2i+\ell}$ for $0 \leq i \leq \ell - 2$ and $x_{i^*} x_{(i+\ell-1)^*}$ for $2\ell - 2 \leq i \leq 3\ell - 3$. The other edges are coloured blue. Note that medium edges also have an odd length. This time an edge is called long if its length is at least ℓ .

Assume that G' contains a red triangle Δ . The same argument we had in the previous case confirms that it cannot have a long edge. Thus all edges are of length at most $\ell - 1$ and these 3 lengths add up to $N = 3\ell - 3$. Thus all edges of Δ are medium. Δ then must have an edge $x_i x_{i+\ell-1}$ for some $1 \leq i \leq \ell - 1$ in which case i is odd and $i + \ell - 1 \leq 2\ell - 2$ is even. If $i + \ell - 1 < 2\ell - 2$, then the edge $x_{i+\ell-1} x_{i+2\ell-2}$ is not red. It follows that the vertices of Δ are $x_{\ell-1}, x_{2\ell-2}$ and x_0 .

Suppose now that C is a noncrossing blue cycle of length ℓ in G' . As in the case when ℓ was odd, we can argue that C cannot have only short edges, by parity reason. Also, every short edge has a length at least 2, thus if it has a non-short edge, then an elementary calculation implies that C has exactly one medium edge

and $\ell - 1$ other edges, each of length 2. It follows from the construction that the endpoints of the blue medium edge are x_{2i} and $x_{2i+\ell-1}$ for some $1 \leq i \leq \ell - 2$. Accordingly, one vertex of C must be x_0 .

Finally, let G be the two-coloured convex geometric graph induced by G' on the vertex set $x_1, x_2, \dots, x_{3\ell-4}$. Were Δ a red triangle in G , it also would be a red triangle in G' , thus it would necessarily contain the vertex x_0 , a contradiction. A similar argument shows that G does not contain a noncrossing blue C_ℓ either. \square

4. NONCROSSING MATCHINGS IN 4-COLOURED GEOMETRIC GRAPHS

Proof of Theorem 6. In view of Theorems 1 and 5, we only have to present a four-colouring of the edges of a complete convex geometric graph on $6k - 5$ vertices such that there are no k (geometrically) disjoint edges of the same colour. Let the vertices of the graph be points on a circle, listed in clockwise order as

$$b_1, \dots, b_{k-1}, r_1, r'_1, \dots, r_{k-1}, r'_{k-1}, g_1, \dots, g_{k-1}, p_k, p'_{k-1}, p_{k-1}, \dots, p'_1, p_1.$$

For simplicity, call the first $k - 1$ vertices blue, the next $2k - 2$ vertices red, the $k - 1$ vertices that follow green, and the last $2k - 1$ vertices purple. We colour the edges of the corresponding complete convex geometric graph using the same set of colours as follows.

First, use red to colour each edge connecting two red vertices, and in addition for each edge that connects a red vertex r'_i to any other vertex. Similarly, every edge between two purple points are coloured purple as well as any edge starting at some vertex p'_i ($1 \leq i \leq k - 1$). Edges of the form $r'_i p'_j$ may be coloured either purple or red. Next, assign blue to each yet uncoloured edge starting at a blue vertex, and similarly, use the green colour for each edge connecting some green vertex to any other vertex except the vertices r'_i, p'_i . Again, edges of the form $b_i g_j$ may be either blue or green. Finally, we colour the edge $p_i r_j$ blue if and only if $i + j \leq k$, else we colour it green. This way we have assigned a colour to each edge of the complete graph.

This graph does not contain k pairwise disjoint purple edges, as it follows directly from the following lemma.

Lemma 15. *Assume that $P = p_1 p'_1 p_2 p'_2 \dots p_n$ is a convex polygon, and $Q = \{q_1, q_2, \dots, q_m\}$ is a set of points lying in the interior of the convex (finite or*

infinite) domain bounded by the line segment p_1p_n , the extension of the line segment $p_1p'_1$ beyond the point p_1 , and the extension of the line segment $p'_{n-1}p_n$ beyond p_n . Consider all the line segments that are either connecting two vertices of P , or connect some point p'_i to a point q_j . Then no n of these line segments can be pairwise disjoint.




Figure 1

Proof. We proceed by induction on n . The claim being obvious for $n = 2$, we assume that $n > 2$, and that we have already proved the statement for smaller values of n . Consider a set S of ℓ pairwise disjoint segments. If each of them has an endpoint in the set $\{p'_1, p'_2, \dots, p'_{n-1}\}$, then clearly $\ell \leq n - 1$. Otherwise one of the segments is of the form $s = p_i p_j$ with $1 \leq i < j \leq n$. Let S_0 be the set of all line segments in S whose both endpoints belong to the set $\{p_i, p'_i, \dots, p_j\}$. Obviously, $s \in S_0$, and $|S_0| \leq j - i$. If $i = 1$ and $j = n$, then clearly $S = S_0$, and we are done. If $i > 1$ but $j = n$, then $S' = S \setminus S_0$ is a set of pairwise disjoint line segments in the configuration determined by the vertices of $P' = p_1 p'_1 p_2 p'_2 \dots p_i$ and the points of $Q' = Q$, just as in the lemma. Since certainly $2 \leq i \leq n - 1$ in this case, it follows from the induction hypothesis that $\ell - |S_0| = |S'| < i$, that is, $\ell < i + |S_0| \leq n$. We can argue similarly if $i = 1$ and $j < n$. Finally, if $1 < i < j < n$, then we may consider $P' = p_1 p'_1 \dots p_i p'_i p_{j+1} p'_{j+1} \dots p_n$, $Q' = Q$ and $S' = S \setminus S_0$ to find, based on the induction hypothesis, that $|S'| < n - (j - i)$. It follows, that

$$|S| = |S'| + |S_0| < (n - j + i) + (j - i) = n,$$

completing the induction step. \square

Similarly, the graph does not contain a noncrossing red M_{2k} either.

Assume next that the graph contains n pairwise disjoint blue edges, and consider only these edges. If they all have a blue endpoint, then clearly $n \leq k - 1$. Otherwise we assume that there are $\ell \geq 1$ edges of the form $p_i r_j$. Since they do not cross, we can list them as $p_{i_1} r_{j_1}, \dots, p_{i_\ell} r_{j_\ell}$ so that $i_1 < \dots < i_\ell$ and $j_1 < \dots < j_\ell$. It follows that

$$i_1 + j_1 + 2(\ell - 1) \leq i_\ell + j_\ell \leq k.$$

The endpoints of the other $n - \ell$ edges must lie in the vertex set

$$\{b_1, b_2, \dots, b_{k-1}, p_1, p_2, \dots, p_{i_1-1}, r_1, r_2, \dots, r_{j_1-1}\}.$$

Consequently,

$$2(n - \ell) \leq (k - 1) + (i_1 - 1) + (j_1 - 1) \leq (k - 1) + k - 2(\ell - 1) - 2 = 2(k - \ell) - 1.$$

It follows that $n < k$ holds in this case, too.

Due to the symmetry in the construction, the existence of a noncrossing green matching on $2k$ vertices can be excluded by a similar argument. \square

The proof of Theorem 8 depends on the following lemma that we found in the pursuit of a possible generalization of the previous arguments.

Lemma 16. *Let r denote any positive integer. If $k \geq 4r$, then the edges of a complete convex geometric graph on $n = (6r + 6)k - (12r + 12)$ vertices can be coloured with $t = 5r + 4$ colours so that it does not contain a monochromatic crossing-free matching of size k .*

Indeed, Theorem 4 and the inequality (1) implies

$$R_c^{(2)}(M_{2k}) = 3k - 1 > \frac{12}{5}k$$

for $k \geq 2$ and

$$R_c^{(3)}(M_{2k}) \geq 4k - 2 > \frac{18}{5}k$$

for $k \geq 8$ (in fact, even for $k \geq 6$). For the values $4 \leq t \leq 8$, the statement of Theorem 8 follows directly from Theorem 7. Finally if $t = 5r + 4 + i$ for some integers $r \geq 1$ and $0 \leq i \leq 4$, then Lemma 16 coupled with relation (1) implies

$$R_c^{(t)}(M_{2k}) \geq (6r + 6 + i)k - (12r + 11 + i) > \frac{6}{5}kt$$

for any $k \geq 6t - 10$. It only remains to prove the lemma.

Proof of Lemma 16. Choose n points on a circle and partition them into $4r + 4$ sets of consecutive points

$$B_i = \{b_{i1}, b_{i2}, \dots, b_{i,k-1}\}, \quad R_i = \{r_{i1}, r'_{i1}, r_{i2}, r'_{i2}, \dots, r_{i,k-2}\}$$

for $1 \leq i \leq 2r + 2$. We choose each numbering in clockwise direction. Thus $|B_i| = k - 1$ and $|R_i| = 2k - 5$. There will be $4r + 4$ corresponding colours that we denote by \mathcal{B}_i and \mathcal{R}_i , respectively.

First, use the colour \mathcal{R}_i for any edge connecting two vertices in R_i , and also for any edge starting at some vertex r'_{ij} , where $1 \leq j \leq k - 3$. Edges of the form $r'_{ij}r'_{uv}$ may belong to either \mathcal{R}_i or \mathcal{R}_u . It follows from Lemma 15 that the colour class \mathcal{R}_i contains at most $k - 3$ pairwise disjoint edges.

Next, we use the colour \mathcal{B}_i to colour each, so far uncoloured edge that has a vertex in B_i . Edges of the form $b_{ij}b_{uv}$ may belong to either \mathcal{B}_i or \mathcal{B}_u . In addition, we assign the colour \mathcal{B}_i to each edge of the form $r_{i^*j}r_{ij'}$ whenever $j' \leq j$ and the number $1 \leq i^* \leq 2r + 2$ satisfies $i^* + 1 \equiv i \pmod{2r + 2}$. As in the proof of Theorem 6, it follows that the colour class \mathcal{B}_i cannot contain k pairwise disjoint edges either.

Finally, we use r further colours $\mathcal{C}_1, \dots, \mathcal{C}_r$ to colour the remaining edges, all of which are of the form $r_{ij}r_{uv}$ for some $i \neq u$. We have $(2r + 2)(k - 2)$ points

$$r_{11}, r_{12}, \dots, r_{1,k-2}, r_{21}, \dots, r_{2,k-2}, r_{2r+2,1}, \dots, r_{2r+2,k-2}$$

positioned along a circle K . Without any loss of generality, we may assume that these are the vertices of a regular polygon $x_1x_2 \dots x_{(2r+2)(k-2)}$ whose center we denote by O . For $2 \leq \ell \leq (r + 1)(k - 2)$, we say that the diagonal x_ix_j is of length ℓ if $j - i \equiv \ell \pmod{(2r + 2)(k - 2)}$. In this case we write $\ell = l(x_ix_j)$. We only have to colour those diagonals whose length is at least $k - 1$.

This is done as follows. For $k - 1 \leq \ell \leq (r + 1)(k - 2)$ and $1 \leq i \leq r$, let a diagonal of length ℓ belong to \mathcal{C}_i if and only if $\ell \equiv 2i - 1 \pmod{2r}$ or $\ell \equiv 2i \pmod{2r}$. To complete the proof of Lemma 16 we only have to show that \mathcal{C}_i does not contain k pairwise disjoint diagonals for $1 \leq i \leq r$.

Thus let $1 \leq i \leq r$, and consider a set \mathcal{M} of pairwise disjoint diagonals in \mathcal{C}_i . The diagonal $e \in \mathcal{M}$ is called *extremal* if \mathcal{M} lies in a closed half plane supported by e . Denote by q the number of such extremal edges. We may suppose that $|\mathcal{M}| > 1$, and thus $q \geq 2$. For any line L through O , fix a pair of disjoint half planes H_L^+ and H_L^- supported by L so that the first one is closed and the other

is open, and introduce

$$\mathcal{H} = \{H_L^+, H_L^- \mid L \text{ is a line through } O\}.$$

For edges $e, f \in \mathcal{M}$, denote by K_{ef} the closed region that e and f cut out of the circle K , it degenerates to a segment when $e = f$. We say that e and f are neighbours, if K_{ef} does not contain any member of \mathcal{M} except e and f . A sequence e_1, e_2, \dots, e_u of pairwise different edges in \mathcal{M} forms a *contiguous* set if they all belong to the same half plane H for some $H \in \mathcal{H}$, and e_j and e_{j+1} are neighbours for every $1 \leq j \leq u-1$. There is a unique partition of \mathcal{M} into maximal contiguous sets $\mathcal{M}_1, \dots, \mathcal{M}_m$.

Claim 17. $m \leq 2q - 2$.

Proof. We say that set \mathcal{M}_j is extremal, if it contains an extremal edge. Since $q \geq 2$, we are done if $m \leq 2$. On the other hand, if $m \geq 2$, then each extremal set has exactly one extremal edge. For each $1 \leq j \leq m$ there is a unique pair of edges $e_j, f_j \in \mathcal{M}_j$ such that \mathcal{M}_j is contained in $K_j = K_{e_j f_j}$. If $e_j \neq f_j$, the shorter of the two diagonals will be denoted by e_j , its midpoint by E_j .

We build up a rooted tree T as follows. The root of the tree is the point O , its further vertices are E_1, \dots, E_m . If by any chance $O = E_j$ for some j (which only can happen if $|\mathcal{M}_j| = 1$), then we extend e_j to a line L and slightly move O around so that it no longer lies in H_L^+ .

First, we connect O to E_j if and only if the segment OE_j only intersects such diagonals in \mathcal{M} that belong to \mathcal{M}_j . This way we get the level one vertices of T . For any E_u not already in T there is a (unique) vertex E_j on level one such that e_j separates \mathcal{M}_u from O . For each level one vertex E_j such that \mathcal{M}_j is not extremal there are at least two different vertices E_u not yet in T such that the segment $E_j E_u$ only intersects such diagonals in \mathcal{M} that belong to \mathcal{M}_u . Connecting each such vertex E_u with E_j we arrive at the level two vertices of T . Iterating this process we arrive at the tree T rooted at O that has $m+1$ vertices, E_j is a leaf if and only if e_j is an extremal edge, and any vertex E_j which is not a leaf has degree at least 3.

Assume first that the root O has a degree at least 2. In this case \mathcal{M}_j is extremal if and only if e_j is extremal, and thus T has exactly q leaves and $m-q$ vertices different from O such that each of them has a degree at least 3. Since T

has exactly m edges, double counting yields

$$2m \geq q + 2 + 3(m - q),$$

and the result follows.

If $\deg(O) = 1$, then let E_j be the unique neighbour of O . In this case \mathcal{M}_j is also extremal, its extremal edge being f_j . This time the number of leaves is $q - 1$, thus T has q degree one vertices, and each further vertex of T has a degree at least 3. It follows that

$$2m \geq q + 3(m + 1 - q),$$

that is, $m \leq 2q - 3$. \square

Write $s_j = |\mathcal{M}_j|$, and denote by n_j the number of vertices x_u that lie in K_j .

Claim 18. *If $1 \leq j \leq m$, then $n_j \geq 2rs_j - (2r - 1)$.*

Proof. List the diagonals d_1, \dots, d_{s_j} in M_j so that $l(d_1) < \dots < l(d_{s_j})$. Note that $l(d_{u+1}) > l(d_u) + 1$ for any $1 \leq u < s_j$. For every u there is an integer k_u such that $l(d_u) = 2rk_u + m_u$, where m_u is either $2i - 1$ or $2i$. It follows that $k_{u+1} > k_u$ for every $1 \leq u < s_j$, thus $l(d_{s_j}) - l(d_1) \geq 2r(s_j - 1) - 1$ and consequently $n_j \geq 2r(s_j - 1) + 1$. \square

Since the regions K_1, K_2, \dots, K_m are pairwise disjoint, each vertex x_u can belong to at most one of them. Moreover, each extremal edge cuts off at least $k - 2$ consecutive vertices that cannot belong to any K_j . Consequently,

$$(2r + 2)(k - 2) \geq \sum_{j=1}^m n_j + q(k - 2) \geq 2r|\mathcal{M}| - m(2r - 1) + q(k - 2),$$

according to Claim 18 and the equality $|\mathcal{M}| = s_1 + \dots + s_m$. It follows that

$$\begin{aligned} 2r(k - |\mathcal{M}|) &\geq (q - 2)(k - 2) - m(2r - 1) + 4r \\ &= (q - 2)(k - 2) - (m - 2)(2r - 1) + 2 \\ &\geq (q - 2)(k - 2 - 2(2r - 1)) + 2 \\ &= (q - 2)(k - 4r) + 2 \\ &> 0. \end{aligned}$$

Thus $|\mathcal{M}| < k$, and the proof of Lemma 16 is complete. \square

REFERENCES

- [1] G. ARAUJO, A. DUMITRESCU, F. HURTADO, M. NOY, AND J. URRUTIA, On the chromatic number of some geometric Kneser graphs, *Comput. Geom. Th. Appl.* **32** (2005) 59–69
- [2] A. BIALOSTOCKI AND H. HARBORTH, Ramsey colorings for diagonals of convex polygons, *Abh. Braunschweig. Wiss. Ges.* **47** (1996) 159–163
- [3] V. CHVÁTAL, Tree – complete graph Ramsey numbers, *J. Graph Theory* **1** (1977) 93
- [4] E.J. COCKAYNE AND P.J. LORIMER, The Ramsey number for stripes, *J. Austral. Math. Soc. A* **19** (1975) 252–256
- [5] R.P. DILWORTH, A decomposition theorem for partially ordered sets, *Annals Math.* **51** (1950) 161–166
- [6] R.L. GRAHAM, B.L. ROTHCHILD, AND J.H. SPENCER, Ramsey Theory, 2nd ed., John Wiley, New York (1990)
- [7] P. GRITZMANN, B. MOHAR, J. PACH, AND R. POLLACK, Embedding a planar triangulation with vertices at specified points (solution to problem E3341), *Amer. Math. Monthly* **98** (1991) 165–166
- [8] H. HARBORTH AND H. LEFMANN, Coloring arcs of convex sets, *Discrete Math.* **220** (2000) 107–117
- [9] GY. KÁROLYI, J. PACH, AND G. TÓTH, Ramsey-type results for geometric graphs. I, *Discrete Comput. Geom.* **18** (1997) 247–255
- [10] GY. KÁROLYI, J. PACH, G. TÓTH, AND P. VALTR, Ramsey-type results for geometric graphs. II, *Discrete Comput. Geom.* **20** (1998) 375–388
- [11] GY. KÁROLYI AND V. ROSTA, Generalized and geometric Ramsey numbers, *Theoretical Comput. Sci.* **263** (2001) 87–98
- [12] J. NEŠETŘIL AND V. RÖDL (EDS.), Mathematics of Ramsey Theory, *Algorithms and Combinatorics* **5**, Springer (1990)
- [13] F.P. RAMSEY, On a problem of formal logic, *Proc. London Math. Soc.* **30** (1930) 264–286
- [14] V. ROSTA, Ramsey theory applications, *Electron. J. Combin.* **11** (2004) Dynamic Survey DS13, 43 pages (electronic)

E-mail address: karolyi@cs.elte.hu ‘‘Gyula Károlyi’’

E-mail address: rosta@renyi.hu ‘‘Vera Rosta’’