

MATHEMATICAL CENTRE TRACTS 121

---

**EIGENVALUE TECHNIQUES  
IN DESIGN  
AND GRAPH THEORY**

W.H. HAEMERS

---

MATHEMATISCH CENTRUM      AMSTERDAM 1980

---

1980 Mathematics subject classification: Primary: 05-02

Secondary: 05B05, 05B20, 05B25,  
05C15, 05C50, 15A18

---


#### ACKNOWLEDGEMENTS

I am grateful to Prof.dr. J.J. Seidel for making me familiar with the application of linear algebra to combinatorics, for stimulating me in writing this monograph and for the many useful suggestions and remarks he gave in realizing it.

I also thank Prof.dr. J.H. van Lint for several important remarks about the manuscript.

I am grateful to E. Baselmans-Weijers for her excellent typing of the manuscript in a very short period of time.

I would like to thank all others who have contributed in one way or another to the realization of the present monograph, and I thank the Mathematical Centre for the opportunity to publish it in their series Mathematical Centre Tracts.



THE SHRIKHANDE GRAPH

## CONTENTS

PREFACE	1
1. MATRICES AND EIGENVALUES	
1.1. Introduction	4
1.2. Interlacing of eigenvalues	6
1.3. More eigenvalue inequalities	10
2. INEQUALITIES FOR GRAPHS	
2.1. Induced subgraphs	15
2.2. Chromatic number	22
3. INEQUALITIES FOR DESIGNS	
3.1. Subdesigns	27
3.2. Intersection numbers	31
4. 4-COLOURABLE STRONGLY REGULAR GRAPHS	
4.1. Introduction	37
4.2. Strongly regular graphs on 40, 50, 56, 64 and 77 vertices	40
4.3. Recapitulation	48
5. GENERALIZED POLYGONS	
5.1. Introduction	50
5.2. An equality for generalized hexagons	53
5.3. Geometric and pseudo-geometric graphs for generalized polygons	60
6. CONSTRUCTIONS	
6.1. Some $2 - (71, 15, 3)$ designs	64
6.2. Some strongly regular graphs	71
APPENDIX I. GRAPHS AND DESIGNS	79
APPENDIX II. TABLES	85
REFERENCES	93
NOTATION	100
INDEX	101



## PREFACE

The application of the theory of matrices and eigenvalues to combinatorics is certainly not new. In a certain sense the study of the eigenvalues of the adjacency matrix of a graph even became a subject of its own, see for instance [B5], [C12], [C13], [H14] and [S2]. Also in the theory of designs, matrix and eigenvalue methods have often been used successfully; see for instance [C6], [C9], [H17] and [R3]. In the present monograph (which is a slightly revised version of the author's thesis) the starting point is a theorem concerning the eigenvalues of partitioned matrices. Applications of this theorem and some known matrix theorems to matrices associated to graphs or designs lead to new results, and new proofs of known results. These concern inequalities of various types, including conclusions for the case of equality. In addition we obtain guiding-principles for constructing strongly regular graphs or 2-designs. Let us give some more details.

Our theorem (Theorem 1.2.3) about eigenvalues and partitionings of matrices, which was announced in [H1], reads as follows:

**THEOREM.** *Let  $A$  be a complete hermitian  $n \times n$  matrix, partitioned into  $m^2$  block matrices, such that all diagonal block matrices are square. Let  $B$  be the  $m \times m$  matrix whose  $i, j$ -th entry equals the average row sum of the  $i, j$ -th block matrix of  $A$  for  $i, j = 1, \dots, m$ . Then the eigenvalues  $\lambda_1(A) \geq \dots \geq \lambda_n(A)$  of  $A$  and the eigenvalues  $\lambda_1(B) \geq \dots \geq \lambda_m(B)$  of  $B$  satisfy*

$$\lambda_i(A) \geq \lambda_i(B) \geq \lambda_{n-m+i}(A) \quad \text{for } i = 1, \dots, m.$$

*Moreover, if for some integer  $k$ ,  $0 \leq k \leq m$ ,  $\lambda_i(A) = \lambda_i(B)$  for  $i = 1, \dots, k$  and  $\lambda_i(B) = \lambda_{n-m+i}(A)$  for  $i = k+1, \dots, m$ , then all the block matrices of  $A$  have constant row and column sums.*

The weaker inequalities  $\lambda_1(A) \geq \lambda_i(B) \geq \lambda_n(A)$  were already observed by C.C. Sims (unpublished), and have been applied successfully by HESTENES & HIGMAN [H10], PAYNE [P4], [P6] and others. They are usually applied under the name Higman-Sims technique. Many proofs by means of this Higman-Sims technique can be shortened by use of our generalization. But, what is more important, our theorem leads to new results, which we shall indicate below.

Suppose  $G$  is a graph on  $n$  vertices, whose  $(0,1)$ -adjacency matrix has eigenvalues  $\lambda_1(G) \geq \dots \geq \lambda_n(G)$ . DELSARTE [D1] proved that, for strongly regular  $G$ , the size of any coclique (independent set) cannot be larger than  $-n\lambda_n(G) / (\lambda_1(G) - \lambda_n(G))$ . A.J. Hoffman (unpublished) proved that this bound holds for any regular graph  $G$ . Using the above theorem we prove that an upper bound for the size of a coclique in any graph  $G$  is provided by

$$-n\lambda_1(G)\lambda_n(G) / (d_{\min}^2 - \lambda_1(G)\lambda_n(G)) ,$$

where  $d_{\min}$  denotes the smallest degree in  $G$ . This generalizes Hoffman's bound, since in case of regularity  $\lambda_1(G) = d_{\min}$  holds. More generally, by use of the same methods we find bounds for the size of an induced subgraph of  $G$  in terms of the average degree of the subgraph (Theorem 2.1.2). Apart from the inequalities of Delsarte and Hoffman we also find inequalities of Bumiller, De Clerck and Payne as corollaries of our result.

By applying the generalization of the Higman-Sims technique (with  $m = 4$ ) to the adjacency matrix of the incidence graph of a design, we obtain bounds for the sizes of a subdesign in terms of the singular values of the incidence matrix (Theorem 3.1.1). For nice designs, such as 2-designs and partial geometries, this result becomes easy to apply, since then the singular values are expressible in the design parameters (for symmetric 2-designs the inequality also appeared in [H4]). Thanks to the second part of our theorem certain conclusions may be drawn easily for the case that the bounds are attained, for the graph case as well as for the design case.

We also prove results concerning the intersection numbers of designs, such as the inequalities of AGRAWAL [A1] (Theorem 3.2.1) and the results of BEKER & HAEMERS [B2] about 2- $(v,k,\lambda)$  designs with an intersection number  $k - \lambda(v - k) / (k - 1)$ .

HOFFMAN [H13] proved that the chromatic number  $\gamma(G)$  of a graph  $G$  satisfies  $\gamma(G) \geq 1 - \lambda_1(G) / \lambda_n(G)$ . To achieve this, Hoffman first proves a generalization of the inequalities of Aronszajn concerning eigenvalues of partitioned matrices. In Section 1.3 we give a new proof of these inequalities using our generalization of the Higman-Sims technique. In Section 2.2 the application of these inequalities yields a generalization of Hoffman's bound (Theorem 2.2.1). For non-trivial strongly regular graphs this leads to  $\gamma(G) \geq \max\{1 - \lambda_1(G) / \lambda_n(G), 1 - \lambda_n(G) / \lambda_2(G)\}$ . In Chapter 4 we use these bounds, and many other results from the previous chapters, in order to determine all 4-colourable strongly regular graphs. This chapter is also



meant to illustrate some applications of the results and techniques obtained in the first three chapters.

Chapter 5 is in the same spirit, but rather independent from the other ones. The main result is the inequality of HAEMERS & ROOS [H3]:  $t \leq s^3$  if  $s \neq 1$  for a generalized hexagon of order  $(s,t)$ , together with some additional regularity for the case of equality. This is proved by rather elementary eigenvalue methods. The same technique applied to generalized quadrangles of order  $(s,t)$  yields the inequality of HIGMAN [H11]:  $t \leq s^2$  if  $s \neq 1$ , the result of BOSE [B7] for the case of equality and a theorem of CAMERON, GOETHALS & SEIDEL [C5] about pseudo-geometric graphs.

Using eigenvalue methods we obtain guiding-principles for the construction of designs and graphs. In Section 6.1 we construct a  $2 - (56,12,3)$  design, for which the framework is provided by Theorem 3.2.4. This design is embeddable in a symmetric  $2 - (71,15,3)$  design. By modifying this design we obtain eight non-isomorphic  $2 - (71,15,3)$  designs. All these designs seem to be new (the construction is also published in [B2]). In Section 6.2 some ideas for the construction of strongly regular graphs are described. We construct strongly regular graphs with parameter sets  $(q^3 + q^2 + q + 1, q^2 + q, q - 1, q + 1)$  and  $(2^{3n} + 2^{2n+1}, 2^{2n} + 2^n - 1, 2^n - 2, 2^n)$  for prime power  $q$  and positive integer  $n$ . Strongly regular graphs with the first parameter set are known; however, our construction yields graphs which are not isomorphic to the known ones. The second family seems to be new. Special attention is given to strongly regular graphs with parameter set  $(40,12,2,4)$ . Several such graphs are constructed with the help of a computer.

In the first appendix we recall some basic concepts and results from the theory of graphs and designs (including finite geometries). The second appendix exhibits explicitly some of the designs and graphs constructed earlier.

CHAPTER I  
MATRICES AND EIGENVALUES

1.1. INTRODUCTION

In this chapter we shall derive some results about eigenvalues of matrices. They provide the main tools for our investigations. We shall assume familiarity with the basic concepts and results from the theory of matrices and eigenvalues. Some general references are [M3], [N1], [W5].

Let  $A$  be a square complex matrix of size  $n$ . The hermitian transpose of  $A$  will be denoted by  $A^*$ . Suppose  $A$  has  $n$  (not necessarily distinct) real eigenvalues, which for instance is the case if  $A$  is hermitian (i.e.  $A = A^*$ ). Then we shall denote these eigenvalues by

$$\lambda_1(A) \geq \dots \geq \lambda_n(A) .$$

If denoted by subscripted variables, eigenvectors will always be ordered according to the ordering of their eigenvalues. Vectors are always column vectors. The linear span of a set of vectors  $u_1, \dots, u_n$  is denoted by  $\langle u_1, \dots, u_n \rangle$ . A basic result, which is important to our purposes, is Rayley's principle (see [N1], [W5]).

1.1.1. RESULT. *Let  $A$  be a hermitian matrix of size  $n$ . For some  $i$ ,  $0 \leq i \leq n$ , let  $u_1, \dots, u_i$  be an orthonormal set of eigenvectors of  $A$  for  $\lambda_1(A), \dots, \lambda_i(A)$ . Then*

i. 
$$\lambda_i(A) \leq \frac{u^* Au}{u^* u} \quad \text{for } u \in \langle u_1, \dots, u_i \rangle, u \neq 0, i \neq 0;$$

*equality holds iff  $u$  is an eigenvector of  $A$  for  $\lambda_i(A)$ .*

ii. 
$$\lambda_{i+1}(A) \geq \frac{u^* Au}{u^* u} \quad \text{for } u \in \langle u_1, \dots, u_i \rangle^\perp, u \neq 0, i \neq n;$$

*equality holds iff  $u$  is an eigenvector of  $A$  for  $\lambda_{i+1}(A)$ .*

For the multiplicity of an eigenvalue we shall always take the geometric one, that is, the maximal number of linearly independent eigenvectors

(to be honest, this agreement is only of influence to the proof of the next lemma, because throughout the remainder of this monograph we shall only consider eigenvalues of diagonalizable matrices). Now we shall prove some easy and well known, but nevertheless useful lemmas.

1.1.2. LEMMA. Let  $M^*$  and  $N$  be complex  $m_1 \times m_2$  matrices. Put

$$A := \begin{bmatrix} 0 & N \\ M & 0 \end{bmatrix}.$$

Then the following are equivalent.

- i.  $\lambda \neq 0$  is an eigenvalue of  $A$  of multiplicity  $f$ ;
- ii.  $-\lambda \neq 0$  is an eigenvalue of  $A$  of multiplicity  $f$ ;
- iii.  $\lambda^2 \neq 0$  is an eigenvalue of  $MN$  of multiplicity  $f$ ;
- iv.  $\lambda^2 \neq 0$  is an eigenvalue of  $NM$  of multiplicity  $f$ .

PROOF.

1. (i)  $\iff$  (ii): let  $AU = \lambda U$  for some matrix  $U$  of rank  $f$ . Write

$$U = \begin{bmatrix} U_1 \\ U_2 \end{bmatrix}, \quad \text{and define } \tilde{U} := \begin{bmatrix} U_1 \\ -U_2 \end{bmatrix},$$

where  $U_i$  has  $m_i$  rows for  $i = 1, 2$ . Then  $NU_2 = \lambda U_1$  and  $MU_1 = \lambda U_2$ . This implies  $A\tilde{U} = -\lambda\tilde{U}$ . Since  $\text{rank } U = \text{rank } \tilde{U}$ , the first equivalence is proved.

2. (iii)  $\iff$  (iv): let  $MNU' = \lambda^2 U'$  for some matrix  $U'$  of rank  $f$ . Then  $NM(NU') = \lambda^2 NU'$ , and  $\text{rank } NU' = \text{rank } U'$ , since

$$\text{rank } \lambda^2 U' = \text{rank } MNU' \leq \text{rank } U',$$

and  $\lambda \neq 0$ . This proves the second equivalence.

3. (i)  $\iff$  (iii): let  $U$  and  $\tilde{U}$  be as in case 1. Put  $n := m_1 + m_2$ . Then

$$A^2 \begin{bmatrix} U \\ \tilde{U} \end{bmatrix} = \lambda^2 \begin{bmatrix} U \\ \tilde{U} \end{bmatrix},$$

and  $\text{rank} \begin{bmatrix} U \\ \tilde{U} \end{bmatrix} = 2f$  if  $\lambda \neq 0$ . On the other hand the multiplicity of an eigenvalue  $\lambda^2 \neq 0$  of  $A^2$  equals

$$\begin{aligned} n - \text{rank}(A^2 - \lambda^2 I) &= n - \text{rank}((A - \lambda I)(A + \lambda I)) \leq \\ &\leq n - (\text{rank}(A - \lambda I) + \text{rank}(A + \lambda I) - n) = 2f. \end{aligned}$$

□

The *singular values* of a complex matrix  $N$ , are the positive eigenvalues of

$$\begin{bmatrix} 0 & N \\ N^* & 0 \end{bmatrix}.$$

By the above lemma we see that they are the same as the square roots of the non-zero eigenvalues of  $NN^*$ .

1.1.3. LEMMA. *Let*

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

*be a complex matrix. If  $A_{11}$  is non-singular, and  $\text{rank } A_{11} = \text{rank } A$ , then*

$$A_{22} = A_{21} A_{11}^{-1} A_{12}.$$

PROOF. For  $i = 1, 2$ , let  $a_{i,j}$  denote the  $j$ -th column of  $A_{i2}$ . From  $\text{rank } A = \text{rank } [A_{11}^* \ A_{21}^*]^*$  it follows

$$\begin{bmatrix} a_{1,j} \\ a_{2,j} \end{bmatrix} = \begin{bmatrix} A_{11} \\ A_{21} \end{bmatrix} u,$$

for some vector  $u$ . But if  $A_{11}$  is non-singular, then  $u = A_{11}^{-1} a_{1,j}$ . Hence  $a_{2,j} = A_{21} A_{11}^{-1} a_{1,j}$ , which proves the lemma.  $\square$

The identity matrix of size  $n$  will be denoted by  $I_n$  or  $I$ . The matrix with all entries equal to one by  $J$ ; a column vector of  $J$  is denoted by  $j$ ;  $J_n$  is a square  $J$  of size  $n$ ; the symbol  $\otimes$  is used for the Kronecker product of matrices.

As a last result in this section we observe that, if  $K := I_n \otimes J_m$ , then

$$\lambda_1(K) = \dots = \lambda_n(K) = m, \quad \lambda_{n+1}(K) = \dots = \lambda_{mn}(K) = 0.$$

## 1.2. INTERLACING OF EIGENVALUES

Suppose  $A$  and  $B$  are square complex matrices of size  $n$  and  $m$ , respectively ( $m \leq n$ ), having only real eigenvalues. If

$$\lambda_i(A) \geq \lambda_i(B) \geq \lambda_{n-m+i}(A)$$

for all  $i = 1, \dots, m$ , then we say that the eigenvalues of  $B$  *interlace* the eigenvalues of  $A$ . If there exists an integer  $k$ ,  $0 \leq k \leq m$ , such that

$$\lambda_i(A) = \lambda_i(B) \quad \text{for } i = 1, \dots, k$$

and

$$\lambda_{n-m+i}(A) = \lambda_i(B) \quad \text{for } i = k+1, \dots, m,$$

then the interlacing will be called *tight*. Our first and fundamental case of interlacing eigenvalues is due to R. Courant (see [C11], p. 28).

**1.2.1. THEOREM.** *Let  $S$  be a complex  $n \times m$  matrix such that  $S^*S = I_m$ . Let  $A$  be a hermitian matrix of size  $n$ . Define  $B := S^*AS$ , and let  $v_1, \dots, v_m$  be an orthonormal set of eigenvectors of  $B$ . Then*

- i. *the eigenvalues of  $B$  interlace the eigenvalues of  $A$ ;*
- ii. *if  $\lambda_i(B) \in \{\lambda_i(A), \lambda_{n-m+i}(A)\}$  for some  $i \in \{1, \dots, m\}$ , then there exists an eigenvector  $v$  of  $B$  for  $\lambda_i(B)$ , such that  $Sv$  is an eigenvector of  $A$  for  $\lambda_i(B)$ ;*
- iii. *if, for some  $\ell \in \{0, \dots, m\}$ ,  $\lambda_i(A) = \lambda_i(B)$  for all  $i = 1, \dots, \ell$ , then  $Sv_i$  is an eigenvector of  $A$  for  $\lambda_i(A)$ , for  $i = 1, \dots, \ell$ ;*
- iv. *if the interlacing is tight, then  $SB = AS$ .*

**PROOF.** Let  $u_1, \dots, u_n$  be an orthonormal set of eigenvectors of  $A$ . For any  $i$ ,  $1 \leq i \leq m$ , take

$$\tilde{v}_i \in \langle v_1, \dots, v_i \rangle \cap \langle S^*u_1, \dots, S^*u_{i-1} \rangle^\perp, \quad \tilde{v}_i \neq 0.$$

Then  $S\tilde{v}_i \in \langle u_1, \dots, u_{i-1} \rangle^\perp$ , hence by 1.1.1

$$\lambda_i(A) \geq \frac{\tilde{v}_i^* S^* A S \tilde{v}_i}{\tilde{v}_i^* S^* S \tilde{v}_i} = \frac{\tilde{v}_i^* B \tilde{v}_i}{\tilde{v}_i^* \tilde{v}_i} \geq \lambda_i(B).$$

Thus also

$$-\lambda_i(B) = \lambda_{m-i+1}(-B) \leq \lambda_{m-i+1}(-A) = -\lambda_{n-m+i}(A).$$

This proves (i).

If  $\lambda_i(B) = \lambda_i(A)$ , then  $\tilde{v}_i$  and  $S\tilde{v}_i$  are eigenvectors of  $B$  and  $A$  respectively for the eigenvalue  $\lambda_i(B) = \lambda_i(A)$ . This, together with the same result applied to  $-B$  and  $-A$  yields (ii).

We shall prove (iii) by induction on  $\ell$ . If  $\ell = 0$ , there is nothing to prove. Suppose  $\ell > 0$ . We have

$$\lambda_\ell(A) = \lambda_\ell(B) = v_\ell^* B v_\ell = \frac{v_\ell^* S^* A S v_\ell}{v_\ell^* S^* S v_\ell}.$$

On the other hand,  $Sv_\ell \in \langle Sv_1, \dots, Sv_{\ell-1} \rangle^\perp$ , and by the induction hypothesis  $Sv_1, \dots, Sv_{\ell-1}$  are orthonormal eigenvectors of  $A$  for  $\lambda_1(A), \dots, \lambda_{\ell-1}(A)$ . Now 1.1.1.ii yields that  $Sv_\ell$  is an eigenvector of  $A$  for  $\lambda_\ell(A)$ . This proves (iii).

Let the interlacing be tight. By applying (iii) to  $A$  with  $\ell = k$  and to  $-A$  with  $\ell = m - k$ , we find that  $Sv_1, \dots, Sv_m$  is an orthonormal set of eigenvectors of  $A$  for  $\lambda_1(B), \dots, \lambda_m(B)$ . Write  $V := [v_1 \dots v_m]$  and  $D := \text{diag}(\lambda_1(B), \dots, \lambda_m(B))$ . Then  $ASV = SVD$ , and  $BV = VD$ . Hence

$$ASV = SBV.$$

Because  $V$  is non-singular, (iv) has now been proved.  $\square$

A direct consequence of the above theorem is the following theorem. This result is known under the name *Cauchy inequalities*, see [H7], [M2], [W5].

**1.2.2. THEOREM.** *Suppose*

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$$

*is a hermitian matrix.*

- i. *The eigenvalues of  $A_{11}$  interlace the eigenvalues of  $A$ .*
- ii. *If the interlacing is tight, then  $A_{12} = 0$ .*

**PROOF.** Let  $m$  be the size of  $A_{11}$ . Define  $S := [I_m \ 0]^*$ , and apply 1.2.1.  $\square$

Another consequence of 1.2.1 is the following result which was announced in [H1] (see also [H2]). This result will often be used in the forthcoming sections.

1.2.3. THEOREM. Let  $A$  be a hermitian matrix partitioned as follows

$$A = \begin{bmatrix} A_{11} & \dots & A_{1m} \\ \vdots & & \vdots \\ A_{m1} & \dots & A_{mm} \end{bmatrix},$$

such that  $A_{ii}$  is square for  $i = 1, \dots, m$ . Let  $b_{ij}$  be the average row sum of  $A_{ij}$ , for  $i, j = 1, \dots, m$ . Define the  $m \times m$  matrix  $B := (b_{ij})$ .

- i. The eigenvalues of  $B$  interlace the eigenvalues of  $A$ .
- ii. If the interlacing is tight, then  $A_{ij}$  has constant row and column sums for  $i, j = 1, \dots, m$ .
- iii. If, for  $i, j = 1, \dots, m$ ,  $A_{ij}$  has constant row and column sums, then any eigenvalue of  $B$  is also an eigenvalue of  $A$  with not smaller a multiplicity.

PROOF. Let  $n_i$  be the size of  $A_{ii}$ . Define

$$\tilde{S}^* := \begin{bmatrix} \underbrace{1 \dots 1}_{n_1} & 0 \dots 0 & \dots & 0 \dots 0 \\ 0 \dots 0 & \underbrace{1 \dots 1}_{n_2} & \dots & 0 \dots 0 \\ 0 \dots 0 & 0 \dots 0 & \dots & 0 \dots 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 \dots 0 & 0 \dots 0 & \dots & \underbrace{1 \dots 1}_{n_m} \end{bmatrix},$$

$D := \text{diag}(\sqrt{n_1}, \dots, \sqrt{n_m})$ , and  $S := \tilde{S} D^{-1}$ . Then  $S^* S = I$  and  $\tilde{S}^* \tilde{S} = D^2$ . We easily see that  $(\tilde{S}^* \tilde{A} \tilde{S})_{ij}$  equals the sum of the entries of  $A_{ij}$ . Hence

$$B = \tilde{S}^* \tilde{A} \tilde{S} D^{-2}.$$

By 1.2.1.i we know that the eigenvalues of  $S^* A S$  interlace the eigenvalues of  $A$ . But  $B$  has the same eigenvalues as  $S^* A S$ , since

$$S^* A S = D^{-1} \tilde{S}^* \tilde{A} \tilde{S} D^{-1} = D^{-1} B D.$$

This proves (i).

It is easily checked that  $A S = S(D^{-1} B D)$  reflects that  $A_{ij}$  has constant row sums for all  $i, j = 1, \dots, m$ . Hence 1.2.1.iv implies (ii).

On the other hand, if  $AS = SD^{-1}BD$  and  $BU = \lambda_i(B)U$  for some matrix  $U$  and integer  $i$ , then  $A(SD^{-1}U) = \lambda_i(B)SD^{-1}U$ , and  $\text{rank } U = \text{rank } SD^{-1}U$ . This proves (iii).  $\square$

As a special case of the above theorem we have that

$$\lambda_1(A) \geq \lambda_i(B) \geq \lambda_n(A) ,$$

for  $i = 1, \dots, m$ . These inequalities are well known and usually applied under the name *Higman-Sims technique*, see [H10], [P4]. We shall also use the name Higman-Sims technique if we apply the more general result 1.2.3. Also 1.2.3.iii is well known, see for instance [C13], [H9] (note that this result remains valid for non-hermitian  $A$ ). We see that 1.2.3.ii gives a sufficient, and that 1.2.3.iii gives a necessary condition for the block matrices of  $A$  to have constant row and column sums. However, neither of these conditions is both necessary and sufficient. This is illustrated by the following partitioned matrices:

$$A := \left( \begin{array}{cc|cc} 0 & 0 & -1 & 1 \\ 0 & 0 & 1 & -1 \\ \hline -1 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 \end{array} \right) , \quad A' := \left( \begin{array}{cc|cc} 0 & 0 & -1 & 1 \\ 0 & 0 & -1 & 1 \\ \hline -1 & -1 & 0 & 0 \\ 1 & 1 & 0 & 0 \end{array} \right) .$$

For both  $A$  and  $A'$  the eigenvalues are  $2, 0, 0, -2$ , and the average row sums of the block matrices are given by the entries of  $B = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$ . The block matrices of  $A$  have constant row sums, whilst the interlacing is not tight. The row sums of the block matrices of  $A'$  are not constant, whilst the eigenvalues of  $B$  are also eigenvalues of  $A$ .

### 1.3. MORE EIGENVALUE INEQUALITIES

In this section we shall use interlacing of eigenvalues in order to prove some known inequalities and equalities, which we shall use in later sections. The first result is due to WEYL [W2] (see also [W5]).

1.3.1. THEOREM. *Let  $A_1$  and  $A_2$  be hermitian matrices of size  $m$ . Then*

$$\lambda_{i-j}(A_1) + \lambda_{1+j}(A_2) \geq \lambda_i(A_1 + A_2) \geq \lambda_{i+j}(A_1) + \lambda_{m-j}(A_2) ,$$

for  $i = 1, \dots, m$ ,  $0 \leq j \leq \min\{i-1, m-i\}$ .



PROOF. Define

$$A := \begin{bmatrix} A_1 - \lambda_{i-j}(A_1)I & 0 \\ 0 & A_2 - \lambda_{1+j}(A_2)I \end{bmatrix}, \quad S := \frac{1}{2}\sqrt{2} \begin{bmatrix} I_m \\ I_m \end{bmatrix}.$$

Then

$$\lambda_{i-j}(A_1 - \lambda_{i-j}(A_1)I) = \lambda_{1+j}(A_2 - \lambda_{1+j}(A_2)I) = 0.$$

Hence  $\lambda_i(A) = \lambda_{i+1}(A) = 0$ . On the other hand

$$S^*AS = \frac{1}{2}(A_1 + A_2 - (\lambda_{i-j}(A_1) + \lambda_{1+j}(A_2))I).$$

With 1.2.1 we now have

$$\lambda_i(A_1 + A_2) - \lambda_{i-j}(A_1) - \lambda_{1+j}(A_2) = 2\lambda_i(S^*AS) \leq 2\lambda_i(A) = 0.$$

If we replace  $A_1$  and  $A_2$  by  $-A_1$  and  $-A_2$ , we get the second inequality.  $\square$

The next theorem is due to HOFFMAN [H13].

1.3.2. THEOREM. Let  $A$  be a hermitian matrix of size  $n$ , partitioned as follows

$$A = \begin{bmatrix} A_{11} & \cdots & A_{1m} \\ \vdots & & \vdots \\ A_{m1} & \cdots & A_{mm} \end{bmatrix},$$

where  $A_{ii}$  is a square matrix of size  $n_i$ , for  $i = 1, \dots, m$ . Let  $j_1, \dots, j_m$  be integers such that  $1 \leq j_i \leq n_i$  for  $i = 1, \dots, m$ . Then

$$\begin{aligned} \lambda_{j_1+\dots+j_m}(A) + \sum_{i=1}^{m-1} \lambda_i(A) &\geq \sum_{i=1}^m \lambda_{j_i}(A_{ii}) \geq \\ &\geq \lambda_{j_1+\dots+j_m-m+1}(A) + \sum_{i=n-m+2}^n \lambda_i(A). \end{aligned}$$

PROOF. Let  $u_1, \dots, u_n$  be an orthonormal set of eigenvectors of  $A$ . Let  $u_{i1}, \dots, u_{in_i}$  be an orthonormal set of eigenvectors of  $A_{ii}$  for  $i = 1, \dots, m$ . Put  $k := j_1 + \dots + j_m$ . Choose a vector  $\tilde{u}_k = [\tilde{u}_{1k}^* \dots \tilde{u}_{mk}^*]^* \neq 0$ , such that

$$\tilde{u}_k \in \langle u_1, \dots, u_{k-1} \rangle^\perp,$$

and

$$\tilde{u}_{ik} \in \langle u_{i1}, \dots, u_{ij_i} \rangle \text{ for } i = 1, \dots, m.$$

It follows from dimension considerations that we can always do so. Now define

$$w_i := \|\tilde{u}_{ik}\|,$$

and

$$\hat{u}_{ik} := \begin{cases} \frac{1}{w_i} \tilde{u}_{ik} & \text{if } w_i \neq 0, \\ u_{ij_i} & \text{if } w_i = 0, \end{cases}$$

for  $i = 1, \dots, m$ . Furthermore put

$$w := (w_1, \dots, w_m)^*,$$

and

$$S := \begin{bmatrix} \hat{u}_{1k} & 0 & \dots & 0 \\ 0 & \hat{u}_{2k} & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & \hat{u}_{mk} \end{bmatrix}.$$

Then we have

$$S^*S = I, \quad Sw = \tilde{u}_k, \quad \text{and } (S^*AS)_{ii} = \hat{u}_{ik}^* A_{ii} \hat{u}_{ik} \text{ for } i = 1, \dots, m.$$

By 1.1.1.i and the choice of  $\hat{u}_{ik}$ , the last formula gives

$$(S^*AS)_{ii} \geq \lambda_{j_i}(A_{ii}) \text{ for } i = 1, \dots, m.$$

Hence

$$(*) \quad \sum_{i=1}^m \lambda_i(S^*AS) = \text{trace } S^*AS \geq \sum_{i=1}^m \lambda_{j_i}(A_{ii}).$$

On the other hand,  $Sw = \tilde{u}_k$ ,  $S^*S = I$  and 1.1.1 yield

$$(**) \quad \lambda_m(S^*AS) \leq \frac{w^* S^* A S w}{w^* w} = \frac{\tilde{u}_k^* \tilde{A} \tilde{u}_k}{\tilde{u}_k^* \tilde{u}_k} \leq \lambda_k(A) .$$

Applying 1.2.1 gives

$$(***) \quad \sum_{i=1}^{m-1} \lambda_i(S^*AS) \leq \sum_{i=1}^{m-1} \lambda_i(A) .$$

Combining (\*), (\*\*) and (\*\*\*) yields the first inequality of our theorem. Again, the second inequality follows by substituting  $-A$  for  $A$  in the first one.  $\square$

If the matrix  $A$  of the above theorem is positive semi-definite and  $m = 2$ , then we have

$$\lambda_{j_1}(A_{11}) + \lambda_{j_2}(A_{22}) \geq \lambda_{j_1+j_2-1}(A) .$$

These are the inequalities of ARONSZAJN [A3] (see also [H7]).

The following consequence of 1.3.2 will turn out to be a useful tool in computations with eigenvalues.

1.3.3. THEOREM. *Suppose*

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$$

*is hermitian of size  $n$ . Suppose  $A$  has just two distinct eigenvalues, that is,*

$$\lambda_1(A) = \dots = \lambda_f(A) > \lambda_{f+1}(A) = \dots = \lambda_n(A)$$

*for some  $f$ ,  $1 \leq f < n$ . Let  $n_1$  and  $n_2$  be the sizes of  $A_{11}$  and  $A_{22}$  respectively. Then*

$$\lambda_i(A_{22}) = \lambda_1(A) + \lambda_n(A) - \lambda_{f-i+1}(A_{11})$$

$$\text{for } \max\{1, f+1-n_1\} \leq i \leq \min\{f, n_2\} ,$$

$$\lambda_i(A_{22}) = \lambda_1(A) \quad \text{for } 1 \leq i < f+1-n_1 ,$$

$$\lambda_i(A_{22}) = \lambda_n(A) \quad \text{for } f < i \leq n_2 .$$

PROOF. By the Cauchy inequalities (1.2.2.i) we have

$$\lambda_i(A) \geq \lambda_i(A_{22}) \geq \lambda_{n_1+i}.$$

This proves the result for  $1 \leq i < f+1-n_1$ , and for  $f < i \leq n_2$ .  
For the remaining values of  $i$ , 1.3.2 gives

$$\lambda_{f+1}(A) + \lambda_i(A) \geq \lambda_{f-i+1}(A_{11}) + \lambda_i(A_{22}) \geq \lambda_f(A) + \lambda_n(A),$$

which proves the required result. □

It is an easy exercise to give a direct proof of the above theorem. The proof then could go analogously to the one of Theorem 5.1 of [C5], where a similar result is stated.

CHAPTER 2  
INEQUALITIES FOR GRAPHS

2.1. INDUCED SUBGRAPHS

In this section we shall derive inequalities for induced subgraphs of graphs, using the results of section 1.2 on interlacing of eigenvalues.

Let  $G$  be a graph on  $n$  vertices. The *eigenvalues* of  $G$  are the eigenvalues of its  $(0,1)$ -adjacency matrix; we denote them by  $\lambda_1(G) \geq \dots \geq \lambda_n(G)$ . Let  $G_1$  be an induced subgraph of  $G$ . Then by 1.2.2 (Cauchy inequalities) the eigenvalues of  $G_1$  interlace the eigenvalues of  $G$ . In particular, if  $G_1$  is a coclique of size  $\alpha$ , then  $\lambda_\alpha(G) \geq \lambda_\alpha(G_1) = 0$ , and  $\lambda_{n-\alpha+1}(G) \leq \lambda_1(G_1) = 0$ . Hence, we have the following result, which was first observed by CVETKOVIĆ [C12] (see also [C13]).

2.1.1. THEOREM. *The size of a coclique of a graph  $G$  cannot exceed the number of nonnegative [nonpositive] eigenvalues of  $G$ .*

Now we shall derive inequalities for induced subgraphs using the Higman-Sims technique (1.2.3). Suppose  $G$  is a graph on  $n$  ( $n \geq 2$ ) vertices of average degree  $d$ . Let the vertex set of  $G$  be partitioned into two non-empty sets, and let  $G_1$  and  $G_2$  be the subgraphs induced by these two sets. For  $i = 1, 2$ , let  $n_i$  be the number of vertices of  $G_i$ , let  $d_i$  be the average degree of  $G_i$ , and let  $\tilde{d}_i$  be the average of the degrees in  $G$  over the vertices of  $G_i$ . Now we can state the following theorem.

2.1.2. THEOREM. *For  $i = 1, 2$*

i. 
$$\lambda_1(G)\lambda_2(G) \geq \frac{d_i d n - \tilde{d}_i^2 n_i}{n - n_i} \geq \lambda_1(G)\lambda_n(G) ,$$

ii. *if equality holds on one of the sides, then  $G_1$  and  $G_2$  are regular, and also the degrees in  $G$  are constant over the vertices of  $G_1$  and  $G_2$ .*

PROOF. If  $G$  is complete we easily see that the theorem is correct. So let us assume that  $G$  is not complete. Let  $A_{11}$ ,  $A_{22}$  and

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$$

be the adjacency matrices of  $G_1$ ,  $G_2$ , and  $G$ , respectively. Put

$$B := \begin{bmatrix} d_1 & \tilde{d}_1 - d_1 \\ \tilde{d}_2 - d_2 & d_2 \end{bmatrix} .$$

Then the entries of  $B$  are the average row sums of the block matrices of  $A$ . By 1.2.3.i we have

$$\lambda_1(A) \geq \lambda_1(B) , \quad \lambda_2(A) \geq \lambda_2(B) , \quad -\lambda_n(A) \geq -\lambda_2(B) .$$

Because  $\text{trace } A = 0$ , we have  $\lambda_n(A) \leq 0$ . From 2.1.1 we know that  $\lambda_2(A) \geq 0$ , since otherwise  $G$  would be complete.  $\text{Trace } B \geq 0$  implies  $\lambda_1(B) \geq 0$ . Hence

$$(*) \quad \lambda_1(A)\lambda_2(A) \geq \lambda_1(B)\lambda_2(B) \geq \lambda_1(A)\lambda_n(A) .$$

On the other hand we know

$$(**) \quad \lambda_1(B)\lambda_2(B) = \det B = d_1\tilde{d}_2 - \tilde{d}_1(\tilde{d}_2 - d_2) .$$

We quickly see

$$dn = \tilde{d}_1n_1 + \tilde{d}_2n_2 , \quad n_2(\tilde{d}_2 - d_2) = n_1(\tilde{d}_1 - d_1) , \quad n = n_1 + n_2 .$$

This yields

$$d_1\tilde{d}_2 - \tilde{d}_1(\tilde{d}_2 - d_2) = (d_1dn - \tilde{d}_1^2n_1) / (n - n_1) .$$

With (\*) and (\*\*) this proves (i).

If equality holds on one of the sides, the interlacing must have been tight. Hence 1.2.3.ii gives (ii). □

Now let us look at the consequences of the above inequalities for some special cases. The size of the largest coclique and clique of  $G$  are denoted by  $\alpha(G)$  and  $\omega(G)$ , respectively.

**2.1.3. THEOREM.** If  $d_{\min}$  and  $d_{\max}$  are the smallest and the largest degree in the graph  $G$ , respectively, then

$$i. \quad \alpha(G) (d_{\min}^2 - \lambda_1(G)\lambda_n(G)) \leq -n\lambda_1(G)\lambda_n(G) ,$$

$$ii. \quad \omega(G) (dn - d_{\max}^2 + \lambda_1(G)\lambda_2(G)) \leq n(d + \lambda_1(G)\lambda_2(G)) .$$

**PROOF.** (i). Substitute  $\alpha(G) = n_1$ ,  $d_1 = 0$  and  $\tilde{d}_1 \geq d_{\min}$  in the right hand inequality of 2.1.2.i.

(ii). Substitute  $\omega(G) = n_1$ ,  $d_1 = \omega(G) - 1$  and  $\tilde{d}_1 \leq d_{\max}$  in the left hand inequality of 2.1.2.i. □

**2.1.4. THEOREM.** If  $G$  is a regular graph on  $n \geq 2$  vertices of degree  $d$ , then

i. any induced subgraph  $G_1$  of  $G$  with  $n_1$  ( $0 < n_1 < n$ ) vertices and average degree  $d_1$  satisfies

$$\lambda_2(G) \geq \frac{nd_1 - n_1d}{n - n_1} \geq \lambda_n(G) ,$$

$$ii. \quad \alpha(G) (d - \lambda_n(G)) \leq -n\lambda_n(G) ,$$

$$iii. \quad \omega(G) (n - d + \lambda_2(G)) \leq n(1 + \lambda_2(G)) .$$

**PROOF.** If  $G$  is regular then  $\lambda_1(G) = d = \tilde{d}_1 = \tilde{d}_2 = d_{\min} = d_{\max}$ . Now 2.1.2.i, 2.1.3.i and 2.1.3.ii give the required results. □

The inequality 2.1.4.ii is an unpublished result of A.J. Hoffman (see [C13], [H2], [L2]). In fact, the inequalities (ii) and (iii) of 2.1.4 (just as the left and the right hand inequality of 2.1.4.i) are equivalent, because either one can be obtained from the other by using  $\omega(\bar{G}) = \alpha(G)$ ,  $\lambda_1(\bar{G}) = n - \lambda_1(G) - 1$  and  $\lambda_i(\bar{G}) = -\lambda_{n-i+2}(G) - 1$  for  $i = 2, \dots, n$  ( $\bar{G}$  is the complement of  $G$ ).

For the graph  $G$  with its subgraphs  $G_1$  and  $G_2$ , we define  $D(G, G_1)$  to be the incidence structure whose points and blocks are the vertices of  $G_1$  and  $G_2$ , respectively, a point and a block being incident if the corresponding vertices are adjacent. If we have equality in any of the inequalities of 2.1.2 - 2.1.4, then 2.1.2.ii yields that  $D(G, G_1)$  is a 1-design, possibly

degenerate. Now let  $G$  be strongly regular. Then by use of 1.2.3.iii it is not difficult to show that equality holds in 2.1.4 iff  $D(G, G_1)$  is a 1-design. If  $G_1$  is a coclique or a clique we have a criterion for  $D(G, G_1)$  to be a 2-design.

**2.1.5. THEOREM.** *If  $G$  is a strongly regular graph on  $n$  vertices of degree  $d$ , then*

- i.  $\alpha(G) \leq 1 + (n-d-1) / (\lambda_2(G) + 1)$  ,
- ii.  $\omega(G) \leq 1 - d / \lambda_n(G)$  ,
- iii. *if equality holds in (i) or (ii), and  $G_1$  is a coclique of size  $\alpha(G)$ , or a clique of size  $\omega(G)$ , respectively, then  $D(G, G_1)$  is a 2-design, possibly degenerate.*

**PROOF.** If  $G$  is strongly regular, we know (see [C5], [S4] or Appendix I)

$$(d - \lambda_2(G))(d - \lambda_n(G)) = n(d + \lambda_2(G)\lambda_n(G)) .$$

From this it follows in a straightforward way that (i) and (ii) are equivalent to 2.1.4.ii and 2.1.4.iii.

From the definition of a strongly regular graph we know that any two points of  $D(G, G_1)$  are incident with a constant number of blocks of  $D(G, G_1)$ .

Furthermore, equality in (i) or (ii) implies that  $D(G, G_1)$  is a 1-design, so in this case  $D(G, G_1)$  is then a 2-design, possibly degenerate.  $\square$

The theorems 2.1.5 and 2.1.4 for strongly regular graphs are known. They are direct consequences of the linear programming bound of DELSARTE [D1] (see also [H2]). They were also proved by BUMILLER [B9].

Applying 2.1.4.i to the point graph of a partial geometry (see Appendix I, or [B6], [T1]) gives the following result of DE CLERCK [C7] (see also [P3] for the case  $\alpha = 1$ , and [B9] for  $t = t_1$ ).

**2.1.6. COROLLARY.** *Let  $P$  be a partial geometry with parameters  $(s, t, \alpha)$ , containing a partial subgeometry  $P_1$  with parameters  $(s_1, t_1, \alpha)$ . Then*

$$s = s_1 \quad \text{or} \quad s \geq s_1 t_1 + \alpha - 1 .$$



PROOF. If  $G$  and  $G_1$  are the point graphs of  $P$  and  $P_1$ , respectively, then (see Appendix I or [T1])  $G_1$  is an induced subgraph of  $G$ , and

$$n = (s+1)(st+\alpha) / \alpha, \quad n_1 = (s_1+1)(s_1t_1+\alpha) / \alpha,$$

$$d = s(t+1), \quad d_1 = s_1(t_1+1), \quad \lambda_2(G) = s - \alpha.$$

Substitution of these values in the left hand inequality of 2.1.4.i leads to

$$(s - s_1)(st + \alpha)(s_1t_1 + \alpha - s - 1) \leq 0.$$

This proves the result. □

The next theorem gives a result in case both Hoffman's bound (2.1.4.ii) and Cvetković's bound (2.1.1) are tight.

2.1.7. **THEOREM.** *Let  $G$  be a strongly regular graph on  $n$  vertices, which is not complete  $\gamma$ -partite (i.e.  $\lambda_2(G) > 0$ ). Let  $f_n(G)$  denote the multiplicity of  $\lambda_n(G)$ . Then*

i.  $\alpha(G) \leq f_n(G),$

ii.  $\alpha(G) \leq 1 + (n - \lambda_1(G) - 1) / (\lambda_2(G) + 1),$

iii. *Let  $G_1$  be a coclique, whose size attains both of these bounds, then  $G_2$ , the subgraph of  $G$  induced by the remaining vertices, is strongly regular with eigenvalues*

$$\lambda_1(G_2) = \lambda_1(G) - \lambda_1(G)\alpha(G) / (n - \alpha(G)), \quad \lambda_2(G_2) = \lambda_2(G),$$

$$\lambda_{n-\alpha(G)}(G_2) = \lambda_2(G) + \lambda_n(G).$$

PROOF. Theorem 2.1.1 implies (i); (ii) is the same as 2.1.5.i. Let  $A$  and  $A_2$  be the adjacency matrices of  $G$  and  $G_2$ , respectively. Then

$$A - \frac{\lambda_1(G) - \lambda_2(G)}{n} J$$

has just two distinct eigenvalues,  $\lambda_2(G)$  and  $\lambda_n(G)$  of multiplicity  $n - f_n(G)$  and  $f_n(G) = \alpha(G)$ , respectively. From 1.3.3 it follows that

$$\tilde{A}_2 := A_2 - \frac{\lambda_1(G) - \lambda_2(G)}{n} J$$

has three distinct eigenvalues,  $\lambda_2(G)$ ,  $\lambda_2(G) + \lambda_n(G)$  and  $\lambda_2(G) + \lambda_n(G) + \alpha(G)(\lambda_1(G) - \lambda_2(G)) / n$ , where the last eigenvalue is simple (has multiplicity one). On the other hand, 2.1.5.iii gives that  $G_2$  is regular of degree  $\lambda_1(G) - \lambda_1(G)\alpha(G) / (n - \alpha(G))$ . This shows that  $\tilde{A}_2$  and  $A_2$  have a common basis of eigenvectors, and that the simple eigenvalue of  $\tilde{A}_2$  belongs to the eigenvector  $j$ . Thus  $A_2$  has the desired eigenvalues, and therefore (see [C6] or Appendix I)  $G_2$  is strongly regular.  $\square$

Using 1.1.3 it is not difficult to show that  $D(G, G_1)$  is a quasi-symmetric 2-design (see Section 3.2), whose block graph is the complement of  $G_2$ . This situation has been studied by SHRIKHANDE [S5].

In proving 2.1.2 we applied interlacing to the product of eigenvalues. We did so in order to get reasonably nice formulas. However, for non-regular graphs the inequality for the product carries less information than the separate inequalities. For this reason, applying the Higman-Sims technique directly to the adjacency matrix of a given non-regular graph, may yield better results than 2.1.2 or 2.1.3. Also, if more is known about the structure of  $G$  or  $G_1$ , it is often possible to get better results by a more detailed application of the Higman-Sims technique. Let us illustrate this by the following result.

**2.1.8. THEOREM.** *Let  $G$  be a regular graph on  $n$  vertices of degree  $d$ , and let the complete bipartite graph  $K_{l,m}$  be an induced subgraph of  $G$ . Let  $x_1$  and  $x_2$ ,  $x_1 \geq x_2$ , be the zeros of*

$$(n - l - m)x^2 + (dl + dm - 2lm)x - lm(n - 2d) .$$

*Then*

$$\lambda_2(G) \geq x_1 \quad \text{and} \quad \lambda_n(G) \leq x_2 .$$

**PROOF.** Without loss of generality, let  $G$  have adjacency matrix

$$A = \begin{bmatrix} 0 & J & A'_{12} \\ J & 0 & A''_{12} \\ A'_{21} & A''_{21} & A_{22} \end{bmatrix} .$$

where the diagonal block matrices are square of sizes  $\ell$ ,  $m$  and  $n_2$ , respectively. Using the Higman-Sims technique (1.2.3) we find that the eigenvalues of

$$B := \begin{pmatrix} 0 & m & d-m \\ \ell & 0 & d-\ell \\ \ell \frac{d-m}{n_2} & m \frac{d-\ell}{n_2} & d - \frac{\ell d + m d - 2\ell m}{n_2} \end{pmatrix}$$

interlace the eigenvalues of  $A$ . Clearly  $\lambda_1(B) = \lambda_1(A) = d$  and hence

$$\lambda_2(B)\lambda_3(B) = (\det B) / d = \ell m(2d-n) / n_2 ,$$

$$\lambda_2(B) + \lambda_3(B) = (\text{trace } B) - d = (2\ell m - \ell d - m d) / n_2 .$$

This yields  $x_1 = \lambda_2(B)$  and  $x_2 = \lambda_3(B)$ . Now the interlacing gives the required result.  $\square$

BUMILLER [B9] showed for strongly regular  $G$  and  $m = 1$  that

$$\ell(\mu + \lambda_2(G)) \leq -d\lambda_n(G) ,$$

where  $\mu = d + \lambda_2(G)\lambda_n(G)$ . Using

$$(d - \lambda_2(G))(d - \lambda_n(G)) = n(d + \lambda_2(G)\lambda_n(G))$$

one easily checks that this follows from the second inequality of the above theorem. PAYNE [P6] proved that

$$(\ell - 1)(m - 1) \leq s^2 ,$$

if  $G$  is the point graph of a generalized quadrangle of order  $(s, t)$  (see Chapter 5 or Appendix I). This follows after substituting

$$n = (s+1)(st+1) , \quad d = s(t+1) , \quad \lambda_2(G) = s-1$$

in the first inequality of the above theorem. It should not be surprising that for this case we obtain the same result as Payne, because he too uses the Higman-Sims technique.

## 2.2. CHROMATIC NUMBER

In this section we shall derive lower bounds for the chromatic number of a graph in terms of its eigenvalues. The main tool is Hoffman's generalization of Aronszajn's inequalities (1.3.2).

Let  $G$  be a non-void graph on  $n$  vertices. Then it follows immediately that  $\gamma(G)$ , the chromatic number of  $G$ , satisfies

$$\gamma(G)\alpha(G) \geq n .$$

Combining this with the upper bounds for  $\alpha(G)$  found in the previous section we obtain lower bounds for  $\gamma(G)$ . For instance, 2.1.3 gives

$$\gamma(G) \geq 1 - d_{\min}^2 / \lambda_1(G)\lambda_n(G) .$$

However, this is not best possible, since HOFFMAN [H13] (see also [B5], [H2], [H14], [L2]) showed that

$$\gamma(G) \geq 1 - \lambda_1(G) / \lambda_n(G) ,$$

which, if  $G$  is not regular, is better than the above bound. If  $G$  is regular, then the two bounds coincide. Taking into account that  $\alpha(G)$  is an integer we get

$$\gamma(G) \geq n / \left\lfloor n \lambda_1(G)\lambda_n(G) / (\lambda_1(G)\lambda_n(G) - d_{\min}^2) \right\rfloor ,$$

which is occasionally better than Hoffman's bound.

HOFFMAN [H13] proves his lower bound by use of the inequalities 1.3.2. We shall use the same technique, but in a more profound way, in order to obtain a generalization of Hoffman's inequality.

**2.2.1. THEOREM.** *Let  $G$  be a graph on  $n$  vertices with chromatic number  $\gamma$ . Let  $k$  be an integer satisfying  $0 \leq k < n/\gamma$ . Then*

- i.  $(\gamma - 1)\lambda_{k+1}(G) \geq -\lambda_{n-k(\gamma-1)}(G) ,$
- ii.  $(\gamma - 1)\lambda_{n-k}(G) \leq -\lambda_{1+k(\gamma-1)}(G) .$

**PROOF.** Let  $A$  be the adjacency matrix of  $G$ . Then without loss of generality

$$A = \begin{bmatrix} A_{11} & \cdots & A_{1\gamma} \\ \vdots & & \vdots \\ A_{\gamma 1} & \cdots & A_{\gamma\gamma} \end{bmatrix},$$

where  $A_{ii}$  is the  $n_i \times n_i$  all-zero matrix, for  $i = 1, \dots, \gamma$ .

First, we assume that  $n_i > k$  for  $i = 1, \dots, \gamma$ . Let  $u_1, \dots, u_n$  denote an orthonormal set of eigenvectors of  $A$ . Define

$$\tilde{A} := A + (\lambda_n(A) - \lambda_1(A)) \sum_{i=1}^k u_i u_i^*.$$

Clearly the matrices  $u_i u_i^*$  and  $A$  have a common basis of eigenvectors. This implies

$$(*) \quad \lambda_i(\tilde{A}) = \lambda_{k+i}(A) \quad \text{for } i = 1, \dots, n-k.$$

For  $i = 1, \dots, \gamma$ , let  $\tilde{A}_{ii}$  be the submatrix of  $\tilde{A}$  corresponding to  $A_{ii}$ . Since  $u_i u_i^*$  is positive semi-definite of rank one,  $\sum_{i=1}^k u_i u_i^*$  is positive semi-definite of rank  $k$ . This yields that  $-\tilde{A}_{ii}$  is positive semi-definite of rank at most  $k$ , hence

$$\lambda_{n_i-k}(\tilde{A}_{ii}) = 0 \quad \text{for } i = 1, \dots, \gamma.$$

Now we apply the left hand inequality of 1.3.2 with  $j_i = n_i - k$ . This gives

$$\lambda_{n-\gamma k}(\tilde{A}) + \sum_{i=1}^{\gamma-1} \lambda_i(\tilde{A}) \geq 0.$$

With (\*) this yields

$$(**) \quad \lambda_{n-k(\gamma-1)}(A) + \sum_{i=1}^{\gamma-1} \lambda_{i+k}(A) \geq 0.$$

Hence

$$(***) \quad (\gamma-1)\lambda_{1+k}(A) \geq -\lambda_{n-k(\gamma-1)}(A).$$

Now suppose that  $n_i \leq k$  for some  $i \in \{1, \dots, \gamma\}$ . Let  $L \subset \{1, \dots, \gamma\}$  be such that  $n_i \leq k$  iff  $i \in L$ . Let  $A'$  be the  $n' \times n'$  submatrix of  $A$ , obtained by discarding all block matrices  $A_{ij}$  with  $i \in L$  or  $j \in L$ . Put  $\ell := |L|$ . From  $k < n/\gamma$  it follows that  $\ell < \gamma$ , hence  $n' > 0$ . Now (\*\*\*) gives

$$(\gamma - \ell - 1)\lambda_{1+k}(A') + \lambda_{n'+k\ell-k(\gamma-1)}(A') \geq 0 .$$

Using  $n' + k\ell \geq n$  and the Cauchy inequalities (1.2.2) we have

$$\lambda_{n'+k\ell-k(\gamma-1)}(A') \leq \lambda_{n-k(\gamma-1)}(A') \leq \lambda_{n-k(\gamma-1)}(A) ,$$

$$\lambda_{1+k}(A') \leq \lambda_{1+k}(A) .$$

Hence

$$(\gamma - \ell - 1)\lambda_{1+k}(A) + \lambda_{n-k(\gamma-1)}(A) \geq 0 .$$

From  $k < n/\gamma$  it follows that  $\lambda_{1+k}(A) \geq \lambda_{n-k(\gamma-1)}(A)$ , hence  $\lambda_{1+k}(A) \geq 0$ .

Thus

$$(\gamma - 1)\lambda_{1+k}(A) \geq -\lambda_{n-k(\gamma-1)}(A) .$$

This proves (i). The proof of (ii) proceeds analogously, but also follows from the above by replacing  $A$  by  $-A$ .  $\square$

We see that the second inequality of the above theorem for  $k = 0$  is Hoffman's bound. In Chapter 4 we shall need a sharpened version of this inequality (see [H13]):

$$-\sum_{i=0}^{\gamma-2} \lambda_{n-i}(G) \geq \lambda_1(G) ,$$

which is in fact just formula (\*\*\*) in the above proof with  $k = 0$ , and  $A$  equal to minus the adjacency matrix of  $G$ .

If  $k > 0$ , the above inequalities are not really bounds for  $\gamma(G)$ , since  $\gamma(G)$  also occurs in an index. However, this does not matter much if we use these inequalities for estimating the chromatic number of a given graph. It is also not difficult to derive proper bounds from these inequalities. The next results illustrate this.

**2.2.2. COROLLARY.** *Let  $f_n(G)$  denote the multiplicity of the eigenvalue  $\lambda_n(G)$ . Then*

$$\gamma(G) \geq \min\{1 + f_n(G), 1 - \lambda_n(G) / \lambda_2(G)\} .$$

**PROOF.** Suppose  $\gamma = \gamma(G) \leq f_n(G)$ . Then  $\lambda_n(G) = \lambda_{n-\gamma+1}(G)$ . Now 2.2.1.i with  $k = 1$  gives  $(\gamma - 1)\lambda_2(G) \geq -\lambda_n(G)$ . This proves the required result.  $\square$

For strongly regular graphs the above results lead to the following theorem.

**2.2.3. THEOREM.** *Let  $G$  be a strongly regular graph on  $n$  vertices. Suppose  $G$  is not the pentagon or a complete  $\gamma$ -partite graph. Then*

$$\gamma(G) \geq \max\{1 - \lambda_1(G) / \lambda_n(G), 1 - \lambda_n(G) / \lambda_2(G)\} .$$

**PROOF.** Due to the above results, it suffices to prove the following claim:

$$f_n(G) \geq -\lambda_n(G) / \lambda_2(G) .$$

To achieve this, we distinguish three cases.

a)  $n \leq 28$ . For this case it follows by straightforward checking that all feasible parameter sets for strongly regular graphs which violate our claim are those of the pentagon and the complete  $\gamma$ -partite graphs.

b)  $\lambda_2(G) < 2$ . If  $G$  has a non-integral eigenvalue, then (see [C6], [C9] or Appendix I)  $\lambda_2(G) = -\frac{1}{2} + \frac{1}{2}\sqrt{n}$ , hence  $n < 25$  and we are in case a. Otherwise  $\lambda_2(G) = 1$ . SEIDEL [S3] determined all strongly regular graphs with  $\lambda_2(G) = 1$ . They satisfy  $n \leq 28$  or  $G$  is a ladder graph (disjoint union of edges), the complement of a lattice graph ( $\overline{L(K_{m,m})}$ ), or the complement of a triangular graph ( $\overline{L(K_m)}$ ). One easily verifies that these three families of graphs satisfy our claim.

c)  $\lambda_2(G) \geq 2$  and  $n > 28$ . If  $G$  is imprimitive ( $G$  is complete  $\gamma$ -partite or the disjoint union of complete graphs), the result is obvious. So assume  $G$  is primitive. Suppose the claim does not hold. Using  $\lambda_2(G) \geq 2$ ,  $\lambda_1(G) < n$  and

$$f_n(G)\lambda_n(G) + (n-1-f_n(G))\lambda_2(G) + \lambda_1(G) = 0$$

we obtain

$$\begin{aligned} f_n^2(G) &< -f_n(G)\lambda_n(G) / \lambda_2(G) = \\ &= n - 1 - f_n(G) + \lambda_1(G) / \lambda_2(G) < \frac{3}{2}n - f_n(G) . \end{aligned}$$

This yields

$$f_n^2(G) + 3f_n(G) < \frac{3}{2}n + 2\sqrt{\frac{3}{2}n} .$$

For primitive  $G$  the absolute bound (see [D2], [S4]) reads

$$n \leq \frac{1}{2}(f_n^2(G) + 3f_n(G)) .$$

Hence  $\frac{1}{2}n < 2\sqrt{\frac{3}{2}n}$ , i.e.  $n < 24$ . This contradicts our assumption, and therefore the theorem is proved.  $\square$

**2.2.4. EXAMPLE.** Let  $G$  be the Schläfli graph, which is drawn in Figure 1; two black or two white vertices are adjacent iff they are on one line, a black vertex is adjacent to a white one iff they are not on a line (see [S3], [H2]). Then  $G$  is strongly regular,  $n = 27$  and

$$\lambda_1(G) = 16, \lambda_2(G) = \dots = \lambda_7(G) = 4, \lambda_8(G) = \dots = \lambda_{27}(G) = -2,$$

$$\lambda_1(\bar{G}) = 10, \lambda_2(\bar{G}) = \dots = \lambda_{21}(\bar{G}) = 1, \lambda_{22}(\bar{G}) = \dots = \lambda_{27}(\bar{G}) = -5,$$

where  $\bar{G}$  denotes the complement of  $G$ . From Figure 1 we see that

$$\alpha(G) \geq 3 \text{ and } \alpha(\bar{G}) \geq 6 .$$

The thin vertical lines partition  $G$  into six cliques, hence  $\gamma(\bar{G}) \leq 6$ . The numbering gives a colouring of  $G$  with nine colours, so  $\gamma(G) \leq 9$ . Using our bounds it follows that equality holds in all these inequalities. Indeed, by 2.1.4.ii or 2.1.5.i we have  $\alpha(G) \leq 3$ ; 2.1.1 yields  $\alpha(\bar{G}) \leq 6$ ;  $\gamma(G) \geq 9$  follows from Hoffman's bound, and  $\gamma(\bar{G}) \geq 6$  follows from our last theorem.




FIGURE I

The chromatic number of strongly regular graphs will be the subject of Chapter 4.



## CHAPTER 3

## INEQUALITIES FOR DESIGNS

3.1. SUBDESIGNS

In this section we shall derive inequalities for subdesigns of designs.

Let  $D$  be a design with incidence matrix  $N$ . It is clear that we cannot apply the Higman-Sims technique (1.2.3) to  $N$ , because  $N$  does not have to be symmetric. Instead, we apply the Higman-Sims technique to

$$A = \begin{bmatrix} 0 & N \\ N^* & 0 \end{bmatrix}.$$

By definition the positive eigenvalues of  $A$  are the singular values of  $N$ . Let  $\sigma_1 \geq \sigma_2 \geq \dots > 0$  denote these singular values. Then we can state the main result of this section.

3.1.1. THEOREM. *Let  $D$  be a  $1 - (v, k, r)$  design with  $b$  blocks. Let  $D_1$  be a possibly degenerate  $1 - (v_1, k_1, r_1)$  subdesign of  $D$  with  $b_1$  blocks. Then*

- i.  $(vr_1 - b_1k)(bk_1 - v_1r) \leq \sigma_2^2(v - v_1)(b - b_1)$ ,
- ii. *if equality holds, then each point [block] of  $D_1$  is incident with a constant number of blocks [points] of  $D_1$ .*

PROOF. Let  $N_1$  and

$$N = \begin{bmatrix} N_1 & N_2 \\ N_3 & N_4 \end{bmatrix}$$

be the incidence matrices of  $D_1$  and  $D$ , respectively. Put

$$A := \begin{bmatrix} 0 & 0 & N_1 & N_2 \\ 0 & 0 & N_3 & N_4 \\ N_1^* & N_3^* & 0 & 0 \\ N_2^* & N_4^* & 0 & 0 \end{bmatrix} \quad \text{and} \quad B := \begin{bmatrix} 0 & 0 & r_1 & r-r_1 \\ 0 & 0 & x & r-x \\ k & k-k_1 & 0 & 0 \\ y & k-y & 0 & 0 \end{bmatrix},$$

where

$$x := b_1(k-k_1)/(v-v_1) \quad \text{and} \quad y := v_1(r-r_1)/(b-b_1).$$

Then the entries of  $B$  are the average row sums of the block matrices of  $A$ .

By 1.1.2 we know

$$\lambda_i(A) = -\lambda_{b+v+1-i}(A), \quad \lambda_j(B) = -\lambda_{5-j}(B),$$

for  $i = 1, \dots, b+v$ ,  $j = 1, \dots, 4$ . We easily have

$$\sigma_1 = \lambda_1(A) = \lambda_1(B) = \sqrt{rk}.$$

From  $\det B = rk(r_1-x)(k_1-y)$  it now follows that

$$\lambda_2(B) = -\lambda_3(B) = \sqrt{(r_1-x)(k_1-y)}.$$

Now 1.2.3.i gives

$$(r_1-x)(k_1-y) \leq \lambda_2^2(A) = \sigma_2^2.$$

With  $b_1k_1 = v_1r_1$  this yields (i).

If equality holds, then the interlacing must be tight. Thus 1.2.3.ii gives (ii).  $\square$

From the above proof it is clear that the result also holds if  $D_1$  is not a 1-design, but then we have to take  $r_1$  and  $k_1$  to be the average row and column sums of  $N_1$ .

For many 1-designs  $\sigma_2^2$  is expressible in terms of the parameters of the design. For instance,  $\sigma_2^2 = r - \lambda$  if  $D$  is a  $2-(v, k, \lambda)$  design, and  $\sigma_2^2 = s + t - \alpha + 1$  if  $D$  is a partial geometry with parameters  $(s, t, \alpha)$  (see Appendix I, or [C6], [T1]).

We shall make explicit two consequences of the above theorem.

3.1.2. COROLLARY. *If a symmetric 2 - (v,k,λ) design contains a symmetric 2 - (v<sub>1</sub>,k<sub>1</sub>,λ<sub>1</sub>) subdesign, possibly degenerate, then*

$$k \geq \left( k_1 - v_1 \frac{k - k_1}{v - v_1} \right)^2 + \lambda .$$

PROOF. Substitute  $b = v$ ,  $k = r$ ,  $b_1 = v_1$ ,  $k_1 = r_1$  and  $\sigma_2^2 = k - \lambda$  in 3.1.1.i. □

3.1.3. COROLLARY. *Let X and Y be a set of points and a set of lines, respectively, of a partial geometry with parameters (s,t,α), such that no point of X is incident with a line of Y. Then*

$$\begin{aligned} (\alpha|X| + (s+t+1-\alpha)(s+1))(\alpha|Y| + (s+t+1-\alpha)(t+1)) &\leq \\ &\leq (s+t+1-\alpha)(s+1)^2(t+1)^2 . \end{aligned}$$

PROOF. Substitute  $k_1 = r_1 = 0$ ,  $b_1 = |Y|$ ,  $v_1 = |X|$ ,  $k = s+1$ ,  $r = t+1$ ,  $v = (s+1)(st+\alpha)/\alpha$ ,  $b = (t+1)(st+\alpha)/\alpha$  and  $\sigma_2^2 = s+t+1-\alpha$  in 3.1.1.i. □

Corollary 3.1.2 appeared in [H4]. A Baer subplane of a projective plane (see [D3]) satisfies 3.1.2 with equality. Other examples which meet this bound (hence where 3.1.1.ii applies) can be found in [H4].

The bound of 3.1.3 can also be tight. For instance, let Q be the partial geometry with parameters (2,4,1) (generalized quadrangle), whose points and lines are the vertices and the triangles of the complement of the Schläfli graph (see Example 2.2.4). There are 15 triangles which do not have a vertex in common with a double six (the black vertices in Figure 1), Thus we have an empty subgeometry (no point and line are incident) of Q having 12 points and 15 lines. This satisfies 3.1.3 with equality.

If  $D_1$  is an empty design ( $k_1 = r_1 = 0$ ), then one easily finds examples which meet the bound of 3.1.1.i. For instance: a projective plane with a maximal arc (see [D3]); a symmetric 2-design containing an oval without tangent blocks (see [A4]); a 2-design having a block repeated  $b/v$  times (see [L1]; here the inequality 3.1.1.i is Mann's inequality [M2]).

Although the results of this section are similar to those of Section 2.1, we did not start with a general inequality for substructures of an incidence structure like we did for subgraphs of a graph in 2.1.2. This has two reasons. Firstly, the formula for an arbitrary incidence structure is more complicated than 2.1.2. The second reason is that there does not seem to be much interest in incidence structures without any additional properties; this is certainly not true for graphs. Yet we shall give one result for an arbitrary incidence structure, namely an inequality for the sizes of an empty substructure.

**3.1.4. THEOREM.** *Let  $D$  be an incidence structure with  $v$  points and  $b$  blocks. Let every point [block] be incident with at least  $r_{\min}$  blocks [ $k_{\min}$  points]. Let  $X$  and  $Y$  be a set of points and a set of blocks, respectively, such that no point of  $X$  is incident with a block of  $Y$ . Then*

$$r_{\min}^2 k_{\min}^2 |X| |Y| \leq \sigma_1^2 \sigma_2^2 (v - |X|) (b - |Y|) ,$$

where  $\sigma_1$  and  $\sigma_2$  denote the two largest singular values of the incidence matrix of  $D$ .

**PROOF.** Let the incidence matrix of  $D$  be

$$N = \begin{bmatrix} 0 & N_2 \\ N_3 & N_4 \end{bmatrix} ,$$

where  $0$  denotes the  $|X| \times |Y|$  all-zero matrix. Let  $r_i$  and  $k_i$  be the average row sums of  $N_i$  and  $N_i^*$ , respectively, for  $i = 2, 3, 4$ . Then by 1.2.3.i the eigenvalues of

$$B := \begin{bmatrix} 0 & 0 & 0 & r_2 \\ 0 & 0 & r_3 & r_4 \\ 0 & k_3 & 0 & 0 \\ k_2 & k_4 & 0 & 0 \end{bmatrix}$$

interlace the eigenvalues of

$$A := \begin{bmatrix} 0 & N \\ N^* & 0 \end{bmatrix} .$$

Now with 1.1.2 it follows that

$$\det B = \lambda_1^2(B)\lambda_2^2(B) \leq \lambda_1^2(A)\lambda_2^2(A) = \sigma_1^2 \sigma_2^2 .$$

On the other hand we have

$$\det B = r_2 r_3 k_2 k_3 = r_2^2 k_3^2 |X| |Y| / ((v - |X|)(b - |Y|)) .$$

Since  $r_2 k_3 \geq r_{\min} k_{\min}$ , the theorem is proved.  $\square$

### 3.2. INTERSECTION NUMBERS

If two distinct blocks of a design  $D$  have exactly  $\rho$  points in common, then  $\rho$  is called an *intersection number* of  $D$ . It is obvious that an intersection number  $\rho$  of a  $1 - (v, k, r)$  design satisfies

$$k \geq \rho \geq \max\{0, 2k - v\} .$$

The next result, which is due to AGRAWAL [A1], gives non-trivial bounds for the intersection numbers of a 1-design. Like in the previous section, the singular values of the incidence matrix of a design  $D$  will be denoted by  $\sigma_1 \geq \sigma_2 \geq \dots > 0$ .

3.2.1. THEOREM. *Let  $D$  be a  $1 - (v, k, r)$  design with  $b$  blocks. Let  $B_1$  and  $B_2$  be distinct blocks of  $D$ . Then*

$$i. \quad |B_1 \cap B_2| \leq 2 \frac{rk - \sigma_2^2}{b} - k + \sigma_2^2 ;$$

*if equality holds then  $|B_1 \cap B_3| + |B_2 \cap B_3| = 2(rk - \sigma_2^2) / b$  for any further block  $B_3$ ,*

$$ii. \quad |B_1 \cap B_2| \geq k - \sigma_2^2 ;$$

*if equality holds then  $|B_1 \cap B_3| = |B_2 \cap B_3|$  for any further block  $B_3$ .*

PROOF. The result is obvious if  $b \leq 3$ , so assume  $b \geq 4$ . Let  $N$  be the incidence matrix of  $D$ , such that the first two columns correspond to the blocks  $B_1$  and  $B_2$ . Define

$$\bar{A} := N^* N .$$

Then the off-diagonal entries of  $A$  are the intersection numbers of  $D$ , and the row and column sums of  $A$  equal  $kr$ . Put  $\rho := |B_1 \cap B_2|$  and consider the following partitioning of  $A$ :

$$A = \begin{bmatrix} \begin{pmatrix} k & \rho \\ \rho & k \end{pmatrix} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} .$$

Define

$$x := 2(kr - k - \rho) / (b - 2) \quad \text{and} \quad B := \begin{pmatrix} k + \rho & kr - k - \rho \\ x & kr - x \end{pmatrix} .$$

Then the entries of  $B$  are the average row sums of the block matrices of  $A$ . Clearly

$$\lambda_1(B) = \lambda_1(A) = rk \quad \text{and} \quad \lambda_2(B) = k + \rho - x .$$

By 1.2.3.i we have

$$\lambda_2(B) \leq \lambda_2(A) = \sigma_2^2 .$$

Hence

$$\sigma_2^2(b - 2) \leq (k + \rho)(b - 2) - 2(kr - k - \rho) .$$

This yields

$$\rho \leq \sigma_2^2 - k + 2(rk - \sigma_2^2) / b .$$

If equality holds, the interlacing is tight and 1.2.3.ii gives that every column sum of  $A_{12}$  equals  $x$ . This proves (i).

To prove (ii) we apply 1.2.1 to  $A$  with

$$S := \begin{pmatrix} -1 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 1 \end{pmatrix}^* \cdot \begin{pmatrix} 2 & 0 \\ 0 & n-2 \end{pmatrix}^{-\frac{1}{2}} .$$

Then

$$B := S^* A S = \begin{pmatrix} k - \rho & 0 \\ 0 & kr - x \end{pmatrix} .$$

It is easily seen that  $kr - x \geq k - \rho$  if  $b \geq 4$ . So  $\lambda_2(B) = k - \rho$ . Hence, by 1.2.1.i

$$k - \rho \leq \lambda_2(A) = \sigma_2^2 .$$

Here equality does not have to imply that the interlacing is tight. Therefore we shall use 1.2.1.ii. If  $\sigma_2^2 = k - \rho = kr - x$ , then  $r = 1$ ,  $\rho = 0$  or  $r = 2$ ,  $\rho = 0$ ,  $b = 4$ , and the result follows immediately by use of (i). If  $\sigma_2^2 = k - \rho < kr - x$ , then 1.2.1.ii implies that  $S(1,0)^* = (-\sqrt{2}, \sqrt{2}, 0, \dots, 0)^*$  is an eigenvector of  $A$  for the eigenvalue  $k - \rho$ . Thus  $A_{12}^* (-1, 1)^* = 0$ . This proves (ii).  $\square$

It is straightforward to verify that equality in (i) or (ii) for a pair of blocks of  $D$  implies also equality for the corresponding blocks of the complement of  $D$ .

Although Agrawal's proof of the above theorem is different from ours, it also uses eigenvalue techniques (in essence the Cauchy inequalities 1.2.2). MAJUMDAR [M1] gives a proof of this theorem for the case that  $D$  is a 2-design, using counting arguments. See also BUSH [B10] and CONNOR [C8] for similar results.

It is clear that our method also leads to inequalities if we consider the intersection pattern of more than two blocks.

**3.2.2. THEOREM.** *Let  $D$  be a  $1 - (v, k, r)$  design with  $b$  blocks. Let  $Y$  be a set of blocks which mutually have  $\rho$  points in common. Then*

- i.  $|Y| (rk - b\rho) \leq b(k - \rho) ,$
- ii.  $|Y| (b\rho - rk + \sigma_2^2) \leq b(\sigma_2^2 - k + \rho) .$

**PROOF.** Let  $N$  be the incidence matrix of  $D$ . We apply the Higman-Sims technique to  $A := N^*N$ , partitioned according to  $Y$  and the other blocks of  $D$ . Put

$$x := |Y| \frac{k(r-1) - \rho(|Y| - 1)}{b - |Y|} .$$

Then

$$B := \begin{pmatrix} k - \rho + \rho|Y| & rk - k + \rho - \rho|Y| \\ x & rk - x \end{pmatrix}$$

carries the average row sums of the block matrices of A. Clearly

$$\lambda_2(B) = k - \rho + \rho |Y| - x = (b(k - \rho) + (b\rho - kr) |Y|) / (b - |Y|) .$$

From 1.2.3.i we have

$$0 \leq \lambda_b(A) \leq \lambda_2(B) \leq \lambda_2(A) = \sigma_2^2 .$$

This lower and upper bound for  $\lambda_2(B)$  yields (i) and (ii), respectively.  $\square$

We define two blocks  $B_1$  and  $B_2$  of a  $1 - (v, k, r)$  design to be *equivalent* if

$$|B_1 \cap B_2| \in \{k, k - \sigma_2^2\} .$$

Then from 3.2.1.ii it is clear that this indeed defines an equivalence relation, and that the number of common points of two blocks only depends on the equivalence classes of these blocks. By the use of 3.2.2 we find bounds for the size of the equivalence classes.

3.2.3. THEOREM. *Let D be a  $1 - (v, k, r)$  design with b blocks. Let Y be an equivalence class of blocks. Then*

- i.  $k$  and  $k - \sigma_2^2$  cannot both be an intersection number of D,
- ii. if  $k - \sigma_2^2$  is an intersection number of D, then

$$|Y| \leq \frac{b\sigma_2^2}{b\sigma_2^2 - bk + rk} ,$$

- iii. if  $k$  is an intersection number of D, then

$$|Y| \leq \frac{b\sigma_2^2}{bk - rk + \sigma_2^2} .$$

PROOF. Assume  $2r \leq b$ ; we may do so because of the remark right after Theorem 3.2.1. Suppose  $k - \sigma_2^2$  is an intersection number. Then  $k - \sigma_2^2 \geq 0$ , hence  $\sigma_2^2 - k + 2(rk - \sigma_2^2) / b \leq 2k(r - 1) / b < k$ . So 3.2.1.i yields that  $k$  cannot be an intersection number. The combination of (i) and (ii) of 3.2.1 yields  $b\sigma_2^2 - bk + rk \geq \sigma_2^2$ . Now formulas (ii) and (iii) follow immediately



from (i) and (ii) of 3.2.2 by substitution of  $\rho = k - \sigma_2^2$  and  $\rho = k$ , respectively.  $\square$

Suppose D is a  $2 - (v, k, \lambda)$  design, so  $\sigma_2^2 = r - \lambda = (bk - rk) / (v - 1)$ . From 3.2.3.iii it follows that D has at most  $b/v$  repeated blocks; this is the inequality of MANN [M2] (see also [L1]). If D has an intersection number  $k - \sigma_2^2 = k - r + \lambda$ , then 3.2.3.ii implies that the size of any equivalence class is at most  $b/(b - v + 1)$ ; this bound appeared in [B2]. This paper also contains the next result (see also [B1]).

A 2-design with just two distinct intersection numbers is called *quasi-symmetric*. Consider the graph G, whose vertices are the blocks of a quasi-symmetric 2-design D, two vertices being adjacent if the number of points which the corresponding blocks have in common equals the larger intersection number. We call G the *block graph* of D. GOETHALS & SEIDEL [G2] (see also [C6]) proved that the block graph of a quasi-symmetric 2-design is strongly regular.

Now suppose D is a  $2 - (v, k, \lambda)$  design with just three distinct intersection numbers  $k - r + \lambda$ ,  $\rho_1$  and  $\rho_2$  ( $\rho_1 > \rho_2$ ). We have already observed that the number of points which two blocks have in common only depends on the equivalence classes of these blocks. For this reason the following definition is legitimate. The *class graph* of D is the graph whose vertices are the equivalence classes, two vertices being adjacent if two blocks representing the corresponding classes have  $\rho_1$  points in common.

**3.2.4. THEOREM.** *Let D be a  $2 - (v, k, \lambda)$  design with just three intersection numbers,  $k - r + \lambda$ ,  $\rho_1$  and  $\rho_2$ . Then the class graph of D is a strongly regular graph on*

$$n := \frac{b(k - r + \lambda - \rho_1)(k - r + \lambda - \rho_2)}{\lambda k^2 - k(r - \lambda) + (r - \lambda)^2 + b\rho_1\rho_2 - \lambda v(\rho_1 + \rho_2)}$$

*vertices, with eigenvalues*

$$\lambda_1(G) = (\lambda v n - b(k - r + \lambda - \rho_2 + \rho_2 n)) / b(\rho_1 - \rho_2) ,$$

$$\lambda_2(G) = (\rho_2 - k + r - \lambda) / (\rho_1 - \rho_2) ,$$

$$\lambda_n(G) = (b(\rho_2 - k + r - \lambda) - n(r - \lambda)) / b(\rho_1 - \rho_2) .$$

PROOF. Let  $N$  be the incidence matrix of  $D$ . Define  $A := N^*N$ . Then

$$(*) \quad A^2 = N^*(NN^*)N = N^*(\lambda J + (r-\lambda)I)N = \lambda k^2 J + (r-\lambda)A .$$

Put  $\rho_0 := k-r+\lambda$ . Let  $x_j$  ( $j = 0,1,2$ ) denote the number of times that  $\rho_j$  occurs in the  $i$ -th row of  $A$ , for some  $i \in \{1, \dots, b\}$ . Then

$$(AJ)_{ii} = k + x_0(k-r+\lambda) + x_1\rho_1 + x_2\rho_2 = rk ,$$

$$(A^2)_{ii} = k^2 + x_0(k-r+\lambda)^2 + x_1\rho_1^2 + x_2\rho_2^2 = \lambda k^2 + k(r-\lambda) ,$$

on using  $(*)$ . Substitute  $x_2 = b-1-x_0-x_1$  and subtract the first equation multiplied by  $(\rho_1+\rho_2)$  from the second one. This yields

$$\begin{aligned} x_0(k-r+\lambda-\rho_1)(k-r+\lambda-\rho_2) &= \\ &= (b-1)\rho_1\rho_2 - k(r-1)(\rho_1+\rho_2) + k(r-\lambda) + k^2(\lambda-1) . \end{aligned}$$

Hence  $x_0$  does not depend on  $i$  and therefore all equivalence classes have size  $x_0+1$ . Now  $n = b/(x_0+1)$  yields the given formula for  $n$ .

Now we partition  $A$  according to the equivalence classes. Let  $\tilde{A}$  denote the adjacency matrix of  $G$ . Then the definition of  $G$  yields that the entries of

$$(**) \quad B := (b/n)((k-r+\lambda-\rho_2)I + (\rho_1-\rho_2)\tilde{A} + \rho_2J) + (r-\lambda)I$$

are the row sums (which are constant) of the block matrices of  $A$ . Since  $A$  has three distinct eigenvalues,  $rk$ ,  $r-\lambda$  and  $0$ , it follows from 1.2.3.iii that each eigenvalue of  $B$  is equal to  $rk$ ,  $r-\lambda$  or  $0$ . We easily check that  $rk$  is a simple eigenvalue of  $B$ , belonging to the all-one vector  $j$ . Now from  $(**)$  the eigenvalues of  $\tilde{A}$  follow. Hence  $\tilde{A}$  has an eigenvector  $j$  and just two distinct eigenvalues not belonging to  $j$ . This implies (see Appendix I or [C6]) that  $G$  is strongly regular.  $\square$

Examples of designs which satisfy the hypothesis of the above theorem can be found in [B1], [B3] or [M8]. For all these examples the class graph is a complete multipartite graph. In Section 6.1 we shall give an example for the above theorem where the class graph is primitive (not complete multipartite or the complement). For other results on 2-designs with an intersection number  $k-r+\lambda$ , see [B2].

## CHAPTER 4

## 4-COLOURABLE STRONGLY REGULAR GRAPHS

4.1. INTRODUCTION

In this chapter we shall illustrate the use of the results and techniques obtained in the previous chapters. The result will be the determination of all 4-colourable strongly regular graphs.

It is obvious that a regular complete  $\gamma$ -partite graph, and a disjoint union of complete graphs on  $\gamma$  vertices are strongly regular graphs with chromatic number  $\gamma$ . Strongly regular graphs, not belonging to one of these two families, are called *primitive*.

Let  $G$  be a strongly regular graph with parameters  $(n, d, p_{11}^1, p_{11}^2)$ . Then (see [C5], [C9] or Appendix I)

$$d = \lambda_1(G) \quad , \quad p_{11}^2 - d = \lambda_2(G)\lambda_n(G) \quad , \quad p_{11}^1 - p_{11}^2 = \lambda_2(G) + \lambda_n(G) \quad ,$$

$$np_{11}^2 = (d - \lambda_2(G))(d - \lambda_n(G)) \quad .$$

Moreover,  $G$  has at most three distinct eigenvalues:

$$\lambda_1(G) \geq \lambda_2(G) = \dots = \lambda_{f+1}(G) \geq 0 \quad , \quad -1 \geq \lambda_{f+2}(G) = \dots = \lambda_n(G) \quad ,$$

where  $f = f_2(G)$ , the multiplicity of  $\lambda_2(G)$ , satisfies

$$f = (-\lambda_n(G)(n-1) - d) / (\lambda_2(G) - \lambda_n(G)) \quad .$$

If  $G$  is primitive, then  $p_{11}^2 > 0$ ,  $\lambda_2(G) > 0$ , and  $\lambda_n(G) < -1$ .

4.1.1. LEMMA. *If  $G$  is a primitive strongly regular graph, not the pentagon, then*

- i.  $d \leq -\lambda_n(G)(\gamma(G) - 1) \quad ,$
- ii.  $-\lambda_n(G) \leq \lambda_2(G)(\gamma(G) - 1) \quad ,$
- iii.  $\lambda_2(G) < \gamma(G) - 1 \quad .$

PROOF. (i) and (ii) are quoted from 2.2.3. Since  $G$  is primitive,  
 $0 < p_{11}^2 = d - \lambda_2(G)\lambda_n(G)$ . Hence (i) gives  $\gamma(G) - 1 \geq -d/\lambda_n(G) > \lambda_2(G)$ .  $\square$

As a direct consequence of this lemma we have the following theorem.

4.1.2. THEOREM. *Given  $\gamma \in \mathbb{N}$ , the number of primitive strongly regular graphs with chromatic number  $\gamma$  is finite.*

PROOF. If the graph  $G$  is primitive, then  $p_{11}^2 \geq 1$  and hence by use of the formulas above

$$n \leq np_{11}^2 = (d - \lambda_2(G))(d - \lambda_n(G)) \leq d(d - \lambda_n(G)) .$$

By Lemma 4.1.1 we have

$$d(d - \lambda_n(G)) < \gamma(G) (\gamma(G) - 1)^5 .$$

This completes the proof.  $\square$

Now let us examine the case  $\gamma(G) \leq 4$ .

4.1.3. LEMMA. *Let  $G$  be a 4-colourable strongly regular graph. Suppose  $G$  has a non-integral eigenvalue. Then  $G$  is the pentagon.*

PROOF. Since  $G$  has a non-integral eigenvalue, we have (see [C9] or Appendix I)

$$\lambda_1(G) = \frac{1}{2}(n-1) , \quad \lambda_2(G) = -\frac{1}{2} + \frac{1}{2}\sqrt{n} , \quad \lambda_n(G) = -\frac{1}{2} - \frac{1}{2}\sqrt{n} ,$$

$$n \equiv 1 \pmod{4} , \quad \sqrt{n} \notin \mathbb{N} .$$

By 2.1.5 we have  $\alpha(G) \leq \sqrt{n}$ , hence

$$4 \geq \gamma(G) \geq n/\alpha(G) \geq n/\lfloor \sqrt{n} \rfloor ,$$

therefore  $n = 16$  or  $n \leq 12$ . Combining the restrictions for  $n$  we have  $n = 5$ , hence  $G$  is the pentagon.  $\square$

4.1.4. LEMMA. *A 4-colourable primitive strongly regular graph has one of the following parameter sets:*

i.	(5,2,0,1) ,	vii.	(16,9,4,6) ,
ii.	(9,4,1,2) ,	viii.	(40,12,2,4),
iii.	(10,3,0,1),	ix	(50,7,0,1) ,
iv.	(15,6,1,3),	x.	(56,10,0,2),
v.	(16,5,0,2),	xi.	(64,18,2,6),
vi	(16,6,2,2),	xii.	(77,16,0,4).

PROOF. Let  $G$  be such a graph. Suppose  $G$  is not the pentagon (which has parameter set (i)). Then by 4.1.3, the eigenvalues of  $G$  are integers. The primitivity of  $G$  yields  $\lambda_2(G) > 0$ ,  $\lambda_n(G) < -1$ . Now 4.1.1.iii gives

$$\lambda_2(G) \in \{1, 2\} .$$

Suppose  $\lambda_2(G) = 1$ . Then by 4.1.1

$$\lambda_n(G) \in \{-2, -3\} , \quad d = \lambda_1(G) \leq 9 .$$

Straightforward computations give that the only feasible parameter sets satisfying these conditions are (ii) - (v), (vii) and (10,6,3,4). However, a graph  $G$  with this last parameter set satisfies  $\alpha(G) \leq 2$ , therefore  $\gamma(G) \geq 5$ . Suppose  $\lambda_2 = 2$ . Then 4.1.1 implies

$$\lambda_n(G) \in \{-2, -3, -4, -5, -6\} , \quad d \leq 18 .$$

With a little more work than for the previous case, this leads to the feasible parameter sets (vi), (viii) - (xii) and (57,14,1,4). However, WILBRINK & BROUWER [W4] proved the nonexistence of a strongly regular graph with this last parameter set.  $\square$

For graphs with parameters (i) - (v) existence and uniqueness is known (see [S3]). Cases (i), (ii) and (iii) are the pentagon, the line graph of  $K_{3,3}$  (also called the lattice graph  $L_2(3)$ ), and the Petersen graph, respectively. It is easily seen that these three graphs have chromatic number three. From 4.1.1 it is clear that none of the other graphs is 3-colourable. Case (iv) is the complement of the line graph of  $K_6$  (also called the complement of the triangular graph  $T(6)$ ), which is easily seen to be 4-colourable. Case (v) is the Clebsch graph (see [S3]). This graph

is given in Figure 2, where two black or two white vertices are adjacent iff they are not on one line, whilst a black vertex is adjacent to a white one iff they are on one line. We almost immediately see that this graph is




FIGURE 2

4-colourable. There are precisely two nonisomorphic strongly regular graphs with parameter set (vi) (see [S6]): the line graph of  $K_{4,4}$  ( $L_2(4)$ ) and the Shrikhande graph (see fly-leaf), both graphs are easily seen to be 4-colourable. Case (vii) is the complementary parameter set of (vi). We quickly see that the complement of the line graph of  $K_{4,4}$  is 4-colourable, however, the complement of the Shrikhande graph is not 4-colourable. Indeed, the size of the largest coclique equals three. The remaining cases are more difficult. They will be treated in the next section.

#### 4.2. STRONGLY REGULAR GRAPHS ON 40, 50, 56, 64 AND 77 VERTICES

In this section we shall study the feasible parameter sets for 4-colourable strongly regular graphs, which remain from the previous section.

The first case is the parameter set  $(40,12,2,4)$ . Although several strongly regular graphs with these parameters are known (see Section 6.2), it will turn out that no such graph has chromatic number four. To prove this we use the following lemma.

4.2.1. LEMMA. *There is no regular bipartite graph on 20 vertices with eigenvalues 4, 2, 0, -2, -4 of multiplicity 1, 6, 6, 6, 1, respectively.*

PROOF. Suppose  $G$  were such a bipartite graph. Let

$$A = \begin{bmatrix} 0 & N \\ N^* & 0 \end{bmatrix}$$

be the adjacency matrix of  $G$ . Then  $N$  is the incidence matrix of a  $1 - (10, 4, 4)$  design,  $D$  say, with singular values  $\sigma_1 = 4, \sigma_2 = \dots = \sigma_7 = 2$ . Let  $B_1$  and  $B_2$  be two distinct blocks of  $D$ . Then 3.2.1.i yields

$$|B_1 \cap B_2| \leq 12/5 .$$

Suppose  $B_1$  and  $B_2$  are disjoint. Let  $x$  and  $y$  be the two points of  $D$  which are not incident with  $B_1$  and  $B_2$ . Let  $B_3$  be a block through  $x$ . Using 3.2.1.ii it follows that

$$|B_1 \cap B_3| = |B_2 \cap B_3| = 1 ,$$

so  $B_3$  is incident with  $y$ . Hence, any block incident with  $x$  is also incident with  $y$ . However, this is not possible, since two points of  $D$  have at most two blocks in common, as follows from 3.2.1.i applied to the dual of  $D$ . So we have

$$|B_1 \cap B_2| \in \{1, 2\} .$$

This implies that  $B := N^*N - J - 3I$  is the adjacency matrix of a (strongly) regular graph with eigenvalues  $-3, 1$  and  $3$  of multiplicity  $3, 6$  and  $1$ , respectively. This is impossible, since

$$30 = \text{trace } B^2 \neq 3 \cdot (-3)^2 + 6 \cdot 1^2 + 1 \cdot 3^2 . \quad \square$$

4.2.2. THEOREM. *There exists no 4-colourable strongly regular graph with parameters  $(40, 12, 2, 4)$ .*

PROOF. Let  $G$  be a strongly regular graph with parameter set  $(40, 12, 2, 4)$ . The eigenvalues of  $A$ , the adjacency matrix of  $G$ , are  $12, 2$  and  $-4$  of multiplicity  $1, 24$  and  $15$ , respectively. Suppose  $G$  is 4-colourable. Then without loss of generality

$$A = \begin{bmatrix} 0 & A_{12} & A_{13} & A_{14} \\ A_{21} & 0 & A_{23} & A_{24} \\ A_{31} & A_{32} & 0 & A_{34} \\ A_{41} & A_{42} & A_{43} & 0 \end{bmatrix}.$$

By 2.1.5.i all block matrices are square of size 10, and by 2.1.5.iii all row and column sums of  $A_{ij}$  are equal to 4, for  $i, j = 1, 2, 3, 4$ ,  $i \neq j$ . Define

$$A_i := \begin{bmatrix} 0 & A_{i2} \\ A_{21} & 0 \end{bmatrix}, \quad A_2 := \begin{bmatrix} 0 & A_{34} \\ A_{43} & 0 \end{bmatrix}, \quad \tilde{A} := A - \frac{1}{4}J, \quad \tilde{A}_i := A_i - \frac{1}{4}J,$$

for  $i = 1, 2$ . Let  $i \in \{1, 2\}$ . Let  $G_i$  be the graph with adjacency matrix  $A_i$ . Now  $\tilde{A}$  has just two distinct eigenvalues 2 and  $-4$  of multiplicity 25 and 15, respectively. Furthermore,  $A_i$  and  $\tilde{A}_i$  have the same eigenvalues, except for the one belonging to the eigenvector  $j$ , which equals 4 for  $A_i$  and  $-1$  for  $\tilde{A}_i$ . Since  $G_i$  is bipartite,  $A_i$  also has an eigenvalue  $-4$ . Now by the Cauchy inequalities (1.2.2) it follows that  $\tilde{A}_i$ , and hence also  $A_i$ , has at least five times the eigenvalue 2. But  $G_i$  is bipartite, therefore  $A_i$ , and hence also  $\tilde{A}_i$ , has at least five times the eigenvalue  $-2$ . Now from 1.3.3 it follows that  $\tilde{A}_{3-i}$ , and hence also  $\tilde{A}_i$  and  $A_i$ , has at least five times the eigenvalue 0. Since  $G_i$  is bipartite on an even number of vertices, the multiplicity of the eigenvalue 0 is even, so at least six. By the same reasoning as above, Theorem 1.3.3 yields that the multiplicities of the eigenvalues 2 and  $-2$  of  $A_i$  are also at least six. Thus  $A_i$  has eigenvalues 4, 2, 0,  $-2$ ,  $-4$  of multiplicity 1, 6, 6, 6, 1, respectively. Now Lemma 4.2.1 finishes the proof.  $\square$

HOFFMAN & SINGLETON [H15] showed the existence and uniqueness of a strongly regular graph with parameters  $(50, 7, 0, 1)$ . So we only have to determine whether this graph is 4-colourable or not. To do so, we shall use a description of the Hoffman-Singleton graph (this description seems to be folklore, since it is well known; however, I could not find a reference) based on the following result, see [B11] or [C10].

**4.2.3. RESULT.** *The thirtyfive lines of  $PG(3, 2)$  can be represented by the thirtyfive triples of a set with seven elements, such that two lines*



*intersect iff the corresponding triples have exactly one element in common.*

REMARK. This result is directly related to the isomorphism of the groups  $\text{PSL}(4,2)$  and the alternating group on eight symbols (see for instance [B4] or [C10]).

Now we construct the Hoffman-Singleton graph as follows. The vertices are the fifteen points and the thirtyfive lines of  $\text{PG}(3,2)$ . Points are mutually non-adjacent; a point is adjacent to a line iff the point is on that line; two lines are adjacent iff the triples, which correspond with these lines according to the above result, are disjoint. It is an easy exercise to check that this construction indeed gives the desired strongly regular graph.

4.2.4. THEOREM. *The Hoffman-Singleton graph has chromatic number four.*

PROOF. Colour the fifteen points red. Fix two elements  $x$  and  $y$  of the 7-set of Result 4.2.3. Colour lines blue, if they correspond to a triple containing  $x$ . Of the remaining lines, colour those yellow, whose corresponding triple contains  $y$ , and colour the other ones green. From our definition it is obvious that this is a correct colouring of the Hoffman-Singleton graph.  $\square$

GEWIRTZ [G1] showed existence and uniqueness of a strongly regular graph with parameters  $(56,10,0,2)$ . Before giving a description of the Gewirtz graph we first prove the following.

4.2.5. PROPOSITION. *If the Gewirtz graph has two disjoint cocliques of size 16, then its chromatic number equals four.*

PROOF. Assume that the Gewirtz graph has adjacency matrix

$$A = \begin{bmatrix} 0 & A_{12} & A_{13} \\ A_{21} & 0 & A_{23} \\ A_{31} & A_{32} & A_{33} \end{bmatrix},$$

where  $A_{12} = A_{21}^*$  is square of size 16. We know that  $A$  has three distinct eigenvalues, 10, 2 and -4. From 2.1.5 it follows that  $A_{12}$ ,  $A_{21}$ ,  $A_{31}$  and  $A_{32}$

have constant row sums equal to 4. Therefore the graph  $G_3$ , whose adjacency matrix is  $A_{33}$ , is a disjoint union of cycles. Suppose one of these cycles has length  $c < 24$ . Partition  $A_{33}$  according to the vertices of this cycle, and the remaining ones. This induces a partition of  $A$  into sixteen block matrices, and the entries of the matrix

$$B := \begin{pmatrix} 0 & 4 & \frac{1}{4}c & 6-\frac{1}{4}c \\ 4 & 0 & \frac{1}{4}c & 6-\frac{1}{4}c \\ 4 & 4 & 2 & 0 \\ 4 & 4 & 0 & 2 \end{pmatrix}$$

are the average row sums of these block matrices. We immediately see that the eigenvalues of  $B$  are

$$\lambda_1(B) = 10, \quad \lambda_2(B) = 2, \quad \lambda_3(B) = \lambda_4(B) = -4.$$

However, we know that

$$\lambda_1(A) = 10, \quad \lambda_2(A) = 2, \quad \lambda_{55}(A) = \lambda_{56}(A) = -4.$$

Hence the eigenvalues of  $B$  interlace the eigenvalues of  $A$  tightly, thus by 1.2.3.ii all block matrices have constant row sums. Therefore,  $\frac{1}{4}c$  is an integer. This proves that each component of  $G_3$  is a cycle of even length. Thus  $G_3$  is bipartite, and therefore the whole graph is 4-colourable.  $\square$

We use the description of the Gewirtz graph given in [G2], where this graph is obtained as the complement of the block graph of a quasi-symmetric  $2 - (21,6,4)$  design with intersection numbers 0 and 2 (see Section 3.2).

**4.2.6. THEOREM.** *The Gewirtz graph has chromatic number four.*

**PROOF.** Let  $D$  be the quasi-symmetric  $2 - (21,6,4)$  design. It is clear that all blocks through a fixed point of  $D$  yield a coclique in our graph of size 16. To see that there is another coclique of the same size, disjoint from this one, we proceed as follows.  $D$  can be obtained from a  $3 - (22,6,1)$  design  $\tilde{D}$  (the extension of  $PG(2,4)$ , see for instance [C6]) by deleting one point and all blocks through that point (i.e.  $D$  is a residual design of  $\tilde{D}$ ). Take a block  $B$  of  $\tilde{D}$ , which is not a block of  $D$ . An elementary counting argument (see [C2]) shows that there are 16 blocks of  $D$  which are disjoint

from B, and which mutually have 2 points in common. Hence these 16 blocks provide a coclique of size 16 in the Gewirtz graph, which, if B has been chosen appropriately, is disjoint from our previous coclique. Application of 4.2.5 completes the proof.  $\square$

Three  $2 - (16,6,2)$  designs on a common point set are called *linked* if any two blocks from distinct designs have 3 or 1 points in common (see [C1] or [M7]). Let  $N_1, N_2$  and  $N_3$  be incidence matrices of three linked  $2 - (16,6,2)$  designs. Then we know that

$$N_i^* J = N_i J = 6J, \quad N_i^* N_i = N_i N_i^* = 2J + 4I,$$

for  $i = 1, 2, 3$ . Moreover, for  $i, j = 1, 2, 3, i \neq j$ , the matrix

$$N_{ij} := \frac{1}{2}(3J - N_i^* N_j)$$

is a  $(0,1)$  matrix by definition. In fact,  $N_{ij}$  is the incidence matrix of a  $2 - (16,6,2)$  design, since

$$N_{ij} N_{ij}^* = \frac{1}{4}(3J - N_i^* N_j)(3J - N_j^* N_i) = 2J + 4I,$$

by use of the above formulas. Define

$$N_{ii} := \frac{1}{2}J - 2I, \quad N_{0j} := N_j, \quad N_{j0} := N_j^*,$$

for  $i = 0, 1, 2, 3, j = 1, 2, 3$ . Then

$$N_{ij} = N_{ji}^*, \quad N_{ij} N_{jk} = 3J - 2N_{ik},$$

for  $i, j, k = 0, 1, 2, 3$ , as follows readily from the formulas above. This implies that

$$A := \begin{bmatrix} 0 & N_{01} & N_{02} & N_{03} \\ N_{10} & 0 & N_{12} & N_{13} \\ N_{20} & N_{21} & 0 & N_{23} \\ N_{30} & N_{31} & N_{32} & 0 \end{bmatrix}$$

is a symmetric matrix, which satisfies

$$A^2 = 18I + 2A + 6(J - I - A).$$

Hence  $A$  is the adjacency matrix of a strongly regular graph with parameters  $(64, 18, 2, 6)$ , which is 4-colourable. We call this graph the *incidence graph* of the three linked designs.

**4.2.7. THEOREM.** *Let  $G$  be a strongly regular graph with parameters  $(64, 18, 2, \epsilon)$  and chromatic number four. Then  $G$  is the incidence graph of three linked  $2 - (16, 6, 2)$  designs.*

**PROOF.** Suppose  $G$  has adjacency matrix

$$A = \begin{bmatrix} 0 & A_{01} & A_{02} & A_{03} \\ A_{10} & 0 & A_{12} & A_{13} \\ A_{20} & A_{21} & 0 & A_{23} \\ A_{30} & A_{31} & A_{32} & 0 \end{bmatrix}.$$

$A$  has eigenvalues 18, 2, and  $-6$  of multiplicity 1, 45 and 18, respectively. By 2.1.5,  $A_{ij}$  is square of size 16, and all row and column sums of  $A_{ij}$  are equal to 6, for  $i, j = 0, 1, 2, 3$ ,  $i \neq j$ . This implies that  $A$  and

$$K := I_4 \otimes J_{16}$$

have a common basis of eigenvectors. Using this, we obtain that

$$\tilde{A} := A - 2I + \frac{1}{2}K$$

has eigenvalues 24, 0 and  $-8$  of multiplicity 1, 48 and 15, respectively. Thus

$$\text{rank } \tilde{A} = 16.$$

For  $i = 0, 1, 2, 3$ , put  $A_{ii} := \frac{1}{2}J_{16} - 2I_{16}$ . Then

$$\text{rank } A_{ii} = 16 \quad \text{and} \quad \tilde{A} = \begin{bmatrix} A_{00} & \cdots & A_{03} \\ \vdots & & \vdots \\ A_{30} & \cdots & A_{33} \end{bmatrix}.$$

From  $\text{rank } A_{00} = \text{rank } \tilde{A}$ , it follows that  $A_{i0} A_{00}^{-1} A_{0j} = A_{ij}$ , for  $i, j = 1, 2, 3$ , on applying 1.1.3. By use of

$$A_{00}^{-1} = (\frac{1}{2}J - 2I)^{-1} = (1/24)J - \frac{1}{2}I \quad \text{and} \quad A_{ij}J = 6J$$

this implies that for  $i, j = 1, 2, 3$

$$(*) \quad \frac{3}{2}J - \frac{1}{2}A_{0i}^*A_{0j} = A_{ij}.$$

This completes the proof. Indeed, if  $i = j$ , then  $(*)$  implies  $A_{0i}A_{0i}^* = 2J + 4I$ , showing that  $A_{0i}$  is the incidence matrix of a  $2 - (16, 6, 2)$  design, and if  $i \neq j$ , then  $(*)$  reflects that the designs represented by  $A_{01}$ ,  $A_{02}$  and  $A_{03}$  are linked, and that  $A_{ij}$  is of the desired form for  $i, j = 1, 2, 3$ .  $\square$

MATHON [M7] proved that there are exactly twelve non-isomorphic triples of linked  $2 - (16, 6, 2)$  designs, which lead to eleven non-isomorphic incidence graphs. Hence there are precisely eleven non-isomorphic 4-colourable strongly regular graphs with parameters  $(64, 18, 2, 6)$ . It is fairly easy to show that one of these graphs is the point graph of the known generalized quadrangle of order  $(3, 5)$  (see Chapter 5; a construction is described in 6.2.3). For completeness we list in Appendix II the systems of three linked  $2 - (16, 6, 2)$  designs, which provide the ten remaining graphs; these systems are taken from Mathon's paper. It is not known whether there are any further strongly regular graphs with these parameters, which are not 4-colourable.

Finally, the next theorem deals with the last set of parameters of Lemma 4.1.4.

**4.2.8. THEOREM.** *There exists no 4-colourable strongly regular graph with parameters  $(77, 16, 0, 4)$ .*

**PROOF.** Let  $G$  be a strongly regular graph with parameters  $(77, 16, 0, 4)$ . Then  $G$  has eigenvalues 16, 2 and  $-6$  of multiplicity 1, 55 and 21, respectively. Suppose  $G$  is 4-colourable, and let  $c$  be the size of the largest colour class. Then  $c \geq \lceil 77/4 \rceil = 20$ . Let

$$A = \begin{bmatrix} 0 & A_{12} \\ A_{21} & A_{22} \end{bmatrix},$$

where  $0$  is square of size  $c$ , be the adjacency matrix of  $G$ . Then  $G_2$ , the graph with adjacency matrix  $A_{22}$ , is 3-colourable. From 2.1.7 it follows that  $c \leq \alpha(G) \leq 21$ , and that  $c = \alpha(G) = 21$  implies that  $G_2$  is the Gewirtz graph.

Since the Gewirtz graph has chromatic number four,  $G$  cannot be coloured with four colours if  $c = 21$  (this shows that  $G$  is 5-colourable if  $\alpha(G) = 21$ , which is the case for the known strongly regular graph with these parameters; see [G2]). Whence  $c = 20$ . Now we apply 1.3.3 to  $A - (2/11)J$ , so as to obtain that

$$\lambda_{57}(A_{22} - \frac{2}{11}J) = -6, \quad \lambda_{56}(A_{22} - \frac{2}{11}J) = -4.$$

Now 1.3.1 gives

$$\lambda_{57}(A_{22}) \geq -6, \quad \lambda_{56}(A_{22}) \geq -4.$$

On the other hand, the average row sum of  $A_{22}$  equals  $592/57$ . Hence by 1.2.3.i (take  $m = 1$ ) we have

$$\lambda_1(A_{22}) \geq 592/57.$$

Now the sharpened version of Hoffman's inequality for the chromatic number (see p. 24), applied to the 3-colourable graph  $G_2$  gives

$$-\lambda_{57}(A_{22}) - \lambda_{56}(A_{22}) \geq \lambda_1(A_{22}).$$

This is a contradiction, proving the theorem.  $\square$

#### 4.3. RECAPITULATION.

All cases of Lemma 4.1.4 have been treated now. The only thing left is to state the main theorem.

4.3.1. THEOREM. *If  $G$  is a 4-colourable strongly regular graph, then one of the following holds:*

- i.  $\gamma(G) = 2$ , and  $G$  is a regular complete bipartite graph, or a disjoint union of edges;
- ii.  $\gamma(G) = 3$ , and  $G$  is a regular complete 3-partite graph, a disjoint union of triangles, the pentagon, the line graph of  $K_{3,3}$ , or the Petersen graph;
- iii.  $\gamma(G) = 4$ , and  $G$  is a regular complete 4-partite graph, a disjoint union of  $K_4$ 's, the complement of the line graph of  $K_6$ , the line graph of  $K_{4,4}$ , or its complement, the Shrikhande graph, the

*Clebsch graph, the Hoffman-Singleton graph, the Gewirtz graph,  
or one of the eleven incidence graphs of three linked 2 - (16,6,2)  
designs.*

CHAPTER 5  
GENERALIZED POLYGONS

5.1. INTRODUCTION

A *generalized n-gon of order (s,t)*,  $s > 0$ ,  $t > 0$ , is a  $1 - (v, s+1, t+1)$  design whose incidence graph has girth  $2n$  and diameter  $n$  (see [T4], [D3], [F1], [H12] or Appendix I). A *generalized polygon* is a generalized  $n$ -gon for some  $n$ .

Generalized  $n$ -gons were introduced by TITS [T4]. An important result is the theorem of FEIT & HIGMAN [F1] (see also [H12], [K2]), which states that a generalized  $n$ -gon of order  $(s,t)$  is an ordinary  $n$ -gon ( $s=t=1$ ) or  $n \in \{2,3,4,6,8,12\}$ .

For a generalized polygon we speak of lines rather than blocks. We shall often omit the adjective "generalized". If  $D$  is a polygon of order  $(s,t)$ , then we immediately see that the dual of  $D$  (points and lines interchanged) is a polygon of order  $(t,s)$ .

Suppose  $N$  is the incidence matrix of the incidence graph of an  $n$ -gon of order  $(s,s)$ . Then  $N$  is the incidence matrix of a  $2n$ -gon of order  $(1,s)$ . Conversely, it can be proved easily that all generalized  $n$ -gons of order  $(1,s)$ ,  $s > 1$ , are of this form. FEIT & HIGMAN [F1] also proved that  $s = 1$  or  $t = 1$  for a 12-gon of order  $(s,t)$ , thus in a sense generalized 12-gons are the same as generalized hexagons of order  $(s,s)$ . Generalized  $n$ -gons of order  $(s,t)$  with  $s > 1$ ,  $t > 1$ , are called *thick*.

A generalized 2-gon is degenerate (every point is incident with every line). It is not difficult to verify that a generalized triangle of order  $(s,t)$  is a  $2 - (s^2+s+1, s+1, 1)$  design, which is the same as a projective plane of order  $s$  (thus  $s = t$ ). So 3-gons of order  $(s,s)$  exist for every prime power  $s$ . For projective planes see DEMBOWSKI [D3] or HUGHES & PIPER [H16].

Thick generalized quadrangles of order  $(s,t)$  are known to exist for  $(s,t)$ ,  $(t,s) = (q,q)$ ,  $(q,q^2)$ ,  $(q^2,q^3)$ ,  $(q-1,q+1)$ , for every prime power  $q$  ( $q \neq 2$  for the last case). Constructions are due to AHRENS & SZEKERES [A2], HALL [H6], KANTOR [K1], PAYNE [P1], [P2] and TITS [T4], see also [D3], [T3]. HIGMAN [H11] showed that .



$$s \leq t^2 \leq s^4$$

for thick quadrangles of order  $(s,t)$ . Several other proofs of this inequality have been found, see [C3], [C5], [H12], [P6] and Section 3 of this chapter; some of these proofs also lead to consequences for the case of equality. There is an extended literature on generalized quadrangles. We mention the survey papers [P4], [T1] and [T3].

Thick generalized hexagons are known to exist for the orders  $(s^3,s)$ ,  $(s,s)$  and  $(s,s^3)$  for prime power  $s$ , see [T4]. A necessary condition for existence of a hexagon of order  $(s,t)$  is that  $st$  be a square, see [F1]. HAEMERS & ROOS [H3] showed that

$$s \leq t^3 \leq s^9$$

for thick hexagons of order  $(s,t)$ . This inequality, together with a result for the case of equality, will be the subject of the next section. For more information about generalized hexagons we refer to [M4], [R1], [S1], [T4], [Y1].

Thick generalized octagons of order  $(s,t)$  are only known to exist for  $(t,s)$ ,  $(s,t) = (2^m, 2^{2m})$ , for odd  $m$ . The construction is due to J. Tits, see [D3]. A necessary condition for existence of an octagon of order  $(s,t)$  is that  $2st$  be a square, see [F1]. HIGMAN [H12] showed that

$$s \leq t^2 \leq s^4$$

for thick octagons. There is hardly any literature about octagons.

Let  $G$  be a connected graph of diameter  $m$ . For vertices  $x$  and  $y$  of  $G$ , let  $\rho(x,y)$  denote the distance between  $x$  and  $y$ . For  $i, j = 0, \dots, m$ , define

$$p_{ij}(x,y) := |\{z \mid \rho(x,z) = i \text{ \& \ } \rho(y,z) = j\}| .$$

If  $p_{ij}(x,y)$  depends on  $i, j$  and  $\rho(x,y)$  only, then  $G$  is called *distance regular* (see [B5]), and we write  $p_{ij}^k := p_{ij}(x,y)$  where  $k := \rho(x,y)$ , and  $d_i := p_{ii}^0$ , for  $i, j = 0, \dots, m$ . The numbers  $p_{ij}^k$  are called the *intersection numbers* of  $G$ . Clearly, a distance regular graph is regular of degree  $d_1$ , and a distance regular graph of diameter 2 is the same as a connected strongly regular graph (in general, a distance regular graph of diameter  $m$  is equivalent to a metric association scheme with  $m$  classes, see [D1]). For a distance regular graph  $G$  of diameter  $m$ , we define the matrices  $A_0, \dots, A_m$ , indexed by the vertices of  $G$ , by

$$(A_i)_{xy} = \begin{cases} 1 & \text{if } \rho(x,y) = i, \\ 0 & \text{otherwise.} \end{cases}$$

Clearly,  $A_0 = I$ ,  $\sum_{i=0}^m A_i = J$  and  $A_1$  is the adjacency matrix of  $G$ . Moreover,

$$(A_i A_j)_{xy} = p_{ij}^{\rho(x,y)}$$

implies

$$A_i A_j = \sum_{k=0}^m p_{ij}^k A_k, \quad A_i J = d_i J, \quad \text{for } i, j = 0, \dots, m.$$

These equations show that  $A_0, \dots, A_m$  generate an  $m+1$  dimensional algebra. This type of algebra turns out to be useful in the study of distance regular graphs and similar configurations, see [B5], [B8], [D1], [H12], [W1].

The *point graph* of a generalized  $n$ -gon  $D$  is the graph whose vertices are the points of  $D$ , two points being adjacent whenever they are on one line of  $D$ . It is well known (see [D3], [Y2]) that the point graph  $G$  of an  $n$ -gon of order  $(s, t)$  is distance regular of diameter  $\lfloor \frac{1}{2}n \rfloor$ , and that the intersection numbers of  $G$  can be expressed in terms of  $s$  and  $t$  (in the forthcoming sections of this chapter we shall exhibit this result for  $n = 6$  and  $n = 4$ ).

A graph  $G$  is called *geometric for an  $n$ -gon* if  $G$  is the point graph of an  $n$ -gon. A graph  $G$  is called *pseudo-geometric for an  $n$ -gon* if  $G$  is distance regular of diameter  $\lfloor \frac{1}{2}n \rfloor$  and its intersection numbers are such that  $G$  could be geometric, that is, there exist integers  $s$  and  $t$ , such that the intersection numbers of  $G$  depend on  $s$  and  $t$  as for geometric graphs.

Let  $D$  be a generalized polygon. An *element* of  $D$  is a point or a line of  $D$ . A sequence of  $\ell + 1$  elements  $e_0, \dots, e_\ell$ , is called a *path of length  $\ell$  between  $e_0$  and  $e_\ell$* , if  $e_i$  is incident with  $e_{i-1}$  for  $i = 1, \dots, \ell$  (thus in  $e_0, \dots, e_\ell$ , points and lines alternate). The *distance* between elements  $e_0$  and  $e_\ell$  of  $D$ , denoted by  $\lambda(e_0, e_\ell)$ , is the length of the shortest path between  $e_0$  and  $e_\ell$ . Thus, if  $e_0$  and  $e_\ell$  are both points, then  $\lambda(e_0, e_\ell)$  is twice the distance between  $e_0$  and  $e_\ell$  in the point graph of  $D$ .

In the next two sections we shall describe a method which for quadrangles and hexagons leads to the inequalities mentioned above, and to the

results in case of equality (unfortunately, this method does not work for octagons). The same method also yields a new proof of a theorem of CAMERON, GOETHALS & SEIDEL [C5], which states that a pseudo-geometric graph for a quadrangle of order  $(s, s^2)$  is geometric.

## 5.2. AN INEQUALITY FOR GENERALIZED HEXAGONS

For  $n = 6$  the definition of a generalized  $n$ -gon is equivalent to the following one:

**5.2.1. DEFINITION.** *A generalized hexagon of order  $(s, t)$  is an incidence structure with points and lines, such that*

- i. *each line has  $s+1$  points,*
- ii. *each point is on  $t+1$  lines,*
- iii. *two distinct lines meet in at most one point,*
- iv. *for any non-incident point-line pair  $x, L$  there is a unique path of length  $< 6$  between  $x$  and  $L$ .*

Throughout this section  $H$  will denote a generalized hexagon of order  $(s, t)$ . By use of the above definition it is straightforward to count the intersection numbers  $p_{ij}^k$  of the point graph of  $H$ . They are exhibited in Table 1. The amount of work in computing these numbers can be reduced by use of the equalities

$$\sum_{i=0}^3 p_{ij}^k = d_j, \quad \sum_{j=0}^3 d_j = v = (s+1)(s^2t^2 + st + 1).$$

This counting also shows that the point graph of a generalized hexagon is distance regular.

TABLE 1

$k$	$p_{11}^k$	$p_{12}^k$	$p_{22}^k$	$p_{13}^k$	$p_{23}^k$	$p_{33}^k$
0	$s(t+1)$	0	$s^2t(t+1)$	0	0	$s^3t^2$
1	$s-1$	$st$	$st(s-1)$	0	$s^2t^2$	$s^2t^2(s-1)$
2	1	$s-1$	$s(t^2+t-1)$	$st$	$st(s-1)(t+1)$	$st(s^2t-st-s+t)$
3	0	$t+1$	$(s-1)(t+1)^2$	$(s-1)(t+1)$	$(t+1)(s^2t-st-s+t)$	$t(t-1)(s^2t-s+t)$

Let  $A_0, \dots, A_3$  be the matrices of the point graph of  $H$ . Define

$$E := A_2 - (s-1)A_1 + (s^2 - s + 1)I - \frac{1}{s+1}J .$$

Then we have the following lemma.

5.2.2. LEMMA. *The eigenvalues of  $E$  are*

$$0 \text{ and } s^2 + st + t^2$$

*of multiplicity*

$$1 + st(s+1)(t+1) \frac{s^2t + st^2 - st + s + t}{s^2 + st + t^2} \text{ and } s^3 \frac{s^2t^2 + st + 1}{s^2 + st + t^2} ,$$

*respectively.*

PROOF. It is clear that  $EJ = 0$ , and that

$$\left( E - (s^2 - s + 1)I + \frac{1}{s+1}J \right)^2 = (s-1)^2 A_1^2 - (s-1)(A_1A_2 + A_2A_1) + A_2^2 .$$

From  $A_iA_j = \sum_{k=0}^3 p_{ij}^k A_k$  and Table 1 we have

$$A_2^2 = s^2t(t+1)I + st(s-1)A_1 + (st^2 + st - s)A_2 + (s-1)(t+1)^2 A_3 ,$$

$$A_1A_2 = A_2A_1 = st A_1 + (s-1)A_2 + (t+1)A_3 ,$$

$$A_1^2 = s(t+1)I + (s-1)A_1 + A_2 .$$

By use of  $A_1 + A_2 + A_3 = J - I$  and  $EJ = 0$ , this leads to

$$E(E - (s^2 + st + t^2)I) = 0 .$$

Hence 0 and  $s^2 + st + t^2$  are the eigenvalues of  $E$ . Finally,

$$\text{trace } E = v(s^2 - s + 1 - (s+1)^{-1}) = s^3(s^2t^2 + st + 1)$$

yields the multiplicities. □

REMARK. In the terminology of DELSARTE [D1], the matrix  $(s^2 + st + t^2)^{-1} E$  is a minimal idempotent in the Bose-Mesner algebra of the association scheme on the points of  $H$ . The underlying theory provides a more elegant

way to prove the previous lemma.

Let  $L_0$  be a line of  $H$ , let  $L_i$  ( $i = 1, 2$ ) denote the set of points at distance  $2i + 1$  from  $L_0$ . Partition  $E$  according to  $L_0, L_1$  and  $L_2$ :

$$E = \begin{bmatrix} E_{00} & E_{01} & E_{02} \\ E_{10} & E_{11} & E_{12} \\ E_{20} & E_{21} & E_{22} \end{bmatrix},$$

so  $E_{ij} = E_{ji}^*$  for  $i, j = 0, 1, 2$ .

5.2.3. LEMMA. *The eigenvalues of  $E_{11}$  are  $0, s^2$  and  $st$  of multiplicity  $st - s + t, t(s^2 - 1)$  and  $s$ , respectively.*

PROOF. Let  $A_{1,11}$  and  $A_{2,11}$  be the submatrices of  $A_1$  and  $A_2$ , respectively, corresponding to  $L_1$ . We easily see that without loss of generality

$$I + A_{1,11} = I_{st+t} \otimes J_s \quad \text{and} \quad I + A_{1,11} + A_{2,11} = I_{s+1} \otimes J_{st}$$

holds, hence

$$E_{11} = I_{s+1} \otimes J_{st} - s(I_{st+t} \otimes J_s) + s^2 I - \frac{1}{s+1} J.$$

From the eigenvalues of  $I_{s+1} \otimes J_{st}$ ,  $I_{st+t} \otimes J_s$ ,  $I$  and  $J$ , and the fact that these four matrices have a common basis of eigenvectors, the eigenvalues of  $E_{11}$  and their multiplicities follow.  $\square$

5.2.4. THEOREM. *A generalized hexagon with  $s+1$  points on a line and  $t+1$  lines through a point satisfies*

- i.  $t \leq s^3$  or  $s = 1$ ,
- ii.  $s \leq t^3$  or  $t = 1$ .

PROOF. From 5.2.2 and 5.2.3 it follows that

$$\text{rank } E = s^3(s^2t^2 + st + 1) / (s^2 + st + t^2), \quad \text{rank } E_{11} = s^2t + s - t.$$

Since  $\text{rank } E_{11} \leq \text{rank } E$ , we have

$$(s^2 t + s - t)(s^2 + st + t^2) \leq s^3 (s^2 t^2 + st + 1) .$$

This yields

$$t^2 (s^2 - 1)(t - s^3) \leq 0 .$$

Thus  $s = 1$  or  $t \leq s^3$ . Applying this result to the dual of H yields (ii).  $\square$

For another proof of this inequality see [H3]. Next we shall derive some additional regularity for hexagons meeting the above bound. To achieve this we need some properties of the matrices  $E_{ij}$ . First we observe that

$$(E_{02})_{xy} = \begin{cases} 1 - 1/(s+1) & \text{if } \lambda(x,y) = 4 , \\ -1/(s+1) & \text{otherwise,} \end{cases}$$

as follows directly from the definition of E and  $E_{02}$ . This implies

$$(E_{02}^* E_{02})_{xy} = \begin{cases} 1 - 1/(s+1) & \text{if } \lambda(x,z) = \lambda(y,z) \text{ for some } z \in L_0 , \\ -1/(s+1) & \text{otherwise.} \end{cases}$$

Hence, without loss of generality

$$E_{02} = j^* \otimes I_{s+1} - \frac{1}{s+1} J_{r(s+1)} ,$$

$$E_{02}^* E_{02} = J_r \otimes I_{s+1} - \frac{1}{s+1} J_{r(s+1)} ,$$

where  $r := s^2 t^2$ . Now the positions of the points of  $L_2$  relative to the points of  $L_0$  give rise to a partition of  $E_{22}$  into  $(s+1)^2$  square block matrices  $F_{ij}$  of size  $s^2 t^2$ :

$$E_{22} = \begin{bmatrix} F_{00} & \cdots & F_{0s} \\ \vdots & & \vdots \\ F_{s0} & \cdots & F_{ss} \end{bmatrix} .$$

It is a matter of straightforward counting to see that  $F_{ij}$  has constant row sums equal to  $t^2(s - 1 + \delta_{ij}) - s^2 t^2 / (s+1)$  for  $i, j = 0, \dots, s$ . This, and the structure of  $E_{02}$  imply that

$$E_{22} E_{02}^* = t^2 E_{02}^* .$$

The following identities are now quickly seen to be true:

$$E_{00} = s^2 \left( I - \frac{1}{s+1} J \right), \quad E_{ij} J = 0 \quad \text{for } i, j = 0, 1, 2,$$

$$E_{00}^2 = s^2 E_{00}, \quad E_{00} E_{02} = s^2 E_{02}, \quad E_{02} E_{02}^* = t^2 E_{00}.$$

5.2.5. LEMMA. *The matrix*

$$\tilde{E}_{22} := E_{22} - s^{-2} E_{02}^* E_{02}$$

*has eigenvalues*

$$0, \quad st+t^2 \quad \text{and} \quad s^2+st+t^2;$$

*the multiplicity of  $s^2+st+t^2$  equals*

$$t^2(s^2-1)(s^3-t) / (s^2+st+t^2).$$

PROOF. Define

$$U := \begin{bmatrix} E_{00} \\ E_{20} \end{bmatrix}, \quad E' := \begin{bmatrix} E_{00} & E_{02} \\ E_{20} & E_{22} \end{bmatrix}.$$

Using the above formulas we obtain

$$(*) \quad UU^*U = s^2(s^2+t^2)U, \quad E'U = (s^2+t^2)U.$$

Since  $\text{rank } U = s$ , the last formula reflects that the columns of  $U$  span an  $s$ -dimensional eigenspace of  $E'$  corresponding to the eigenvalue  $s^2+t^2$ .

Thanks to 5.2.2 and 5.2.3 the eigenvalues of  $E$  and  $E_{11}$  are known. By use of 1.3.3 we obtain that the non-zero eigenvalues of  $E'$  are

$$s^2+t^2, \quad st+t^2 \quad \text{and} \quad s^2+st+t^2,$$

of multiplicity

$$s, \quad t(s^2-1) \quad \text{and} \quad t^2(s^2-1)(s^3-t) / (s^2+st+t^2),$$

respectively. Now from (\*) it follows that

$$E' - s^{-2} U^* U$$

has just two distinct non-zero eigenvalues  $st+t^2$  and  $s^2+st+t^2$ , with the

same multiplicities as before. On the other hand one easily verifies that

$$E' - s^{-2} U^* U = \begin{bmatrix} 0 & 0 \\ 0 & \tilde{E}_{22} \end{bmatrix},$$

which proves the lemma. □

The important thing in the last lemma is that the eigenvalue  $s^2 + st + t^2$  disappears if  $t = s^3$ . In order to give a combinatorial interpretation of this phenomenon, we need two definitions. For a line  $L$  and points  $x$  and  $y$  of a generalized hexagon we define:

$$P_{ijk}(L, x, y) := |\{z \mid \lambda(z, L) = 2i + 1, \lambda(z, x) = 2j, \lambda(z, y) = 2k\}|,$$

for  $i = 0, 1, 2, j, k = 0, 1, 2, 3$ ; the configuration *induced* by  $L, x$  and  $y$  is the configuration formed by the points and the lines, which are on a shortest path between  $L$  and  $x, L$  and  $y$ , or  $x$  and  $y$ . For example, Figure 3 gives all possible configurations induced by  $L, x$  and  $y$  if  $\lambda(L, x) = \lambda(L, y) = 5$  and  $\lambda(x, y) = 4$ .




FIGURE 3

**5.2.6. THEOREM.** *If a generalized hexagon has order  $(s, s^3)$ , then  $P_{ijk}(L, x, y)$  only depends on  $i, j, k$  and the configuration induced by  $L, x$  and  $y$ .*

**PROOF.** First observe that



$$\sum_{i=0}^2 p_{ijk}(L, x, y), \quad \sum_{j=0}^3 p_{ijk}(L, x, y) \quad \text{and} \quad \sum_{k=0}^3 p_{ijk}(L, x, y)$$

only depend on  $i, j, k$  and the configuration induced by  $L, x$  and  $y$  (in fact,  $\sum_{i=0}^2 p_{ijk}(L, x, y) = p_{jk}^{\rho}(x, y)$ ). Subsequently, we verify (this is an easy but tiresome job), that the theorem is true if  $i = 0$ , or  $j \leq 1$ , or  $k \leq 1$ , and also if  $\lambda(L, x) < 5$  or  $\lambda(L, y) < 5$ . Thus it suffices to prove the theorem for  $i = j = k = 2$ ,  $\lambda(L, x) = \lambda(L, y) = 5$ . From the definitions of  $E, E_{22}$  and  $p_{ijk}(L, x, y)$  it follows that

$$\begin{aligned} \left( (E_{22} - \frac{1}{s+1} J)^2 \right)_{xy} &= (s^2 - s + 1)^2 p_{200}(L_0, x, y) + \\ &- 2(s-1)(s^2 - s + 1)p_{201}(L_0, x, y) + 2(s^2 - s + 1)p_{202}(L_0, x, y) + \\ &+ (s-1)^2 p_{211}(L_0, x, y) - (s-1)(p_{212}(L_0, x, y) + p_{221}(L_0, x, y)) + \\ &+ p_{222}(L_0, x, y). \end{aligned}$$

Now take  $t = s^3$ . Lemma 5.2.5 implies that  $\tilde{E}_{22}$  has just two distinct eigenvalues 0 and  $s^4(1+s^2)$ . Hence

$$\tilde{E}_{22}^2 = s^4(1+s^2)\tilde{E}_{22}.$$

Using the formulas for the matrices  $E_{ij}$  this yields

$$\left( E_{22} - \frac{1}{s+1} J \right)^2 = s^4(1+s^2)E_{22} - s^2 E_{02}^* E_{02} + s^8(s+1)^{-1} J.$$

This implies that  $\left( (E_{22} - (s+1)^{-1} J)^2 \right)_{xy}$  only depends on the configuration induced by  $L_0, x$  and  $y$ . Combination with the previous steps yields that for  $x, y \in L_2$ ,  $p_{222}(L_0, x, y)$  only depends on the configuration induced by  $L_0, x$  and  $y$ . This completes the proof.  $\square$

With the available formulas the values of  $p_{ijk}(L, x, y)$  are readily computed. For example, in Figure 3 we give  $p_{222}(L, x, y)$  for the given configurations.

RONAN [R1], [R2] and THAS [T2], give sufficient conditions for a generalized hexagon to be one of the known ones. One hopes of course that a result like the one above will imply such a sufficient condition. Unfor-


tunately, the gap between the condition we have and the condition we need, still seems to be too large to close up. It is worthwhile to remark that the known hexagons of order  $(s,t)$  with  $t \neq s^3$  do not satisfy the condition of the above theorem.


Finally we remark that similar techniques yield the inequality (see MATHON [M6])


$$t \leq s^3 + t_2(s^2 - s + 1)$$

for a regular near hexagon with parameters  $(s,t,t_2)$ , as introduced by SHULT & YANUSHKA [S7] (see also [Y2]). If  $t_2 = 0$ , then a near hexagon is the same as a generalized hexagon.

### 5.3. GEOMETRIC AND PSEUDO-GEOMETRIC GRAPHS FOR GENERALIZED POLYGONS

In this section we deal with the question whether a pseudo-geometric graph is geometric for a generalized  $n$ -gon. It is clear that for  $n \in \{2,3\}$  the point graph of an  $n$ -gon is the complete graph. Assume  $G$  is the point graph of an  $n$ -gon  $D$  with  $n > 3$ . Then three points of  $D$  which form a triangle in  $G$ , must lie on one line of  $D$ . This implies that the graph  cannot be an induced subgraph of  $G$ . The next result states that the converse is also true.


5.3.1. LEMMA. *For a generalized  $n$ -gon with  $n > 3$ , a pseudo-geometric graph  $G$  is geometric iff  is not an induced subgraph of  $G$ .*

PROOF. Only the "if" part remains to be proved. Take  $n$  even (the case  $n$  odd is not difficult, but superfluous because of the Feit-Higman theorem). Let  $D$  be the incidence structure whose points are the vertices of  $G$ , and whose lines are the cliques (= complete subgraphs) of  $G$  of size  $p_{11}^1 + 2$ . For two adjacent vertices of  $G$ , there are  $p_{11}^1$  vertices adjacent to both, but all these vertices are mutually adjacent since otherwise  occurs. This means that every edge of  $G$  determines a unique line of  $D$ . This proves that  $D$  is a  $1 - (v, s+1, t+1)$  design, where  $s = p_{11}^1 + 1$  and  $t+1 = d_1/s$ . Let  $G'$  be the incidence graph of  $D$ . Suppose  $G$  has a  $c$ -cycle as an induced subgraph. Then from  $p_{1,k-1}^k = 1$  and  $p_{1,k}^k = s - 1$  for  $1 \leq k < \frac{1}{2}n$  it follows that  $c \geq n$  for  $c$  even, and  $c = 3$  or  $c > n$  for  $c$  odd. Therefore, since each triangle of  $G$  is on a line of  $D$ , we have that  $G'$  has girth at least  $2n$ . Now  $t+1 = p_{1, \frac{1}{2}n-1}^{\frac{1}{2}n}$

implies that the distance between a point and a line of  $D$  (regarded as vertices of  $G'$ ) is at most  $n-1$ . Hence  $G'$  has diameter  $n$  and girth  $2n$ . This proves that  $D$  is an  $n$ -gon, whose point graph is  $G$ .  $\square$

A direct consequence of this lemma (which was pointed out to me by D.E. Taylor) is the following.

5.3.2. COROLLARY. *For a generalized  $n$ -gon with  $n > 4$ , a pseudo-geometric graph is geometric.*

PROOF. If  $n > 4$ , then  $p_{11}^2 = 1$ . Hence  does not occur in a pseudo-geometric graph. Now 5.3.1 gives the result.  $\square$

What remains to be studied are generalized quadrangles. A very easy counting argument shows that the point graph  $G$  of a quadrangle of order  $(s,t)$  is strongly regular with intersection numbers  $p_{11}^1 = s-1$ ,  $p_{11}^2 = t+1$ ,  $p_{11}^0 = d_1 = s(t+1)$  (this proves that the point graph of an  $n$ -gon is distance regular, in the case  $n = 4$ ). This implies (see [C5], [T1] or Appendix I) that the eigenvalues of  $G$ , and hence the eigenvalues of any pseudo-geometric graph for a quadrangle of order  $(s,t)$ , are

$$s(t+1), \quad s-1 \quad \text{and} \quad -t-1$$

of multiplicity

$$1, \quad s^2(st+1)/(s+t) \quad \text{and} \quad st(s+1)(t+1)/(s+t),$$

respectively.

There exist many pseudo-geometric graphs for quadrangles, which are not geometric. The Shrikhande graph (see fly-leaf) is one of them. More examples (including an infinite family) are given in Section 6.2. The following theorem, which is due to CAMERON, GOETHALS & SEIDEL [C5], gives a sufficient condition for a pseudo-geometric graph to be geometric, as well as the extension of Higman's inequality to pseudo-geometric graphs for generalized quadrangles.

5.3.3. THEOREM. *Let  $G$  be a pseudo-geometric graph for a thick generalized quadrangle of order  $(s,t)$ . Then*

- i.  $t \leq s^2$ ,
- ii. *if equality holds, then  $G$  is geometric,*

iii. equality implies that all subconstituents of  $G$  are strongly regular.

PROOF. Let  $A$  be the adjacency matrix of  $G$ . Define

$$E := - (s+1)A + (s^2 - 1)I + J .$$

Then from the eigenvalues of  $A$  it follows that  $E$  has just one non-zero eigenvalue  $(s+1)(s+t)$  of multiplicity  $s^2(st+1)/(s+t)$ . Hence

$$(*) \quad \text{rank } E = s^2(st+1)/(s+t) .$$

Let  $x$  be a vertex of  $G$ . Partition  $A$  and  $E$  according to  $x$ , the vertices adjacent to  $x$ , and the vertices not adjacent to  $x$ :

$$A = \begin{bmatrix} 0 & j^* & 0 \\ j & A_{11} & A_{12} \\ 0 & A_{21} & A_{22} \end{bmatrix} , \quad E = \begin{bmatrix} s^2 & -sj^* & j^* \\ -sj & E_{11} & E_{12} \\ j & E_{21} & E_{22} \end{bmatrix} ,$$

where  $A_{12} = A_{21}^*$  and  $E_{12} = E_{21}^*$ . For  $i = 1, 2$ , let  $G_i$  be the graph with adjacency matrix  $A_{ii}$  (so  $G_i$  is a subconstituent of  $G$ ). Then  $G_i$  has  $s(t+1)$  vertices, and is regular of degree  $p_{11}^1 = s-1$ . Hence  $s-1$  is an eigenvalue of  $A_{11}$  of multiplicity  $c$ , say. It is known (see [B5], [C12] or Appendix I) that  $c$  equals the number of components of  $G_i$ . Clearly each component has at least  $s$  vertices. Hence

$$(**) \quad cs \leq s(t+1) .$$

The matrices  $A_{11}$ ,  $I$  and  $J$  have a common basis of eigenvectors. Using this it follows that  $E_{11}$  has an eigenvalue  $0$  of multiplicity  $c-1$  (one of the eigenvalues  $s-1$  of  $A_{11}$  corresponding to the eigenvector  $j$  leads to the eigenvalue  $s(t+1)$  of  $E_{11}$ ). Hence


$$\text{rank } E_{11} = s(t+1) - (c-1) .$$

Now using  $(*)$  and  $(**)$  we have

$$s(t+1) - t \leq \text{rank } E_{11} \leq \text{rank } E = s^2(st+1)/(s+t) .$$

This yields

$$t(s-1)(t-s^2) \leq 0,$$

proving (i). Suppose equality holds. Then we must have equality in (\*\*), which means that  $G_1$  is a disjoint union of complete graphs on  $s$  vertices. Since  $x$  is arbitrary, this yields that  does not occur in  $G$ . Now (ii) follows on applying 5.3.1.

Since the disjoint union of complete graphs of the same size is strongly regular, it only remains to be proved that  $G_2$  is strongly regular. This we shall prove analogously to the proof of 5.2.5. We know that the eigenvalues of  $E_{11}$  are  $0$ ,  $s(s+1)$  and  $s(s^2+1)$  of multiplicity  $s^2$ ,  $(s-1)(s^2+1)$  and  $1$ , respectively. Now using 1.3.3 and the eigenvalues of  $E$ , we obtain that the matrix

$$E' := \begin{bmatrix} s^2 & j^* \\ j & E_{22} \end{bmatrix}$$

has eigenvalues  $0$ ,  $s^2(s+1)$  and  $2s^2$ , where  $2s^2$  is a simple eigenvalue with eigenvector  $[s^2 \ j^*]^*$ . Hence  $E_{22}$ , and also  $A_{22}$ , has just two distinct eigenvalues not belonging to the eigenvector  $j$ . This proves that  $G_2$  is strongly regular.  $\square$

From (iii) of the above theorem it follows that the number of points adjacent to three mutually non-adjacent points of a quadrangle of order  $(s, s^2)$  is constant. This result was first proved by BOSE [B7].

A quadrangle of order  $(s, t)$  is the same as a partial geometry with parameters  $(s, t, 1)$  (see [H11], [T1] or Appendix I). For pseudo-geometric graphs for a partial geometry with parameters  $(s, t, \alpha)$ , where  $\alpha > 1$ , a result like Lemma 5.3.1 does not hold anymore. Therefore the question in the beginning of this section is much more difficult to answer for these geometries.

CHAPTER 6  
CONSTRUCTIONS

6.1. SOME 2 - (71,15,3) DESIGNS

In this section we shall construct eight non-isomorphic 2 - (71,15,3) designs. First we shall construct a 2 - (56,12,3) design  $D$ , which satisfies the hypothesis of Theorem 3.2.4. Next we show that  $D$  is embeddable in a 2 - (71,15,3) design. A less extensive treatment of this construction appeared in [B2]. Designs with these parameters seem to be new (see [C4] p.104, or [H5] p.297).

The most important ingredient for our construction is  $\mathbb{F}_8$ , the field with eight elements. Let  $G$  be the group  $\text{AFL}(1,8)$ , that is, the group of order 168 defined by  $x \mapsto ax^{2^i} + b$ ,  $a, b \in \mathbb{F}_8$ ,  $a \neq 0$ ,  $i \in \mathbb{Z}$ . We shall identify  $\mathbb{F}_8$  with  $\text{AG}(3,2)$ , the 3-dimensional affine space over  $\mathbb{F}_2$ . Although  $G$  is not the full automorphism group of  $\text{AG}(3,2)$ ,  $G$  acts transitively on the elements (we reserve the word points for points of a design), the lines (i.e. unordered pairs of elements), the planes (sets of the form  $\{a, b, c, a+b+c\}$ ), and the sets of four elements which do not form a plane. Moreover, the stabilizer of a line  $L$  has four orbits on lines:  $L$  itself, the lines intersecting  $L$ , the lines parallel to  $L$ , and the lines skew to  $L$ .

Now we shall define the incidence structure  $D$ . The points of  $D$  are the fifty-six ordered pairs of distinct elements of  $\mathbb{F}_8$ . The blocks of  $D$  are the seventy 4-subsets of  $\mathbb{F}_8$ . Let  $\alpha$ , satisfying  $\alpha^3 = \alpha + 1$ , be a primitive element of  $\mathbb{F}_8$ . The point  $(0,1)$  is defined to be incident with the following blocks:

$$\begin{aligned}
 & \{0,1,\alpha,\alpha^3\}, \{0,1,\alpha,\alpha^2\}, \{0,1,\alpha^2,\alpha^3\}, \{\alpha,\alpha^2,\alpha^3,\alpha^4\}, \{\alpha^2,\alpha^3,\alpha^4,\alpha^5\}, \\
 (*) & \{0,1,\alpha^2,\alpha^6\}, \{0,1,\alpha^2,\alpha^4\}, \{0,1,\alpha^4,\alpha^6\}, \{\alpha,\alpha^2,\alpha^4,\alpha^6\}, \{\alpha,\alpha^3,\alpha^4,\alpha^6\}, \\
 & \{0,1,\alpha^4,\alpha^5\}, \{0,1,\alpha,\alpha^4\}, \{0,1,\alpha,\alpha^5\}, \{\alpha,\alpha^2,\alpha^4,\alpha^5\}, \{\alpha,\alpha^2,\alpha^5,\alpha^6\}.
 \end{aligned}$$

Now we let  $G$  act on  $D$ . This defines  $D$ , because  $G$  acts transitively on the points of  $D$ , and because the map  $x \mapsto x^2$ , which fixes the point  $(0,1)$ , also fixes the set of blocks incident with  $(0,1)$ . A point  $(x,y)$  is called

equivalent to a point  $(w,z)$  if  $\{x,y\} = \{w,z\}$ . Two blocks  $b \subset \mathbb{F}_8$  and  $b' \subset \mathbb{F}_8$  are called *equivalent* if  $b = b'$ , or  $b \cap b' = \emptyset$  (i.e.  $b \cup b' = \mathbb{F}_8$ ). We know that  $G$  has two orbits on the blocks of  $D$ . The first orbit contains fifty-six blocks, they are the 4-subsets of  $\mathbb{F}_8$  which do not form a plane; these blocks will be called blocks of *type I*. The second orbit consists of the fourteen 4-subsets of  $\mathbb{F}_8$  which are planes; these blocks are of *type II*. It is clear that equivalent blocks are of the same type.

**6.1.1. LEMMA.** *Let  $(x,y)$  be a point of  $D$ . Let  $b \subset \mathbb{F}_8$  and  $b' \subset \mathbb{F}_8$  be distinct equivalent blocks of  $D$ .*

- i. *If  $(x,y)$  is incident with  $b$ , then  $\{x,y\} \subset b$  or  $\{x,y\} \subset b'$ .*
- ii. *If  $\{x,y\} \subset b$ ,  $b$  of type I, then exactly one of the following two statements is true.*
  1.  *$(x,y)$  is incident with  $b$  and not with  $b'$ , and  $(y,x)$  is incident with  $b'$  and not with  $b$ ;*
  2.  *$(x,y)$  is incident with  $b'$  and not with  $b$ , and  $(y,x)$  is incident with  $b$  and not with  $b'$ .*
- iii. *If  $b$  is of type II, then  $(x,y)$  is incident with  $b$  iff  $\{x,y\} \subset b$ .*
- iv.  *$b$  and  $b'$  have no points of  $D$  in common.*

**PROOF.** Without loss of generality take  $(x,y) = (0,1)$ . We may do so, because  $G$  is transitive on the points of  $D$ . Blocks incident with  $(0,1)$  are given in (\*). On applying the map  $x \mapsto x+1$  to (\*) we find that the blocks incident with  $(1,0)$  are the following ones:

$$\begin{aligned}
 & \{0,1,\alpha,\alpha^3\}, \{0,1,\alpha^3,\alpha^6\}, \{0,1,\alpha,\alpha^6\}, \{\alpha,\alpha^3,\alpha^5,\alpha^6\}, \{\alpha,\alpha^4,\alpha^5,\alpha^6\}, \\
 (**) \quad & \{0,1,\alpha^2,\alpha^6\}, \{0,1,\alpha^5,\alpha^6\}, \{0,1,\alpha^2,\alpha^5\}, \{\alpha^2,\alpha^3,\alpha^5,\alpha^6\}, \{\alpha,\alpha^2,\alpha^3,\alpha^5\}, \\
 & \{0,1,\alpha^4,\alpha^5\}, \{0,1,\alpha^3,\alpha^5\}, \{0,1,\alpha^3,\alpha^4\}, \{\alpha^3,\alpha^4,\alpha^5,\alpha^6\}, \{\alpha^2,\alpha^3,\alpha^4,\alpha^6\}.
 \end{aligned}$$

The first column of (\*) and (\*\*) consists of blocks of type II, all other blocks are of type I. Now (i), (ii) and (iii) are just a matter of verification. (iv) follows immediately from (ii) for blocks of type I, and from (iii) for blocks of type II.  $\square$

From (ii) and (iii) of this lemma we conclude the following. If each point of  $D$  is replaced by its equivalent partner and each block of  $D$  of type I by its equivalent partner, then incidence is not changed. Hence the permutation of the points of  $D$ , which interchanges the equivalent pairs of points, is an automorphism of  $D$ . This automorphism is different from, and commutes with, the automorphisms provided by the group  $G$ .

6.1.2. THEOREM.  $D$  is a 2 - (56,12,3) design.

PROOF. First we prove that  $D$  is a 1-design. Since every point is incident with fifteen blocks, the average number of points incident with a block equals twelve. By 6.1.1.iii a block of type II is incident with exactly twelve points. Now, since  $G$  acts transitively on blocks of the same type, also the blocks of type I are incident with exactly twelve points. We have seen that  $G$  has three orbits on the (unordered) pairs of lines of  $\mathbb{F}_8$  (intersecting, parallel, skew), for which the following pairs are representatives:

$$\{(0,1),\{0,\alpha\}\}, \quad \{(0,1),\{\alpha,\alpha^3\}\}, \quad \{(0,1),\{\alpha^2,\alpha^4\}\}.$$

From this it follows that the group  $2 \times G$ , which is an automorphism group of  $D$ , has seven orbits on the (unordered) pairs of points of  $D$ , for which the following ones are representatives:

$$\begin{aligned} &\{(0,1),\{0,\alpha\}\}, \quad \{(0,1),\{\alpha,\alpha^3\}\}, \quad \{(0,1),\{\alpha^2,\alpha^4\}\}, \\ &\{(1,0),\{0,\alpha\}\}, \quad \{(1,0),\{\alpha,\alpha^3\}\}, \quad \{(1,0),\{\alpha^2,\alpha^4\}\}, \\ &\{(1,0),\{0,1\}\}. \end{aligned}$$

Blocks incident with  $(0,1)$  and  $(1,0)$  are given in (\*) and (\*\*), respectively. Using the maps  $x \mapsto \alpha x$ ,  $x \mapsto x + \alpha$  and  $x \mapsto \alpha(x + \alpha)$ , we obtain the blocks incident with  $(0,\alpha)$ ,  $(\alpha,\alpha^3)$  and  $(\alpha^2,\alpha^4)$ . The blocks incident with  $(0,\alpha)$  are

$$\begin{aligned} &\{0,\alpha,\alpha^2,\alpha^4\}, \{0,\alpha,\alpha^2,\alpha^3\}, \{0,\alpha,\alpha^3,\alpha^4\}, \{\alpha^2,\alpha^3,\alpha^4,\alpha^5\}, \{\alpha^3,\alpha^4,\alpha^5,\alpha^6\}, \\ &\{0,1,\alpha,\alpha^3\}, \{0,\alpha,\alpha^3,\alpha^5\}, \{0,1,\alpha,\alpha^5\}, \{1,\alpha^2,\alpha^3,\alpha^5\}, \{1,\alpha^2,\alpha^4,\alpha^5\}, \\ &\{0,\alpha,\alpha^5,\alpha^6\}, \{0,\alpha,\alpha^2,\alpha^5\}, \{0,\alpha,\alpha^2,\alpha^6\}, \{\alpha^2,\alpha^3,\alpha^5,\alpha^6\}, \{1,\alpha^2,\alpha^3,\alpha^6\}. \end{aligned}$$

The blocks incident with  $(\alpha,\alpha^3)$  are



$$\begin{aligned} &\{0,1,\alpha,\alpha^3\}, \{0,\alpha,\alpha^3,\alpha^4\}, \{1,\alpha,\alpha^3,\alpha^4\}, \{0,1,\alpha^2,\alpha^4\}, \{1,\alpha^2,\alpha^4,\alpha^6\}, \\ &\{\alpha,\alpha^3,\alpha^4,\alpha^5\}, \{\alpha,\alpha^2,\alpha^3,\alpha^4\}, \{\alpha,\alpha^2,\alpha^3,\alpha^5\}, \{0,\alpha^2,\alpha^4,\alpha^5\}, \{0,1,\alpha^2,\alpha^5\}, \\ &\{\alpha,\alpha^2,\alpha^3,\alpha^6\}, \{0,\alpha,\alpha^2,\alpha^3\}, \{0,\alpha,\alpha^3,\alpha^6\}, \{0,\alpha^2,\alpha^4,\alpha^6\}, \{0,\alpha^4,\alpha^5,\alpha^6\}. \end{aligned}$$

The blocks incident with  $(\alpha^2, \alpha^4)$  are

$$\begin{aligned} &\{0,\alpha,\alpha^2,\alpha^4\}, \{0,\alpha^2,\alpha^4,\alpha^5\}, \{0,\alpha,\alpha^2,\alpha^4,\alpha^5\}, \{0,\alpha,\alpha^3,\alpha^5\}, \{1,\alpha,\alpha^3,\alpha^5\}, \\ &\{\alpha^2,\alpha^4,\alpha^5,\alpha^6\}, \{\alpha^2,\alpha^3,\alpha^4,\alpha^5\}, \{\alpha^2,\alpha^3,\alpha^4,\alpha^6\}, \{0,\alpha^3,\alpha^5,\alpha^6\}, \{0,\alpha,\alpha^3,\alpha^6\}, \\ &\{1,\alpha^2,\alpha^3,\alpha^4\}, \{0,\alpha^2,\alpha^3,\alpha^4\}, \{0,1,\alpha^2,\alpha^4\}, \{0,1,\alpha^3,\alpha^5\}, \{0,1,\alpha^5,\alpha^6\}. \end{aligned}$$

Blocks incident with  $(0,1)$  are marked by  $\{$ . Blocks incident with  $(1,0)$  are marked by  $\}$ . We see that for each of the seven pairs of points there are exactly three blocks incident with both points. Hence  $D$  is a  $2 - (56,12,3)$  design.  $\square$

Next we shall see that  $D$  satisfies the hypothesis of 3.2.4. The *line graph* of a geometry is the graph whose vertices are the lines, two vertices being adjacent iff the lines intersect.

**6.1.3. THEOREM.**  $D$  has just three intersection numbers 3, 2 and 0 ( $= k - r + \lambda$ ). The class graph of  $D$  is the complement of the line graph of  $PG(3,2)$ .

**PROOF.** Let  $b_1 \subset \mathbb{F}_8$  and  $b_2 \subset \mathbb{F}_8$  be non-equivalent blocks of  $D$ . Let  $b_1'$  and  $b_2'$  be the equivalent partners of  $b_1$  and  $b_2$ , respectively. Let  $B_1, B_2, B_1', B_2'$  be the sets of points incident with  $b_1, b_2, b_1', b_2'$ , respectively. Then by 6.1.1

$$|B_1 \cap B_1'| = |B_2 \cap B_2'| = 0 = k - r + \lambda.$$

Hence by 3.2.1.ii

$$|B_1 \cap B_2| = |B_1 \cap B_2'| = |B_1' \cap B_2| = |B_1' \cap B_2'|.$$

This implies

$$|(B_1 \cup B_1') \cap (B_2 \cup B_2')| = 4|B_1 \cap B_2|.$$

From (ii) and (iii) of 6.1.1 it follows that

$$\begin{aligned}
& |(B_1 \cup B_1') \cap (B_2 \cup B_2')| = \\
& = |\{(x,y) \in \mathbb{F}_8^2 \mid x \neq y, \{x,y\} \subset b_1 \text{ or } \{x,y\} \subset b_1', \\
& \qquad \qquad \qquad \{x,y\} \subset b_2 \text{ or } \{x,y\} \subset b_2'\}| = \\
& = \begin{cases} 8 & \text{if } |b_1 \cap b_2| = 2, \\ 12 & \text{if } |b_1 \cap b_2| \in \{1,3\}. \end{cases}
\end{aligned}$$

Hence 3, 2 and 0 are the intersection numbers of  $D$ . It is clear that the thirtyfive equivalence classes of blocks of  $D$  can be represented by the 4-subsets of  $\mathbb{F}_8$  containing 0. Therefore, they can also be represented by all 3-subsets of  $\mathbb{F}_8 - \{0\}$ . Now using Result 4.2.3 we see that the class graph of  $D$  is indeed the complement of the line graph of  $PG(3,2)$ .  $\square$

By the above theorem, there exists a 2-1 correspondence between the blocks of  $D$  and the lines of  $PG(3,2)$ , such that two blocks have no point in common iff they correspond to the same line of  $PG(3,2)$ , two blocks have two points in common iff they correspond to intersecting lines and two blocks have three points in common iff they correspond to skew lines.

6.1.4. THEOREM.  $D$  is embeddable in a symmetric  $2 - (71,15,3)$  design.

PROOF. We extend  $D$  to  $D_1$  with fifteen points (called *new points*), being the points of  $PG(3,2)$  and one block (*new block*). The points incident with the new block are precisely the new points. We define a new point to be incident with an old block (block of  $D$ ) iff the line of  $PG(3,2)$  corresponding to that block contains that point. Now it is easily seen that  $D_1$  is a  $1 - (71,15,15)$  design, and that any two distinct blocks of  $D_1$  have three points in common. This proves that the dual of  $D_1$ , and therefore  $D_1$  itself, is a symmetric  $2 - (71,15,3)$  design.  $\square$

From 6.1.1.iii it follows that the seven equivalence classes of blocks of type II of  $D$  correspond to seven mutually intersecting lines of  $PG(3,2)$ . For these lines we may take seven lines through one point or seven lines in one plane. Let  $D_1$  be the embedding of  $D$  in which blocks of type II correspond to lines through one point, and let  $D_2$  be the other embedding of  $D$ . Define  $D_1^*$  and  $D_2^*$  to be the dual of  $D_1$  and  $D_2$ , respectively. We shall show

that these four  $2 - (71,15,3)$  designs are non-isomorphic. To achieve this we define a block  $B$  of a  $2 - (71,15,3)$  design to be *special* if the derived design with respect to  $B$  (i.e. the  $2 - (15,3,2)$  subdesign of the  $2 - (71,15,3)$  design, formed by the points of  $B$  and the blocks distinct from  $B$ ) consists of two identical copies of a  $2 - (15,3,1)$  design. It is not difficult to see that the residual design with respect to a special block  $B$  (i.e. the  $2 - (56,12,3)$  subdesign of the  $2 - (71,15,3)$  design, formed by the points off  $B$  and the blocks distinct from  $B$ ) satisfies the hypothesis of Theorem 3.2.4 and that its class graph is isomorphic to the complement of the block graph of the  $2 - (15,3,1)$  design associated to  $B$  (which clearly is quasi-symmetric, since  $\lambda = 1$ ). From the proof of 6.1.4 it follows that the new blocks of  $D_1$  and  $D_2$  are special. Moreover, 6.1.1.iii implies that the new point of  $D_1$  which is incident with all old blocks of type II, is a special block of  $D_1^*$ . By verification it turns out that these are the only special blocks of  $D_1$ ,  $D_1^*$  and  $D_2$ . However,  $D_2^*$  has seven special blocks. They are the seven new points of  $D_2$ , which lie in the plane of  $PG(3,2)$  corresponding to the old blocks of type II. This already shows that

$$D_2 \not\cong D_2^*, D_2 \not\cong D_1, D_2 \not\cong D_1^*, D_2^* \not\cong D_1, D_2^* \not\cong D_1^* .$$

We know that any  $2 - (15,3,1)$  design associated to a special block of  $D_1$  or  $D_2$  is the design formed by the points and lines of  $PG(3,2)$ . By use of 6.1.1.iii it follows that also the  $2 - (15,3,1)$  design associated to the special block of  $D_1^*$  is the design which comes from  $PG(3,2)$ . Let  $D^*$  be the residual design of  $D_1^*$  with respect to the special block. Then the class graph of  $D^*$  is again the complement of the line graph of  $PG(3,2)$ . This means that, similarly as for the design  $D$ , interchanging points and planes of  $PG(3,2)$  yields a second embedding of  $D^*$  into a  $2 - (71,15,3)$  design. Let  $D_3^*$  be this  $2 - (71,15,3)$  design and let  $D_3$  be the dual of  $D_3^*$ . By verification it follows that  $D_3^*$  has just one special block (the one we started with), but that  $D_3$  has precisely seven special blocks. This shows that  $D_1^* \not\cong D_3^*$ . By further investigation it turns out that the  $2 - (15,3,1)$  design associated to any of the seven special blocks of  $D_3$  is again the design obtained from  $PG(3,2)$ , however, the seven special blocks of  $D_2^*$  give other  $2 - (15,3,1)$  designs (in fact all seven of them give the second design in the list of WHITE, COLE & CUMMINGS [W3]). This proves that  $D_2^* \not\cong D_3$ , and therefore  $D_1 \not\cong D_1^*$ . Hence  $D_1, D_1^*, D_2, D_2^*, D_3$  and  $D_3^*$  are all non-isomorphic. But there is still more. As remarked before, the  $2 - (15,3,1)$  design associ-

ated to one of the seven special blocks of  $D_3$  (these seven special blocks form an orbit under the automorphism group of  $D_3$ ) is the design formed by the points and lines of  $PG(3,2)$ . This implies that once again we can make another  $2 - (71,15,3)$  design by taking away the (two identical)  $2 - (15,3,1)$  designs and putting them back again after having interchanged points and planes. Call this new design  $D_4$ , and let  $D_4^*$  be its dual. It turns out that  $D_4$  has one special block, and that  $D_4^*$  has seven special blocks. Hence  $D_4$  is not isomorphic to  $D_1, D_1^*, D_2^*, D_3$  and  $D_4^*$ . In addition,  $D_4$  is not isomorphic to  $D_2$  and  $D_3^*$ , since otherwise  $D_3$  would have been isomorphic to  $D_1$  or  $D_1^*$ .

Because of time considerations we did not check whether it is possible to produce still more  $2 - (71,15,3)$  designs by playing once again the same game with respect to a special block of  $D_4^*$ . Thus we have the following result.

6.1.5. THEOREM. *There exist at least eight  $2 - (71,15,3)$  designs.*

The designs  $D_1^*, D_2^*, D_3^*$  and  $D_4^*$  are given explicitly in Appendix II. By taking residual designs with respect to various special blocks we obtain (at least) four non-isomorphic  $2 - (56,12,3)$  designs which satisfy the hypothesis of Theorem 3.2.4. One of these designs has a class graph which is non-isomorphic to the class graph of the other ones.

An *oval* in a  $2 - (71,15,3)$  design is a set  $S$  of six points such that any block has two or no points with  $S$  in common, see [A4]. Let  $S$  be an oval. It is clear that exactly twenty-six blocks do not meet  $S$ . Therefore, by 3.1.1 an oval of a  $2 - (71,15,3)$  design is equivalent to an empty subdesign with six points and twenty-six blocks. From 3.1.1.ii (see also [A4]) it follows that the subdesign of the  $2 - (71,15,3)$  design formed by the points off  $S$  and the blocks not meeting  $S$  is the dual of a  $2 - (26,6,3)$  design. By verification it follows that the following blocks of  $D$  provide an oval in  $D_1^*$ :

$$\{1, \alpha, \alpha^2, \alpha^3\}, \{1, \alpha, \alpha^2, \alpha^5\}, \{0, 1, \alpha, \alpha^5\}$$

$$\{1, \alpha^2, \alpha^4, \alpha^5\}, \{\alpha, \alpha^2, \alpha^4, \alpha^5\}, \{0, \alpha^3, \alpha^4, \alpha^6\}.$$

We conclude this section with a remark about automorphism groups. The group  $2 \times G$  of order 336 is an automorphism group of  $D_1, D_2, D_3$  and their

duals. The designs  $D_4$  and  $D_4^*$  have an automorphism group of order 48, viz. the stabilizer of a special block of  $D_3$ . F.C. Bussemaker has verified by use of a computer that the groups mentioned above are the full automorphism groups.

ACKNOWLEDGEMENT. I thank H.J. Beker, F.C. Bussemaker, R.H.F. Denniston and M. Hall Jr. for various contributions to this section.

## 6.2. SOME STRONGLY REGULAR GRAPHS

Suppose  $A$  is the adjacency matrix of a strongly regular graph  $G$  on  $n$  vertices of degree  $d$ . Furthermore, assume that  $A$  admits the following structure:

$$A = \begin{bmatrix} A_{11} & \cdots & A_{1m} \\ \vdots & & \vdots \\ A_{m1} & \cdots & A_{mm} \end{bmatrix},$$

where  $A_{ij}$  is a square matrix of size  $c := n/m$  having constant row sums equal to  $b_{ij}$ , say, for  $i, j = 1, \dots, m$ . From 1.2.3.iii it follows that the eigenvalues of the matrix  $B := (b_{ij})$  satisfy

$$(*) \quad \lambda_1(B) = \lambda_1(A) = d, \quad \lambda_i(B) \in \{\lambda_2(A), \lambda_n(A)\} \quad \text{for } i = 2, \dots, m.$$

Hence (\*) yields directives for the construction of strongly regular graphs, whose adjacency matrix admits this block structure. Let us consider two special cases of this structure.

CASE 1: All diagonal entries of  $B$  are equal to  $r_0$ , say, and all off-diagonal entries of  $B$  are equal to  $r_1$ , say, that is,

$$B = (r_0 - r_1)I + r_1 J.$$

Hence by (\*)

$$(m-1)r_1 + r_0 = d \quad \text{and} \quad r_0 - r_1 \in \{\lambda_2(A), \lambda_n(A)\}.$$

This implies

$$r_0 = (\lambda_2(A)(m-1) + d) / m, \quad r_1 = (d - \lambda_2(A)) / m,$$

or

$$r_0 = (\lambda_n(A)(m-1) + d) / m, \quad r_1 = (d - \lambda_n(A)) / m.$$

**6.2.1. EXAMPLE.** We wish to construct a strongly regular graph  $G$  with parameter set  $(q^3 + q^2 + q + 1, q^2 + q, q - 1, q + 1)$ , admitting the block structure of case 1 with  $m = q^2 + 1$  and  $c = q + 1$ . Then (see Appendix I)  $\lambda_2(G) = q - 1$ ,  $\lambda_n(G) = -q - 1$  and the formulas above yield  $r_0 = q$ ,  $r_1 = 1$ . It is indeed possible to construct  $G$ , by use of this framework. To do so we define the permutation matrices  $P$  and  $Q$  of size  $q + 1$  by

$$P := \begin{bmatrix} 0 & I_q \\ 1 & 0 \end{bmatrix} \quad \text{and} \quad (Q)_{ij} := \begin{cases} 1 & \text{if } i + j = q + 2, \\ 0 & \text{otherwise.} \end{cases}$$

It easily follows that

$$P^{q+1} = I, \quad (P^k)^* = P^{q+1-k} = P^{-k}, \quad Q^2 = I,$$

$$P^k Q = (P^k Q)^* = Q P^{-k}, \quad \sum_{i=0}^q P^i = J.$$

For  $k = 1, \dots, q$ , define  $R_k := P^k Q = Q P^{-k}$ . Then

$$R_k^* = R_k, \quad R_k R_\ell = P^{k-\ell}, \quad \sum_{i=0}^q R_i = J, \quad \sum_{i=0}^q \sum_{j=0}^q R_i R_j = (q+1)J,$$


for  $k, \ell = 1, \dots, q$ . Let  $p_1, \dots, p_v$  ( $v := q^2$ ) be the points, and let  $C_1, \dots, C_{q+1}$  be the parallel classes of an affine plane of order  $q$ . For  $i, j = 1, \dots, q^2 + 1$ , define the  $(q+1) \times (q+1)$  matrices

$$A_{ij} := \begin{cases} J - I & \text{if } i = j, \\ I & \text{if } i \neq j, i = q^2 + 1 \text{ or } j = q^2 + 1, \\ R_k & \text{if } i \neq j, i \leq q^2, j \leq q^2, \text{ and } C_k \text{ contains the} \\ & \text{line through } p_i \text{ and } p_j. \end{cases}$$

By use of the formulas above it is straightforward to verify that the square matrix  $A$  of size  $q^3 + q^2 + q + 1$ , built up with these block matrices  $A_{ij}$ , satisfies

$$(A + I)^2 = (q+1)J + q^2 I.$$

Therefore,  $A$  is the adjacency matrix of the desired strongly regular graph.

A strongly regular graph with the same parameters is provided by the point graph of a generalized quadrangle of order  $(q, q)$  (see Section 5.1). However, for  $q > 4$  the graphs constructed above need not be geometric. Indeed, we can order the points and the parallel classes of an affine plane of order  $q$  in such a way that the graph we obtained by our construction has  as an induced subgraph. Then by 5.3.1, this graph is not geometric, and therefore non-isomorphic to any of those which come from generalized quadrangles.

As a second example of graphs admitting the block structure of case 1 we mention the eleven incidence graphs of three linked  $2 - (16, 6, 2)$  designs (see Section 4.2).

CASE 2: All diagonal entries of  $B$  are equal to  $r_0$ , say. The off-diagonal entries of  $B$  take exactly two values  $r_1$  and  $r_2$ , say ( $r_1 > r_2$ ). Then by (\*) the  $(0, 1)$  matrix

$$\tilde{B} := \frac{1}{r_1 - r_2} (B - r_2(J - I) - r_0 I)$$

has just three distinct eigenvalues, one of which is simple and belongs to the eigenvector  $j$ . Hence  $\tilde{B}$  is the adjacency matrix of a strongly regular graph  $G'$  with eigenvalues

$$d' := \lambda_1(G') = (d - r_2 m + r_2 - r_0) / (r_1 - r_2) ,$$

$$\lambda_2(G') = (\lambda_2(A) + r_2 - r_0) / (r_1 - r_2) ,$$

$$\lambda_m(G') = (\lambda_n(A) + r_2 - r_0) / (r_1 - r_2) .$$

6.2.2. EXAMPLE. We wish to construct a strongly regular graph  $G$  with parameters  $(40, 12, 2, 4)$ , admitting the block structure of case 2 with  $m = 10$ ,  $c = 4$ ,  $r_0 = r_2 = 0$ ,  $r_1 = 2$ . From  $\lambda_2(G) = 2$ ,  $\lambda_{40}(G) = -4$  and the above formulas it follows that  $d' = 6$ ,  $\lambda_2(G') = 1$ ,  $\lambda_{10}(G') = -2$ . Hence  $G'$  is the complement of the Petersen graph. For  $h = 0, 1$  and  $i, j = 2, 3, 4$ , define the square  $(0, 1)$  matrices  $T_{hij}$  of size four, by

$$(T_{hij})_{kl} := \begin{cases} h & \text{if } (k, l) \in \{(1, 1), (1, l), (k, 1), (k, l)\} , \\ 1 - h & \text{otherwise.} \end{cases}$$

Then we have for  $h, h' = 0, 1; i, j, i', j' = 2, 3, 4$ ,

$$\begin{aligned} T_{hij} &= T_{hji}^* , & T_{hij} J &= 2J , & T_{hij} + T_{(1-h)ij} &= J , \\ T_{hij} T_{h'i'j'}^* &= J \text{ if } j \neq j' , & T_{hij} T_{hi'j}^* &= 2T_{1ii'} , \\ T_{hij} T_{(1-h)i'j}^* &= 2T_{0ii'} . \end{aligned}$$

With the help of these properties it is relatively easy to check that the following two matrices are adjacency matrices of strongly regular graphs with parameters  $(40, 12, 2, 4)$ .

$$\begin{bmatrix} 0 & T_{044} & T_{034} & T_{044} & T_{033} & T_{122} & T_{024} & 0 & 0 & 0 \\ T_{044} & 0 & 0 & T_{044} & T_{134} & 0 & T_{122} & T_{034} & 0 & T_{122} \\ T_{043} & 0 & 0 & T_{133} & T_{043} & T_{024} & 0 & T_{133} & T_{024} & 0 \\ T_{044} & T_{044} & T_{133} & 0 & 0 & T_{123} & 0 & T_{033} & 0 & T_{024} \\ T_{033} & T_{143} & T_{034} & 0 & 0 & 0 & T_{123} & T_{144} & T_{122} & 0 \\ T_{122} & 0 & T_{042} & T_{132} & 0 & 0 & T_{124} & 0 & T_{044} & T_{134} \\ T_{042} & T_{122} & 0 & 0 & T_{132} & T_{142} & 0 & 0 & T_{032} & T_{022} \\ 0 & T_{043} & T_{133} & T_{033} & T_{144} & 0 & 0 & 0 & T_{123} & T_{123} \\ 0 & 0 & T_{042} & 0 & T_{122} & T_{044} & T_{023} & T_{132} & 0 & T_{033} \\ 0 & T_{122} & 0 & T_{042} & 0 & T_{143} & T_{022} & T_{132} & T_{033} & 0 \end{bmatrix} \begin{bmatrix} 0 & T_{022} & T_{032} & T_{042} & T_{022} & T_{032} & T_{042} & 0 & 0 & 0 \\ T_{022} & 0 & T_{033} & T_{043} & T_{022} & 0 & 0 & T_{032} & T_{042} & 0 \\ T_{023} & T_{033} & 0 & T_{044} & 0 & T_{022} & 0 & T_{032} & 0 & T_{042} \\ T_{024} & T_{034} & T_{044} & 0 & 0 & 0 & T_{022} & 0 & T_{023} & T_{024} \\ T_{022} & T_{022} & 0 & 0 & 0 & T_{033} & T_{043} & T_{033} & T_{033} & 0 \\ T_{023} & 0 & T_{022} & 0 & T_{033} & 0 & T_{044} & T_{033} & 0 & T_{043} \\ T_{024} & 0 & 0 & T_{022} & T_{034} & T_{044} & 0 & 0 & T_{033} & T_{043} \\ 0 & T_{023} & T_{023} & 0 & T_{033} & T_{033} & 0 & 0 & T_{044} & T_{044} \\ 0 & T_{024} & 0 & T_{023} & T_{033} & 0 & T_{033} & T_{044} & 0 & T_{044} \\ 0 & 0 & T_{024} & T_{024} & 0 & T_{034} & T_{034} & T_{044} & T_{044} & 0 \end{bmatrix}$$

MATHON [M5] used the block structure of case 2 for the construction of strongly regular graphs with parameters  $(pq^2, \frac{1}{2}(pq^2-1), \frac{1}{4}(pq^2-5), \frac{1}{4}(pq^2-1))$  for prime powers  $p$  and  $q$ ,  $p \equiv 1, q \equiv -1 \pmod{4}$ . The strongly regular graph  $G'$ , which provides the framework for Mathon's construction has parameters  $(p, \frac{1}{2}(p-1), \frac{1}{4}(p-5), \frac{1}{4}(p-1))$ .

Any graph constructed in one of the examples above has the property that  $p_{11}^1 + 2 = p_{11}^2$ . This implies that its adjacency matrix  $A$  satisfies

$$(A+I)^2 = p_{11}^2 J + (d+1-p_{11}^2)I ,$$

where  $d$  denotes the degree of the graph. This yields the well-known fact that  $A+I$  is the incidence matrix of a symmetric  $2-(n, d+1, p_{11}^2)$  design. Similarly, if a strongly regular graph satisfies  $p_{11}^1 = p_{11}^2$ , then its adjacency matrix itself is the incidence matrix of a symmetric  $2-(n, d, p_{11}^1)$  design. This phenomenon is behind the next example, where we derive a strongly regular graph with  $p_{11}^1 + 2 = p_{11}^2$  from one with  $p_{11}^1 = p_{11}^2$ .



**6.2.3. EXAMPLE.** We start with a description of the generalized quadrangle  $Q$  of order  $(2^{\ell}-1, 2^{\ell}+1)$ ,  $\ell \in \mathbb{N}$ , due to HALL [H6]. Consider  $AG(3, q)$ , the three dimensional affine geometry over  $\mathbb{F}_q$ , with  $q = 2^{\ell}$ . Let  $S$  be a set of  $m := q + 2$  lines from  $AG(3, q)$  passing through one point, such that no three lines lie in one plane. Such a set exists, because it corresponds to a complete oval in the projective plane  $PG(2, q)$ , which exists iff  $q$  is even, see [D3]. It is easy to prove that each plane of  $AG(3, q)$  contains two or no lines from  $S$ . The points of  $Q$  are the points of  $AG(3, q)$ ; the lines of  $Q$  are the lines of  $S$ , and the lines of  $AG(3, q)$  which are parallel to a line of  $S$ ; a point and a line are incident in  $Q$ , iff they are incident in  $AG(3, q)$ . Now it is easy to prove that  $Q$  is a generalized quadrangle of order  $(q-1, q+1)$ . Let us partition the adjacency matrix  $A$  of the line graph of  $Q$  (i.e. the point graph of the dual of  $Q$ ) into  $m^2$  square block matrices of size  $q^2$ , according to the  $m$  parallel classes in  $Q$ :

$$A = \begin{bmatrix} A_{11} & \cdots & A_{1m} \\ \vdots & & \vdots \\ A_{m1} & \cdots & A_{mm} \end{bmatrix}.$$

From the structure of  $Q$  it follows that we may arrange the lines such that

$$A_{ij} = 0 \quad \text{if } i = j;$$

$$A_{ij} = I_q \otimes J_q \quad \text{if } i \text{ odd and } j = i+1, \text{ or } j \text{ odd and } i = j+1;$$

$$A_{ij} = \begin{bmatrix} P_{11} & \cdots & P_{1q} \\ \vdots & & \vdots \\ P_{q1} & \cdots & P_{qq} \end{bmatrix},$$

for all other values of  $i, j$ , where  $P_{kl}$  is a permutation matrix of size  $q$ , for  $k, l = 1, \dots, q$ . Now we derive a new matrix  $\tilde{A}$  from  $A$  by replacing

$$\begin{bmatrix} A_{ii} & A_{i,i+1} \\ A_{i+1,i} & A_{i+1,i+1} \end{bmatrix} \quad \text{by} \quad \begin{bmatrix} A_{i,i+1} & A_{ii} \\ A_{i+1,i+1} & A_{i+1,i} \end{bmatrix},$$

for  $i = 1, 3, \dots, m-1$ . Then it follows that

$$\tilde{A}^2 = A^2 = qJ + q^2I.$$

Since  $\tilde{A}$  is symmetric with all diagonal entries equal to one, the matrix

$\tilde{A} - I$  is the adjacency matrix of a strongly regular graph with parameters  $(q^2(q+2), q^2+q-1, q-2, q)$ . For  $q = 2$  this graph is the Clebsch graph (see Section 4.1, Figure 2), but for all other values of  $q = 2^l$  these strongly regular graphs seem to be new.

We remark that in the above example  $A$  has the block structure of case 1, whilst  $\tilde{A}$  has the block structure of case 2 with the cocktailparty graph on  $m$  vertices (complete  $\frac{1}{2}m$ -partite graph) as the underlying strongly regular graph.

The remainder of this section will be devoted to strongly regular graphs with parameters  $(40, 12, 2, 4)$ . For convenience we call such graphs *40-graphs*. Examples 6.2.1 ( $q = 3$ ) and 6.2.2 provide 40-graphs. The point graph of a generalized quadrangle of order  $(3, 3)$  is a 40-graph. PAYNE [P5] proved that there are exactly two generalized quadrangles of order  $(3, 3)$  (one being the dual of the other). In fact, these two geometric 40-graphs are the graph of Example 6.2.1 with  $q = 3$ , and the second graph of Example 6.2.2. From  $p_{11}^1 = 2$  it follows that a subgraph of a 40-graph induced by all vertices adjacent to a given vertex, is regular of degree two, so a disjoint union of cycles. But we can say more.

**6.2.4. LEMMA.** *Let  $G$  be a 40-graph. Let  $x$  be a vertex of  $G$  and let  $G_x$  be the subgraph of  $G$  induced by the vertices adjacent to  $x$ . Then  $G_x$  is one of the following graphs:*

- i. *a 12-cycle;*
- ii. *the disjoint union of a 9-cycle and a triangle;*
- iii. *the disjoint union of two 6-cycles;*
- iv. *the disjoint union of a 6-cycle and two triangles;*
- v. *the disjoint union of four triangles.*

**PROOF.** We only have to prove that the number of vertices of any component of  $G_x$  is divisible by three. If  $G_x$  is connected, there is nothing to prove. Suppose  $G_x$  has a component  $C$  of size  $c < 12$ . We partition  $A$  into sixteen block matrices according to: the vertex  $x$ , the vertices of  $C$ , the remaining vertices of  $G_x$ , and the vertices not adjacent to  $x$ . Then the entries of

$$B := \begin{pmatrix} 0 & c & 12-c & 0 \\ 1 & 2 & 0 & 9 \\ 1 & 0 & 2 & 9 \\ 0 & c/3 & 4-c/3 & 8 \end{pmatrix}$$

are the average row sums of the block matrices of A. It is easy to see that

$$\lambda_1(B) = 12, \quad \lambda_2(B) = \lambda_3(B) = 2, \quad \lambda_4(B) = -4.$$

On the other hand we know

$$\lambda_1(A) = 12, \quad \lambda_2(A) = \lambda_3(A) = 2, \quad \lambda_{40}(A) = -4.$$

So the eigenvalues of B interlace the eigenvalues of A tightly. Hence, by 1.2.3.ii the row sums of the block matrices are constant, so  $c/3$  is an integer.  $\square$

We associate with a 40-graph a 5-tuple  $(a_1, \dots, a_5)$ , where  $a_1, \dots, a_5$  denote the number of vertices  $x$  for which  $G_x$  has the form  $(i), \dots, (v)$ , respectively, of the above lemma. Using 5.3.1 we observe that a 40-graph is the point graph of a generalized quadrangle iff its 5-tuple is  $(0, 0, 0, 0, 40)$ . The first graph of Example 6.2.2 has 5-tuple  $(0, 0, 4, 24, 12)$ . R. Mathon (private communication) constructed a 40-graph with 5-tuple  $(0, 0, 0, 36, 4)$ .

WEISFEILER [W1] describes an algorithm for generating strongly regular graphs with a given parameter set, based on the principle of backtracking. By use of this algorithm we wrote a computer program (in Algol 60) for the construction of 40-graphs. Weisfeiler's algorithm rejects isomorphism only partially. This means that some of the produced 40-graphs may be isomorphic. We had our program run for about ten minutes. It turned out that, although we obtained about twohundred (not necessarily non-isomorphic) 40-graphs, the process of finding all 40-graphs still was in the beginning phase. For this reason there was no hope for completing the whole search. It seems that there are thousands of 40-graphs. Still we wanted to test the few hundred 40-graphs we found on isomorphism. A complete test on isomorphisms would have been too expensive. Therefore we just computed the 5-tuple of each 40-graph. It turned out that twentyone of these 40-graphs had different 5-tuples. So we found at least twentyone non-isomorphic 40-graphs. One of these is the first graph of Example 6.2.2. But none of these graphs has the

5-tuple of a geometric 40-graph or Mathon's 40-graph. So we have the following result.

6.2.5. THEOREM. *There exist at least twentyfour strongly regular graphs with parameters  $(40,12,2,4)$ .*

These 40-graphs are given in Appendix II, except for the three 40-graphs that are already exhibited in the Examples 6.2.1 and 6.2.2. The first graph in the list is Mathon's 40-graph.

We noticed already that a 40-graph gives a  $2 - (40,13,4)$  design. There is no reason why non-isomorphic graphs should lead to non-isomorphic designs. However, it has been checked that our twentyfour 40-graphs do produce twentyfour non-isomorphic  $2 - (40,13,4)$  designs.

An *oval* in a  $2 - (40,13,4)$  design is a set  $S$  of four points, such that any block has at most two points in common with  $S$ , see [A4]. Easy counting arguments give that twelve blocks are disjoint from  $S$  and four blocks have exactly one point in common with  $S$ . Suppose we have a 40-graph with a coclique of size four, such that any vertex is adjacent to two or to no vertices of that coclique. Then this coclique of the 40-graph produces an oval in the corresponding  $2 - (40,13,4)$  design. Conversely, it can be proved that any oval in a  $2 - (40,13,4)$  design, obtained from a 40-graph, corresponds to such a coclique. We see that the two 40-graphs of Example 6.2.2 produce  $2 - (40,13,4)$  designs with ten disjoint ovals. Also the last six 40-graphs of Appendix II supply designs with ovals. The remaining sixteen 40-graphs have no ovals.

In 4.2.2 we saw that the chromatic number of any 40-graph is at least five. Since the complement of the Petersen graph is 5-colourable, it follows that the two 40-graphs of Example 6.2.2 are 5-colourable as well.

APPENDIX I  
GRAPHS AND DESIGNS

This appendix contains the basic concepts and results from the theory of graphs and designs, which are used in the present monograph. Some general references are [B5], [C13], [H8], [W6] for graphs, [D3], [H5], [H17], [R3] for designs, and [B7], [C6] for both. We shall assume knowledge of Section 1.1.

A *graph* consists of a finite non-empty set of *vertices* together with a set of *edges*, where each edge is an unordered pair of vertices (so our graphs are finite, undirected and without loops or multiple edges). The two vertices of an edge are called *adjacent* (or *joint*). A graph is *complete* if every pair of vertices is an edge. The complete graph on  $n$  vertices is denoted by  $K_n$ . A graph without edges is called *void* (or *null*). The *complement* of a graph  $G$  is the graph  $\bar{G}$  on the same vertex-set as  $G$ , where any two vertices are adjacent whenever they are not adjacent in  $G$ . The *disjoint union* of a collection of graphs  $G_1, \dots, G_m$  on disjoint vertex sets is the graph whose vertex-set is the union of all vertex-sets, and whose edge-set is the union of all edge-sets of  $G_1, \dots, G_m$ . A graph is *disconnected* if it is the disjoint union of two or more graphs. Any graph  $G$  is the disjoint union of one or more connected (= not disconnected) graphs, called the *components* of  $G$ .

Let  $G$  be a graph on  $n$  vertices. A sequence of distinct vertices  $x_0, \dots, x_\ell$  of  $G$  is a *path* of length  $\ell$  between  $x_0$  and  $x_\ell$  if  $\{x_{i-1}, x_i\}$  is an edge for  $i = 1, \dots, \ell$ . The *distance*  $\rho(x, y)$  between two vertices  $x$  and  $y$  is the length of the shortest path between  $x$  and  $y$  ( $\rho(x, y) = \infty$  if  $x$  and  $y$  are in distinct components of  $G$ ). The *diameter* of  $G$  is the largest distance in  $G$ . A sequence of vertices  $x_0, \dots, x_\ell$  is a *circuit* of length  $\ell$  if  $x_1, \dots, x_\ell$  are distinct,  $x_0 = x_\ell$ ,  $\ell > 2$  and  $\{x_{i-1}, x_i\}$  is an edge for  $i = 1, \dots, \ell$ . The *girth* of  $G$  is the length of the shortest circuit in  $G$ . The *adjacency matrix* of  $G$  is the  $n \times n$  matrix  $A$ , indexed by the vertices of  $G$ , defined by

$$(A)_{xy} = \begin{cases} 1 & \text{if } \{x, y\} \text{ is an edge,} \\ 0 & \text{otherwise.} \end{cases}$$

Obviously,  $\bar{G}$  has adjacency matrix  $J - A - I$ . The *eigenvalues* of  $G$  are the eigenvalues of  $A$ ; they are denoted by  $\lambda_1(G) \geq \dots \geq \lambda_n(G)$  (the eigenvalues are real, because  $A$  is symmetric). We easily have that

$$\lambda_1(K_n) = n - 1, \quad \lambda_2(K_n) = \dots = \lambda_n(K_n) = -1.$$

The *incidence matrix*  $N$  of  $G$ , whose rows are indexed by the vertices and whose columns are indexed by the edges, is defined by

$$(N)_{x,E} = \begin{cases} 1 & \text{if } x \in E, \\ 0 & \text{otherwise.} \end{cases}$$

The graph with adjacency matrix  $N^*N - 2I$  is called the *line graph* of  $G$ , denoted by  $L(G)$ . The *subgraph* of  $G$  induced by a set  $S$  of vertices of  $G$  is the graph with vertex-set  $S$ , where two vertices are adjacent whenever they are adjacent in  $G$  (a subgraph is always an induced subgraph). Note that the adjacency matrix of a subgraph of  $G$  is a principal submatrix of  $A$ . A *clique* is a complete subgraph; a *coclique* (or *independent set of vertices*) is a void subgraph. The size of the largest clique and coclique is denoted by  $\omega(G)$  and  $\alpha(G)$ , respectively. A *colouring* of  $G$  is a colouring of the vertices, such that adjacent vertices have different colours (i.e. a partition of the vertices into cocliques). Vertices which are coloured with the same colour form a *colour class*.  $G$  is *k-colourable* if  $G$  admits a colouring with  $k$  colours; the smallest possible value of  $k$  is the *chromatic number* of  $G$ , denoted by  $\gamma(G)$ . It easily follows that

$$\gamma(G) \geq \omega(G), \quad \gamma(G)\alpha(G) \geq n.$$

If  $\gamma(G) = 2$ , then  $G$  is *bipartite*. By use of 1.1.2 it follows that if  $G$  is bipartite, then

$$\lambda_i(G) = -\lambda_{n+1-i}(G) \quad \text{for } i = 1, \dots, n.$$

Conversely,  $\lambda_1(G) = -\lambda_n(G)$  implies that  $G$  is void or bipartite; this follows from the Perron-Frobenius theorem on non-negative matrices (see [C13], [M3]). If  $G$  can be coloured with  $\gamma$  colours, such that all pairs of differently coloured vertices are edges, then  $G$  is *complete  $\gamma$ -partite* (i.e. the complement of the disjoint union of complete graphs). The complete bipartite graph ( $\gamma = 2$ ) is denoted by  $K_{\ell,m}$ , where  $\ell$  and  $m$  are the sizes of the two colour classes. By use of 1.1.2 it follows that

$$\lambda_1(K_{\ell,m}) = -\lambda_{\ell+m}(K_{\ell,m}) = \sqrt{\ell m}, \quad \lambda_i(K_{\ell,m}) = 0 \quad \text{for } i = 2, \dots, \ell+m-1.$$

The *degree* (or *valency*) of a vertex is the number of vertices adjacent to that vertex. A graph is *regular* (of degree  $d$ ) if all its vertices have the same degree (equal to  $d$ ).

Let  $G$  be regular of degree  $d$ . Then  $Aj = dj$ , hence  $d$  is an eigenvalue of  $G$  with eigenvector  $j$ . Moreover, the matrices  $A$ ,  $J$  and  $I$  have a common basis of eigenvectors. By use of this we obtain

$$\lambda_1(\bar{G}) = n - d - 1, \quad \lambda_i(\bar{G}) = -\lambda_{n+2-i}(G) - 1 \quad \text{for } i = 2, \dots, n.$$

If  $G$  is connected, then the Perron-Frobenius theorem yields that  $d$  is the largest eigenvalue of  $G$  with multiplicity one. Hence,  $d = \lambda_1(G)$  and its multiplicity equals the number of components of  $G$  (see [B5] for an elementary proof). If  $G$  is connected and  $d = 2$ , then  $G$  is called an *n-cycle* (or *circuit*).  $G$  is *strongly regular* if  $G$  is not void or complete and the adjacency matrix  $A$  satisfies

$$(1) \quad AJ = dJ, \quad (A - \lambda_2(G)I)(A - \lambda_n(G)I) = p_{11}^2 J,$$

for some number  $p_{11}^2$ . This is equivalent to requiring that  $A$  has precisely two distinct eigenvalues not belonging to the eigenvector  $j$ .

Now let  $G$  be strongly regular. Then also  $\bar{G}$  is strongly regular. Computation of the diagonal entries and the row sums of both sides of the second equality of (1) yields

$$(2) \quad d + \lambda_2(G)\lambda_n(G) = p_{11}^2, \quad (d - \lambda_2(G))(d - \lambda_n(G)) = p_{11}^2 n.$$

Define  $p_{11}^1 := \lambda_2(G) + \lambda_n(G) + p_{11}^2$ . Then (1) becomes

$$AJ = dJ, \quad A^2 = dI + p_{11}^1 A + p_{11}^2 (J - A - I).$$

This reflects that  $G$  satisfies the following three properties:  $G$  is regular of degree  $d$ ; for any pair of adjacent vertices there are exactly  $p_{11}^1$  vertices adjacent to both; for any pair of non-adjacent vertices there are exactly  $p_{11}^2$  vertices adjacent to both. The integers  $n$ ,  $d$ ,  $p_{11}^1$ ,  $p_{11}^2$  are the *parameters* of  $G$ . Let  $f$  be the multiplicity of the eigenvalue  $\lambda_2(G)$ . Then

$$0 = \text{trace } A = d + f\lambda_2(G) + (n - f - 1)\lambda_n(G),$$

whence

$$f = \frac{d + (n-1)\lambda_n(G)}{\lambda_n(G) - \lambda_2(G)}, \quad \lambda_n(G)(n-2f-1) = -d - f(p_{11}^1 - p_{11}^2).$$

So, if  $n \neq 2f+1$ , then  $\lambda_n(G)$  is rational, and therefore  $\lambda_2(G)$  and  $\lambda_n(G)$  are integers. It is an easy exercise (see [C6]) to show that  $n = 2f+1$  implies

$$n-1 = 2d = 4p_{11}^2 = 4(p_{11}^1 + 1), \quad \lambda_2(G) = -\lambda_n(G) - 1 = -\frac{1}{2} + \frac{1}{2}\sqrt{n}.$$

By use of the above results and trace  $A = 0$ , it follows that  $\lambda_2(G) \geq 0$  and  $\lambda_n(G) \leq -1$ . From (2) we have that  $\lambda_2(G) = 0$  iff  $p_{11}^2 = d$ . It is easily seen that  $p_{11}^2 = d$  reflects that  $G$  is a complete  $\gamma$ -partite graph. If  $\lambda_n(G) = -1$ , then  $\lambda_2(\bar{G}) = 0$ , hence  $G$  is the disjoint union of complete graphs and  $p_{11}^2 = 0$ . Conversely, by (1)  $p_{11}^2 = 0$  implies  $d = \lambda_2(G)$  and  $\lambda_n(G) = -1$ . These two families of strongly regular graphs are called *imprimitive*. Let  $x$  be a vertex of  $G$ . The two *subconstituents* of  $G$  with respect to  $x$  are the subgraphs of  $G$  induced by the vertices adjacent to  $x$  and by the vertices non-adjacent to  $x$ . The subconstituents are regular of degree  $p_{11}^1$  and  $d - p_{11}^2$ , respectively. Examples of primitive (= not imprimitive) strongly regular graphs are: the pentagon (5-cycle),  $L(K_m)$  for  $m \geq 5$  (= *triangular graph*),  $L(K_{m,m})$  for  $m \geq 3$  (= *lattice graph*) and their complements. The *Petersen graph* is the complement of  $L(K_5)$  and has parameters  $(10,3,0,1)$ .

An *incidence structure* consists of a finite non-empty set  $V_1$  of *points* and a finite non-empty set  $V_2$  of *blocks*, together with a subset of  $V_1 \times V_2$  of *flags*. A point and a block are *incident* if they form a flag. Often blocks are identified with the sets of points with which they are incident; if so, blocks are denoted by capitals, otherwise we use small types. An incidence structure without flags is called *empty*.

Let  $D$  be an incidence structure with  $v$  points and  $b$  blocks. An incidence structure  $D'$  formed by points and blocks of  $D$  is a *substructure* of  $D$  whenever a point and a block are incident in  $D'$  iff they are incident in  $D$ . The *incidence matrix*  $N$  of  $D$ , whose rows are indexed by the points and whose columns are indexed by the blocks, is defined by

$$(N)_{x,B} = \begin{cases} 1 & \text{if } x \in B, \\ 0 & \text{otherwise.} \end{cases}$$



The incidence structures with incidence matrices  $N^*$  and  $J - N$  are called the *dual* and the *complement* of  $D$ , respectively. The graph with adjacency matrix  $\begin{bmatrix} 0 & N \\ N^* & 0 \end{bmatrix}$  is the *incidence graph* of  $D$ . Clearly  $D$  and its dual have the same incidence graph. We call  $D$  a  $t - (v, k, \lambda)$  *design* (or  $t$ -*design* with *parameters*  $(v, k, \lambda)$ ) if all blocks have size  $k$ , and if any set of  $t$  points is contained in exactly  $\lambda$  blocks. A  $t - (v, k, \lambda)$  design with  $k < t$  or  $v - k < t$  is *degenerate*. A *design* is a  $t$ -design for some  $t$ . A *subdesign* is a substructure which is a design. Note that we allow repeated blocks (two or more blocks incident with exactly the same points).

Let  $D$  be a non-degenerate  $t - (v, k, \lambda)$  design. The complement of  $D$  is also a non-degenerate  $t$ -design. By elementary counting we see that for  $t \geq 1$   $D$  is also a  $(t-1) - (v, k, \lambda(v-t+1)/(k-t+1))$  design. In particular,  $D$  is a  $1 - (v, k, r)$  design, where

$$r := \binom{v-1}{t-1} / \binom{k-1}{t-1} ;$$

$r$  equals the number of blocks incident with any point. Counting flags yields  $bk = vr$ . If  $D$  is a  $1 - (v, k, r)$  design, then  $rk$  is the largest eigenvalue of  $NN^*$  and hence by 1.1.2  $\sqrt{rk}$  is the largest singular value of  $N$ . A 2-design is also called (*balanced incomplete*) *block design*. Now let  $D$  be a non-degenerate 2- $(v, k, \lambda)$  design. In terms of the incidence matrix  $N$  this means

$$(3) \quad N^*J = kJ, \quad NJ = rJ, \quad NN^* = \lambda J + (r - \lambda)I.$$

$NN^*$  has eigenvalues  $\lambda v + r - \lambda = rk$  and  $r - \lambda$  of multiplicity 1 and  $v - 1$ , respectively. By 1.1.2 these eigenvalues are the squares of the singular values of  $N$ . From  $rk \neq 0$ ,  $r - \lambda \neq 0$ , it follows that  $v = \text{rank } NN^* = \text{rank } N$ , hence  $b \geq v$  (Fisher's inequality). If  $b = v$  (i.e.  $r = k$ ),  $D$  is called *symmetric*. Formula (3) yields  $N(N^* - (\lambda/r)J) = (r - \lambda)I$ . If  $D$  is symmetric,  $N$  is square, hence  $(N^* - (\lambda/r)J)N = (r - \lambda)I$ , i.e.  $N^*N = \lambda J + (r - \lambda)I$ , and therefore the dual of  $D$  is a symmetric 2- $(v, k, \lambda)$  design as well. Let  $D$  be symmetric, and let  $B$  be a block of  $D$ . The subdesign formed by the points incident with  $B$  and the blocks distinct from  $B$  is a 2- $(k, \lambda, \lambda - 1)$  design (possibly degenerate), called the *derived design* of  $D$  with respect to  $B$ . Similarly, the subdesign formed by the points not incident with  $B$  and the blocks distinct from  $B$  is a 2- $(v - k, k - \lambda, \lambda)$  design (possibly degenerate), called the *residual design* of  $D$  with respect to  $B$ . A 2- $(v - k, k - \lambda, \lambda)$  design  $D'$  is *embeddable* in  $D$  if  $D'$  is a residual design of  $D$ . For example, the

points and lines of  $PG(2,q)$ , the projective plane of order  $q$ , form a symmetric  $2 - (q^2+q+1, q+1, 1)$  design; the degenerate  $2 - (q+1, 1, 0)$  design is a derived design, and the affine plane of order  $q$ , which is a  $2 - (q^2, q, 1)$  design, is a residual design.

A *partial geometry* with parameters  $(s, t, \alpha)$ ,  $s, t, \alpha \in \mathbb{N}$ , is a  $1 - (v, s+1, t+1)$  design satisfying the following two conditions:

- any two blocks have at most one point in common;
- (4) for any non-incident point-block pair  $(x, B)$  the number of blocks incident with  $x$  and intersecting  $B$  equals  $\alpha$ .

For a partial geometry we speak of lines rather than blocks. Let  $D$  denote a partial geometry with parameters  $(s, t, \alpha)$  and incidence matrix  $N$ . The graph with adjacency matrix  $NN^* - (t+1)I$  is the *point graph* of  $D$ . The *line graph* of  $D$  is the point graph of the dual of  $D$ . From the definition it readily follows that the point graph  $G$  of  $D$  is strongly regular with parameters

$$(v, s(t+1), t(\alpha-1) + s - 1, \alpha(t+1)) .$$

By use of our identities for strongly regular graphs we obtain

$$\lambda_1(G) = s(t+1) , \quad \lambda_2(G) = s - \alpha , \quad \lambda_n(G) = -t - 1 ,$$

$$v = (s+1)(st+\alpha)/\alpha , \quad b = (t+1)(st+\alpha)/\alpha .$$

Hence  $NN^*$  has the eigenvalues  $(s+1)(t+1)$ ,  $0$  and  $s+t+1-\alpha$  of multiplicity  $1$ ,  $s(s+1-\alpha)(st+\alpha)/\alpha(s+t+1-\alpha)$  and  $st(s+1)(t+1)/\alpha(s+t+1-\alpha)$ , respectively. By 1.1.2 the square roots of these eigenvalues are the singular values of  $N$ . A *partial subgeometry* is a substructure, which itself is a partial geometry. Let  $D'$  be a partial subgeometry of  $D$  with parameters  $(s', t', \alpha)$ . Then the point graph of  $D'$  is an induced subgraph of the point graph of  $D$ . This can be seen as follows. Let  $x$  and  $y$  be two points of  $D'$  and suppose there exists a line  $L$  of  $D$ , which is not a line of  $D'$ , incident with  $x$  and  $y$ . Let  $M$  be a line of  $D'$  incident with  $x$ . By (4) there are  $\alpha$  lines of  $D'$  incident with  $x$  and intersecting  $M$ . Hence there are at least  $\alpha+1$  such lines in  $D$ . This is a contradiction. So two points are on a line of  $D'$  iff they are on a line of  $D$ , which proves the claim. A partial geometry with parameters  $(s, t, 1)$  is the same as a *generalized quadrangle of order*  $(s, t)$ .

## APPENDIX II

## TABLES

First we list the twentyone strongly regular graphs with parameters  $(40,12,2,4)$ , as promised in Section 6.2. Together with the ones of Examples 6.2.1 and 6.2.2 they form the twentyfour 40-graphs of Theorem 6.2.5. The respective 5-tuples of the 40-graphs listed below are:

$(0,0,0,36,4)$	$(8,18,2,9,3)$	$(16,14,4,5,1)$
$(12,18,6,3,1)$	$(4,20,4,10,2)$	$(18,18,0,3,1)$
$(12,12,0,14,2)$	$(9,18,0,9,4)$	$(18,20,0,0,2)$
$(6,22,6,3,3)$	$(18,12,9,0,1)$	$(27,12,0,0,1)$
$(0,32,0,0,8)$	$(0,12,0,18,10)$	$(0,36,0,0,4)$
$(0,24,0,12,4)$	$(8,8,0,20,4)$	$(6,12,12,10,0)$
$(2,8,12,16,2)$	$(10,8,18,4,0)$	$(0,16,6,12,6)$

2 3 4 5 6 7 8 9 10 11 12 13	9 10 15 16 17 21 22 24 28 32 35 38	9 10 15 16 17 21 22 24 28 32 35 38
1 3 4 14 15 16 17 18 19 20 21 22	6 8 9 10 14 19 23 24 32 33 34 39	6 8 9 10 14 19 23 24 32 33 34 39
1 2 4 23 24 25 26 27 28 29 30 31	4 7 12 15 16 19 23 24 30 31 35 39	4 7 12 15 16 19 23 24 30 31 35 39
1 2 3 32 33 34 35 36 37 38 39 40	3 8 10 13 17 18 20 21 30 32 37 39	3 8 10 13 17 18 20 21 30 32 37 39
1 6 7 14 17 20 23 27 31 32 37 39	7 9 12 14 18 20 21 22 28 31 34 39	7 9 12 14 18 20 21 22 28 31 34 39
1 5 7 15 18 21 24 28 29 33 35 40	2 9 12 13 15 17 18 23 31 36 37 38	2 9 11 15 17 18 20 23 31 35 36 37
1 5 6 16 19 22 25 26 30 34 36 38	3 5 10 11 14 16 18 24 33 36 37 38	3 5 10 11 14 16 18 24 33 36 37 38
1 9 10 14 17 20 25 26 30 33 35 40	2 4 11 15 18 20 22 24 29 34 35 36	2 4 12 13 15 18 22 24 29 34 36 38
1 8 10 15 18 21 23 27 31 34 36 38	1 2 5 6 11 16 20 25 30 35 37 39	1 2 5 6 12 13 16 25 30 37 38 39
1 8 9 16 19 22 24 28 29 32 37 39	1 2 4 7 12 13 22 25 29 33 38 39	1 2 4 7 11 20 22 25 29 33 35 39
1 12 13 14 17 20 24 28 29 34 36 38	7 8 9 13 15 19 21 25 28 30 33 36	6 7 10 13 15 19 21 25 28 30 34 36
1 11 13 15 18 21 25 26 30 32 37 39	3 5 6 10 17 19 21 25 29 34 35 36	3 5 8 9 17 19 21 25 29 33 35 36
1 11 12 16 19 22 23 27 31 33 35 40	4 6 10 11 16 19 22 26 28 31 34 37	4 8 9 11 16 19 22 26 28 31 33 37
2 5 8 11 15 16 23 28 30 32 36 40	2 5 7 15 17 19 22 26 29 30 32 37	2 5 7 15 17 19 22 26 29 30 32 37
2 6 9 12 14 16 24 26 31 33 37 38	1 3 6 8 11 14 21 26 29 31 38 39	1 3 6 8 11 14 21 26 29 31 38 39
2 7 10 13 14 15 25 27 29 34 35 39	1 3 7 9 13 20 23 26 29 32 34 36	1 3 7 9 13 20 23 26 29 32 34 36
2 5 8 11 18 19 24 26 31 34 35 39	1 4 6 12 14 20 24 26 28 30 33 36	1 4 6 12 14 20 24 26 28 30 33 36
2 6 9 12 17 19 25 27 29 32 36 40	4 5 6 7 8 23 25 26 28 32 35 38	4 5 6 7 8 23 25 26 28 32 35 38
2 7 10 13 17 18 23 28 30 33 37 38	2 3 11 12 13 14 20 27 28 32 35 38	2 3 11 12 13 14 20 27 28 32 35 38
2 5 8 11 21 22 25 27 29 33 37 38	4 5 8 9 16 17 19 27 29 31 33 38	4 5 6 10 16 17 19 27 29 31 34 38
2 6 9 12 20 22 23 28 30 34 35 39	1 4 5 11 12 15 23 27 32 33 34 37	1 4 5 11 12 15 23 27 32 33 34 37
2 7 10 13 20 21 24 26 31 32 36 40	1 5 8 10 13 14 23 27 30 31 35 36	1 5 8 10 13 14 23 27 30 31 35 36
3 5 9 13 14 19 21 24 25 32 35 38	2 3 6 16 18 21 22 27 28 29 30 33	2 3 6 16 18 21 22 27 28 29 30 33
3 6 10 11 15 17 22 23 25 33 36 39	1 2 3 7 8 17 25 27 28 31 34 37	1 2 3 7 8 17 25 27 28 31 34 37
3 7 8 12 16 18 20 23 24 34 37 40	9 10 11 12 18 24 26 27 29 30 31 32	9 10 11 12 18 24 26 27 29 30 31 32
3 5 9 13 16 18 20 26 28 33 36 39	13 14 15 16 17 18 25 27 33 34 35 39	13 14 15 16 17 18 25 27 33 34 35 39
3 6 10 11 14 19 21 26 27 34 37 40	19 20 21 22 23 24 25 26 36 37 38 39	19 20 21 22 23 24 25 26 36 37 38 39
3 5 9 13 16 18 20 26 28 33 36 39	1 5 11 13 17 18 19 23 24 29 39 40	1 5 11 13 17 18 19 23 24 29 39 40
3 6 10 11 16 18 20 30 31 32 35 38	8 10 12 14 15 16 20 23 25 28 37 40	8 10 12 14 15 16 20 23 25 28 37 40
3 7 8 12 14 19 21 29 31 33 36 39	3 4 9 11 14 17 22 23 25 34 38 40	3 4 9 11 14 17 22 23 25 34 38 40
3 5 9 13 15 17 22 29 30 34 37 40	3 5 6 13 15 20 22 24 25 32 33 40	3 5 6 13 15 20 22 24 25 32 33 40
4 5 10 12 14 18 22 23 26 29 33 34	1 2 4 14 16 18 19 21 25 31 36 40	1 2 4 14 16 18 19 21 25 31 36 40
4 6 8 13 15 19 20 24 27 30 32 34	2 7 10 11 17 20 21 23 26 31 35 40	2 7 10 12 13 17 21 23 26 31 38 40
4 7 9 11 16 17 21 25 28 31 32 33	2 5 8 12 13 16 21 24 26 30 38 40	2 5 8 11 16 20 21 24 26 30 35 40
4 6 8 13 16 17 21 23 26 29 36 37	1 3 8 9 12 18 19 22 26 33 37 40	1 3 6 10 12 18 19 22 26 34 37 40
4 7 9 11 14 18 22 24 27 30 35 37	6 7 8 11 12 16 17 22 27 32 39 40	6 7 8 11 12 16 17 22 27 32 39 40
4 5 10 12 15 19 20 25 28 31 35 36	4 6 7 9 13 14 21 24 27 29 35 40	4 6 7 9 13 14 21 24 27 29 35 40
4 7 9 11 15 19 20 23 26 29 39 40	1 6 7 10 15 18 19 20 27 30 34 40	1 7 8 9 15 18 19 20 27 30 33 40
4 5 10 12 16 17 21 24 27 30 38 40	2 3 4 5 9 10 15 26 27 28 36 40	2 3 4 5 9 10 15 26 27 28 36 40
4 6 8 13 14 18 22 25 28 31 38 39	28 29 30 31 32 33 34 35 36 37 38 39	28 29 30 31 32 33 34 35 36 37 38 39

9 10 15 16 17 21 22 24 28 32 35 38  
 7 8 9 14 15 19 23 24 30 32 33 39  
 4 6 10 12 16 19 23 24 31 34 35 39  
 3 8 10 13 17 18 20 21 30 32 37 39  
 7 9 12 14 18 20 21 22 28 31 34 39  
 3 9 11 14 15 17 18 23 31 36 37 38  
 2 5 10 12 13 16 18 24 33 36 37 38  
 2 4 11 15 18 20 22 24 29 34 35 36  
 1 2 5 6 11 16 20 25 30 35 37 39  
 1 3 4 7 11 14 22 25 29 33 38 39  
 6 8 9 10 13 19 21 25 28 33 34 36  
 3 5 7 15 17 19 21 25 29 30 35 36  
 4 7 11 15 16 19 22 26 28 30 31 37  
 2 5 6 10 17 19 22 26 29 32 34 37  
 1 2 6 8 12 13 21 26 29 31 38 39  
 1 3 7 9 13 20 23 26 29 32 34 36  
 1 4 6 12 14 20 24 26 28 30 33 36  
 4 5 6 7 8 23 25 26 28 32 35 38  
 2 3 11 12 13 14 20 27 28 32 35 38  
 4 5 8 9 16 17 19 27 29 31 33 38  
 1 4 5 11 12 15 23 27 32 33 34 37  
 1 5 8 10 13 14 23 27 30 31 35 36  
 2 3 6 16 18 21 22 27 28 29 30 33  
 1 2 3 7 8 17 25 27 28 31 34 37  
 9 10 11 12 18 24 26 27 29 30 31 32  
 13 14 15 16 17 18 25 27 33 34 35 39  
 19 20 21 22 23 24 25 26 36 37 38 39  
 1 5 11 13 17 18 19 23 24 29 39 40  
 8 10 12 14 15 16 20 23 25 28 37 40  
 2 4 9 12 13 17 22 23 25 34 38 40  
 3 5 6 13 15 20 22 24 25 32 33 40  
 1 2 4 14 16 18 19 21 25 31 36 40  
 2 7 10 11 17 20 21 23 26 31 35 40  
 3 5 8 11 14 16 21 24 26 30 38 40  
 1 3 8 9 12 18 19 22 26 33 37 40  
 6 7 8 11 12 16 17 22 27 32 39 40  
 4 6 7 9 13 14 21 24 27 29 35 40  
 1 6 7 10 15 18 19 20 27 30 34 40  
 2 3 4 5 9 10 15 26 27 28 36 40  
 28 29 30 31 32 33 34 35 36 37 38 39  
  
 9 10 15 16 17 21 22 24 28 32 35 38  
 6 8 9 10 12 17 23 24 33 34 35 39  
 4 7 14 15 16 19 23 24 30 31 32 39  
 3 8 10 12 13 17 18 21 30 37 38 39  
 7 9 12 14 18 20 21 22 28 31 34 39  
 2 9 13 15 17 18 20 23 31 32 36 39  
 3 5 10 11 16 18 20 24 33 35 36 37  
 2 4 11 14 15 18 22 24 29 34 36 38  
 1 2 5 6 11 14 16 25 30 37 38 39  
 1 2 4 7 13 20 22 25 29 32 33 39  
 7 8 9 13 15 19 21 25 28 30 33 36  
 2 4 5 16 17 19 21 25 29 31 35 36  
 4 6 10 11 16 19 22 26 28 31 34 37  
 3 5 8 9 17 19 22 26 29 32 33 37  
 1 3 6 8 11 20 21 26 29 31 35 39  
 1 4 6 12 14 20 24 26 28 30 33 36  
 4 5 6 7 8 23 25 26 28 32 35 38  
 2 3 11 12 13 14 20 27 28 32 35 38  
 5 6 7 10 15 17 19 27 29 30 34 38  
 1 4 5 11 12 15 23 27 32 33 34 37  
 1 5 8 10 13 14 23 27 30 31 35 36  
 2 3 6 16 18 21 22 27 28 29 30 33  
 1 2 3 7 8 17 25 27 28 31 34 37  
 9 10 11 12 18 24 26 27 29 30 31 32  
 13 14 15 16 17 18 25 27 33 34 35 39  
 19 20 21 22 23 24 25 26 36 37 38 39  
 1 5 11 13 17 18 19 23 24 29 39 40  
 8 10 12 14 15 16 20 23 25 28 37 40  
 2 4 9 12 13 17 22 23 25 34 38 40  
 3 5 6 13 15 20 22 24 25 32 33 40  
 1 2 4 14 16 18 19 21 25 31 36 40  
 2 7 10 11 17 20 21 23 26 31 35 40  
 3 5 8 11 14 16 21 24 26 30 38 40  
 1 3 8 9 12 18 19 22 26 33 37 40  
 6 7 8 11 12 16 17 22 27 32 39 40  
 4 6 7 9 13 14 21 24 27 29 35 40  
 1 6 7 10 15 18 19 20 27 30 34 40  
 2 3 4 5 9 10 15 26 27 28 36 40  
 28 29 30 31 32 33 34 35 36 37 38 39  
  
 10 12 14 16 17 20 22 24 28 32 35 38  
 4 7 9 10 15 16 23 24 32 33 37 39  
 4 8 11 12 14 19 23 24 30 34 35 39  
 2 3 11 13 17 18 21 22 31 32 38 39  
 7 8 10 12 13 17 18 20 30 36 37 39  
 10 11 14 15 18 20 21 23 28 31 34 37  
 2 5 14 16 18 19 21 24 29 31 35 36  
 3 5 9 15 18 20 22 24 29 33 34 38  
 2 8 10 11 13 14 19 25 28 33 36 38  
 1 2 5 6 9 21 22 25 30 34 35 39  
 3 4 6 9 16 17 20 25 29 35 36 37  
 1 3 5 13 14 21 23 25 29 32 33 37  
 4 5 9 12 16 19 22 26 28 31 34 37  
 1 3 6 7 9 12 15 26 31 36 38 39  
 2 6 8 14 17 19 22 26 29 30 32 37  
 1 2 7 11 13 20 23 26 29 30 34 38  
 1 4 5 11 15 21 24 26 28 30 33 36  
 4 5 6 7 8 23 25 26 28 32 35 38  
 3 7 9 13 15 20 21 27 28 30 32 35  
 1 5 6 8 11 16 19 27 31 32 33 39  
 4 6 7 10 12 17 19 27 29 33 34 38  
 1 4 8 10 13 15 23 27 29 31 35 36  
 2 3 6 12 16 18 22 27 28 30 33 36  
 1 2 3 7 8 17 25 27 28 31 34 37  
 9 10 11 12 18 24 26 27 29 30 31 32  
 13 14 15 16 17 18 25 27 33 34 35 39  
 19 20 21 22 23 24 25 26 36 37 38 39  
 1 6 9 13 17 18 19 23 24 29 39 40  
 7 8 11 12 15 16 21 22 25 28 39 40  
 3 5 10 15 16 17 19 23 25 31 38 40  
 4 6 7 13 14 20 22 24 25 30 33 40  
 1 2 4 12 15 18 19 20 25 34 36 40  
 2 8 9 12 17 20 21 23 26 31 35 40  
 3 6 8 10 13 16 21 24 26 32 36 40  
 1 3 7 10 11 18 19 22 26 33 37 40  
 5 7 9 11 14 17 22 23 27 32 34 40  
 2 5 6 11 12 13 15 24 27 35 38 40  
 1 4 8 9 14 16 18 21 27 30 37 40  
 2 3 4 5 10 14 20 26 27 28 29 40  
 28 29 30 31 32 33 34 35 36 37 38 39  
  
 10 12 14 16 17 20 22 24 28 32 35 38  
 5 7 9 10 15 22 23 24 32 33 34 39  
 4 8 11 12 14 19 23 24 30 34 35 39  
 3 8 10 13 17 18 21 22 31 32 36 39  
 2 7 11 12 13 17 18 20 30 37 38 39  
 10 11 14 15 18 20 21 23 28 31 34 37  
 2 5 14 16 18 19 21 24 29 31 35 36  
 3 4 9 15 16 18 20 24 29 33 37 38  
 2 8 10 11 13 14 19 25 28 33 36 38  
 1 2 4 6 9 16 21 25 30 35 37 39  
 3 5 6 9 17 20 22 25 29 34 35 36  
 1 3 5 13 14 21 23 25 29 32 33 37  
 4 5 9 12 16 19 22 26 28 31 34 37  
 1 3 6 7 9 12 15 26 31 36 38 39  
 2 6 8 14 17 19 22 26 29 30 32 37  
 1 7 8 10 13 20 23 26 29 30 34 36  
 1 4 5 11 15 21 24 26 28 30 33 36  
 4 5 6 7 8 23 25 26 28 32 35 38  
 3 7 9 13 15 20 21 27 28 30 32 35  
 1 5 6 8 11 16 19 27 31 32 33 39  
 4 6 7 10 12 17 19 27 29 33 34 38  
 1 2 4 11 13 15 23 27 29 31 35 38  
 2 3 6 12 16 18 22 27 28 30 33 36  
 1 2 3 7 8 17 25 27 28 31 34 37  
 9 10 11 12 18 24 26 27 29 30 31 32  
 13 14 15 16 17 18 25 27 33 34 35 39  
 19 20 21 22 23 24 25 26 36 37 38 39  
 1 6 9 13 17 18 19 23 24 29 39 40  
 7 8 11 12 15 16 21 22 25 28 39 40  
 3 5 10 15 16 17 19 23 25 31 38 40  
 4 6 7 13 14 20 22 24 25 30 33 40  
 1 2 4 12 15 18 19 20 25 34 36 40  
 2 8 9 12 17 20 21 23 26 31 35 40  
 3 6 8 10 13 16 21 24 26 32 36 40  
 1 3 7 10 11 18 19 22 26 33 37 40  
 4 7 9 11 14 17 22 23 27 32 34 40  
 5 6 8 10 12 13 15 24 27 35 36 40  
 1 5 8 9 14 18 21 27 30 37 40  
 2 3 4 5 10 14 20 26 27 28 29 40  
 28 29 30 31 32 33 34 35 36 37 38 39

10 12 14 16 17 20 22 24 28 32 35 38 10 12 14 16 17 20 22 24 28 32 35 38 10 12 14 16 17 20 22 24 28 32 35 38  
 5 7 10 15 16 19 23 24 30 32 34 39 5 7 10 15 16 19 23 24 30 31 35 39 5 7 10 15 16 19 23 24 31 32 33 39  
 4 8 11 12 14 19 23 24 31 33 35 39 4 8 11 12 14 19 23 24 32 33 34 39 4 8 11 12 14 19 23 24 30 34 35 39  
 3 8 10 13 17 18 21 22 30 32 37 39 3 8 10 13 17 18 21 22 30 31 38 39 3 8 10 13 17 18 21 22 31 32 36 39  
 2 7 11 12 13 17 18 20 31 36 38 39 2 7 11 12 13 17 18 20 32 36 37 39 2 7 11 12 13 17 18 20 30 37 38 39  
 10 11 14 15 18 20 21 23 28 31 34 37 10 11 14 15 18 20 21 23 28 31 34 37 10 11 14 15 18 20 21 23 28 31 34 37  
 2 5 9 14 18 21 22 24 29 33 35 37 2 5 9 14 18 21 22 24 29 33 34 38 2 5 9 14 18 21 22 24 29 34 35 36  
 3 4 9 15 16 18 20 24 29 34 36 38 3 4 9 15 16 18 20 24 29 35 36 37 3 4 9 15 16 18 20 24 29 33 37 38  
 7 8 10 13 14 19 20 25 28 30 33 36 7 8 10 13 14 19 20 25 28 30 33 36 7 8 10 13 14 19 20 25 28 30 33 36  
 1 2 4 6 9 12 15 25 35 36 37 39 1 2 4 6 9 11 16 25 34 36 38 39 1 2 4 6 9 12 15 25 35 36 37 39  
 3 5 6 14 16 17 19 25 29 32 36 37 3 5 6 10 17 19 22 25 29 33 35 37 3 5 6 14 16 17 19 25 29 32 36 37  
 1 3 5 10 13 21 23 25 29 33 34 38 1 3 5 13 14 15 23 25 29 31 36 38 1 3 5 10 13 21 23 25 29 33 34 38  
 4 5 9 12 16 19 22 26 28 31 34 37 4 5 9 12 16 19 22 26 28 31 34 37 4 5 9 12 16 19 22 26 28 31 34 37  
 1 3 6 7 9 11 22 26 30 34 38 39 1 3 6 7 9 12 21 26 30 35 37 39 1 3 6 7 9 11 22 26 31 33 38 39  
 2 6 8 10 17 19 22 26 29 31 33 38 2 6 8 12 17 19 20 26 29 30 34 38 2 6 8 10 17 19 22 26 29 30 34 38  
 1 2 8 11 13 20 23 26 29 30 35 37 1 2 8 10 13 21 23 26 29 32 33 37 1 2 8 11 13 20 23 26 29 31 35 36  
 1 4 5 11 15 21 24 26 28 30 33 36 1 4 5 11 15 21 24 26 28 30 33 36 1 4 5 11 15 21 24 26 28 30 33 36  
 4 5 6 7 8 23 25 26 28 32 35 38 4 5 6 7 8 23 25 26 28 32 35 38 4 5 6 7 8 23 25 26 28 32 35 38  
 2 3 9 11 13 15 21 27 28 32 35 38 2 3 9 11 13 15 21 27 28 32 35 38 2 3 9 11 13 15 21 27 28 32 35 38  
 1 5 6 8 9 16 21 27 31 32 33 39 1 5 6 8 9 15 22 27 31 32 33 39 1 5 6 8 9 16 21 27 30 32 34 39  
 4 6 7 12 17 19 20 27 29 30 34 35 4 6 7 14 16 17 19 27 29 31 32 36 4 6 7 12 17 19 20 27 29 31 33 35  
 1 4 7 13 14 15 23 27 29 31 32 36 1 4 7 11 13 20 23 27 29 30 34 35 1 4 7 13 14 15 23 27 29 30 32 37  
 2 3 6 12 16 18 22 27 28 30 33 36 2 3 6 12 16 18 22 27 28 30 33 36 2 3 6 12 16 18 22 27 28 30 33 36  
 1 2 3 7 8 17 25 27 28 31 34 37 1 2 3 7 8 17 25 27 28 31 34 37 1 2 3 7 8 17 25 27 28 31 34 37  
 9 10 11 12 18 24 26 27 29 30 31 32 9 10 11 12 18 24 26 27 29 30 31 32 9 10 11 12 18 24 26 27 29 30 31 32  
 13 14 15 16 17 18 25 27 33 34 35 39 13 14 15 16 17 18 25 27 33 34 35 39 13 14 15 16 17 18 25 27 33 34 35 39  
 19 20 21 22 23 24 25 26 36 37 38 39 19 20 21 22 23 24 25 26 36 37 38 39 19 20 21 22 23 24 25 26 36 37 38 39  
 1 6 9 13 17 18 19 23 24 29 39 40 1 6 9 13 17 18 19 23 24 29 39 40 1 6 9 13 17 18 19 23 24 29 39 40  
 7 8 11 12 15 16 21 22 25 28 39 40 7 8 11 12 15 16 21 22 25 28 39 40 7 8 11 12 15 16 21 22 25 28 39 40  
 2 4 9 14 16 17 21 23 25 31 38 40 2 4 9 14 15 17 22 23 25 32 37 40 2 4 9 14 16 17 21 23 25 31 38 40  
 3 5 6 13 15 20 22 24 25 30 35 40 3 5 6 12 13 20 21 24 25 33 35 40 3 5 6 13 15 20 22 24 25 30 35 40  
 1 2 4 11 18 19 20 22 25 33 34 40 1 2 4 11 16 17 20 23 26 31 38 40 1 2 4 11 18 19 20 22 25 33 34 40  
 3 7 9 12 15 17 20 23 26 32 37 40 3 7 9 12 15 17 20 23 26 31 38 40 3 7 9 12 15 17 20 23 26 32 37 40  
 2 6 8 12 13 14 21 24 26 32 36 40 2 6 8 12 13 14 21 24 26 32 36 40 2 6 8 12 13 14 21 24 26 32 36 40  
 1 3 7 10 16 18 19 21 26 31 36 40 1 2 8 11 14 18 19 22 26 31 36 40 1 3 7 10 16 18 19 21 26 31 36 40  
 5 8 9 10 11 17 22 23 27 34 35 40 5 8 9 10 12 17 21 23 27 34 35 40 5 8 9 10 11 17 22 23 27 34 35 40  
 4 6 7 10 11 13 16 24 27 33 38 40 4 6 8 11 13 14 16 24 27 30 38 40 4 6 7 10 11 13 16 24 27 33 38 40  
 1 5 8 12 14 15 18 19 27 30 37 40 1 4 7 10 12 15 18 19 27 33 37 40 1 5 8 12 14 15 18 19 27 30 37 40  
 2 3 4 5 10 14 20 26 27 28 29 40 2 3 4 5 10 14 20 26 27 28 29 40 2 3 4 5 10 14 20 26 27 28 29 40  
 28 29 30 31 32 33 34 35 36 37 38 39 28 29 30 31 32 33 34 35 36 37 38 39 28 29 30 31 32 33 34 35 36 37 38 39

5 9 10 12 17 20 21 23 26 27 30 31  
5 6 8 12 18 20 22 24 25 27 29 32  
7 9 10 11 18 19 22 23 25 28 29 31  
5 6 7 11 17 19 21 24 26 28 30 32  
2 4 10 12 14 15 22 23 26 28 33 35  
2 4 9 11 13 16 18 20 30 31 33 35  
3 4 8 9 14 15 17 20 29 32 34 35  
1 2 7 10 13 16 21 24 25 28 34 35  
1 3 6 7 14 16 22 24 26 27 33 36  
1 3 5 8 13 15 18 19 30 31 33 36  
3 4 6 12 13 15 21 23 25 27 34 36  
1 2 5 11 14 16 17 19 29 31 34 36  
6 8 10 11 17 22 26 29 35 36 37 38  
5 7 9 12 18 21 25 30 35 36 37 38  
5 7 10 11 20 24 27 31 33 34 37 38  
6 8 9 12 19 23 28 32 33 34 37 38  
1 4 7 12 13 22 25 31 32 33 37 39  
2 3 6 10 14 21 26 31 32 34 37 39  
3 4 10 12 16 24 27 29 30 35 37 39  
1 2 6 7 15 23 28 29 30 36 37 39  
1 4 8 11 14 18 27 28 29 33 37 40  
2 3 5 9 13 17 27 28 30 34 37 40  
1 3 5 11 16 20 25 26 32 35 37 40  
2 4 8 9 15 19 25 26 31 36 37 40  
2 3 8 11 14 17 23 24 30 33 38 39  
1 4 5 9 13 18 23 24 29 34 38 39  
1 2 9 11 15 19 21 22 32 35 38 39  
3 4 5 8 16 20 21 22 31 36 38 39  
2 3 7 12 13 19 20 21 26 33 38 40  
1 4 6 10 14 19 20 22 25 34 38 40  
1 3 6 12 15 17 18 24 28 35 38 40  
2 4 7 10 16 17 18 23 27 36 38 40  
5 6 9 10 15 16 17 21 25 29 39 40  
7 8 11 12 15 16 18 22 26 30 39 40  
5 6 7 8 13 14 19 23 27 31 39 40  
9 10 11 12 13 14 20 24 28 32 39 40  
13 14 15 16 17 18 19 20 21 22 23 24  
13 14 15 16 25 26 27 28 29 30 31 32  
17 18 19 20 25 26 27 28 33 34 35 36  
21 22 23 24 29 30 31 32 33 34 35 36

8 9 10 12 17 20 21 23 26 27 30 31  
5 6 11 12 18 20 22 24 25 27 29 31  
7 8 9 10 18 19 22 23 25 28 29 32  
5 6 7 11 17 19 21 24 26 28 30 32  
2 4 10 12 14 15 22 23 26 28 33 35  
2 4 8 9 13 16 18 20 30 32 33 35  
3 4 9 11 14 15 17 20 29 31 34 35  
1 3 6 12 13 15 21 24 25 28 34 35  
1 3 6 7 14 16 22 24 26 27 33 36  
1 3 5 11 13 15 18 19 30 31 33 36  
2 4 7 10 13 16 21 23 25 27 34 36  
1 2 5 8 14 16 17 19 29 32 34 36  
6 8 10 11 17 22 26 29 35 36 37 38  
5 7 9 12 18 21 25 30 35 36 37 38  
5 7 8 10 20 24 27 32 33 34 37 38  
6 9 11 12 19 23 28 31 33 34 37 38  
1 4 7 12 13 22 25 31 32 33 37 39  
2 3 6 10 14 21 26 31 32 34 37 39  
3 4 10 12 16 24 27 29 30 35 37 39  
1 2 6 7 15 23 28 29 30 36 37 39  
1 4 8 11 14 18 27 28 29 33 37 40  
2 3 5 9 13 17 27 28 30 34 37 40  
1 3 5 11 16 20 25 26 32 35 37 40  
2 4 8 9 15 19 25 26 31 36 37 40  
2 3 8 11 14 17 23 24 30 33 38 39  
1 4 5 9 13 18 23 24 29 34 38 39  
1 2 9 11 15 19 21 22 32 35 38 39  
3 4 5 8 16 20 21 22 31 36 38 39  
2 3 7 12 13 19 20 21 26 33 38 40  
1 4 6 10 14 19 20 22 25 34 38 40  
1 3 6 12 15 17 18 24 28 35 38 40  
2 4 7 10 16 17 18 23 27 36 38 40  
5 6 9 10 15 16 17 21 25 29 39 40  
7 8 11 12 15 16 18 22 26 30 39 40  
5 6 7 8 13 14 19 23 27 31 39 40  
9 10 11 12 13 14 20 24 28 32 39 40  
13 14 15 16 17 18 19 20 21 22 23 24  
13 14 15 16 25 26 27 28 29 30 31 32  
17 18 19 20 25 26 27 28 33 34 35 36  
21 22 23 24 29 30 31 32 33 34 35 36

8 9 10 12 17 20 22 23 25 27 29 31  
5 7 8 10 18 20 21 24 26 27 29 32  
6 9 11 12 18 19 21 23 26 28 29 31  
5 6 7 11 17 19 22 24 25 28 30 32  
2 4 10 12 14 15 22 23 26 28 33 35  
3 4 8 9 13 15 18 20 30 32 33 35  
2 4 9 11 14 16 17 20 29 31 34 35  
1 2 6 12 13 16 21 24 25 28 34 35  
1 3 6 7 14 16 22 24 26 27 33 36  
1 2 5 11 13 16 18 19 30 31 33 36  
3 4 7 10 13 15 21 23 25 27 34 36  
1 3 5 8 14 15 17 19 29 32 34 36  
6 8 10 11 17 22 26 29 35 36 37 38  
5 7 9 12 18 21 25 30 35 36 37 38  
5 6 11 12 20 24 27 31 33 34 37 38  
7 8 9 10 19 23 28 32 33 34 37 38  
1 4 7 12 13 21 26 31 32 33 37 39  
2 3 6 10 14 22 25 31 32 34 37 39  
3 4 10 12 16 24 27 29 30 35 37 39  
1 2 6 7 15 23 28 29 30 36 37 39  
2 3 8 11 14 17 27 28 30 34 37 40  
1 3 5 11 16 20 25 26 32 35 37 40  
2 4 8 9 15 19 25 26 31 36 37 40  
1 4 8 11 14 18 23 24 29 33 38 39  
2 3 5 9 13 17 23 24 29 34 38 39  
1 2 9 11 15 19 21 22 32 35 38 39  
3 4 5 8 16 20 21 22 31 36 38 39  
2 3 7 12 13 19 20 21 26 33 38 40  
1 4 6 10 14 19 20 22 25 34 38 40  
1 3 7 10 15 17 18 24 28 35 38 40  
2 4 6 12 16 17 18 23 27 36 38 40  
5 6 9 10 15 16 17 21 25 29 39 40  
7 8 11 12 15 16 18 22 26 30 39 40  
5 6 7 8 13 14 19 23 27 31 39 40  
9 10 11 12 13 14 20 24 28 32 39 40  
13 14 15 16 17 18 19 20 21 22 23 24  
13 14 15 16 25 26 27 28 29 30 31 32  
17 18 19 20 25 26 27 28 33 34 35 36  
21 22 23 24 29 30 31 32 33 34 35 36

8 9 10 12 17 20 22 23 25 27 29 31  
5 7 10 11 17 20 21 24 26 27 30 31  
6 8 9 12 17 19 21 23 26 28 30 32  
5 6 7 11 18 19 21 24 26 28 29 32  
2 4 10 12 14 15 22 23 26 28 33 35  
3 4 9 11 13 15 18 20 30 31 33 35  
2 4 8 9 14 16 17 20 29 32 34 35  
1 3 7 10 13 15 21 24 25 28 34 35  
1 3 6 7 14 16 22 24 26 27 33 36  
1 2 5 8 13 16 18 19 30 32 33 36  
2 4 6 12 13 16 21 23 25 27 34 36  
1 3 5 11 14 15 17 19 29 31 34 36  
6 8 10 11 17 22 26 29 35 36 37 38  
5 7 9 12 18 21 25 30 35 36 37 38  
5 6 11 12 20 24 27 32 33 34 37 38  
7 8 9 10 19 23 28 32 33 34 37 38  
1 4 7 12 13 21 26 31 32 33 37 39  
2 3 6 10 14 22 25 31 32 34 37 39  
3 4 10 12 16 24 27 29 30 35 37 39  
1 2 6 7 15 23 28 29 30 36 37 39  
2 3 8 11 14 17 27 28 30 34 37 40  
1 3 5 11 16 20 25 26 32 35 37 40  
2 4 8 9 15 19 25 26 31 36 37 40  
1 4 8 11 14 18 23 24 29 33 38 39  
2 3 5 9 13 17 23 24 29 34 38 39  
1 2 9 11 15 19 21 22 32 35 38 39  
3 4 5 8 16 20 21 22 31 36 38 39  
2 3 7 12 13 19 20 21 26 33 38 40  
1 4 6 10 14 19 20 22 25 34 38 40  
1 3 7 10 15 17 18 24 28 35 38 40  
2 4 6 12 16 17 18 23 27 36 38 40  
5 6 9 10 15 16 17 21 25 29 39 40  
7 8 11 12 15 16 18 22 26 30 39 40  
5 6 7 8 13 14 19 23 27 31 39 40  
9 10 11 12 13 14 20 24 28 32 39 40  
13 14 15 16 17 18 19 20 21 22 23 24  
13 14 15 16 25 26 27 28 29 30 31 32  
17 18 19 20 25 26 27 28 33 34 35 36  
21 22 23 24 29 30 31 32 33 34 35 36

Here are listed the ten systems of three linked 2-(16,6,2) designs, as promised in Section 4.2. The incidence graphs of these systems, together with the point graph of the generalized quadrangle of order (3,5) form the eleven 4-colourable strongly regular graphs with parameters (64,18,2,6) of Theorem 4.3.1.

2 3 4 5 9 13	2 3 4 5 9 13	2 3 4 5 9 13	1 2 3 7 15 16	1 2 3 7 15 16
1 3 4 6 10 14	1 3 4 6 10 14	1 3 4 6 10 14	1 2 5 9 10 11	1 2 5 9 10 11
1 2 4 7 11 15	1 2 4 7 11 15	1 2 4 7 11 15	1 3 5 6 8 14	1 3 5 6 8 14
1 2 3 8 12 16	1 2 3 8 12 16	1 2 3 8 12 16	1 4 6 7 9 13	1 4 6 7 9 13
1 6 7 8 9 13	1 6 7 8 9 13	1 6 7 8 9 13	1 4 10 12 14 16	1 4 10 12 14 16
2 5 7 8 10 14	2 5 7 8 10 14	2 5 7 8 10 14	1 8 11 12 13 15	1 8 11 12 13 15
3 5 6 8 11 15	3 5 6 8 11 15	3 5 6 8 11 15	2 3 4 5 12 13	2 3 4 5 12 13
4 5 6 7 12 16	4 5 6 7 12 16	4 5 6 7 12 16	2 4 6 11 14 15	2 4 6 11 14 15
1 5 10 11 12 13	1 5 10 11 12 13	1 5 10 11 12 13	2 6 8 10 13 16	2 6 8 10 13 16
2 6 9 11 12 14	2 6 9 11 12 14	2 6 9 11 12 14	2 7 8 9 12 14	2 7 8 9 12 14
3 7 9 10 12 15	3 7 9 10 12 15	3 7 9 10 12 15	3 4 8 9 10 15	3 4 8 9 10 15
4 8 9 10 11 16	4 8 9 10 11 16	4 8 9 10 11 16	3 6 9 11 12 16	3 6 9 11 12 16
1 5 9 14 15 16	1 5 9 14 15 16	1 5 9 14 15 16	3 7 10 11 13 14	3 7 10 11 13 14
2 6 10 13 15 16	2 6 10 13 15 16	2 6 10 13 15 16	4 5 7 8 11 16	4 5 7 8 11 16
3 7 11 13 14 16	3 7 11 13 14 16	3 7 11 13 14 16	5 6 7 10 12 15	5 6 7 10 12 15
4 8 12 13 14 15	4 8 12 13 14 15	4 8 12 13 14 15	5 9 13 14 15 16	5 9 13 14 15 16
2 3 4 6 11 16	2 3 4 6 11 16	2 3 4 6 11 16	1 2 3 6 11 13	1 2 3 6 11 13
1 3 4 5 12 15	1 3 4 5 12 15	1 3 4 5 12 15	1 2 5 12 14 15	1 2 5 12 14 15
1 2 4 8 9 14	1 2 4 8 9 14	1 2 4 8 9 14	1 3 4 5 7 10	1 3 4 5 7 10
1 2 3 7 10 13	1 2 3 7 10 13	1 2 3 7 10 13	1 4 9 11 15 16	1 4 9 11 15 16
1 5 7 8 11 16	1 5 6 8 10 16	1 5 6 8 10 16	1 6 7 8 12 16	1 6 7 8 12 16
2 6 7 8 12 15	2 5 9 10 11 15	2 5 9 10 11 15	1 8 9 10 13 14	1 8 9 10 13 14
3 5 6 7 9 14	3 8 10 11 12 14	3 8 10 11 12 14	2 3 5 8 9 16	2 3 5 8 9 16
4 5 6 8 10 13	4 5 7 8 11 13	4 5 7 8 11 13	2 4 6 9 10 12	2 4 6 9 10 12
1 6 9 10 12 16	1 7 9 11 12 16	1 7 9 11 12 16	2 4 7 13 14 16	2 4 7 13 14 16
2 5 9 10 11 15	2 6 7 8 12 15	2 6 7 8 12 15	2 7 8 10 11 15	2 7 8 10 11 15
3 8 10 11 12 14	3 5 6 7 9 14	3 5 6 7 9 14	3 4 8 11 12 14	3 4 8 11 12 14
4 7 9 11 12 13	4 6 9 10 12 13	4 6 9 10 12 13	3 6 10 14 15 16	3 6 10 14 15 16
1 6 11 13 14 15	1 6 11 13 14 15	1 6 11 13 14 15	3 7 9 12 13 15	3 7 9 12 13 15
2 5 12 13 14 16	2 8 9 13 14 16	2 8 9 13 14 16	4 5 6 8 13 15	4 5 6 8 13 15
3 8 9 13 15 16	3 5 12 13 15 16	3 5 12 13 15 16	5 6 7 9 11 14	5 6 7 9 11 14
4 7 10 14 15 16	4 7 10 14 15 16	4 7 10 14 15 16	5 10 11 12 13 16	5 10 11 12 13 16
2 3 4 7 12 14	2 3 5 6 10 12	2 3 5 7 11 12	1 2 6 7 10 14	1 2 6 9 14 16
1 3 4 8 11 13	1 4 6 8 11 12	1 4 5 6 9 11	1 2 9 12 13 16	1 2 7 10 12 13
1 2 4 5 10 16	1 2 9 12 13 15	1 2 10 11 14 16	4 6 8 9 14 16	4 6 7 8 10 14
1 2 3 6 9 15	1 2 6 7 14 16	1 2 5 8 13 15	4 7 8 10 12 13	4 8 9 12 13 16
1 5 6 8 12 14	1 4 5 7 9 10	1 4 7 8 10 12	1 3 4 6 12 15	1 3 4 13 14 15
2 5 6 7 11 13	2 3 7 8 9 11	2 3 6 8 9 10	1 4 5 11 13 14	1 4 5 6 11 12
3 6 7 8 10 16	3 4 6 7 13 15	3 4 5 8 14 16	2 3 8 13 14 15	2 3 6 8 12 15
4 5 7 8 9 15	3 4 9 12 14 16	3 4 10 11 13 15	2 5 6 8 11 12	2 5 8 11 13 14
1 7 9 10 11 14	1 3 10 11 15 16	1 3 6 7 15 16	1 3 7 8 9 11	1 3 8 10 11 16
2 8 9 10 12 13	2 4 5 8 15 16	2 4 9 12 15 16	1 5 8 10 15 16	1 5 7 8 9 15
3 5 9 11 12 16	5 6 9 11 13 16	5 6 10 12 14 15	2 3 4 10 11 16	2 3 4 7 9 11
4 6 10 11 12 15	5 7 11 12 14 15	6 8 11 12 13 16	2 4 5 7 9 15	2 4 5 10 15 16
1 7 12 13 15 16	1 3 5 8 13 14	1 3 9 12 13 14	3 5 6 9 10 13	3 5 6 7 13 16
2 8 11 14 15 16	2 4 10 11 13 14	2 4 6 7 13 14	3 5 7 12 14 16	3 5 9 10 12 14
3 5 10 13 14 15	6 8 9 10 14 15	5 7 9 10 13 16	6 7 11 13 15 16	6 9 10 11 13 15
4 6 9 13 14 16	7 8 10 12 13 16	7 8 9 11 14 15	9 10 11 12 14 15	7 11 12 14 15 16

1 2 3 7 15 16	1 2 3 7 15 16	1 2 3 7 15 16	1 2 3 7 15 16	2 3 4 5 9 13
1 2 5 9 10 11	1 2 5 9 10 11	1 2 5 9 10 11	1 2 5 9 10 11	1 3 4 6 10 14
1 3 5 6 8 14	1 3 5 6 8 14	1 3 5 6 8 14	1 3 5 6 8 14	1 2 4 7 11 15
1 4 6 7 9 13	1 4 6 7 9 13	1 4 6 7 9 13	1 4 6 7 9 13	1 2 3 8 12 16
1 4 10 12 14 16	1 4 10 12 14 16	1 4 10 12 14 16	1 4 10 12 14 16	1 6 7 8 9 13
1 8 11 12 13 15	1 8 11 12 13 15	1 8 11 12 13 15	1 8 11 12 13 15	2 5 7 8 10 14
2 3 4 5 12 13	2 3 4 5 12 13	2 3 4 5 12 13	2 3 4 5 12 13	3 5 6 8 11 15
2 4 6 11 14 15	2 4 6 11 14 15	2 4 6 11 14 15	2 4 6 11 14 15	4 5 6 7 12 16
2 6 8 10 13 16	2 6 8 10 13 16	2 6 8 10 13 16	2 6 8 10 13 16	1 5 10 11 12 13
2 7 8 9 12 14	2 7 8 9 12 14	2 7 8 9 12 14	2 7 8 9 12 14	2 6 9 11 12 14
3 4 8 9 10 15	3 4 8 9 10 15	3 4 8 9 10 15	3 4 8 9 10 15	3 7 9 10 12 15
3 6 9 11 12 16	3 6 9 11 12 16	3 6 9 11 12 16	3 6 9 11 12 16	4 8 9 10 11 16
3 7 10 11 13 14	3 7 10 11 13 14	3 7 10 11 13 14	3 7 10 11 13 14	1 5 9 14 15 16
4 5 7 8 11 16	4 5 7 8 11 16	4 5 7 8 11 16	4 5 7 8 11 16	2 6 10 13 15 16
5 6 7 10 12 15	5 6 7 10 12 15	5 6 7 10 12 15	5 6 7 10 12 15	3 7 11 13 14 16
5 9 13 14 15 16	5 9 13 14 15 16	5 9 13 14 15 16	5 9 13 14 15 16	4 8 12 13 14 15
1 2 3 6 11 13	1 2 3 6 11 13	1 2 3 6 11 13	1 2 3 6 11 13	2 3 4 6 11 16
1 2 5 12 14 15	1 2 5 12 14 15	1 2 5 12 14 15	1 2 5 12 14 15	1 3 4 8 9 15
1 3 4 5 7 10	1 3 4 5 7 10	1 3 4 5 7 10	1 3 4 5 7 10	1 2 4 5 12 14
1 4 9 11 15 16	1 4 9 11 15 16	1 4 9 11 15 16	1 4 9 11 15 16	1 2 3 7 10 13
1 6 7 8 12 16	1 6 7 8 12 16	1 6 7 8 12 16	1 6 7 8 12 16	1 5 6 8 10 16
1 8 9 10 13 14	1 8 9 10 13 14	1 8 9 10 13 14	1 8 9 10 13 14	2 5 9 10 11 15
2 3 5 8 9 16	2 3 5 8 9 16	2 3 5 8 9 16	2 3 5 8 9 16	3 8 10 11 12 14
2 4 6 9 10 12	2 4 6 9 10 12	2 4 6 9 10 12	2 4 6 9 10 12	4 5 7 8 11 13
2 4 7 13 14 16	2 4 7 13 14 16	2 4 7 13 14 16	2 4 7 13 14 16	1 7 9 11 12 16
2 7 8 10 11 15	2 7 8 10 11 15	2 7 8 10 11 15	2 7 8 10 11 15	2 6 7 8 12 15
3 4 8 11 12 14	3 4 8 11 12 14	3 4 8 11 12 14	3 4 8 11 12 14	3 5 6 7 9 14
3 6 10 14 15 16	3 6 10 14 15 16	3 6 10 14 15 16	3 6 10 14 15 16	4 6 9 10 12 13
3 7 9 12 13 15	3 7 9 12 13 15	3 7 9 12 13 15	3 7 9 12 13 15	1 6 11 13 14 15
4 5 6 8 13 15	4 5 6 8 13 15	4 5 6 8 13 15	4 5 6 8 13 15	2 8 9 13 14 16
5 6 7 9 11 14	5 6 7 9 11 14	5 6 7 9 11 14	5 6 7 9 11 14	3 5 12 13 15 16
5 10 11 12 13 16	5 10 11 12 13 16	5 10 11 12 13 16	5 10 11 12 13 16	4 7 10 14 15 16
1 2 6 7 10 14	1 2 6 9 14 16	1 2 6 7 10 14	1 2 6 9 14 16	2 3 7 8 9 11
1 2 9 12 13 16	1 2 7 10 12 13	1 2 9 12 13 16	1 2 7 10 12 13	1 4 5 6 9 11
4 6 8 9 14 16	4 6 7 8 10 14	4 6 8 9 14 16	4 6 7 8 10 14	1 2 10 11 14 16
4 7 8 10 12 13	4 8 9 12 13 16	4 7 8 10 12 13	4 8 9 12 13 16	1 2 9 12 13 15
1 3 4 6 12 15	1 3 4 13 14 15	1 3 4 6 12 15	1 3 4 6 12 15	1 4 7 8 10 12
1 4 5 11 13 14	1 4 5 6 11 12	1 4 5 11 13 14	1 4 5 6 11 12	2 3 5 6 10 12
2 3 8 13 14 15	2 3 6 8 12 15	2 3 8 13 14 15	2 3 6 8 12 15	3 4 9 12 14 16
2 5 6 8 11 12	2 5 8 11 13 14	2 5 6 8 11 12	2 5 8 11 13 14	3 4 10 11 13 15
1 3 8 10 11 16	1 3 7 8 9 11	1 3 8 10 11 16	1 3 8 10 11 16	1 3 5 8 13 14
1 5 7 8 9 15	1 5 8 10 15 16	1 5 7 8 9 15	1 5 8 10 15 16	2 4 6 7 13 14
2 3 4 7 9 11	2 3 4 10 11 16	2 3 4 7 9 11	2 3 4 10 11 16	5 7 9 10 13 16
2 4 5 10 15 16	2 4 5 7 9 15	2 4 5 10 15 16	2 4 5 7 9 15	5 7 11 12 14 15
3 5 6 9 10 13	3 5 6 7 13 16	3 5 6 9 10 13	3 5 6 7 13 16	1 3 6 7 15 16
3 5 7 12 14 16	3 5 9 10 12 14	3 5 7 12 14 16	3 5 9 10 12 14	2 4 5 8 15 16
6 7 11 13 15 16	6 9 10 11 13 15	6 7 11 13 15 16	6 9 10 11 13 15	6 8 9 10 14 15
9 10 11 12 14 15	7 11 12 14 15 16	9 10 11 12 14 15	7 11 12 14 15 16	6 8 11 12 13 16



Here we give the 2 - (71,15,3) designs  $D_1^*$ ,  $D_2^*$ ,  $D_3^*$  and  $D_4^*$ , which are constructed in Section 6.1. Together with their duals they form the eight 2 - (71,15,3) designs of Theorem 6.1.5.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	2	3	16	17	18	19	20	21	22	23	24	25	26	27	1	2	3	16	17	18	19	20	21	22	23	24	25	26	27
1	2	3	28	29	30	31	32	33	34	35	36	37	38	39	1	2	3	28	29	30	31	32	33	34	35	36	37	38	39
1	4	5	16	17	28	29	40	41	42	43	44	45	46	47	1	4	5	16	17	28	29	40	41	42	43	44	45	46	47
1	4	5	18	19	30	31	48	49	50	51	52	53	54	55	1	4	5	18	19	30	31	48	49	50	51	52	53	54	55
1	6	7	16	17	30	31	56	57	58	59	60	61	62	63	1	6	7	16	17	30	31	56	57	58	59	60	61	62	63
1	6	7	18	19	28	29	64	65	66	67	68	69	70	71	1	6	7	18	19	28	29	64	65	66	67	68	69	70	71
1	8	9	20	21	32	33	40	41	48	49	56	57	64	65	1	8	9	20	21	32	33	40	41	48	49	56	57	64	65
1	8	9	22	23	34	35	42	43	50	51	58	59	66	67	1	8	9	22	23	34	35	42	43	50	51	58	59	66	67
1	10	11	20	21	34	35	44	45	52	53	60	61	68	69	1	10	11	20	21	34	35	44	45	52	53	60	61	68	69
1	10	11	22	23	32	33	46	47	54	55	62	63	70	71	1	10	11	22	23	32	33	46	47	54	55	62	63	70	71
1	12	13	24	25	36	37	40	41	50	51	60	61	70	71	1	12	13	24	25	36	37	40	41	50	51	60	61	70	71
1	12	13	26	27	38	39	42	43	48	49	62	63	68	69	1	12	13	26	27	38	39	42	43	48	49	62	63	68	69
1	14	15	24	25	38	39	44	45	54	55	56	57	66	67	1	14	15	24	25	38	39	44	45	54	55	56	57	66	67
1	14	15	26	27	36	37	46	47	52	53	58	59	64	65	1	14	15	26	27	36	37	46	47	52	53	58	59	64	65
2	4	6	20	22	36	38	40	42	52	54	56	58	68	70	2	4	6	20	22	36	38	40	42	52	54	56	58	68	70
2	4	6	21	23	37	39	41	43	53	55	57	59	69	71	2	4	6	21	23	37	39	41	43	53	55	57	59	69	71
2	5	7	20	22	37	39	44	46	48	50	60	62	64	66	2	5	7	20	22	37	39	44	46	48	50	60	62	64	66
2	5	7	21	23	36	38	45	47	49	51	61	63	65	67	2	5	7	21	23	36	38	45	47	49	51	61	63	65	67
2	8	10	24	26	28	30	40	46	48	54	59	61	67	69	2	8	10	24	26	28	30	40	46	48	54	59	61	67	69
2	8	10	25	27	29	31	41	47	49	55	58	60	66	68	2	8	10	25	27	29	31	41	47	49	55	58	60	66	68
2	9	11	24	26	29	31	43	45	51	53	56	62	64	70	2	9	11	24	26	29	31	43	45	51	53	56	62	64	70
2	9	11	25	27	28	30	42	44	50	52	57	63	65	71	2	9	11	25	27	28	30	42	44	50	52	57	63	65	71
2	12	14	16	18	32	34	40	44	51	55	59	63	64	68	2	12	14	16	18	32	34	40	44	51	55	59	63	64	68
2	12	14	17	19	33	35	41	45	50	54	58	62	65	69	2	12	14	17	19	33	35	41	45	50	54	58	62	65	69
2	13	15	16	18	33	35	42	46	49	53	57	61	66	70	2	13	15	16	18	33	35	42	46	49	53	57	61	66	70
2	13	15	17	19	32	34	43	47	48	52	56	60	67	71	2	13	15	17	19	32	34	43	47	49	53	56	60	66	70
3	4	7	24	27	32	35	40	42	53	55	60	62	65	67	3	4	7	24	27	32	35	40	42	53	55	60	62	65	67
3	4	7	25	26	33	34	41	43	52	54	61	63	64	66	3	4	7	25	26	33	34	41	43	52	54	60	62	65	67
3	5	6	24	27	33	34	44	46	49	51	56	58	69	71	3	5	6	24	27	33	34	44	46	49	51	56	59	68	70
3	5	6	25	26	32	35	45	47	48	50	57	59	68	70	3	5	6	25	26	32	35	45	47	48	50	56	58	69	71
3	8	11	16	19	37	38	41	44	49	52	59	62	67	70	3	8	11	16	19	37	38	41	44	48	52	59	63	66	70
3	8	11	17	18	36	39	40	45	48	53	58	63	66	71	3	8	11	17	18	36	39	43	47	50	54	57	61	64	68
3	9	10	16	19	36	39	43	46	51	54	57	60	65	68	3	9	10	16	19	36	39	40	44	49	53	58	62	67	71
3	9	10	17	18	37	38	42	47	50	55	56	61	64	69	3	9	10	17	18	37	38	42	46	51	55	58	63	67	71
3	12	15	20	23	29	30	40	47	51	52	57	62	66	69	3	12	15	20	23	29	30	43	44	48	55	58	61	65	70
3	12	15	21	22	28	31	41	46	50	53	56	63	67	68	3	12	15	21	22	28	31	41	46	50	53	56	63	67	68
3	13	14	20	23	28	31	43	44	48	55	58	61	65	70	3	13	14	20	23	28	31	40	47	51	52	57	62	66	69
3	13	14	21	22	29	30	42	45	49	54	59	60	64	71	3	13	14	21	22	29	30	42	45	49	54	59	60	64	71
4	8	13	18	23	25	30	34	37	45	46	56	62	65	68	4	8	12	18	21	27	31	33	38	44	47	58	60	66	71
4	8	13	19	22	24	31	35	36	44	47	57	63	64	69	4	8	12	19	23	24	30	34	36	45	46	56	62	64	69
4	9	12	18	21	26	30	33	39	44	47	58	60	67	70	4	9	13	18	20	26	31	32	39	45	46	56	61	67	70
4	9	12	19	20	27	31	32	38	45	46	59	61	66	71	4	9	13	19	22	25	30	35	37	44	47	57	63	65	68
4	10	15	16	21	25	28	35	38	48	51	58	62	64	71	4	10	14	16	23	26	29	33	37	49	50	56	61	66	68
4	10	15	17	20	24	29	34	39	49	50	59	63	65	70	4	10	14	17	21	25	28	34	39	48	51	58	63	64	70
4	11	14	16	23	26	28	32	36	49	50	56	60	66	69	4	11	15	16	22	27	29	32	36	48	51	57	60	67	69
4	11	14	17	22	27	29	33	37	48	51	57	61	67	68	4	11	15	17	20	24	28	35	38	49	50	59	62	65	71
5	8	12	16	22	26	29	35	39	52	55	56	61	65	71	5	8	13	16	22	27	28	34	39	52	55	56	61	65	71
5	8	12	17	23	27	28	34	38	53	54	57	60	64	70	5	8	13	17	20	24	29	33	37	53	54	58	63	67	69
5	9	13	16	20	25	29	32	37	53	54	58	63	67	69	5	9	12	16	23	26	28	35	38	53	54	57	60	64	70
5	9	13	17	21	24	28	33	36	52	55	59	62	66	68	5	9	12	17	21	25	29	32	36	52	55	59	62	66	68
5	10	14	18	20	26	31	34	36	41	42	57	62	67	71	5	10	15	18	20	26	30	34	36	41	42	57	63	66	71
5	10	14	19	21	27	30	35	37	40	43	56	63	66	70	5	10	15	19	22	25	31	33	38	40	43	59	61	64	69
5	11	15	18	22	25	31	33	38	40	43	59	60	65	69	5	11	14	18	21	27	30	35	37	40	43	56	62	67	70
5	11	15	19	23	24	30	32	39	41	42	58	61	64	68	5	11	14	19	23	24	31	32	39	41	42	58	60	65	68
6	8	15	18	21	27	29	32	36	43	44	50	54	61	62	6	8	15	18	23	25	29	35	39	40	46	49	52	60	63
6	8	15	19	20	26	28	33	37	42	45	51	55	60	63	6	8	15	19	21	26	28	32	37	42	44	51	54	61	62
6	9	14	18	23	24	29	35	38	41	46	48	52	60	63	6	9	14	18	22	24	29	34							

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15  
1 2 3 16 17 18 19 20 21 22 23 24 25 26 27  
1 2 3 28 29 30 31 32 33 34 35 36 37 38 39  
1 4 5 16 17 28 29 40 41 42 43 44 45 46 47  
1 4 5 18 19 30 31 48 49 50 51 52 53 54 55  
1 6 7 16 17 30 31 56 57 58 59 60 61 62 63  
1 6 7 18 19 28 29 64 65 66 67 68 69 70 71  
1 8 9 20 21 32 33 40 41 48 49 56 57 64 65  
1 8 9 22 23 34 35 42 43 50 51 58 59 66 67  
1 10 11 20 21 34 35 44 45 52 53 60 61 68 69  
1 10 11 22 23 32 33 46 47 54 55 62 63 70 71  
1 12 13 24 25 36 37 42 43 48 49 62 63 68 69  
1 12 13 26 27 38 39 40 41 50 51 60 61 70 71  
1 14 15 24 25 38 39 46 47 52 53 58 59 64 65  
1 14 15 26 27 36 37 44 45 54 55 56 57 66 67  
2 4 6 24 26 32 34 44 46 48 50 56 58 68 70  
2 4 6 25 27 33 35 45 47 49 51 57 59 69 71  
2 5 7 24 26 33 35 40 42 52 54 60 62 64 66  
2 5 7 25 27 32 34 41 43 53 55 61 63 65 67  
2 8 10 16 18 36 38 41 45 51 55 58 62 64 68  
2 8 10 17 19 37 39 43 47 49 53 56 60 66 70  
2 9 11 16 18 37 39 40 44 50 54 59 63 65 69  
2 9 11 17 19 36 38 42 46 48 52 57 61 67 71  
2 12 14 20 22 28 30 42 45 51 52 56 63 65 70  
2 12 14 21 23 29 31 40 47 49 54 58 61 67 68  
2 13 15 20 22 29 31 41 46 48 55 59 60 66 69  
2 13 15 21 23 28 30 43 44 50 53 57 62 64 71  
3 4 7 20 23 36 39 44 46 49 51 60 62 65 67  
3 4 7 21 22 37 38 45 47 48 50 61 63 64 66  
3 5 6 20 23 37 38 40 42 53 55 56 58 69 71  
3 5 6 21 22 36 39 41 43 52 54 57 59 68 70  
3 8 11 24 27 28 31 42 47 50 55 57 60 65 68  
3 8 11 25 26 29 30 40 45 48 53 59 62 67 70  
3 9 10 24 27 29 30 41 44 49 52 58 63 66 71  
3 9 10 25 26 28 31 43 46 51 54 56 61 64 69  
3 12 15 16 19 32 35 41 47 50 52 56 62 67 69  
3 12 15 17 18 33 34 43 45 48 54 58 60 65 71  
3 13 14 16 19 33 34 40 46 51 53 57 63 66 68  
3 13 14 17 18 32 35 42 44 49 55 59 61 64 70  
4 8 12 16 20 24 28 32 36 53 54 59 61 66 71  
4 8 12 17 23 26 29 34 39 52 55 57 63 64 69  
4 9 13 16 21 25 28 33 37 52 55 58 60 67 70  
4 9 13 17 22 27 29 35 38 53 54 56 62 65 68  
4 10 14 18 22 24 30 32 38 40 43 57 60 67 69  
4 10 14 19 21 26 31 34 37 41 42 59 62 65 71  
4 11 15 18 23 25 30 33 39 41 42 56 61 66 68  
4 11 15 19 20 27 31 35 36 40 43 58 63 64 70  
5 8 13 18 21 24 31 35 39 45 46 56 63 67 71  
5 8 13 19 22 26 30 33 36 44 47 58 61 65 69  
5 9 12 18 20 25 31 34 38 44 47 57 62 66 70  
5 9 12 19 23 27 30 32 37 45 46 59 60 64 68  
5 10 15 16 23 24 29 35 37 48 51 57 61 65 70  
5 10 15 17 20 26 28 33 38 49 50 59 63 67 68  
5 11 14 16 22 25 29 34 36 49 50 56 60 64 71  
5 11 14 17 21 27 28 32 39 48 51 58 62 66 69  
6 8 14 18 20 27 29 33 37 43 46 50 52 61 62  
6 8 14 19 23 25 28 35 38 41 44 48 54 60 63  
6 9 15 18 21 26 29 32 36 42 47 51 53 60 63  
6 9 15 19 22 24 28 34 39 40 45 49 55 61 62  
6 10 12 16 22 27 31 33 39 42 44 48 53 64 67  
6 10 12 17 21 25 30 35 36 40 46 50 55 65 66  
6 11 13 16 23 26 31 32 38 43 45 49 52 65 66  
6 11 13 17 20 24 30 34 37 41 47 51 54 64 67  
7 8 15 16 21 27 30 34 38 42 46 49 54 69 70  
7 8 15 17 22 25 31 32 37 40 44 51 52 68 71  
7 9 14 16 20 26 30 35 39 43 47 48 55 68 71  
7 9 14 17 23 24 31 33 36 41 45 50 53 69 70  
7 10 13 18 23 27 28 34 36 40 47 48 52 56 59  
7 10 13 19 20 25 29 32 39 42 45 50 54 57 58  
7 11 12 18 22 26 28 35 37 41 46 49 53 57 58  
7 11 12 19 21 24 29 33 38 43 44 51 55 56 59

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15  
1 2 3 16 17 18 19 20 21 22 23 24 25 26 27  
1 2 3 28 29 30 31 32 33 34 35 36 37 38 39  
1 4 5 16 17 28 29 40 41 42 43 44 45 46 47  
1 4 5 18 19 30 31 48 49 50 51 52 53 54 55  
1 6 7 16 17 30 31 56 57 58 59 60 61 62 63  
1 6 7 18 19 28 29 64 65 66 67 68 69 70 71  
1 8 9 20 21 32 33 40 41 48 49 56 57 64 65  
1 8 9 22 23 34 35 42 43 50 51 58 59 66 67  
1 10 11 20 21 34 35 44 45 52 53 60 61 68 69  
1 10 11 22 23 32 33 46 47 54 55 62 63 70 71  
1 12 13 24 25 36 37 42 43 48 49 62 63 68 69  
1 12 13 26 27 38 39 40 41 50 51 60 61 70 71  
1 14 15 24 25 38 39 46 47 52 53 58 59 64 65  
1 14 15 26 27 36 37 44 45 54 55 56 57 66 67  
2 4 6 24 26 32 34 44 46 48 50 56 58 68 70  
2 4 6 25 27 33 35 45 47 49 51 57 59 69 71  
2 5 7 20 24 34 36 40 47 51 54 56 58 68 70  
2 5 7 21 25 35 37 41 46 50 55 57 59 69 71  
2 8 10 16 18 37 38 42 45 51 55 58 62 64 71  
2 8 10 17 19 36 39 43 44 50 53 57 63 65 70  
2 9 11 16 27 28 38 40 48 53 55 58 63 66 69  
2 9 11 17 26 29 39 41 49 52 54 59 62 67 68  
2 12 14 18 22 30 32 40 42 44 54 59 61 65 69  
2 12 14 19 23 31 33 41 43 45 55 58 60 64 68  
2 13 15 20 22 28 30 43 47 48 52 57 60 67 71  
2 13 15 21 23 29 31 42 46 49 53 56 61 66 70  
3 4 7 22 26 32 39 43 45 49 53 56 58 69 71  
3 4 7 23 27 33 38 42 44 48 52 57 59 68 70  
3 5 6 20 22 36 39 41 42 52 55 61 63 64 66  
3 5 6 21 23 37 38 40 43 53 54 60 62 65 67  
3 8 11 24 26 29 31 42 47 51 55 57 60 65 69  
3 8 11 25 27 28 30 43 46 50 54 56 61 64 68  
3 9 10 18 25 30 37 41 44 47 49 58 60 66 70  
3 9 10 19 24 31 36 40 45 46 48 59 61 67 71  
3 12 15 16 18 32 34 41 46 51 53 57 63 67 68  
3 12 15 17 19 33 35 40 47 50 52 56 62 66 69  
3 13 14 16 20 28 34 45 49 50 55 59 62 65 70  
3 13 14 17 21 29 35 44 48 51 54 58 63 64 71  
4 8 12 16 20 25 29 33 36 53 54 59 60 66 71  
4 8 12 17 21 24 28 32 37 52 55 58 61 67 70  
4 9 13 16 19 23 34 37 39 47 54 57 61 64 69  
4 9 13 17 18 22 35 36 38 46 55 56 60 65 68  
4 10 14 20 23 24 29 30 38 41 50 56 63 67 69  
4 10 14 21 22 25 28 31 39 40 51 57 62 66 68  
4 11 15 18 20 26 31 35 37 40 43 59 63 64 70  
4 11 15 19 21 27 30 34 36 41 42 58 62 65 71  
5 8 13 18 20 27 31 33 39 44 46 58 62 67 69  
5 8 13 19 21 26 30 32 38 45 47 59 63 66 68  
5 9 12 22 24 27 29 30 35 45 53 57 62 64 70  
5 9 12 23 25 26 28 31 34 44 52 56 63 65 71  
5 10 15 16 22 26 29 33 37 48 50 58 61 65 68  
5 10 15 17 23 27 28 32 36 49 51 59 60 64 69  
5 11 14 16 19 25 32 35 39 42 48 56 60 67 70  
5 11 14 17 18 24 33 34 38 43 49 57 61 66 71  
6 8 15 16 23 24 30 35 39 40 44 49 55 68 71  
6 8 15 17 22 25 31 34 38 41 45 48 54 69 70  
6 9 14 16 21 26 30 33 36 43 46 51 52 69 70  
6 9 14 17 20 27 31 32 37 42 47 50 53 68 71  
6 10 12 18 21 27 29 34 39 43 47 48 55 56 59  
6 10 12 19 20 26 28 35 38 42 46 49 54 57 58  
6 11 13 18 23 25 29 32 36 40 45 50 52 57 58  
6 11 13 19 22 24 28 33 37 41 44 51 53 56 59  
7 8 14 18 23 26 28 35 36 41 47 48 53 61 62  
7 8 14 19 22 27 29 34 37 40 46 49 52 60 63  
7 9 15 18 21 24 28 33 39 42 45 50 54 60 63  
7 9 15 19 20 25 29 32 38 43 44 51 55 61 62  
7 10 13 16 24 27 31 32 35 41 43 52 54 65 66  
7 10 13 17 25 26 30 33 34 40 42 53 55 64 67  
7 11 12 16 21 22 31 36 38 44 47 49 50 64 67  
7 11 12 17 20 23 30 37 39 45 46 48 51 65 66

## REFERENCES

- [A1] H. AGRAWAL, *On the bounds of the number of common treatments between blocks of certain two associate PBIB designs*, Calcutta Statist. Assoc. Bull. 13 (1964), 76-79.
- [A2] R.W. AHRENS & G. SZEKERES, *On a combinatorial generalization of the 27 lines associated with a cubic surface*, J. Austral. Math. Soc. 10 (1969), 485-492.
- [A3] N. ARONSZAJN, *Rayleigh-Ritz and A. Weinstein method for approximation of eigenvalues. I. Operators in a Hilbert space*, Proc. American Nat. Acad. Sci. 34 (1948), 474-480.
- [A4] E.F. ASSMUS, Jr. & J.H. van LINT, *Ovals in projective designs*, J. Combinatorial Theory A 27 (1979), 307-324.
- [B1] H.J. BEKER, *Constructions and decompositions of designs*, Ph.D. Thesis, University of London, 1976.
- [B2] H.J. BEKER & W. HAEMERS, *2-designs having an intersection number  $k - n$* , J. Combinatorial Theory A 28 (1980), 64-81.
- [B3] H.J. BEKER & C.J. MITCHEL, *A construction method for point divisible designs*, J. Statist. Planning and Inference 2 (1978), 293-306.
- [B4] N. BIGGS, *Finite groups of automorphisms*, London Math. Soc. Lecture Notes No. 6, Cambridge University Press, Cambridge, 1971.
- [B5] N. BIGGS, *Algebraic Graph Theory*, Cambridge tracts in math. No. 67, Cambridge University Press, Cambridge, 1974.
- [B6] R.C. BOSE, *Strongly regular graphs, partial geometries and partially balanced designs*, Pacific J. Math. 13 (1963), 389-419.
- [B7] R.C. BOSE, *Graphs and designs*, in "Finite geometric structures and their applications" (A. Barlotti ed.), Cremonese, Rome, 1973, pp. 3-104.
- [B8] R.C. BOSE & D.M. MESNER, *On linear associative algebras corresponding to association schemes of partially balanced designs*, Ann. Math. Statist. 30 (1959), 21-38.

- [B9] C. BUMILLER, *Pointwise products of incidence matrices of subgraphs* (preprint).
- [B10] K.A. BUSH, *Bounds for the number of common symbols in balanced and certain partially balanced designs*, J. Combinatorial Theory A 23 (1977), 46-51.
- [B11] F.C. BUSSEMAKER & J.J. SEIDEL, *Symmetric Hadamard matrices of order 36*, Ann. N.Y. Acad. Sci. 175 (1970), 66-79; also: Report Techn. Univ. Eindhoven 70-WSK-02.
- [C1] P.J. CAMERON, *On groups with several doubly-transitive permutation representations*, Math. Z. 128 (1972), 1-14.
- [C2] P.J. CAMERON, *Extending symmetric designs*, J. Combinatorial Theory A 14 (1973), 215-220.
- [C3] P.J. CAMERON, *Partial quadrangles*, Quart. J. Math. Oxford (3) 25 (1974), 1-13.
- [C4] P.J. CAMERON, *Parallelisms of Complete Designs*, London Math. Soc. Lecture Notes 23, Cambridge University Press, Cambridge, 1976.
- [C5] P.J. CAMERON, J.M. GOETHALS & J.J. SEIDEL, *Strongly regular graphs having strongly regular subconstituents*, J. Algebra 55 (1978), 257-280.
- [C6] P.J. CAMERON & J.H. van LINT, *Graph theory, coding theory and block designs*, London Math. Soc. Lecture Notes No. 19, Cambridge University Press, Cambridge, 1975.
- [C7] F. de CLERCK, *Partial geometries: a combinatorial survey*, Bull. Belgisch Wiskundig Genootschap (to appear).
- [C8] W.S. CONNOR, *On the structure of balanced incomplete block designs*, Ann. Math. Statist. 23 (1952), 57-71.
- [C9] W.S. CONNOR & W.H. CLATWORTHY, *Some theorems for partially balanced designs*, Ann. Math. Statist. 25 (1954), 100-112.
- [C10] G.M. CONWELL, *The 3-space PG(3,2) and its group*, Ann. Math. 11 (1910), 60-76.
- [C11] R. COURANT & D. HILBERT, *Methoden der Mathematischen Physik*, Vol. 1, Springer Verlag, Berlin, 1924.
- [C12] D.M. CVETKOVIĆ, *Graphs and their spectra*, Thesis, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. 354-356 (1971), 1-50.

- [C13] D.M. CVETKOVIĆ, M. DOOB & H. SACHS, *Spectra of Graphs - Theory and Applications*, V.E.B. Deutscher Verlag der Wissenschaften, Berlin, 1980.
- [D1] P. DELSARTE, *An algebraic approach to the association schemes of coding theory*, Philips Res. Rep. Suppl. 10 (1973).
- [D2] P. DELSARTE, J.M. GOETHALS & J.J. SEIDEL, *Spherical codes and designs*, Geometriae Dedicata 6 (1977), 363-388.
- [D3] P. DEMBOWSKI, *Finite Geometries*, Springer Verlag, Berlin-Heidelberg-New York, 1968.
- [F1] W. FEIT & G. HIGMAN, *The non-existence of certain generalized polygons*, J. Algebra 1 (1964), 114-131.
- [G1] A. GEWIRTZ, *The uniqueness of  $g(2,2,10,56)$* , Trans. New York Acad. Sci. 31 (1969), 656-675.
- [G2] J.M. GOETHALS & J.J. SEIDEL, *Strongly regular graphs derived from combinatorial designs*, Canad. J. Math. 22 (1970), 597-614.
- [H1] W. HAEMERS, *A generalization of the Higman-Sims technique*, Proc. Kon. Nederl. Akad. Wetensch. A 81 (= Indag. Math. 40) (1978), 445-447.
- [H2] W. HAEMERS, *Eigenvalue methods*, in: "Packing and Covering in Combinatorics" (A. Schrijver ed.), Tract 106, Mathematical Centre, Amsterdam, 1979, pp. 15-38.
- [H3] W. HAEMERS & C. ROOS, *An inequality for generalized hexagons*, Geometriae Dedicata (to appear).
- [H4] W. HAEMERS & M.S. SHRIKHANDE, *Some remarks on subdesigns of symmetric designs*, J. Statist. Planning and Inference (to appear).
- [H5] M. HALL, Jr., *Combinatorial Theory*, Blaisdell, Waltham-Toronto-London, 1967.
- [H6] M. HALL, Jr., *Affine generalized quadrilaterals*, in: "Studies in pure mathematics" (L. Mirsky ed.), Academic Press, 1971, pp. 113-116.
- [H7] H.L. HAMBURGER & M.E. GRIMSHAW, *Linear transformations in  $n$ -dimensional vector space*, Cambridge University Press, London-New York, 1951.
- [H8] F. HARARY, *Graph Theory*, Addison-Wesley, Reading, Mass., 1969.
- [H9] E.V. HAYNSWORTH, *Applications of a theorem on partitioned matrices*, J. Res. Nat. Bur. Standards B 63 (1959), 73-78.

- [H10] M.D. HESTENES & D.G. HIGMAN, *Rank 3 groups and strongly regular graphs*, in: "Computers in Algebra and Number Theory" (G. Birkhoff & M. Hall, Jr., eds.), SIAM-AMS Proc. IV, Amer. Math. Soc., Providence, R.I., 1971, pp. 141-159.
- [H11] D.G. HIGMAN, *Partial geometries, generalized quadrangles and strongly regular graphs*, in: *Atti Convegno di Geometria e sue Applicazioni*", Perugia, 1971, pp. 263-293.
- [H12] D.G. HIGMAN, *Invariant relations, coherent configurations and generalized polygons*, in: "Combinatorics; part 3: combinatorial group theory" (M. Hall, Jr. & J.H. van Lint, eds.), Tract 57, Mathematical Centre, Amsterdam, 1974, pp. 27-43
- [H13] A.J. HOFFMAN, *On eigenvalues and colourings of graphs*, in: "Graph Theory and its Applications" (B. Harris, ed.), Acad. Press, New York, 1970, pp. 79-91.
- [H14] A.J. HOFFMAN, *Eigenvalues of graphs*, in: "Studies in Graph Theory; part II" (D.R. Fulkerson, ed.), M.A.A. Studies in Math., 1975, pp. 225-245.
- [H15] A.J. HOFFMAN & R.R. SINGLETON, *On Moore graphs of diameters 2 and 3*, I.B.M. J. Res. Develop. 4 (1960), 497-504.
- [H16] D.R. HUGHES & F.C. PIPER, *Projective Planes*, Springer Verlag, Berlin-Heidelberg-New York, 1973.
- [H17] D.R. HUGHES & F.C. PIPER, *Block Designs* (in preparation).
- [K1] W.M. KANTOR, *Generalized quadrangles associated with  $G_2(q)$*  (preprint).
- [K2] R. KILMOYER & L. SOLOMON, *On the theorem of Feit-Higman*, J. Combinatorial Theory A 15 (1973), 310-322.
- [L1] J.H. van LINT & H.J. RYSER, *Block designs with repeated blocks*, Discrete Math. 3 (1972), 381-396.
- [L2] L. LOVÁSZ, *On the Shannon capacity of a graph*, IEEE Trans. Information Theory 25 (1979), 1-7.
- [M1] K.N. MAJUMDAR, *On some theorems in combinatorics relating to incomplete block designs*, Ann. Math. Statist. 24 (1953), 379-389.

- [M2] H.B. MANN, *A note on balanced incomplete block designs*, Ann. Math. Statist. 40 (1969), 679-680.
- [M3] M. MARCUS & H. MINC, *A survey of matrix theory and matrix inequalities*, Allyn & Bacon, Boston, 1964.
- [M4] R. MATHON, *3-class association schemes*, in: "Proc. Conf. on Algebraic Aspects of Combinatorics", (D.G. Corneil & E. Mendelsohn, eds.), Congressus Numerantium XIII, Utilitas, Winnipeg, 1975, pp. 123-155.
- [M5] R. MATHON, *Symmetric conference matrices of order  $pq^2 + 1$* , Canad. J. Math. 30 (1978), 321-331.
- [M6] R. MATHON, *On primitive association schemes with three classes* (in preparation).
- [M7] R. MATHON, *The systems of linked (16,6,2) designs* (submitted to Ars Combinatoria).
- [M8] C.J. MITCHELL, *An infinite family of symmetric designs*, Discrete Math. 26 (1979), 247-250.
- [N1] B. NOBLE & J.W. DANIEL, *Applied Linear Algebra* (second edition), Prentice-Hall, Englewood Cliffs, 1977.
- [P1] S.E. PAYNE, *Nonisomorphic generalized quadrangles*, J. Algebra 18 (1971), 201-212.
- [P2] S.E. PAYNE, *Generalized quadrangles of order (s-1,s+1)*, J. Algebra 22 (1972), 97-119.
- [P3] S.E. PAYNE, *A restriction on the parameters of a subquadrangle*, Bull. Amer. Math. Soc. 79 (1973), 747-748.
- [P4] S.E. PAYNE, *Finite generalized quadrangles: a survey*, in: "Proc. Int. Conf. on Projective Planes", Wash. State Univ. Press, 1973, pp. 291-261.
- [P5] S.E. PAYNE, *All generalized quadrangles of order 3 are known*, J. Combinatorial Theory A 18 (1975), 203-206.
- [P6] S.E. PAYNE, *An inequality for generalized quadrangles*, Proc. Amer. Math. Soc. 71 (1978), 147-152.

- [R1] M.A. RONAN, *A geometric characterization of Moufang hexagons*, Ph. D. Thesis, University of Illinois, 1978.
- [R2] M.A. RONAN, *A combinatorial characterization of the dual Moufang hexagon* (preprint).
- [R3] H.J. RYSER, *Combinatorial Mathematics*, Carus Mathematical Monograph 14, Wiley, New York, 1963.
- [S1] G.J. SCHELLEKENS, *On a hexagonal structure I & II*, Proc. Kon. Nederl. Akad. Wetensch. A 65 (= Indag. Math. 24) (1962), 201-217, 218-234.
- [S2] A.J. SCHWENK & R.J. WILSON, *Eigenvalues of graphs*, in: "Selected Topics in Graph Theory" (L.W. Beineke & R.J. Wilson, eds.), Academic Press, London-New York-San Francisco, 1978, pp. 307-336.
- [S3] J.J. SEIDEL, *Strongly regular graphs with  $(-1,1,0)$  adjacency matrix having eigenvalue 3*, Linear Algebra and Appl. 1 (1968), 281-298.
- [S4] J.J. SEIDEL, *Strongly regular graphs*, in: "Surveys in Combinatorics" (B. Bollobás, ed.), London Math. Soc. Lecture Notes No. 38, Cambridge University Press, Cambridge, 1979, pp. 157-180.
- [S5] M.S. SHRIKHANDE, *Strongly regular graphs and quasi-symmetric designs*, Utilitas Mathematica 3 (1973), 297-309.
- [S6] S.S. SHRIKHANDE, *The uniqueness of the  $L_2$  association scheme*, Ann. Math. Statist. 30 (1959), 781-798.
- [S7] E.E. SHULT & A. YANUSHKA, *Near  $n$ -gons and line systems*, Geometriae Dedicata (to appear).
- [T1] J.A. THAS, *Combinatorics of partial geometries and generalized quadrangles*, in: "Higher Combinatorics", (M. Aigner, ed.), Reidel, Dordrecht, 1977, pp. 183-199.
- [T2] J.A. THAS, *A remark on the theorem of Yanushka-Ronan characterizing the generalized hexagon  $H(q)$  arising from the group  $G_2(q)$*  (preprint).
- [T3] J.A. THAS & S.E. PAYNE, *Classical finite generalized quadrangles: a combinatorial study*, Ars Combinatoria 2 (1976), 57-110.
- [T4] J. TITS, *Sur le triality et certains groupes qui s'en déduisent*, Publ. Math. I.H.E.S. 2 (1959), 14-60.



- [W1] B. WEISFEILER, *On Construction and Identification of Graphs*, Lecture Notes in Math. No. 558, Springer Verlag, Berlin-Heidelberg-New York, 1976.
- [W2] H. WEYL, *Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung)*, Math. Ann. 71 (1912), 441-479.
- [W3] H.S. WHITE, F.N. COLE & L.D. CUMMINGS, *Complete classification of the triad systems on fifteen elements*, Memoir Nat. Acad. Sci. 14, second memoir (1925), 1-89.
- [W4] H.A. WILBRINK & A.E. BROUWER, *A (57,14,1) strongly regular graph does not exist*, Report ZW 121/78, Math. Centre Amsterdam, 1978.
- [W5] J.H. WILKINSON, *The Algebraic Eigenvalue Problem*, Clarendon, Oxford, 1965.
- [W6] R.J. WILSON, *Introduction to Graph Theory* (second edition), Longman, London, 1979.
- [Y1] A. YANUSHKA, *Generalized hexagons of order (t,t)*, Israel. J. Math. 23 (1976), 309-324.
- [Y2] A. YANUSHKA, *A sufficient condition for the existence of an association scheme* (preprint).

## NOTATION

$A^*$	hermitian transpose of matrix $A$ .
$(A)_{ij}$	$ij$ -th entry of matrix $A$
$(b_{ij})$	matrix with entries $b_{ij}$ .
$\lambda_1(A) \geq \dots \geq \lambda_n(A)$	eigenvalues (if real) of matrix $A$ .
$A \otimes B$	Kronecker product of matrices $A$ and $B$ .
$\text{diag}(a_1, \dots, a_n)$	diagonal matrix with diagonal entries $a_1, \dots, a_n$ .
$I$ or $I_n$	$n \times n$ identity matrix.
$J$	all-one matrix.
$J_n$	$n \times n$ all-one matrix.
$j$	all-one vector.
$\ u\ $	length of vector $u$ .
$\langle u_1, \dots, u_n \rangle$	linear span of vectors $u_1, \dots, u_n$ .
$\langle u_1, \dots, u_n \rangle^\perp$	orthogonal complement of $\langle u_1, \dots, u_n \rangle$ .
$\bar{G}$	complement of graph $G$ .
$\alpha(G)$	size of largest coclique of $G$ .
$\gamma(G)$	chromatic number of $G$ .
$\omega(G)$	size of largest clique of $G$ .
$\lambda_1(G) \geq \dots \geq \lambda_n(G)$	eigenvalues of $G$ .
$f_i(G)$	multiplicity of eigenvalue $\lambda_i(G)$ .
$L(G)$	line graph of $G$ .
$D(G, G_1)$	incidence structure formed by $G$ and $G_1$ , see p. 17.
$K_n$	complete graph on $n$ vertices.
$K_{\ell, m}$	complete bipartite graph on $\ell + m$ vertices.
$\rho(x, y)$	distance between vertices $x$ and $y$ , see p. 79.
$p_{ij}(x, y)$ or $p_{ij}^k$	intersection numbers of a graph, see p. 51 and p. 81.
$\lambda(e_1, e_2)$	distance between elements of an $n$ -gon, see p. 52.
$p_{ijk}(L, x, y)$	see p. 58.
$\mathbb{Z}$	set of integers.
$\mathbb{N}$	set of positive integers.
$\mathbb{F}_q$	field with $q$ elements.
$\text{PG}(n, q)$	$n$ -dimensional projective geometry over $\mathbb{F}_q$ .
$\text{AG}(n, q)$	$n$ -dimensional affine geometry over $\mathbb{F}_q$ .
$\delta_{ij}$	1 if $i = j$ ; 0 if $i \neq j$ .
$\lfloor x \rfloor$	lower integer part of real number $x$ .
$\lceil x \rceil$	upper integer part of real number $x$ .

## INDEX

adjacency matrix	79	distance regular graph	51
adjacent vertices	79	dual of an incidence structure	83
balanced incomplete block design	83	edge	79
bipartite graph	80	eigenvalues of a graph	15,80
block	82	element of a generalized n-gon	52
block design	83	embeddable	83
block graph	35	empty incidence structure	82
Cauchy inequalities	8	equivalent blocks	34,65
chromatic number	80	equivalent points	65
circuit	79,81	flag	82
class graph	35	(generalized) n-gon	50
Clebsch graph	39	(generalized) hexagon	53
clique	80	(generalized) polygon	50
coclique	80	(generalized) quadrangle	84
colour class	80	geometric graph	52
colourable, k-	80	Gewirtz graph	42
colouring	80	girth	79
complement of a graph	79	graph	79
complement of an inc. structure	83	graph, 40-	76
complete graph	79	Higman-Sims technique	10
complete $\gamma$ -partite graph	80	Hoffman-Singleton graph	42
component	79	imprimitive str. reg. graph	37,82
connected graph	79	incidence	82
cycle, n-	81	incidence graph	46,83
degenerate design	83	incidence matrix	80,82
degree	81	incidence structure	82
derived design	83	induced configuration	58
design	83	induced subgraph	80
design, t-	83	independent set of vertices	80
design, t - (v,k, $\lambda$ )	83	interlacing eigenvalues	7
diameter	79	intersection numbers of a design	31
disconnected graph	79	intersection numbers of a graph	51
disjoint union of graphs	79	joint vertices	79
distance in a graph	79	lattice graph	82
distance in a generalized n-gon	52	length of a circuit	79

length of a path in a graph	79	quasi-symmetric 2-design	35
length of a path in a gen. n-gon	52	regular graph	81
line graph	67,84	residual design	83
linked 2 - (16,6,2) designs	45	Schläfli graph	26
new block	68	Shrikhande graph	fly-leaf
new point	68	singular values	5
null graph	79	special block	69
order of a generalized n-gon	50	strongly regular graph	81
oval	70,78	subconstituent	82
parameters of a t-design	83	subdesign	83
parameters of a partial geometry	84	subgraph	80
parameters of a str. reg. graph	81	substructure	82
partial geometry	84	symmetric 2-design	83
partial subgeometry	84	thick generalized n-gon	50
path in a generalized n-gon	52	tight interlacing	7
path in a graph	79	triangular graph	82
Petersen graph	82	type I, II block	65
point	82	valency	81
point graph	52,84	vertex	79
primitive str. reg. graph	37,82	void graph	79
pseudo-geometric graph	52		

## OTHER TITLES IN THE SERIES MATHEMATICAL CENTRE TRACTS

A leaflet containing an order-form and abstracts of all publications mentioned below is available at the Mathematisch Centrum, Tweede Boerhaavestraat 49, Amsterdam, 1091AL The Netherlands. Orders should be sent to the same address.

---

- MCT 1 T. VAN DER WALT, *Fixed and almost fixed points*, 1963. ISBN 90 6196 002 9.
- MCT 2 A.R. BLOEMENA, *Sampling from a graph*, 1964. ISBN 90 6196 003 7.
- MCT 3 G. DE LEVE, *Generalized Markovian decision processes, part I: Model and method*, 1964. ISBN 90 6196 004 5.
- MCT 4 G. DE LEVE, *Generalized Markovian decision processes, part II: Probabilistic background*, 1964. ISBN 90 6196 005 3.
- MCT 5 G. DE LEVE, H.C. TIJMS & P.J. WEEDA, *Generalized Markovian decision processes, Applications*, 1970. ISBN 90 6196 051 7.
- MCT 6 M.A. MAURICE, *Compact ordered spaces*, 1964. ISBN 90 6196 006 1.
- MCT 7 W.R. VAN ZWET, *Convex transformations of random variables*, 1964. ISBN 90 6196 007 X.
- MCT 8 J.A. ZONNEVELD, *Automatic numerical integration*, 1964. ISBN 90 6196 008 8.
- MCT 9 P.C. BAAYEN, *Universal morphisms*, 1964. ISBN 90 6196 009 6.
- MCT 10 E.M. DE JAGER, *Applications of distributions in mathematical physics*, 1964. ISBN 90 6196 010 X.
- MCT 11 A.B. PAALMAN-DE MIRANDA, *Topological semigroups*, 1964. ISBN 90 6196 011 8.
- MCT 12 J.A.TH.M. VAN BERCKEL, H. BRANDT CORSTIUS, R.J. MOKKEN & A. VAN WIJNGAARDEN, *Formal properties of newspaper Dutch*, 1965. ISBN 90 6196 013 4.
- MCT 13 H.A. LAUWERIER, *Asymptotic expansions*, 1966, out of print; replaced by MCT 54 and 67.
- MCT 14 H.A. LAUWERIER, *Calculus of variations in mathematical physics*, 1966. ISBN 90 6196 020 7.
- MCT 15 R. DOORNBOS, *Slippage tests*, 1966. ISBN 90 6196 021 5.
- MCT 16 J.W. DE BAKKER, *Formal definition of programming languages with an application to the definition of ALGOL 60*, 1967. ISBN 90 6196 022 3.
- MCT 17 R.P. VAN DE RIET, *Formula manipulation in ALGOL 60, part 1*, 1968. ISBN 90 6196 025 8.
- MCT 18 R.P. VAN DE RIET, *Formula manipulation in ALGOL 60, part 2*, 1968. ISBN 90 6196 038 X.
- MCT 19 J. VAN DER SLOT, *Some properties related to compactness*, 1968. ISBN 90 6196 026 6.
- MCT 20 P.J. VAN DER HOUWEN, *Finite difference methods for solving partial differential equations*, 1968. ISBN 90 6196 027 4.

- MCT 21 E. WATTEL, *The compactness operator in set theory and topology*, 1968. ISBN 90 6196 028 2.
- MCT 22 T.J. DEKKER, *ALGOL 60 procedures in numerical algebra, part 1*, 1968. ISBN 90 6196 029 0.
- MCT 23 T.J. DEKKER & W. HOFFMANN, *ALGOL 60 procedures in numerical algebra, part 2*, 1968. ISBN 90 6196 030 4.
- MCT 24 J.W. DE BAKKER, *Recursive procedures*, 1971. ISBN 90 6196 060 6.
- MCT 25 E.R. PAERL, *Representations of the Lorentz group and projective geometry*, 1969. ISBN 90 6196 039 8.
- MCT 26 EUROPEAN MEETING 1968, *Selected statistical papers, part I*, 1968. ISBN 90 6196 031 2.
- MCT 27 EUROPEAN MEETING 1968, *Selected statistical papers, part II*, 1969. ISBN 90 6196 040 1.
- MCT 28 J. OOSTERHOFF, *Combination of one-sided statistical tests*, 1969. ISBN 90 6196 041 X.
- MCT 29 J. VERHOEFF, *Error detecting decimal codes*, 1969. ISBN 90 6196 042 8.
- MCT 30 H. BRANDT CORSTIUS, *Excercises in computational linguistics*, 1970. ISBN 90 6196 052 5.
- MCT 31 W. MOLENAAR, *Approximations to the Poisson, binomial and hypergeometric distribution functions*, 1970. ISBN 90 6196 053 3.
- MCT 32 L. DE HAAN, *On regular variation and its application to the weak convergence of sample extremes*, 1970. ISBN 90 6196 054 1.
- MCT 33 F.W. STEUTEL, *Preservation of infinite divisibility under mixing and related topics*, 1970. ISBN 90 6196 061 4.
- MCT 34 I. JUHÁSZ, A. VERBEEK & N.S. KROONENBERG, *Cardinal functions in topology*, 1971. ISBN 90 6196 062 2.
- MCT 35 M.H. VAN EMDEN, *An analysis of complexity*, 1971. ISBN 90 6196 063 0.
- MCT 36 J. GRASMAN, *On the birth of boundary layers*, 1971. ISBN 90 6196 064 9.
- MCT 37 J.W. DE BAKKER, G.A. BLAAUW, A.J.W. DUIJVESTIJN, E.W. DIJKSTRA, P.J. VAN DER HOUWEN, G.A.M. KAMSTEEG-KEMPER, F.E.J. KRUSEMAN ARETZ, W.L. VAN DER POEL, J.P. SCHAAP-KRUSEMAN, M.V. WILKES & G. ZOUTENDIJK, *MC-25 Informatica Symposium*, 1971. ISBN 90 6196 065 7.
- MCT 38 W.A. VERLOREN VAN THEMAAT, *Automatic analysis of Dutch compound words*, 1971. ISBN 90 6196 073 8.
- MCT 39 H. BAVINCK, *Jacobi series and approximation*, 1972. ISBN 90 6196 074 6.
- MCT 40 H.C. TIJMS, *Analysis of (s,S) inventory models*, 1972. ISBN 90 6196 075 4.
- MCT 41 A. VERBEEK, *Superextensions of topological spaces*, 1972. ISBN 90 6196 076 2.
- MCT 42 W. VERVAAT, *Success epochs in Bernoulli trials (with applications in number theory)*, 1972. ISBN 90 6196 077 0.
- MCT 43 F.H. RUYMGAART, *Asymptotic theory of rank tests for independence*, 1973. ISBN 90 6196 081 9.
- MCT 44 H. BART, *Meromorphic operator valued functions*, 1973. ISBN 90 6196 082 7.

- MCT 45 A.A. BALKEMA, *Monotone transformations and limit laws*, 1973.  
ISBN 90 6196 083 5.
- MCT 46 R.P. VAN DE RIET, *ABC ALGOL, A portable language for formula manipulation systems, part 1: The language*, 1973. ISBN 90 6196 084 3.
- MCT 47 R.P. VAN DE RIET, *ABC ALGOL, A portable language for formula manipulation systems, part 2: The compiler*, 1973. ISBN 90 6196 085 1.
- MCT 48 F.E.J. KRUSEMAN ARETZ, P.J.W. TEN HAGEN & H.L. OUDSHOORN, *An ALGOL 60 compiler in ALGOL 60, Text of the MC-compiler for the EL-X8*, 1973. ISBN 90 6196 086 X.
- MCT 49 H. KOK, *Connected orderable spaces*, 1974. ISBN 90 6196 088 6.
- MCT 50 A. VAN WIJNGAARDEN, B.J. MAILLOUX, J.E.L. PECK, C.H.A. KOSTER, M. SINTZOFF, C.H. LINDSEY, L.G.L.T. MEERTENS & R.G. FISHER (Eds), *Revised report on the algorithmic language ALGOL 68*, 1976. ISBN 90 6196 089 4.
- MCT 51 A. HORDIJK, *Dynamic programming and Markov potential theory*, 1974.  
ISBN 90 6196 095 9.
- MCT 52 P.C. BAAYEN (ed.), *Topological structures*, 1974. ISBN 90 6196 096 7.
- MCT 53 M.J. FABER, *Metrizability in generalized ordered spaces*, 1974.  
ISBN 90 6196 097 5.
- MCT 54 H.A. LAUWERIER, *Asymptotic analysis, part 1*, 1974. ISBN 90 6196 098 3.
- MCT 55 M. HALL JR. & J.H. VAN LINT (Eds), *Combinatorics, part 1: Theory of designs, finite geometry and coding theory*, 1974.  
ISBN 90 6196 099 1.
- MCT 56 M. HALL JR. & J.H. VAN LINT (Eds), *Combinatorics, part 2: graph theory, foundations, partitions and combinatorial geometry*, 1974. ISBN 90 6196 100 9.
- MCT 57 M. HALL JR. & J.H. VAN LINT (Eds), *Combinatorics, part 3: Combinatorial group theory*, 1974. ISBN 90 6196 101 7.
- MCT 58 W. ALBERS, *Asymptotic expansions and the deficiency concept in statistics*, 1975. ISBN 90 6196 102 5.
- MCT 59 J.L. MIJNHEER, *Sample path properties of stable processes*, 1975.  
ISBN 90 6196 107 6.
- MCT 60 F. GÖBEL, *Queueing models involving buffers*, 1975. ISBN 90 6196 108 4.
- \* MCT 61 P. VAN EMDE BOAS, *Abstract resource-bound classes, part 1*.  
ISBN 90 6196 109 2.
- \* MCT 62 P. VAN EMDE BOAS, *Abstract resource-bound classes, part 2*.  
ISBN 90 6196 110 6.
- MCT 63 J.W. DE BAKKER (ed.), *Foundations of computer science*, 1975.  
ISBN 90 6196 111 4.
- MCT 64 W.J. DE SCHIPPER, *Symmetric closed categories*, 1975. ISBN 90 6196 112 2.
- MCT 65 J. DE VRIES, *Topological transformation groups 1 A categorical approach*, 1975. ISBN 90 6196 113 0.
- MCT 66 H.G.J. PIJLS, *Locally convex algebras in spectral theory and eigenfunction expansions*, 1976. ISBN 90 6196 114 9.

- \* MCT 67 H.A. LAUWERIER, *Asymptotic analysis, part 2.*  
ISBN 90 6196 119 x.
- MCT 68 P.P.N. DE GROEN, *Singularly perturbed differential operators of second order*, 1976. ISBN 90 6196 120 3.
- MCT 69 J.K. LENSTRA, *Sequencing by enumerative methods*, 1977.  
ISBN 90 6196 125 4.
- MCT 70 W.P. DE ROEVER JR., *Recursive program schemes: semantics and proof theory*, 1976. ISBN 90 6196 127 0.
- MCT 71 J.A.E.E. VAN NUNEN, *Contracting Markov decision processes*, 1976.  
ISBN 90 6196 129 7.
- MCT 72 J.K.M. JANSEN, *Simple periodic and nonperiodic Lamé functions and their applications in the theory of conical waveguides*, 1977.  
ISBN 90 6196 130 0.
- MCT 73 D.M.R. LEIVANT, *Absoluteness of intuitionistic logic*, 1979.  
ISBN 90 6196 122 x.
- MCT 74 H.J.J. TE RIELE, *A theoretical and computational study of generalized aliquot sequences*, 1976. ISBN 90 6196 131 9.
- MCT 75 A.E. BROUWER, *Treelike spaces and related connected topological spaces*, 1977. ISBN 90 6196 132 7.
- MCT 76 M. REM, *Associations and the closure statement*, 1976. ISBN 90 6196 135 1.
- MCT 77 W.C.M. KALLENBERG, *Asymptotic optimality of likelihood ratio tests in exponential families*, 1977 ISBN 90 6196 134 3.
- MCT 78 E. DE JONGE, A.C.M. VAN ROOIJ, *Introduction to Riesz spaces*, 1977.  
ISBN 90 6196 133 5.
- MCT 79 M.C.A. VAN ZUIJLEN, *Empirical distributions and rankstatistics*, 1977.  
ISBN 90 6196 145 9.
- MCT 80 P.W. HEMKER, *A numerical study of stiff two-point boundary problems*, 1977. ISBN 90 6196 146 7.
- MCT 81 K.R. APT & J.W. DE BAKKER (Eds), *Foundations of computer science II*,  
part 1, 1976. ISBN 90 6196 140 8.
- MCT 82 K.R. APT & J.W. DE BAKKER (Eds), *Foundations of computer science II*,  
part 2, 1976. ISBN 90 6196 141 6.
- MCT 83 L.S. VAN BENTEM JUTTING, *Checking Landau's "Grundlagen" in the AUTOMATH system*, 1979 ISBN 90 6196 147 5.
- MCT 84 H.L.L. BUSARD, *The translation of the elements of Euclid from the Arabic into Latin by Hermann of Carinthia (?) books vii-xii*, 1977.  
ISBN 90 6196 148 3.
- MCT 85 J. VAN MILL, *Supercompactness and Wallman spaces*, 1977.  
ISBN 90 6196 151 3.
- MCT 86 S.G. VAN DER MEULEN & M. VELDHORST, *Torrix I*, 1978.  
ISBN 90 6196 152 1.
- \* MCT 87 S.G. VAN DER MEULEN & M. VELDHORST, *Torrix II*,  
ISBN 90 6196 153 x.
- MCT 88 A. SCHRIJVER, *Matroids and linking systems*, 1977.  
ISBN 90 6196 154 8.



- MCT 89 J.W. DE ROEVER, *Complex Fourier transformation and analytic functionals with unbounded carriers*, 1978.  
ISBN 90 6196 155 6.
- \* MCT 90 L.P.J. GROENEWEGEN, *Characterization of optimal strategies in dynamic games*, . ISBN 90 6196 156 4.
- MCT 91 J.M. GEYSEL, *Transcendence in fields of positive characteristic*, 1979. ISBN 90 6196 157 2.
- MCT 92 P.J. WEEDA, *Finite generalized Markov programming*, 1979.  
ISBN 90 6196 158 0.
- MCT 93 H.C. TIJMS (ed.) & J. WESSELS (ed.), *Markov decision theory*, 1977.  
ISBN 90 6196 160 2.
- MCT 94 A. BIJLSMA, *Simultaneous approximations in transcendental number theory*, 1978 . ISBN 90 6196 162 9.
- MCT 95 K.M. VAN HEE, *Bayesian control of Markov chains*, 1978 .  
ISBN 90 6196 163 7.
- \* MCT 96 P.M.B. VITÁNYI, *Lindenmayer systems: structure, languages, and growth functions*, . ISBN 90 6196 164 5.
- \* MCT 97 A. FEDERGRUEN, *Markovian control problems; functional equations and algorithms*, . ISBN 90 6196 165 3.
- MCT 98 R. GEEL, *Singular perturbations of hyperbolic type*, 1978.  
ISBN 90 6196 166 1
- MCT 99 J.K. LENSTRA, A.H.G. RINNOOY KAN & P. VAN EMDE BOAS, *Interfaces between computer science and operations research*, 1978.  
ISBN 90 6196 170 X.
- MCT 100 P.C. BAAYEN, D. VAN DULST & J. OOSTERHOFF (Eds), *Proceedings bicentennial congress of the Wiskundig Genootschap, part 1*, 1979.  
ISBN 90 6196 168 8.
- MCT 101 P.C. BAAYEN, D. VAN DULST & J. OOSTERHOFF (Eds), *Proceedings bicentennial congress of the Wiskundig Genootschap, part 2*, 1979.  
ISBN 90 9196 169 6.
- MCT 102 D. VAN DULST, *Reflexive and superreflexive Banach spaces*, 1978.  
ISBN 90 6196 171 8.
- MCT 103 K. VAN HARN, *Classifying infinitely divisible distributions by functional equations*, 1978 . ISBN 90 6196 172 6.
- MCT 104 J.M. VAN WOUWE, *Go-spaces and generalizations of metrizability*, 1979.  
ISBN 90 6196 173 4.
- \* MCT 105 R. HELMERS, *Edgeworth expansions for linear combinations of order statistics*, . ISBN 90 6196 174 2.
- MCT 106 A. SCHRIJVER (Ed.), *Packing and covering in combinatorics*, 1979.  
ISBN 90 6196 180 7.
- MCT 107 C. DEN HEIJER, *The numerical solution of nonlinear operator equations by imbedding methods*, 1979. ISBN 90 6196 175 0.
- MCT 108 J.W. DE BAKKER & J. VAN LEEUWEN (Eds), *Foundations of computer science III, part 1*, 1979. ISBN 90 6196 176 9.

- MCT 109 J.W. DE BAKKER & J. VAN LEEUWEN (Eds), *Foundations of computer science III*, part 2, 1979. ISBN 90 6196 177 7.
- MCT 110 J.C. VAN VLIET, *ALGOL 68 transput*, part I: *Historical Review and Discussion of the Implementation Model*, 1979. ISBN 90 6196 178 5.
- MCT 111 J.C. VAN VLIET, *ALGOL 68 transput*, part II: *An implementation model*, 1979. ISBN 90 6196 179 3.
- MCT 112 H.C.P. BERBEE, *Random walks with stationary increments and Renewal theory*, 1979. ISBN 90 6196 182 3.
- MCT 113 T.A.B. SNIJDERS, *Asymptotic optimality theory for testing problems with restricted alternatives*, 1979. ISBN 90 6196 183 1.
- MCT 114 A.J.E.M. JANSSEN, *Application of the Wigner distribution to harmonic analysis of generalized stochastic processes*, 1979. ISBN 90 6196 184 x.
- MCT 115 P.C. BAAYEN & J. VAN MILL (Eds), *Topological Structures II*, part 1, 1979. ISBN 90 6196 185 5.
- MCT 116 P.C. BAAYEN & J. VAN MILL (Eds), *Topological Structures II*, part 2, 1979. ISBN 90 6196 186 6.
- MCT 117 P.J.M. KALLENBERG, *Branching processes with continuous state space*, 1979. ISBN 90 6196 188 2.
- MCT 118 P. GROENEBOOM, *Large deviations and Asymptotic efficiencies*, 1980. ISBN 90 6196 190 4.
- MCT 119 F. PETERS, *Sparse matrices and substructures*, 1980. ISBN 90 6196 192 0.
- MCT 120 W.P.M. DE RUYTER, *On the Asymptotic Analysis of Large Scale Ocean Circulation*, 1980. ISBN 90 6196 192 9.
- MCT 121 W.H. HAEMERS, *Eigenvalue techniques in design and graph theory*, 1980. ISBN 90 6196 194 7.

AN ASTERISK BEFORE THE NUMBER MEANS "TO APPEAR"