

MATHEMATICAL CENTRE TRACTS

4



MATHEMATICAL CENTRE TRACTS

4

**GENERALIZED MARKOVIAN  
DECISION PROCESSES**

PART II

PROBABILISTIC BACKGROUND

BY

G. DE LEVE

2<sup>nd</sup> Edition

MATHEMATISCH CENTRUM AMSTERDAM

1970

EERSTE DRUK IN 1964

## CONTENTS

### PART II

	page
Preface to the second edition	I
CHAPTER 1 THE FUNDAMENTAL STOCHASTIC PROCESS	1
1. General properties	1
2. Random losses	25
3. Stationary strong Markov processes	35
4. Stationary Markov processes and random losses	40
CHAPTER 2 THE DECISION PROCESS	59
1. The basic probability space	59
2. The probabilistic foundation of the decision process	73
3. Properties of the decision process	82
4. A new foundation of the decision process	109
5. Stationary strong Markovian decision processes	114
References	117
Errata and addenda	118
List of symbols	130



Preface to the second edition,

This volume reports on the results obtained in the study of the probabilistic foundation of the Markovian decision processes. The first printing of this book was defaced by a rather large number of printing errors and obscurities. Since a complete revision was out of question the text has been improved by adding a list of addenda and errata and a list of symbols at the end of the book.

The author is very grateful to Mr. A. Hordijk, who read the first printing, suggested improvements, made a list of symbols and posed questions which are still difficult to answer. Some of his comments are included in the list of addenda and errata.

Amsterdam, 1969

dL





## CHAPTER 1

### The fundamental stochastic process

#### 1. General properties

In this chapter we shall consider a class of stochastic processes with a common state space  $X^*$ .

The state space  $X^*$  with points  $x$  is an  $M$ -dimensional Borel set. Since  $X^*$  is also the parameter set of the class of stochastic processes considered, we denote the latter by  $\{S_x^* ; x \in X^*\}$ .

The stochastic processes  $S_x^*$  are defined by means of the following tools:

- 1) the state space  $X^*$  with points  $x$ ;
- 2) a space  $\Omega^*$  with points  $\omega$  ;
- 3) a family of  $\omega$ -functions  $\{x_t^*(\omega) ; t \in [0, \infty)\}$ , defined on  $\Omega^*$ , such that for each  $t \in [0, \infty)$  the  $\omega$ -function  $x_t^*(\omega)$  maps  $\Omega^*$  into  $X^*$ ;
- 4) the  $\sigma$ -field  $G^*$  of  $M$ -dimensional Borel sets in  $X^*$ ;
- 5) the smallest  $\sigma$ -field  $H^*$  with respect to which the  $\omega$ -functions  $\{x_t^*(\omega) ; t \in [0, \infty)\}$  are measurable.
- 6) the function  $P^*[K; x]$  of sets  $K \in H^*$  and points  $x \in X^*$ , satisfying the properties:
  - a) for each  $x \in X^*$ , the set function  $P^*[K; x]$  assigns a probability measure to the sets  $K \in H^*$ ;
  - b) for each  $K \in H^*$ , the  $x$ -function  $P^*[K; x]$  is measurable with respect to  $G^*$ .

A stochastic process  $S_{x_0}^*$  is defined by a family of stochastic variables  $\{\underline{x}_{-t; x_0}^* ; t \in [0, \infty)\}$ , the probability distributions of which are given by

$$\text{Prob} \{ \underline{x}_{-t; x_0}^* \in A \} = P^* [ \Lambda_{t; A}^* ; x_0 ] , \quad (1.1)$$

where  $\Lambda_{t; A}^*$  is defined by

$$\Lambda_{t; A}^* \stackrel{\text{def}}{=} \{ \omega | x_t^*(\omega) \in A \} \quad (1.2)$$

and  $A \in G^*$ .

For each  $x$  the set function  $P^*[K;x]$  represents a probability measure  $P^*$  defined on  $H^*$ . Consequently, for each  $x \in X^*$  we have a triple  $\{\Omega^*, H^*, P^*\}$ . Such a triple is called a probability space. The stochastic processes  $S_x^*$  are defined by means of probability spaces with identical  $\Omega^*$  and  $H^*$ , but with different probability measures  $P^*$ .

The points  $x \in X^*$  and  $\omega \in \Omega^*$  are called the states and the realizations of the stochastic processes respectively. The space  $\Omega^*$  is called the sample space, while the functions  $x_t^*(\omega)$  are named sample functions. Finally, the points  $t \in [0, \infty)$  represent points of time.

Usually in the theory of stochastic processes the  $\sigma$ -field  $H^*$  is completed with all subsets of sets of probability measure 0. In this section, however, we consider various probability measures  $P^*[K;x]$ ; one for each  $x \in X^*$ . So, if we want an "x-free" extension of  $H^*$ , we need to be more selective in completing the  $\sigma$ -field  $H^*$ .

Let  $\Lambda_0^*$  be an  $\omega$ -set with the following properties:

- 1) for each  $\omega \in \Lambda_0^*$ , the t-function  $x_t^*(\omega)$  is continuous from the right;
- 2) in each bounded time interval in  $[0, \infty)$  and for each  $\omega \in \Lambda_0^*$ , the t-function  $x_t^*(\omega)$  has only a finite number of discontinuities.

#### Assumption 1

For each  $x \in X^*$ , a set  $K_x \in H^*$  can be found such that

- a)  $\Lambda_0^* \subset K_x$ ;
- b)  $P^*[K_x;x] = 0$ .

The  $\sigma$ -field  $F^*$  is the smallest  $\sigma$ -field of  $\omega$ -sets that contains  $H^*$  and includes all subsets of  $\Lambda_0^*$ .

The domain of definition of the set function  $P^*[K;x]$  is from now on regarded as extended to  $F^*$ . This extension is unique (cf. [1] p.90).

#### Lemma 1.1

For each  $K \in F^*$ , the x-function  $P^*[K;x]$  is measurable with respect to  $G^*$ .

Proof:

If  $K \in H^*$ , the  $x$ -function  $P^*[K;x]$  is measurable with respect to  $G^*$  (cf. tool 6 of  $S_x^*$ ). Further, if  $K \in \Lambda_0^*$ , we have  $P^*[K;x] = 0$  for all  $x \in X^*$ .

Let  $J$  be the class of  $\omega$ -sets  $K \in F^*$  for which the assertion is true.

We have now proved that

- a)  $J \supset H^*$ ;
- b)  $K \in J$  if  $K \in \Lambda_0^*$ .

The following points can easily be verified:

- c)  $\bar{K} \in J$  if  $K \in J$ ;
- d)  $\bigcup_{j=1}^{\infty} K_j \in J$  if  $K_j \in J$  and if  $K_j \subset K_{j+1}$  ( $j=1,2,\dots$ ).

These properties of  $J$  imply that  $J$  is a  $\sigma$ -field, which contains  $H^*$  and includes all subsets of  $\Lambda_0^*$  ([2] p.599). Hence,  $J=F^*$ .

Now we are in a position to prove the following lemma:

Lemma 1.2.1

If  $I$  is any open time interval and if  $B$  is a closed set in  $X^*$ , then for

$$M_{I;B} \stackrel{\text{def}}{=} \{\omega \mid \forall_{t \in I} x_t^*(\omega) \in B\} \quad (1.3)$$

we have

$$M_{I;B} \in F^*. \quad (1.4)$$

Proof:

Let  $\{t_j; j=1,2,\dots\}$  be the set of all rational numbers in  $[0, \infty)$ . Obviously, we have

$$\bigcap_{t_j \in I} \Lambda_{t_j;B} \supset M_{I;B} \quad (1.5)$$

and thus

$$\bar{\Lambda}_0^* \cap \bigcap_{t_j \in I} \Lambda_{t_j;B} \supset \bar{\Lambda}_0^* \cap M_{I;B}. \quad (1.6)$$

The left hand sides of (1.5) and (1.6) belong to  $F^*$ . We now prove the converse of (1.6).

For each  $t \in I$  a monotone decreasing subsequence  $\{s_m; m=1, 2, \dots\}$  of  $\{t_j; j=1, 2, \dots\}$  can be found such that

$$\lim_{m \rightarrow \infty} s_m = t. \quad (1.7)$$

Hence, if  $\omega \in \overline{\Lambda}_0^* \cap \bigcap_{t_j \in I} \Lambda_{t_j; B}$ , we find (B is closed)

$$x_t^*(\omega) = \lim_{s_m \downarrow t} x_{s_m}^*(\omega) \in B. \quad (1.8)$$

This result can be obtained for each  $t \in I$  and therefore

$$\overline{\Lambda}_0^* \cap \bigcap_{t_j \in I} \Lambda_{t_j; B} \stackrel{1)}{\subset} \overline{\Lambda}_0^* \cap M_{I; B}. \quad (1.9)$$

From (1.6) and (1.9) it follows that

$$\overline{\Lambda}_0^* \cap M_{I; B} = \overline{\Lambda}_0^* \cap \bigcap_{t_j \in I} \Lambda_{t_j; B} \in F^*. \quad (1.10)$$

Since  $F^*$  includes all subsets of  $\overline{\Lambda}_0^*$  we have

$$M_{I; B} \in F^*. \quad (1.4)$$

This terminates the proof.

Let  $i_1$  and  $i_2$  be the left and the right boundary point of an open interval  $I$  respectively. If  $B$  is a closed set in  $X^*$ , we have

$$1) M_{\{i_1\} \cup I; B} = \Lambda_{i_1; B} \cap M_{I; B} \in F^*; \quad (1.11)$$

$$2) M_{I \cup \{i_2\}; B} = M_{I; B} \cap \Lambda_{i_2; B} \in F^*; \quad (1.12)$$

$$3) M_{\{i_1\} \cup I \cup \{i_2\}; B} = \Lambda_{i_1; B} \cap M_{I; B} \cap \Lambda_{i_2; B} \in F^*. \quad (1.13)$$

---

1) The identity (1.10) implies the separability of the stochastic processes  $\{S_x^*; x \in X^*\}$  with respect to the class of all closed sets (cf. [2] p.51).

So we have proved the following lemma:

Lemma 1.2

If  $B$  is a closed set in  $X^*$  and if  $I$  is any interval in  $[0, \infty)$ , then

$$M_{I;B} \in F^*. \quad (1.14)$$

Lemma 1.3

If  $B$  is a closed set in  $X^*$ , there exists a sequence of open sets  $\{B_n; n=1,2,\dots\}$  and a sequence of closed sets  $\{B_n^*; n=1,2,\dots\}$  satisfying:

$$1) B_{n-1} \supset B_n^* \supset B_n \supset B_{n+1}^*; \quad (1.15)$$

$$2) \bigcap_{n=1}^{\infty} B_n = \bigcap_{n=1}^{\infty} B_n^* = B. \quad (1.16)$$

Proof :

Let  $B_n$  and  $B_n^*$  be defined by

$$B_n \stackrel{\text{def}}{=} \{x \mid \exists x_1 \in B; |x-x_1| < \frac{1}{n}\} \quad (1.17)$$

and

$$B_n^* \stackrel{\text{def}}{=} \{x \mid \exists x_1 \in B; |x-x_1| \leq \frac{1}{n}\} \quad (1.18)$$

respectively.

The assertion will now be obvious.

Lemma 1.4.1

If  $B$  is any open set in  $X^*$  and if  $I$  is any interval in  $[0, \infty)$ , for

$$\Lambda_{I;B} \stackrel{\text{def}}{=} \{\omega \mid \exists t \in I \ x_t^{**}(\omega) \in B\} \quad (1.19)$$

we have

$$\Lambda_{I;B} \in F^*. \quad (1.20)$$

Proof :

By lemma 1.2

$$\Lambda_{I;B} = \overline{M_{I;B}} \in F^*. \quad (1.21)$$

Lemma 1.4.2

If  $B$  is any closed set in  $X^*$  and  $I$  is a bounded closed interval

$[i_1, i_2]$  in  $[0, \infty)$ , then

$$\Lambda_{I;B} \in F^*. \quad (1.22)$$

Proof:

We consider the sequence  $T = \{t_j; j=1,2,\dots\}$  consisting of

- a) the points  $i_1$  and  $i_2$ ;
- b) the rational points in  $I$ .

If

$$\omega \in \overline{\Lambda}_0^* \cap \Lambda_{I;B}, \quad (1.23)$$

then

$$\exists_{t \in I} x_t^*(\omega) \in B. \quad (1.24)$$

Since the  $t$ -function  $x_t^*(\omega)$  is continuous from the right, the following statements are true:

$$\forall_k \exists_{t \in I} \exists_m \forall_s \in [t, t + \frac{2}{m}] x_t^*(\omega) \in B \text{ \& } x_s^*(\omega) \in B_k^* \quad (1.25)$$

$$\forall_k \exists_m \forall_n \exists_{t_j \in T} \forall_s \in [t_j, t_j + \frac{1}{m}] x_{t_j}^*(\omega) \in B_n \text{ \& } x_s^*(\omega) \in B_k^*. \quad (1.26)$$

Hence, (1.23) implies

$$\omega \in \overline{\Lambda}_0^* \cap \bigcap_{k=1}^{\infty} \bigcup_{m=1}^{\infty} \bigcap_{n=1}^{\infty} \bigcup_{t_j \in T} \Lambda_{t_j; B_n} \cap M[t_j, t_j + \frac{1}{m}]; B_k^*. \quad (1.27)$$

Thus,

$$\overline{\Lambda}_0^* \cap \Lambda_{I;B} \subset \overline{\Lambda}_0^* \cap \bigcap_{k=1}^{\infty} \bigcup_{m=1}^{\infty} \bigcap_{n=1}^{\infty} \bigcup_{t_j \in T} \Lambda_{t_j; B_n} \cap M[t_j, t_j + \frac{1}{m}]; B_k^*. \quad (1.28)$$

We shall now prove the converse of (1.28).

If (1.27) is true, then

$$\left[ \bigcap_{k=1}^m \{m_k \geq 2m_{k-1}\} \right] \left[ \bigcap_{kn} \{t_{kn} \in T\} \right] \forall_s \in [t_{kn}, t_{kn} + \frac{1}{2^{k-1}}] x_{t_{kn}}^*(\omega) \in B_n \text{ \& } x_s^*(\omega) \in B_k^*. \quad (1.29)$$

For each  $k$  we consider the sequence of points  $\{t_{kn}; n=1,2,\dots\}$ .

If  $n \geq n_0$ , we find

$$x_{t_{kn}}^*(\omega) \in B_{n_0} \subset B_{n_0}^*. \quad (1.30)$$

Since  $I$  is closed and bounded, the points of accumulation  $\{t_k^\alpha; \alpha=1,2,\dots\}$  of  $\{t_{kn}; n=1,2,\dots\}$  belong to  $I$ . If (1.27) is true, one of the following cases will arise:

- a) At least one of the points  $\{t_k^\alpha; \alpha=1,2,\dots\}$ , say  $t_k^1$ , is a point of continuity of the  $t$ -function  $x_t^*(\omega)$ ;
- b) All points  $\{t_k^\alpha; \alpha=1,2,\dots\}$  are points of discontinuity of the  $t$ -function  $x_t^*(\omega)$ .

In case a) it follows from (1.30) that for the point  $\omega$  considered we find

$$x_{t_k^1}^*(\omega) \in \bigcap_{n_0=1}^{\infty} B_{n_0}^* = B. \quad (1.31)$$

Hence,

$$\omega \in \Lambda_{I;B} \cap \overline{\Lambda}_0^*. \quad (1.32)$$

In case b), because of assumption 1, the number of accumulation points must be finite ( $\omega \in \overline{\Lambda}_0^*$ ).

If  $s_k^\alpha$  is defined by

$$s_k^\alpha = t_k^\alpha + \frac{1}{2^{m_k}}; \quad k=1,2,\dots, \quad (1.33)$$

then, since  $\frac{1}{2^{m_k-1}} > 0$ , for each  $\alpha$  an integer  $n_\alpha$  can be found such that

$$s_k^\alpha \in \left[ t_{kn_\alpha}^\alpha, t_{kn_\alpha}^\alpha + \frac{1}{2^{m_k-1}} \right]. \quad (1.34)$$

Consequently,

$$x_{s_k^\alpha}^*(\omega) \in B_k^*; \quad k=1,2,\dots \quad (1.35)$$

If  $t_k'$  denotes the superior of  $\{t_k^\alpha; \alpha=1,2,\dots\}$  and if we consider the sequence  $\{t_k'; k=1,2,\dots\}$ , then we can easily verify that this sequence runs through a finite number of points in  $I$  (points of discontinuity of  $x_t^*(\omega)$ ). So a subsequence of  $\{t_k'; k=1,2,\dots\}$ , say  $\{t_{k(h)}'; h=1,2,\dots\}$ ,

exists that satisfies

$$t'_{k(h)} = t' = \liminf \{t'_k; k=1,2,\dots\}. \quad (1.36)$$

Now let  $s_k$  be defined by

$$s_k = t'_k + \frac{1}{2^m k} ; k=1,2,\dots \quad (1.37)$$

Since  $k(h) \geq h$  and thus  $B_{k(h)}^* \subset B_h^*$ , it follows from (1.35) that

$$x_{s_{k(h)}}^*(\omega) \in B_h^* ; h=1,2,3,\dots \quad (1.38)$$

Further, we can easily verify that  $\lim_{h \rightarrow \infty} s_{k(h)} = t'$  and thus for each  $h_0$  ( $\omega \in \overline{\Lambda}_0^*$ )

$$x_{t'}^*(\omega) = \lim_{h \rightarrow \infty} x_{s_{k(h)}}^*(\omega) \in B_{h_0}^*. \quad (1.39)$$

Consequently,

$$x_{t'}^*(\omega) \in \bigcap_{h_0=1}^{\infty} B_{h_0}^* = B. \quad (1.40)$$

Hence,

$$\omega \in \Lambda_{I;B} \cap \overline{\Lambda}_0^*. \quad (1.41)$$

We have now proved that both case a) and case b) lead to (1.41).

This implies that the converse of (1.28) is also true.

Thus,

$$\overline{\Lambda}_0^* \cap \Lambda_{I;B} = \overline{\Lambda}_0^* \cap \bigcap_{k=1}^{\infty} \bigcup_{m=1}^{\infty} \bigcap_{n=1}^{\infty} \bigcup_{t_j \in I} \Lambda_{t_j; B_n} \cap M_{[t_j, t_j + \frac{1}{m}]; B_k}^* \in F^*. \quad (1.42)$$

and therefore

$$\Lambda_{I;B} \in F^*. \quad (1.43)$$

This ends the proof.

The following lemma can easily be proved (cf. (1.11), (1.12) and (1.13)):

Lemma 1.4.3

If  $I$  is any interval in  $[0, \infty)$  and if  $B$  is a closed set, then

$$\Lambda_{I;B} \in F^*. \quad (1.44)$$



Lemmas 1.2, 1.4.1 and 1.4.3 imply:

Lemma 1.4

If  $I$  is any interval in  $[0, \infty)$  and if  $B$  is either closed or open, then

$$\Lambda_{I;B} \in F^* \quad (1.45)$$

and

$$M_{I;B} = \overline{\Lambda_{I;B}} \in F^*. \quad (1.46)$$

If  $C$  is a closed set in  $X^*$  and if  $\omega$  is a realization of a stochastic process  $S_x^*$ , let  $t(\omega; C)$  be the moment that the system is for the first time in  $C$ . If the initial state of the stochastic process  $S_x^*$  belongs to  $C$ , then  $t(\omega; C) = 0$ .

This point of time can also be defined by

$$t(\omega; C) \stackrel{\text{def}}{=} \begin{cases} \inf \{t | x_t^*(\omega) \in C\}, & \text{if } x_t^*(\omega) \in C \text{ for some finite } t. \\ \infty, & \text{otherwise.} \end{cases} \quad (1.47)$$

Let the  $\omega$ -set  $\Xi_{I;C}$  be defined by

$$\Xi_{I;C} \stackrel{\text{def}}{=} \{\omega | t(\omega; C) \in I\}, \quad (1.48)$$

where  $I$  is an interval in  $[0, \infty)$ .

Lemma 1.5.1

For any interval  $I$  in  $[0, \infty)$  and for each closed set  $C$  we have

$$\Xi_{I;C} \in F^*. \quad (1.49)$$

(Thus,  $t(\omega; C)$  is measurable with respect to  $F^*$ .)

Proof:

Let us consider a closed interval  $I = [i_1, i_2]$ . It can easily be verified that for this choice of  $I$  the  $\omega$ -set  $\overline{\Lambda}_0^* \cap \Xi_{I;C}$  is given by

$$\overline{\Lambda}_0^* \cap \Xi_{I;C} = \overline{\Lambda}_0^* \cap M_{[0, i_1)}; \overline{C} \cap \Lambda_{[i_1, i_2]}; C \in F^*. \quad (1.50)$$

Hence,  $\Xi_{I;C} \in F^*$ .

The proofs for other types of intervals are obvious. This ends the proof.

Let us introduce the  $\omega$ -functions  $x^*(\omega; C)$ , defined by

$$x^*(\omega; C) \stackrel{\text{def}}{=} \begin{cases} x_t^*(\omega; C), & \text{if } t(\omega; C) < \infty. \\ x_0^*(\omega), & \text{if } t(\omega; C) = \infty. \end{cases} \quad (1.51)$$

Note that by this definition the state at the end of the period  $[0, t(\omega; C)]$  is given by  $x^*(\omega; C)$  if  $t(\omega; C) < \infty$ .

The function  $x^*(\omega; C)$  is defined for each  $\omega \in \Omega^*$ .

Let the  $\omega$ -set  $\Delta_{B; C}$  be defined by

$$\Delta_{B; C} \stackrel{\text{def}}{=} \{\omega \mid x^*(\omega; C) \in B\}. \quad (1.52)$$

Lemma 1.5.2

For each  $B \in G^*$  and for each closed set  $C$  we have

$$\Delta_{B; C} \in F^*. \quad (1.53)$$

(Thus,  $x^*(\omega; C)$  is an  $\omega$ -function which is measurable with respect to  $F^*$ .)

Proof:

Let for a fixed  $m$  the  $\omega$ -function  $x_{(m)}^*(\omega)$  be defined by ( $k=1, 2, \dots$ )

$$x_{(m)}^*(\omega) \stackrel{\text{def}}{=} \begin{cases} x_0^*(\omega), & \text{if } \omega \in \Lambda_0^* \cup \overline{\Xi [0, \infty); C} \\ x_{\frac{k}{2^m}}^*(\omega), & \text{if } \omega \in \Xi \left[ \frac{k-1}{2^m}, \frac{k}{2^m} \right); C \cap \overline{\Lambda_0^*}. \end{cases} \quad (1.54)$$

That is

$$x_{(m)}^*(\omega) = \sum_{k=0}^{\infty} \chi_{(m); k}(\omega) \frac{x_k^*(\omega)}{2^m}, \quad (1.55)$$

where

$$\chi_{(m); 0}(\omega) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \omega \in \Lambda_0^* \cup \Xi [0, \infty); C \\ 0, & \text{otherwise} \end{cases} \quad (1.56)$$

and for  $k=1, 2, \dots$

$$X_{(m);k}(\omega) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \omega \in \Xi \left[ \frac{k-1}{2^m}, \frac{k}{2^m} \right); C \cap \overline{\Lambda}_0^* \\ 0, & \text{otherwise} \end{cases} \quad (1.57)$$

Obviously, the  $\omega$ -functions  $\{X_{(m);k}(\omega); k=0,1,\dots\}$  are measurable with respect to  $F^*$ .

Consequently, the  $\omega$ -functions  $x_{(m)}^*(\omega)$  are measurable with respect to  $F^*$ .

It can easily be verified that for  $m \rightarrow \infty$  the sequence of  $\omega$ -functions  $\{x_{(m)}^*(\omega); m=1,2,\dots\}$  converges everywhere to an  $\omega$ -function, let us say  $x_{(\infty)}^*(\omega)$ . From this it follows that the  $\omega$ -function  $x_{(\infty)}^*(\omega)$  is measurable with respect to  $F^*$ .

Since for  $\omega \in \overline{\Lambda}_0^*$  the  $t$ -function  $x_t^*(\omega)$  is continuous from the right, we find for these points

$$x_{(\infty)}^*(\omega) = x^*(\omega; C). \quad (1.58)$$

Consequently, if  $B \in G^*$ ,

$$\Delta_{B;C} \cap \overline{\Lambda}_0^* = \{\omega \mid x_{(\infty)}^*(\omega) \in B\} \cap \overline{\Lambda}_0^* \in F^*. \quad (1.59)$$

Thus,

$$\Delta_{B;C} \in F^*. \quad (1.60)$$

This ends the proof.

Let us assume that the set  $C$  has been chosen in such a way that for each  $x \in X^*$  we have

$$P^* \left[ \Xi_{[0,\infty)}; C; x \right] = 1. \quad (1.61)$$

Since each combination of a measurable  $\omega$ -function and the probability space  $\{\Omega^*; F^*; P^*\}$  generates a stochastic variable, the  $\omega$ -functions  $t(\omega; C)$  and  $x^*(\omega; C)$  lead us to the stochastic variables  $\underline{t}_{C;x}$  and  $\underline{x}_{C;x}^*$ . The probability distributions of these variables are given by

$$\text{Prob}\{\underline{t}_{C;x} \in I\} \stackrel{\text{def}}{=} P^* \left[ \Xi_{I;C}; x \right] \quad (1.62)$$

and

$$\text{Prob}\{\underline{x}_{C;x}^* \in B\} \stackrel{\text{def}}{=} P^*[\Delta_{B;C;x}] \quad (1.63)$$

respectively.

The stochastic variable  $\underline{t}_{C;x}$  represents the length of the time period preceding the moment at which the system first is in C, while  $\underline{x}_{C;x}^*$  denotes the state at the end of this period if (1.61) is true.

Summarizing:

Lemma 1.5

If assumption 1 and condition (1.61) are satisfied, the probability distribution of the length  $\underline{t}_{C;x}$  of the period preceding the moment at which the system first is in C and that of the state  $\underline{x}_{C;x}^*$  at that point of time are defined. They are given by (1.62) and (1.63) respectively.

Let B be a closed set in  $X^*$  and let us define a family of  $\omega$ -functions  $\{x_t^*(\omega; B); t \in [0, \infty)\}$  by

$$x_t^*(\omega; B) \stackrel{\text{def}}{=} \begin{cases} x_{t(\omega; B)+t}^*(\omega), & \text{if } t(\omega; B) < \infty \\ x_t^*(\omega), & \text{if } t(\omega; B) = \infty. \end{cases} \quad (1.64)$$

Lemma 1.6

The  $\omega$ -functions  $\{x_t^*(\omega; B); t \in [0, \infty)\}$  are measurable with respect to  $F^*$ .

Proof:

Let us consider the  $\omega$ -functions  $x_{(m);t}^*(\omega)$ , for  $k=1, 2, \dots$ , defined by

$$x_{(m);t}^*(\omega) \stackrel{\text{def}}{=} \begin{cases} x_t^*(\omega), & \text{if } \omega \in \Lambda_0^* \cup \overline{\Xi}[0, \infty); B \\ x_{t+\frac{k}{2^m}}^*(\omega), & \text{if } \omega \in \overline{\Lambda}_0^* \cap \Xi\left[\frac{k-1}{2^m}, \frac{k}{2^m}\right); B \end{cases} \quad (1.65)$$

Then

$$x_{(m);t}^*(\omega) = \sum_{k=0}^{\infty} \chi_k(\omega) x_{t+\frac{k}{2^m}}^*(\omega), \quad (1.66)$$

where

$$\chi_0(\omega) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \omega \in \Lambda_0^* \cup \overline{\Xi} [0, \infty); B \\ 0, & \text{if } \omega \in \overline{\Lambda_0^*} \cap \Xi [0, \infty); B \end{cases} \quad (1.67)$$

and for  $k=1, 2, \dots$

$$\chi_k(\omega) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \omega \in \overline{\Lambda_0^*} \cap \Xi \left[ \frac{k-1}{2^m}, \frac{k}{2^m} \right); B \\ 0, & \text{otherwise.} \end{cases} \quad (1.68)$$

The remainder of the proof is identical with that of lemma 1.5.2 and is therefore omitted.

This ends the proof.

It follows from (1.64) that for each  $\omega \in \overline{\Lambda_0^*}$

- 1) the  $t$ -function  $x_t^*(\omega; B)$  is continuous from the right;
- 2) in each finite time interval in  $[0, \infty)$  the  $t$ -function  $x_t^*(\omega; B)$  has only a finite number of discontinuities.

If  $B$  and  $C$  are closed sets, let us introduce the  $\omega$ -functions  $t(\omega; B; C)$  and  $x^*(\omega; B; C)$ , defined by

$$t(\omega; B; C) \stackrel{\text{def}}{=} \begin{cases} \inf \{ t | x_t^*(\omega; B) \in C \}, & \text{if } x_t^*(\omega; B) \in C \text{ for some} \\ & \text{finite } t \\ \infty, & \text{otherwise} \end{cases} \quad (1.69)$$

and

$$x^*(\omega; B; C) \stackrel{\text{def}}{=} \begin{cases} x_t^*(\omega; B; C)(\omega; B), & \text{if } t(\omega; B; C) < \infty \\ x_0^*(\omega; B), & \text{if } t(\omega; B; C) = \infty \end{cases} \quad (1.70)$$

respectively.

#### Lemma 1.7

If  $B$  and  $C$  are closed sets in  $X^*$ , the  $\omega$ -functions  $t(\omega; B; C)$  and  $x^*(\omega; B; C)$ , defined by (1.69) and (1.70), are measurable with respect to  $F^*$ .

Proof:

The function  $x_t^*(\omega; B)$  has the same properties as the function  $x_t^*(\omega)$ . Therefore, lemma 1.7 is a direct consequence of lemmas 1.5.1 and 1.5.2.

If  $C$  is a closed set in  $X^*$ , a sequence of open sets  $\{\bar{B}_n; n=1, 2, \dots\}$  can be found such that (cf. lemma 1.3)

$$\bar{B}_n \supset \bar{B}_{n+1} \supset \dots \supset C \quad (1.71)$$

and

$$\bigcap_{n=1}^{\infty} \bar{B}_n = C. \quad (1.72)$$

Consequently, the sequence of closed sets  $\{B_n; n=1, 2, \dots\}$  satisfies

$$B_n \subset B_{n+1} \subset \dots \subset \bar{C} \quad (1.73)$$

and

$$\bigcup_{n=1}^{\infty} B_n = \bar{C}. \quad (1.74)$$

If  $C$  is a closed set in  $X^*$  and if  $\omega$  is a realization of a stochastic process  $S_x^*$ , let  $t(\omega; [C])$  be the moment that the system enters into  $C$  for the first time.

If the initial state of the stochastic process  $S_x^*$  belongs to  $\bar{C}$ , then we obviously have

$$t(\omega; [C]) = t(\omega; C). \quad (1.75)$$

If the initial state is an element of  $C$ , then  $t(\omega; C) = 0$  but the first entry in  $C$  does not occur before a state of  $\bar{C}$  has been assumed.

Let us consider the sequence  $\{t_n(\omega); n=1, 2, \dots\}$ , defined by

$$t_n(\omega) \stackrel{\text{def}}{=} t(\omega; B_n) + t(\omega; B_n; C). \quad (1.76)$$

Obviously, the  $\omega$ -functions  $t_n(\omega)$  are measurable with respect to  $F^*$ . The function  $t_n(\omega)$  represents the time needed for being first in  $B_n$  and then in  $C$ . Consequently, by (1.73)

$$t(\omega; [C]) \leq t_n(\omega). \quad (1.77)$$

Since

$$0 \leq t_n(\omega) \leq t_{n-1}(\omega), \quad (1.78)$$

we can define an  $\omega$ -function  $t_\infty(\omega)$  by

$$t_\infty(\omega) = \begin{cases} \lim_{n \rightarrow \infty} t_n(\omega), & \text{if } t_n(\omega) < \infty \text{ for some } n. \\ \infty, & \text{otherwise.} \end{cases} \quad (1.79)$$

It can easily be verified that  $t_\infty(\omega)$  is measurable with respect to  $F^{**}$ .

It follows from the definition of  $t(\omega; [C])$  that for each  $\delta > 0$  and for some  $t \in [0, t(\omega; [C]) + \delta)$  we have

$$x_t^*(\omega) \in \bar{C}. \quad (1.80)$$

Thus for some  $t \in [0, t(\omega; [C]) + \delta)$  and a sufficient large  $n$

$$x_t^*(\omega) \in B_n. \quad (1.81)$$

Hence, for each  $\delta > 0$  and a sufficient large  $n$

$$t(\omega; [C]) \leq t(\omega; B_n) + t(\omega; B_n; C) + \delta. \quad (1.82)$$

Thus, by (1.77) and (1.82)

$$t(\omega; [C]) = t_\infty(\omega). \quad (1.83)$$

So we have proved the following lemma:

Lemma 1.8.1

The  $\omega$ -function  $t(\omega; [C])$  is measurable with respect to  $F^{**}$ .

Let us introduce the  $\omega$ -function  $x^*(\omega; [C])$ , defined by

$$x^*(\omega; [C]) \stackrel{\text{def}}{=} \begin{cases} x_{t(\omega; [C])}^*(\omega), & \text{if } t(\omega; [C]) < \infty \\ x_0^*(\omega), & \text{if } t(\omega; [C]) = \infty. \end{cases} \quad (1.84)$$

Note, that by this definition the state at the end of the period

$[0, t(\omega; [C])]$  is given by  $x^*(\omega; [C])$  unless  $t(\omega; [C]) = \infty$ .

We shall now demonstrate that the  $\omega$ -function  $x^*(\omega; [C])$  is measurable with respect to  $F^{**}$ . To this end we introduce the sequence of  $\omega$ -

functions  $\{x_{(n)}^*(\omega); n=1,2,\dots\}$ , where

$$x_{(n)}^*(\omega) = \chi(\omega) x_{(n)}^*(\omega; B_n; C) + (1 - \chi(\omega)) x_0^*(\omega) \quad (1.85)$$

with

$$\chi(\omega) = \begin{cases} 1, & \text{if } t(\omega; [C]) < \infty \text{ and } \omega \in \overline{\Lambda}_0^* . \\ 0, & \text{if } t(\omega; [C]) = \infty \text{ or if } \omega \in \Lambda_0^* . \end{cases} \quad (1.86)$$

It can easily be verified that the  $\omega$ -functions  $\{x_n^*(\omega); n=1,2,\dots\}$  are measurable with respect to  $F^*$ .

Since

$$t(\omega; [C]) = \lim_{n \rightarrow \infty} (t(\omega; B_n) + t(\omega; B_n; C)) , \quad (1.82)$$

we find for  $\omega \in \overline{\Lambda}_0^*$

$$x^*(\omega; [C]) = x_{(n)}^*(\omega) . \quad (1.87)$$

This implies that for all  $\omega$  the sequences  $\{x_{(n)}^*(\omega); n=1,2,\dots\}$  converge to a limit, say  $x_\infty^*(\omega)$ .

The  $\omega$ -function  $x_\infty^*(\omega)$  is measurable with respect to  $F^*$ .

Obviously, we have for  $\omega \in \overline{\Lambda}_0^*$

$$x^*(\omega; [C]) = x_\infty^*(\omega) . \quad (1.88)$$

So we have proved the following lemma:

Lemma 1.8.2

The  $\omega$ -function  $x^*(\omega; [C])$  is measurable with respect to  $F^*$ .

Let us introduce the  $\omega$ -sets  $\Xi_{I; [C]}$  and  $\Delta_{B; [C]}$ , defined by

$$\Xi_{I; [C]} \stackrel{\text{def}}{=} \{\omega \mid t(\omega; [C]) \in I\} \quad (1.89)$$

and

$$\Delta_{B; [C]} \stackrel{\text{def}}{=} \{\omega \mid x^*(\omega; [C]) \in B\} \quad (1.90)$$

respectively.

We now assume that the closed set  $C$  is chosen in such a way that for each  $x$



$$P^* [\Xi_{[0, \infty)}; [C]; x] = 1. \quad (1.91)$$

The  $\omega$ -functions  $t(\omega; [C])$  and  $x(\omega; [C])$  together with the probability spaces  $\{\Omega^*; F^*; P^*\}$  generate the stochastic variables  $\underline{t}_{[C]; x}$  and  $\underline{x}_{[C]; x}^*$ ; the corresponding probability distributions are given by

$$\text{Prob} \{ \underline{t}_{[C]; x} \in I \} \stackrel{\text{def}}{=} P^* [\Xi_I; [C]; x] \quad (1.92)$$

and

$$\text{Prob} \{ \underline{x}_{[C]; x}^* \in B \} \stackrel{\text{def}}{=} P^* [\Delta_B; [C]; x] \quad (1.93)$$

respectively.

The stochastic variable  $\underline{t}_{[C]; x}$  represents the length of the time period preceding the first entry in C, while  $\underline{x}_{[C]; x}^*$  denotes the state at the end of that period if (1.91) is true.

Summarizing:

Lemma 1.8

If assumption 1 and condition (1.86) are satisfied, the probability distribution of the length  $\underline{t}_{[C]; x}$  of the period preceding the first entry in C and that of the state  $\underline{x}_{[C]; x}^*$  at that point of time are defined. They are given by (1.92) and (1.93) respectively.

We now consider the  $\omega$ -functions  $x_t^*(\omega; [C])$ , defined by

$$x_t^*(\omega; [C]) \stackrel{\text{def}}{=} \begin{cases} x_{t(\omega; [C]) + t}^*(\omega), & \text{if } t(\omega; [C]) < \infty. \\ x_t^*(\omega), & \text{if } t(\omega; [C]) = \infty. \end{cases} \quad (1.94)$$

Repeating the arguments made in the proof of lemma 1.6 we can prove:

Lemma 1.9

The  $\omega$ -functions  $\{x_t^*(\omega; [C]); t \in [0, \infty)\}$  are measurable with respect to  $F^*$ .

Our future discussions are based on the following assumption:

Assumption 2

If  $x^*(t)$  is any mapping of the time axis  $[0, \infty)$  into the state space  $X^*$ , one and only one point  $\omega$  can be found such that

$$x_t^*(\omega) = x^*(t). \quad (1.95)$$

We introduce the following notation:

$$x_t^*(\omega; t_0) \stackrel{\text{def}}{=} x_{t+t_0}^*(\omega). \quad (1.96)$$

Lemma 1.10

For each  $\omega \in \Omega^*$  and  $t_0 \in [0, \infty)$ , one and only one point  $\omega_1 \in \Omega^*$  can be found such that for  $t \geq 0$

$$x_t^*(\omega_1) = x_t^*(\omega; t_0). \quad (1.97)$$

Proof:

If we write  $x^*(t) = x_t^*(\omega; t_0)$  the assertion follows at once from assumption 2.

The point transformation, defined by (1.97), will be denoted by

$$\omega_1 = T_{t_0}(\omega). \quad (1.98)$$

The point transformation (1.98) also introduces a transformation of  $\omega$ -sets.

The  $\omega_1$ -set  $K_1$  will be called the  $t_0$ -image of  $K$  if

$$K_1 = \{\omega_1 \mid \omega_1 = T_{t_0}(\omega), \omega \in K\}. \quad (1.99)$$

We write:

$$K_1 = T_{t_0}(K). \quad (1.100)$$

Conversely, we can define a set transformation  $K = T_{t_0}^{-1}(K_1)$  by

$$T_{t_0}^{-1}(K_1) \stackrel{\text{def}}{=} \{\omega \mid \omega_1 = T_{t_0}(\omega), \omega_1 \in K_1\}. \quad (1.101)$$

If  $K_1 \in F^*$ , let us introduce the class

$$F_{t_0}^* \stackrel{\text{def}}{=} \{T_{t_0}^{-1}(K_1) \mid K_1 \in F^*\}. \quad (1.102)$$

Lemma 1.11

The set transformation  $K = T_{t_0}^{-1}(K_1)$  generates an isomorphism of  $F^*$  with  $F_{t_0}^*$ .

Proof:

We first prove that  $F_{t_0}^*$  is a  $\sigma$ -field. This can easily be done by verifying the following properties:

a)  $\Omega^* \in F_{t_0}^*$

b) if  $K = T_{t_0}^{-1}(K_1) \in F_{t_0}^*$ , then

$$\bar{K} = \Omega^* - T_{t_0}^{-1}(K_1) = T_{t_0}^{-1}(\Omega^* - K_1) \in F_{t_0}^* ;$$

c) if  $K_i = T_{t_0}^{-1}(K_{i;1}) \in F_{t_0}^*$  ( $i=1,2,\dots$ ), we also have

$$\bigcup_{i=1}^{\infty} K_i = \bigcup_{i=1}^{\infty} T_{t_0}^{-1}(K_{i;1}) = T_{t_0}^{-1}\left(\bigcup_{i=1}^{\infty} K_{i;1}\right) \in F_{t_0}^* .$$

Consequently,  $F_{t_0}^*$  is a  $\sigma$ -field.

Since

$$\begin{aligned} T_{t_0}(T_{t_0}^{-1}(K_1)) &= \{\omega_1 \mid \omega_1 = T_{t_0}(\omega); \omega \in T_{t_0}^{-1}(K_1)\} = \\ &= \{\omega_1 \mid \omega_1 \in K_1\} = K_1, \end{aligned} \quad (1.103)$$

the set transformation  $K = T_{t_0}^{-1}(K_1)$  generates an isomorphism of  $F^*$  with  $F_{t_0}^*$ .

This proves the lemma completely.

Lemma 1.12

The  $\sigma$ -field  $F_{t_0}^*$  satisfies  $F_{t_0}^* \subset F^*$ .

Proof:

Let  $J$  be the class of sets  $K$  belonging to both  $F^*$  and  $F_{t_0}^*$ . Obviously,  $J$  is a  $\sigma$ -field.

Let  $J_1$  be the class of sets  $K_1 \in F^*$  satisfying  $T_{t_0}^{-1}(K_1) \in J$ .

The following properties of  $J_1$  can easily be verified:

- a) If  $K_1 \in J_1$ , then  $T_{t_0}^{-1}(\overline{K_1}) = \overline{T_{t_0}^{-1}(K_1)} \in J$ . Thus,  $\overline{K_1} \in J_1$ ;
- b) If  $K_{1;i} \in J_1$ , then  $T_{t_0}^{-1}(\bigcup_{i=1}^{\infty} K_{1;i}) = \bigcup_{i=1}^{\infty} T_{t_0}^{-1}(K_{1;i}) \in J$ .  
Thus,  $\bigcup_{i=1}^{\infty} K_{1;i} \in J_1$ .

Hence,  $J_1$  is a  $\sigma$ -field.

- c) If  $K_1 \subset \Lambda_o^*$ , then  $T_{t_0}^{-1}(K_1) \subset \Lambda_o^*$  and thus  $T_{t_0}^{-1}(K_1) \in J$ .  
Consequently,  $K_1 \in J_1^o$ ;
- d) If  $K_1 = \Lambda_{t;B}$  with  $B \in G^*$ , then  $T_{t_0}^{-1}(K_1) = \Lambda_{t+t_0;B} \in J$ . Thus,  
 $\Lambda_{t;B} \in J_1$ .

Hence,  $J_1$  is a  $\sigma$ -field that contains the sets  $\Lambda_{t;B}$  and the subsets of  $\Lambda_o^*$ . Thus,  $J_1 = F^*$ . Consequently,  $J = F_{t_0}^* \subset F^*$ .

This ends the proof.

Lemma 1.13

For each  $\omega \in \Omega^*$  and for each closed set  $C \in G^*$  one and only one point  $\omega_1 \in \Omega^*$  can be found such that

$$x_t^*(\omega_1) = x_t^*(\omega; [C]); \quad t \geq 0. \quad (1.104)$$

Proof :

If we write  $x^*(t) = x_t^*(\omega; [C])$ , the assertion follows at once from assumption 2.

The point transformation, defined by (1.104) will be denoted by

$$\omega_1 = T_{[C]}(\omega). \quad (1.105)$$

This point transformation also introduces a transformation of  $\omega$ -sets in  $\Omega^*$ . The  $\omega_1$ -set  $K_1$  will be called the  $[C]$ -image of  $K$  if

$$K_1 = \{\omega_1 \mid \omega_1 = T_{[C]}(\omega); \omega \in K\}. \quad (1.106)$$

Lemma 1.14

If  $\Omega^*_{[C]}$  is defined by

$$\Omega^*_{[C]} \stackrel{\text{def}}{=} T_{[C]}(\Omega^*), \quad (1.107)$$

then

$$\Omega^*_{[C]} = \overline{\Xi}_{(0, \infty); [C]} \cup \Xi_{0; C}. \quad (1.108)$$

Proof:

If we have either  $t(\omega; [C]) = 0$  or  $t(\omega; [C]) = \infty$ , by (1.94) and (1.104) we find  $\omega_1 = T_{[C]}(\omega) = \omega$ . Consequently,  $t(\omega_1; [C]) = t(\omega; [C])$ .

Hence,  $\omega_1 \in \overline{\Xi}_{(0, \infty); [C]}$ .

If  $0 < t(\omega; [C]) < \infty$ , then  $t(\omega_1; C) = 0$ . Therefore,  $\omega_1 \in \Xi_{0; C}$ .

So we have proved that

$$T_{[C]}(\Omega^*) \subset \overline{\Xi}_{(0, \infty); [C]} \cup \Xi_{0; C}. \quad (1.109)$$

We shall now demonstrate that the converse of (1.109) is also true.

If  $\omega' \in \overline{\Xi}_{(0, \infty); [C]}$ , then

$$\omega' = T_{[C]}(\omega') \quad (1.110)$$

and thus

$$\omega' \in T_{[C]}(\Omega^*). \quad (1.111)$$

If  $\omega' \in \Xi_{0; C}$ , if  $\omega''$  satisfies

$$t(\omega''; [C]) > 0 \quad (1.112)$$

and if  $\omega'''$  is given by

$$x_t^*(\omega''') = \begin{cases} x_t^*(\omega''), & \text{if } t < t(\omega''; [C]) \\ x_{t-t(\omega''; [C])}^*(\omega'), & \text{if } t \geq t(\omega''; [C]) \end{cases} \quad (1.113)$$

then

$$\omega' = T_{[C]}(\omega'''). \quad (1.114)$$

Consequently, if  $\omega' \in \Xi_{0;C}$ , we also have

$$\omega' \in T_{[C]}(\Omega^*). \quad (1.115)$$

Summarizing, if  $\omega' \in \overline{\Xi}_{(0,\infty);[C]} \cup \Xi_{0;C}$ , we find

$$\omega' \in T_{[C]}(\Omega^*) \quad (1.116)$$

and thus

$$T_{[C]}(\Omega^*) \supset \overline{\Xi}_{(0,\infty);[C]} \cup \Xi_{0;C}. \quad (1.117)$$

The relations (1.109) and (1.117) imply

$$T_{[C]}(\Omega^*) = \overline{\Xi}_{(0,\infty);[C]} \cup \Xi_{0;C}. \quad (1.118)$$

This ends the proof.

Conversely, we can define a set transformation  $K = T_{[C]}^{-1}(K_1)$  by

$$T_{[C]}^{-1}(K_1) \stackrel{\text{def}}{=} \{\omega \mid \omega_1 = T_{[C]}(\omega); \omega_1 \in K_1\}. \quad (1.119)$$

Let us consider the following classes:

$$F^*[C] \stackrel{\text{def}}{=} \{K_1 \cap \Omega^*_{[C]} \mid K_1 \in F^*\} \quad (1.120)$$

$$F^*_{[C]} \stackrel{\text{def}}{=} \{T_{[C]}^{-1}(K_1) \mid K_1 \in F^*[C]\}. \quad (1.121)$$

Repeating the arguments made in the proofs of lemmas 1.11 and 1.12 we find:

Lemma 1.15

The set transformation  $K = T_{[C]}^{-1}(K_1)$  generates an isomorphism of  $F^*[C]$  with  $F^*_{[C]}$ .

Lemma 1.16

The  $\sigma$ -field  $F^*_{[C]}$  satisfies  $F^*_{[C]} \subset F^*$ .

In future we shall use the point transformations  $\omega_1 = T^j_{[C]}(\omega)$ , defined by

$$T_{[C]}^1(\omega) = T_{[C]}(\omega) \quad (1.122)$$

and

$$T_{[C]}^j(\omega) = T_{[C]}(T_{[C]}^{j-1}(\omega)); \quad j=2,3,\dots \quad (1.123)$$

The point transformation  $\omega_1 = T_{[C]}^j(\omega)$  introduces a transformation of  $\omega$ -sets.

We write

$$K_1 = T_{[C]}^j(K) \quad (1.124)$$

if

$$K_1 = \{\omega_1 \mid \omega_1 = T_{[C]}^j(\omega), \omega \in K\}. \quad (1.125)$$

We obviously have

$$T_{[C]}^1(K) = T_{[C]}(K). \quad (1.126)$$

Conversely, we can define a set transformation  $K = T_{[C]}^{-j}(K_1)$  by

$$T_{[C]}^{-j}(K_1) = T_{[C]}^{-1}(T_{[C]}^{-j+1}(K_1)); \quad j=1,2,\dots \quad (1.127)$$

with

$$T_{[C]}^0(K_1) = K_1. \quad (1.128)$$

By means of lemmas 1.15 and 1.16 we can easily verify:

Lemma 1.17

$$\text{If } K \in F^*, \text{ then } T_{[C]}^j(K) \in F^*. \quad (1.129)$$

$$\text{If } K_1 \in F^*, \text{ then } T_{[C]}^{-j}(K_1) \in F^*. \quad (1.130)$$

Consider a closed set  $C$  in  $X^*$ , satisfying the following assumption:

For each  $j \geq 1$  and for each  $x \in X^*$  we have

$$P^* [T_{[C]}^{-j+1}(\exists [0, \infty); [C]); x] = 1. \quad (1.131)$$

Let us define the  $\omega$ -function  $t_j(\omega; [C])$  by

$$t_j(\omega; [C]) = t(T_{[C]}^{j-1}(\omega); [C]). \quad (1.132)$$

By lemma 1.17 the  $\omega$ -function  $t_j(\omega; [C])$  is measurable with respect to  $F^*$ .

By (1.131) the length of the period between the  $(j-1)^{st}$  and the  $j^{th}$  entry in C is almost surely defined and equal to  $t_j(\omega; [C])$ .

Let us define the  $\omega$ -function  $x_j^*(\omega; [C])$  by

$$x_j^*(\omega; [C]) = x^*(T_{[C]}^{j-1}(\omega); [C]). \quad (1.133)$$

By lemma 1.17 the  $\omega$ -function  $x_j(\omega; [C])$  is measurable with respect to  $F^*$ .

By (1.131) the state at the  $j^{th}$  entry in C is almost surely defined and equal to  $x_j(\omega; [C])$ .

Summarizing:

Lemma 1.18

The  $\omega$ -functions  $t_j(\omega; [C])$  and  $x_j^*(\omega; [C])$  ( $j=1,2,\dots$ ), defined by (1.132) and (1.333) respectively, are measurable with respect to  $F^*$ .

The  $\omega$ -functions  $t_j(\omega; [C])$  and  $x_j^*(\omega; [C])$  together with the probability spaces  $\{\Omega^*; F^*; P^*\}$  generate the stochastic variables  $\underline{t}_{[C]}; x; j$  and  $\underline{x}_{[C]}^*; x; j$ ; the corresponding probability distributions are given by

$$\text{Prob} \{ \underline{t}_{[C]}; x; j \in I \} \stackrel{\text{def}}{=} P^* [ T_{[C]}^{-j+1} (\underline{\varepsilon}_I; [C]); x ] \quad (1.134)$$

and

$$\text{Prob} \{ \underline{x}_{[C]}^*; x; j \in B \} \stackrel{\text{def}}{=} P^* [ T_{[C]}^{-j+1} (\Delta_B; [C]); x ] . \quad (1.135)$$

The stochastic variable  $\underline{t}_{[C]}; x; j$  represents almost surely the length of the period between the  $(j-1)^{st}$  and the  $j^{th}$  entry in C, while  $\underline{x}_{[C]}^*; x; j$  denotes the state at the  $j^{th}$  entry.

So we have proved the following lemma:

Lemma 1.19

If the assumptions 1 and 2 and the condition (1.131) are satisfied, the probability distributions of the lengths  $\underline{t}_{[C]}; x; j$  of the



periods between successive entries in  $C$  and those of the entry states  $\underline{x}^*_{[C]}; x; j$  are defined. They are given by (1.134) and (1.135).

## 2. Random losses

The considerations in this section do not longer start from the assumption that almost all  $t$ -functions  $x_t^*(\omega)$  are continuous from the right. On the other hand we still assume that almost all  $t$ -functions  $x_t^*(\omega)$  have only a finite number of discontinuities in a finite interval. Moreover, the assertions, stated in lemmas 1.5 ff., are supposed to be true. In chapter 2 of this part we shall show that in a special case these lemmas can be proved without the continuity assumption.

A stochastic process  $S_x^*$  is also called a random walk in  $X^*$ . Let us assume that losses are incurred during walks in  $X^*$ . We distinguish the following types of losses:

- a) The "first type" loss is defined by means of a closed set  $A$  and a bounded real valued function  $\gamma_{\text{disc}}(x)$ , which is measurable with respect to  $G^*$ . If the initial state  $x_0^*(\omega)$  of the random walk belongs to  $A$ , a loss  $\gamma_{\text{disc}}(x_0^*(\omega))$  is incurred at the start. Moreover, each entry  $x_j^*(\omega; [A])$  in  $A$  costs  $\gamma_{\text{disc}}(x_j^*(\omega; [A]))$ . In our future discussions we shall make use of a constant  $\gamma_d$ , that satisfies for each  $x \in X^*$

$$|\gamma_{\text{disc}}(x)| \leq \gamma_d < \infty; \quad (1.136)$$

- b) The "second type" loss is defined by means of a bounded continuous function  $\gamma_{\text{cont}}(x)$ . The "second type" loss incurred during the period  $[s_1, s_2)$  is then given by the Riemann integral

$$\int_{s_1}^{s_2} \gamma_{\text{cont}}(x_t^*(\omega)) dt. \quad (1.137)$$

In our future discussions we shall make use of a constant  $\gamma_c$  that satisfies for each  $x \in X^*$

$$|\gamma_{\text{cont}}(x)| \leq \gamma_c < \infty. \quad (1.138)$$

In this section we consider random losses, which will be incurred in the periods  $[0, t_0)$ ,  $[0, t(\omega; B)]$  and  $[0, t(\omega; [C]))$ .

Let the  $\omega$ -functions  $\{\hat{t}_n(\omega; [A]); n=1, 2, \dots\}$  be defined by

$$\hat{t}_n(\omega; [A]) \stackrel{\text{def}}{=} \sum_{j=1}^n t_j(\omega; [A]). \quad (1.139)$$

Note that set A has been used in the definition of the "first type" loss.

We now assume that the closed set A satisfies for each x

$$\lim_{n \rightarrow \infty} P^* [\Xi_{t_0; [A]; n; x}] = 0, \quad (1.140)$$

where

$$\Xi_{t_0; [A]; n} \stackrel{\text{def}}{=} \{\omega \mid \hat{t}_n(\omega; [A]) < t_0\}.$$

Let  $n(\omega; t_0; [A])$  be the number of entries in A during the period  $[0, t_0)$ .

According to this definition

$$n(\omega; t_0; [A]) = n, \text{ if } \hat{t}_n(\omega; [A]) < t_0 \leq \hat{t}_{n+1}(\omega; [A]). \quad (1.141)$$

Obviously, the following lemma is true:

Lemma 1.20

The  $\omega$ -function  $n(\omega; t_0; [A])$  is measurable with respect to  $F^*$ .

We now start our discussion with the losses of the first type.

A real valued  $\omega$ -function  $k_{\text{disc}}(\omega; t_0)$  is defined by

$$k_{\text{disc}}(\omega; t_0) = \begin{cases} \sum_{j=1}^{n(\omega; t_0; [A])} \gamma_{\text{disc}}(x_j^*(\omega; [A])), & \text{if } \omega \in \Xi_{0; \bar{A}} \text{ and } n(\omega; t_0; [A]) < \infty. \\ \gamma_{\text{disc}}(x_0^*(\omega)) + \sum_{j=1}^{n(\omega; t_0; [A])} \gamma_{\text{disc}}(x_j^*(\omega; [A])), & \text{if} \\ \quad \omega \in \Xi_{0; A} \text{ and } n(\omega; t_0; [A]) < \infty. & (1.142) \\ 0, & \text{otherwise.} \end{cases}$$

Lemma 1.21

The  $\omega$ -function  $k_{\text{disc}}(\omega; t_0)$  is measurable with respect to  $F^*$ .

Proof:

Since

- a)  $\gamma_{\text{disc}}(x)$  is Borel measurable with respect to  $G^*$ ,
- b)  $x_j^*(\omega; [A])$  are measurable with respect to  $F^*$  ( $j=1,2,\dots$ ),

we find that both ( $i=1,2,\dots$ )

$$\sum_{j=1}^i \gamma_{\text{disc}}(x_j^*(\omega; [A])) \text{ and } \gamma_{\text{disc}}(x_0^*(\omega))$$

are measurable with respect to  $F^*$ .

Let us introduce the  $\omega$ -functions  $\chi_0(\omega)$  and  $\chi_i(\omega; t_0)$ , defined by

$$\chi_0(\omega) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \omega \in E_{0;A} \\ 0, & \text{otherwise} \end{cases} \quad (1.143)$$

and

$$\chi_i(\omega; t_0) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \hat{t}_i(\omega; [A]) < t_0 \leq \hat{t}_{i+1}(\omega; [A]) \\ 0, & \text{otherwise} \end{cases} \quad (1.144)$$

respectively.

It can easily be verified that the  $\omega$ -functions

$$\left\{ \sum_{i=1}^m \chi_i(\omega; t_0) [\chi_0(\omega) \gamma_{\text{disc}}(x_0^*(\omega)) + \sum_{j=1}^i \gamma_{\text{disc}}(x_j^*(\omega; [A]))] \right\};$$

;  $m=1,2,\dots$  } (1.145)

are measurable with respect to  $F^*$ .

Since the sequence (1.145) converges everywhere to  $k_{\text{disc}}(\omega; t_0)$ , this  $\omega$ -function is measurable with respect to  $F^*$ .

This ends the proof.

By (1.140) and (1.142)  $k_{\text{disc}}(\omega; t_0)$  represents almost surely the "first type" loss incurred in the period  $[0, t_0)$ .

We now consider a closed set B, satisfying

$$\lim_{n \rightarrow \infty} P^* [\Xi_B; [A]; n; x] = 0, \quad (1.146)$$

where

$$\Xi_B; [A]; n \stackrel{\text{def}}{=} \{\omega \mid \hat{t}_n(\omega; [A]) \leq t(\omega; B)\}. \quad (1.147)$$

Let  $n(\omega; B; [A])$  be the number of entries in A during the period  $[0, t(\omega; B)]$ .

According to this definition

$$n(\omega; B; [A]) = n, \text{ if } \hat{t}_n(\omega; [A]) \leq t(\omega; B) < \hat{t}_{n+1}(\omega; [A]). \quad (1.148)$$

The proof of the following lemma is obvious.

Lemma 1.22

The  $\omega$ -function  $n(\omega; B; [A])$  is measurable with respect to  $F^*$ .

A real valued  $\omega$ -function  $k_{\text{disc}}(\omega; B)$  is defined by

$$k_{\text{disc}}(\omega; B) = \begin{cases} \sum_{j=1}^{n(\omega; B; [A])} \gamma_{\text{disc}}(x_j^*(\omega; [A])), & \text{if } \omega \in \Xi_{O; A}^- \text{ and} \\ & n(\omega; B; [A]) < \infty. \\ \sum_{j=1}^{n(\omega; B; [A])} \gamma_{\text{disc}}(x_j^*(\omega; [A])) + \gamma_{\text{disc}}(x_0^*(\omega)), & \text{if} \\ & \omega \in \Xi_{O; A} \text{ and } n(\omega; B; [A]) < \infty. \\ 0, & \text{otherwise.} \end{cases} \quad (1.149)$$

The following lemma can easily be proved (cf. lemma 1.21):

Lemma 1.23

The  $\omega$ -function  $k_{\text{disc}}(\omega; B)$  is measurable with respect to  $F^*$ .

By (1.146) and (1.149)  $k_{\text{disc}}(\omega; B)$  represents almost surely the "first type" loss incurred in the period  $[0, t(\omega; B)]$ .

Next we consider a closed set C, satisfying

$$\lim_{n \rightarrow \infty} P^* [\Xi[C]; [A]; n; x] = 0, \quad (1.150)$$

where

$$\Xi [C]; [A]; n \stackrel{\text{def}}{=} \{\omega | \hat{t}_n(\omega; [A]) < t(\omega; [C])\}. \quad (1.151)$$

Let  $n(\omega; [C]; [A])$  be the number of entries in A during the period  $[0, t(\omega; [C]))$ .

According to this definition

$$n(\omega; [C]; [A]) = n, \text{ if } \hat{t}_n(\omega; [A]) < t(\omega; [C]) \leq \hat{t}_{n+1}(\omega; [A]). \quad (1.152)$$

The proof of the following lemma is obvious.

Lemma 1.24

The  $\omega$ -function  $n(\omega; [C]; [A])$  is measurable with respect to  $F^*$ .

A real valued  $\omega$ -function  $k_{\text{disc}}(\omega; [C])$  is now defined by

$$k_{\text{disc}}(\omega; [C]) \stackrel{\text{def}}{=} \begin{cases} \sum_{j=1}^{n(\omega; [C]; [A])} \gamma_{\text{disc}}(x_j^*(\omega; [A])), & \text{if } n(\omega; [C]; [A]) < \infty \text{ and} \\ & \omega \in \Xi_{O; \bar{A}}. \\ \sum_{j=1}^{n(\omega; [C]; [A])} \gamma_{\text{disc}}(x_j^*(\omega; [A])) + \gamma_{\text{disc}}(x_0^*(\omega)), & \text{if} \\ & n(\omega; [C]; [A]) < \infty \text{ and } \omega \in \Xi_{O; A}. \\ 0, & \text{otherwise.} \end{cases} \quad (1.153)$$

The following lemma can easily be proved (cf. lemma 1.21):

Lemma 1.25

The  $\omega$ -function  $k_{\text{disc}}(\omega; [C])$  is measurable with respect to  $F^*$ .

By (1.150) and (1.153)  $k_{\text{disc}}(\omega; [C])$  represents almost surely the "first type" loss incurred in the period  $[0, t(\omega; [C]))$ .

Let us introduce the  $\omega$ -function  $x_t^{***}(\omega)$ , defined by

$$x_t^{***}(\omega) \stackrel{\text{def}}{=} \begin{cases} \lim_{n \rightarrow \infty} x_{t + \frac{1}{n}}^*(\omega), & \text{if } \omega \in \bar{\Lambda}_O^*. \\ x_0^*(\omega), & \text{if } \omega \in \Lambda_O^*. \end{cases} \quad (1.154)$$

Lemma 1.26

The  $\omega$ -functions  $\{x_t^{***}(\omega); t \in [0, \infty)\}$  are measurable with respect to  $F^*$ .

The  $t$ -functions  $\{x_t^{***}(\omega); \omega \in \Omega^*\}$  are continuous from the right and have almost surely in each finite interval only a finite number of discontinuities.

Proof:

Consider the sequence of  $\omega$ -functions  $\{x_{n;t}^{***}(\omega); n=1,2,\dots\}$ , defined by

$$x_{n;t}^{***}(\omega) \stackrel{\text{def}}{=} \begin{cases} x_{t+\frac{1}{n}}^*(\omega), & \text{if } \omega \in \bar{\Lambda}_0^* \\ x_0^*(\omega), & \text{if } \omega \in \Lambda_0^* \end{cases} \quad (1.155)$$

The  $\omega$ -functions  $x_{n;t}^{***}(\omega)$  are measurable with respect to  $F^*$ . It can easily be verified that the sequence converges everywhere to  $x_t^{***}(\omega)$ . Consequently, the  $\omega$ -function  $x_t^{***}(\omega)$  is measurable with respect to  $F^*$ . Since  $x_t^*(\omega)$  has only a finite number of discontinuities in a finite interval, the second part of the assertion is obvious.

This ends the proof.

Lemma 1.27

The  $\omega$ -functions  $\gamma_{\text{cont}}(x_t^{***}(\omega))$  are measurable with respect to  $F^*$ .

The  $t$ -functions  $\gamma_{\text{cont}}(x_t^{***}(\omega))$  are continuous from the right and have almost surely only a finite number of discontinuities in a finite interval.

Proof:

Since  $\gamma_{\text{cont}}(x)$  is a continuous function, the assertions are immediate.

Lemma 1.28

The Riemann integral

$$k_{\text{cont}}(\omega; s) \stackrel{\text{def}}{=} \int_0^s \gamma_{\text{cont}}(x_t^{***}(\omega)) dt \quad (1.156)$$

exists for each  $s < \infty$  and represents an  $\omega$ -function which is measurable with respect to  $F^*$ .

Proof:

By lemma 1.27 we obviously have

$$\int_0^s \gamma_{\text{cont}}(x_t^{***}(\omega)) dt = \lim_{n \rightarrow \infty} \frac{1}{2^n} \sum_{j=1}^{2^n} \gamma_{\text{cont}}\left(\frac{x_{j/s}^{***}(\omega)}{2^n}\right) \leq s \gamma_c < \infty. \quad (1.157)$$

Consequently,  $k_{\text{cont}}(\omega; s)$  exists and is measurable with respect to  $F^*$ .

This ends the proof.

The  $\omega$ -function  $k_{\text{cont}}(\omega; t_0)$  represents almost surely the "second type" costs incurred in the period  $[0, t_0)$ .

Next we introduce the  $\omega$ -functions  $k_{\text{cont}}(\omega; B)$  and  $k_{\text{cont}}(\omega; [C])$ , defined by

$$k_{\text{cont}}(\omega; B) \stackrel{\text{def}}{=} \begin{cases} k(\omega; t(\omega; B)), & \text{if } t(\omega; B) < \infty \\ 0, & \text{otherwise} \end{cases} \quad (1.158)$$

and

$$k_{\text{cont}}(\omega; [C]) \stackrel{\text{def}}{=} \begin{cases} k(\omega; t(\omega; [C])), & \text{if } t(\omega; [C]) < \infty \\ 0, & \text{otherwise} \end{cases} \quad (1.159)$$

respectively.

Lemma 1.29

The  $\omega$ -functions  $k_{\text{cont}}(\omega; B)$  and  $k_{\text{cont}}(\omega; [C])$  are measurable with respect to  $F^*$ .

Proof:

Let us introduce the  $\omega$ -functions  $\{w_n(\omega); n=1, 2, \dots\}$ , defined for each  $j$  by

$$w_n(\omega) \stackrel{\text{def}}{=} \begin{cases} k(\omega; \frac{j}{n}), & \text{if } \frac{j-1}{n} \leq t(\omega; B) < \frac{j}{n} \\ 0, & \text{otherwise} \end{cases} \quad (1.160)$$

Obviously, the  $\omega$ -functions  $\{w_n(\omega); n=1, 2, \dots\}$  are measurable with respect to  $F^*$ .

The sequence  $\{w_n(\omega); n=1, 2, \dots\}$  converges everywhere to the  $\omega$ -function  $k_{\text{cont}}(\omega; B)$ . Consequently, the  $\omega$ -function  $k_{\text{cont}}(\omega; B)$  is measurable with respect to  $F^*$ .

The proof for  $k_{\text{cont}}(\omega; [C])$  goes along the same lines and is therefore

omitted.

This ends the proof.

If the sets B and C satisfy for each x

$$P^* [\bar{E}_{[0, \infty)}; B; x] = 1 \quad (1.161)$$

and

$$P^* [\bar{E}_{[0, \infty)}; [C]; x] = 1 \quad (1.162)$$

the  $\omega$ -function  $k_{\text{cont}}(\omega; B)$  represents almost surely the "second type" loss incurred in the period  $[0, t(\omega; B)]$ , while with regard to the period  $[0, t(\omega; [C])]$  this loss almost surely is given by  $k_{\text{cont}}(\omega; [C])$ .

Under (1.140), (1.146), (1.150), (1.161) and (1.162) the total costs incurred in the period  $[0, t_0)$ ,  $[0, t(\omega; B)]$  and  $[0, t(\omega; [C])]$  are almost surely given by

$$k(\omega; t_0) \stackrel{\text{def}}{=} k_{\text{disc}}(\omega; t_0) + k_{\text{cont}}(\omega; t_0), \quad (1.163)$$

$$k(\omega; B) \stackrel{\text{def}}{=} k_{\text{disc}}(\omega; B) + k_{\text{cont}}(\omega; B) \quad (1.164)$$

and

$$k(\omega; [C]) \stackrel{\text{def}}{=} k_{\text{disc}}(\omega; [C]) + k_{\text{cont}}(\omega; [C]) \quad (1.165)$$

respectively.

Remark that

$$\text{a) } |k(\omega; t_0)| \leq t_0 \gamma_c + n(\omega; t_0; [A]) \gamma_d; \quad (1.166)$$

$$\text{b) } |k(\omega; B)| \leq t(\omega; B) \gamma_c + n(\omega; B; [A]) \gamma_d; \quad (1.167)$$

$$\text{c) } |k(\omega; [C])| \leq t(\omega; [C]) \gamma_c + n(\omega; [C]; [A]) \gamma_d. \quad (1.168)$$

Obviously, we have:

Lemma 1.30

The  $\omega$ -functions  $k(\omega; t_0)$ ,  $k(\omega; B)$  and  $k(\omega; [C])$  are measurable with respect to  $F^*$ .

The  $\omega$ -functions  $k(\omega; t_0)$ ,  $k(\omega; B)$ ,  $k(\omega; [C])$ ,  $n(\omega; t_0; [A])$ ,  $n(\omega; B; [A])$  and  $n(\omega; [C]; [A])$  together with the probability spaces  $\{\Omega^*; F^*; P^*\}$  generate the stochastic variables  $\underline{k}_{t_0; x}$ ,  $\underline{k}_{B; x}$ ,  $\underline{k}_{[C]; x}$ ,



$\underline{n}_{t_0;x}, \underline{n}_{B;x}$  and  $\underline{n}_{[C];x}$ ; the corresponding probability distributions are given by

$$\text{Prob} \{ \underline{k}_{t_0;x} \in I \} \stackrel{\text{def}}{=} P^* [K_{I;t_0;x}] , \quad (1.169)$$

$$\text{Prob} \{ \underline{k}_{B;x} \in I \} \stackrel{\text{def}}{=} P^* [K_{I;B;x}] , \quad (1.170)$$

$$\text{Prob} \{ \underline{k}_{[C];x} \in I \} \stackrel{\text{def}}{=} P^* [K_{I;[C];x}] , \quad (1.171)$$

$$\text{Prob} \{ \underline{n}_{t_0;x} = n \} \stackrel{\text{def}}{=} P^* [N_{n;t_0;x}] , \quad (1.172)$$

$$\text{Prob} \{ \underline{n}_{B;x} = n \} \stackrel{\text{def}}{=} P^* [N_{n;B;x}] , \quad (1.173)$$

and

$$\text{Prob} \{ \underline{n}_{[C];x} = n \} \stackrel{\text{def}}{=} P^* [N_{n;[C];x}] , \quad (1.174)$$

where

$$K_{I;t_0} \stackrel{\text{def}}{=} \{ \omega \mid k(\omega; t_0) \in I \} , \quad (1.175)$$

$$K_{I;B} \stackrel{\text{def}}{=} \{ \omega \mid k(\omega; B) \in I \} , \quad (1.176)$$

$$K_{I;[C]} \stackrel{\text{def}}{=} \{ \omega \mid k(\omega; [C]) \in I \} , \quad (1.177)$$

$$N_{n;t_0} \stackrel{\text{def}}{=} \{ \omega \mid n(\omega; t_0; [A]) = n \} , \quad (1.178)$$

$$N_{n;B} \stackrel{\text{def}}{=} \{ \omega \mid n(\omega; B; [A]) = n \} , \quad (1.179)$$

$$N_{n;[C]} \stackrel{\text{def}}{=} \{ \omega \mid n(\omega; [C]; [A]) = n \} , \quad (1.180)$$

and  $I$  is any interval in  $(-\infty, +\infty)$ .

So we have proved:

Lemma 1.31

Under (1.140), (1.146), (1.150), (1.161) and (1.162) the probability distributions of the random losses  $\underline{k}_{t_0;x}, \underline{k}_{B;x}$  and  $\underline{k}_{[C];x}$  incurred in the periods  $[0, t_0)$ ,  $[0, t_{B;x}]$  and  $[0, t_{[C];x})$  respectively as well as those of the number of entries  $\underline{n}_{t_0;x}, \underline{n}_{B;x}$  and  $\underline{n}_{[C];x}$  in  $A$  during the same periods are defined. They are given by (1.169) through (1.174).

Finally, let us define the  $\omega$ -functions  $n_j(\omega; t_0; [A])$ ,  $n_j(\omega; [C]; [A])$ ,  $k_j(\omega; t_0)$  and  $k_j(\omega; [C])$  by

$$n_j(\omega; t_0; [A]) = n(T_{(j-1)t_0}(\omega); t_0; [A]) \quad (1.181)$$

$$n_j(\omega; [C]; [A]) = n(T_{[C]}^{j-1}(\omega); [C]; [A]) \quad (1.182)$$

$$k_j(\omega; t_0) = k(T_{(j-1)t_0}(\omega); t_0) \quad (1.183)$$

and

$$k_j(\omega; [C]) = k(T_{[C]}^{j-1}(\omega); [C]). \quad (1.184)$$

By means of lemmas 1.11 and 1.17 we can easily verify that the  $\omega$ -functions  $n_j(\omega; t_0; [A])$ ,  $n_j(\omega; [C]; [A])$ ,  $k_j(\omega; t_0)$  and  $k_j(\omega; [C])$  are measurable with respect to  $F^*$ .

We now assume that for each  $j \geq 1$  and for each  $x$  we have

$$\lim_{n \rightarrow \infty} P^* [T_{(j-1)t_0}^{-1}(\Xi_{t_0; [A]; n}; x) = 0 \quad (1.185)$$

$$\lim_{n \rightarrow \infty} P^* [T_{[C]}^{-j+1}(\Xi_{[C]; [A]; n}; x) = 0 \quad (1.186)$$

and

$$P^* [T_{[C]}^{-j+1}(\Xi_{[0, \infty); [C]}; x) = 1. \quad (1.187)$$

The  $\omega$ -functions  $n_j(\omega; t_0; [A])$  and  $n_j(\omega; [C]; [A])$  represent the number of entries in A during the periods  $[(j-1)t_0, jt_0)$  and  $[T_{(j-1)}^{-1}(\omega; [C]), T_j(\omega; [C]))$  respectively.

By (1.185) the costs incurred in the period  $[(j-1)t_0, jt_0)$  are almost surely given by  $k_j(\omega; t_0)$ . By (1.186) and (1.187) the costs incurred between the  $(j-1)^{\text{st}}$  and the  $j^{\text{th}}$  entry in C are almost surely given by  $k_j(\omega; [C])$ .

The  $\omega$ -functions  $n_j(\omega; t_0; [A])$ ,  $n_j(\omega; [C]; [A])$ ,  $k_j(\omega; t_0)$  and  $k_j(\omega; [C])$  together with the probability spaces  $\{\Omega^*; F^*; P^*\}$  generate the stochastic variables  $\underline{n}_{t_0; x; j}$ ,  $\underline{n}_{[C]; x; j}$ ,  $\underline{k}_{t_0; x; j}$  and  $\underline{k}_{[C]; x; j}$ ; the corresponding probability distributions are given by

$$\text{Prob} \{ \underline{n}_{t_0; x; j} = n \} \stackrel{\text{def}}{=} P^* [T_{(j-1)t_0}^{-1} (N_n; t_0); x] \quad (1.188)$$

$$\text{Prob} \{ \underline{n}_{[C]; x; j} = n \} \stackrel{\text{def}}{=} P^* [T_{[C]}^{-j+1} (N_n; [C]); x] \quad (1.189)$$

$$\text{Prob} \{ \underline{k}_{t_0; x; j} \in I \} \stackrel{\text{def}}{=} P^* [T_{(j-1)t_0}^{-1} (K_I; t_0); x] \quad (1.190)$$

and

$$\text{Prob} \{ \underline{k}_{[C]; x; j} \in I \} \stackrel{\text{def}}{=} P^* [T_{[C]}^{-j+1} (K_I; [C]); x] \quad (1.191)$$

Let the stochastic variables  $\{\hat{t}_{[C]; x; n} ; n=1,2,\dots\}$  be defined by

$$\hat{t}_{[C]; x; n} = \sum_{j=1}^n t_{[C]; x; j} ; j=1,2,\dots \quad (1.192)$$

The stochastic variables  $\underline{n}_{t_0; x; j}$  and  $\underline{n}_{[C]; x; j}$  represent the number of entries in A during the periods  $[(j-1)t_0, jt_0)$  and

$[\hat{t}_{[C]; x; j-1}, \hat{t}_{[C]; x; j})$  respectively.

The stochastic variables  $\underline{k}_{t_0; x; j}$  and  $\underline{k}_{[C]; x; j}$  represent almost surely the costs incurred in the periods  $[(j-1)t_0, jt_0)$  and

$[\hat{t}_{[C]; x; j-1}, \hat{t}_{[C]; x; j})$  respectively.

So we have proved the following lemma:

Lemma 1.32

Under (1.185), (1.186) and (1.187) the probability distributions of  $\underline{n}_{t_0; x; j}$ ,  $\underline{n}_{[C]; x; j}$ ,  $\underline{k}_{t_0; x; j}$  and  $\underline{k}_{[C]; x; j}$  are defined; they are given by (1.188) through (1.191).

3. Stationary strong Markov processes

Let us consider the  $\omega$ -functions  $\hat{x}_t^*(\omega; t_0)$  and  $\hat{x}_t^*(\omega; [C])$  defined by

$$\hat{x}_t^*(\omega; t_0) \stackrel{\text{def}}{=} \begin{cases} x_t^*(\omega), & \text{if } t < t_0 \\ x_{t_0}^*(\omega), & \text{if } t \geq t_0 \end{cases} \quad (1.193)$$

and

$$\hat{x}_t^*(\omega; [C]) \stackrel{\text{def}}{=} \begin{cases} x_t^*(\omega), & \text{if } t < t(\omega; [C]) \\ x_{t(\omega; [C])}^*(\omega; [C]), & \text{if } t \geq t(\omega; [C]) \end{cases} \quad (1.194)$$

We can easily prove the following lemma:

Lemma 1.33

If  $C$  is a closed set in  $X^*$ , the  $\omega$ -functions  $\hat{x}_t^*(\omega; t_0)$  and  $\hat{x}_t^*(\omega; [C])$  are measurable with respect to  $F^*$ .

We now introduce the following notation:

The class of  $\omega$ -sets  $\hat{H}_t^*$  is the smallest  $\sigma$ -field with respect to which the  $\omega$ -functions  $\{\hat{x}_t^*(\omega; t_0); t \in [0, \infty)\}$  are measurable.

The class of  $\omega$ -sets  $\hat{H}_{[C]}^*$  is the smallest  $\sigma$ -field with respect to which the  $\omega$ -functions  $\{\hat{x}_t^*(\omega; [C]); t \in [0, \infty)\}$  are measurable.

The class of  $\omega$ -sets  $\hat{H}_t^*$  is the smallest  $\sigma$ -field with respect to which the  $\omega$ -functions  $\{\hat{x}_t^*(\omega; t_0); t \in [0, \infty)\}$  are measurable.

The class of  $\omega$ -sets  $\hat{H}_{[C]}^*$  is the smallest  $\sigma$ -field with respect to which the  $\omega$ -functions  $\{\hat{x}_t^*(\omega; [C]); t \in [0, \infty)\}$  are measurable.

Let  $F_1^*$  be a  $\sigma$ -field of  $\omega$ -sets in  $\Omega^*$  that satisfies

$$F_1^* \subset F^*. \quad (1.195)$$

Let  $\hat{y}(\omega)$  be a measurable ( $F^*$ ) and integrable  $\omega$ -function satisfying for some  $K \in F_1^*$  and for each  $\Lambda \in F_1^*$

$$P^* [K \cap \Lambda; x] = \int_{\Lambda} P^* [d\omega; x] \hat{y}(\omega). \quad (1.196)$$

Then the conditional probability of  $K$  relative to  $F_1^*$ , denoted by

$$P^* [K; x | F_1^*], \quad (1.197)$$

is defined as any  $\omega$ -function  $y(\omega)$  which is almost surely equal to  $\hat{y}(\omega)$ .

By the Radon-Nicodym theorem ([1], p.132) a family of such  $\omega$ -functions exists of which

- a) each one is measurable with respect to  $F_1^*$ ;
- b) each two are identical except for an  $\omega$ -set of probability measure 0.

Note that the expression (1.197) is an  $\omega$ -function which is measurable with respect to  $F_1^*$ .

The  $\omega$ -function  $P^* [K; x | F_1^*]$  is called a regular conditional probability measure, if

- 1) for each  $\omega \in \Omega^*$  the set function  $P^* [K; x | F_1^*]$  is a probability measure defined on  $F^*$ ;
- 2) for each  $K \in F^*$  the  $\omega$ -function  $P^* [K; x | F_1^*]$  is measurable with respect to  $F_1^*$ .

In this book the probability space  $\{\Omega^*; F^*; P^*\}$  will be called strongly Markovian if and only if

- 1) for each  $t_0 \in [0, \infty)$ , for each  $K \in H_{t_0}^*$  and for each  $x \in X^*$  we have

$$P^* [K; x | \hat{H}_{t_0}^*] = P^* [T_{t_0}(K); x_{t_0}^*(\omega)]; \quad (1.198)$$

- 2) for each  $x \in X^*$ , for each closed set  $C$  in  $X^*$  satisfying

$$P^* [E_{[0, \infty)}; [C]; x] = 1, \quad (1.199)$$

for each  $K \in H_{[C]}^*$  we have

$$P^* [K; x | \hat{H}_{[C]}^*] = P^* [T_{[C]}(K); x^*(\omega; [C])]. \quad (1.200)$$

If  $\{\Omega^*; F^*; P^*\}$  is a strongly Markovian probability space, the basic stochastic process  $S_x^*$  is called a "stationary strong Markov process".

In [1] on p.577 and in [5] on p.91 condition (1.200) is replaced by a more stringent one.

The equations (1.198) and (1.200) are equivalent to

$$P^* [K_1 \cap \Lambda_1; x] = \int_{\Lambda_1} P^* [d\omega; x] P^* [T_{t_0}(K_1); x_{t_0}^*(\omega)] \quad (1.201)$$

and

$$P^* [K_2 \cap \Lambda_2; x] = \int_{\Lambda_2} P^* [d\omega; x] P^* [T_{[C]}(K_2); x^*(\omega; [C])], \quad (1.202)$$

where  $\Lambda_1 \in \hat{H}_{t_0}^*$ ,  $K_1 \in H_{t_0}^*$ ,  $\Lambda_2 \in \hat{H}_{[C]}^*$  and  $K_2 \in H_{[C]}^*$ .

Let the class of  $\omega$ -sets  $\hat{F}_{t_0}^*$  be the smallest  $\sigma$ -field of  $\omega$ -sets containing  $\hat{H}_{t_0}^*$  and including all subsets of  $\Lambda_{t_0}^*$ .

Let the class of  $\omega$ -sets  $\hat{F}_{[C]}^*$  be the smallest  $\sigma$ -field of  $\omega$ -sets containing  $\hat{H}_{[C]}^*$  and including all subsets of  $\Lambda_{[C]}^*$ .

The following lemma can easily be proved:

Lemma 1.34

If the probability space is strongly Markovian, then for each  $x \in X^*$ ,  $t_0 \in [0, \infty)$  and closed set  $C$  satisfying (1.199), we have

$$P^* [K_1; x | \hat{F}_{t_0}^*] = P^* [T_{t_0}(K_1); x_{t_0}^*(\omega)] \quad (1.203)$$

and

$$P^* [K_2; x | \hat{F}_{[C]}^*] = P^* [T_{[C]}(K_2); x^*(\omega; [C])], \quad (1.204)$$

where  $K_1 \in F_{t_0}^*$  and  $K_2 \in F_{[C]}^*$ .

Let  $y_{t_0}(\omega_1)$  and  $y(\omega)$  be two  $\omega$ -functions, satisfying

$$a) \quad 0 \leq y_{t_0}(\omega_1) \leq 1 \quad (1.205)$$

$$b) \quad y(\omega) = y_{t_0}(T_{t_0}(\omega)). \quad (1.206)$$

Lemma 1.35

- 1) If  $y(\omega)$  is measurable with respect to  $F_{t_0}^*$ , then  $y_{t_0}(\omega_1)$  is measurable with respect to  $F^*$ .
- 2) If  $y_{t_0}(\omega_1)$  is measurable with respect to  $F^*$ , then  $y(\omega)$  is measurable with respect to  $F_{t_0}^*$ .
- 3) If  $\Lambda \in \hat{F}_{t_0}^*$ , if  $y(\omega)$  is measurable with respect to  $F_{t_0}^*$  and if  $\{\Omega^*; F^*; P^*\}$  is strongly Markovian, then

$$\int_{\Lambda} P^* [d\omega; x] y(\omega) = \int_{\Lambda} P^* [d\omega; x] \int_{\Omega^*} P^* [d\omega_1; x_{t_0}^*(\omega)] y_{t_0}(\omega_1). \quad (1.207)$$

Proof :

We first consider the cases 1) and 2). If  $M_r$  and  $M_r'$  are defined

by 
$$M_r \stackrel{\text{def}}{=} \{ \omega | y(\omega) \leq r \} \quad (1.208)$$

and 
$$M'_r \stackrel{\text{def}}{=} \{ \omega | y_{t_0}(\omega_1) \leq r \} \quad (1.209)$$

respectively, then we can easily verify that

$$M'_r = T_{t_0}(M_r) \quad (1.210)$$

and

$$M_r = T_{t_0}^{-1}(M'_r). \quad (1.211)$$

The assertions are now a simple consequence of lemma 1.11.

We consider the third case.

Let the sets  $M_{k;m}$  and  $M'_{k;m}$  be defined by

$$M_{k;m} \stackrel{\text{def}}{=} \{ \omega | \frac{k-1}{2^m} \leq y(\omega) < \frac{k}{2^m} \} \quad (1.212)$$

and

$$M'_{k;m} \stackrel{\text{def}}{=} \{ \omega_1 | \frac{k-1}{2^m} \leq y_{t_0}(\omega_1) < \frac{k}{2^m} \} \quad (1.213)$$

respectively.

We can easily verify that

$$M'_{k;m} = T_{t_0}(M_{k;m}) \quad (1.214)$$

and

$$M_{k;m} = T_{t_0}^{-1}(M'_{k;m}). \quad (1.215)$$

Thus,

$$M'_{k;m} \in F^*. \quad (1.216)$$

Moreover, by lemma 1.34

$$\begin{aligned} \int_{\Lambda} P^* [d\omega; x] y(\omega) &= \lim_{n \rightarrow \infty} \sum_{k=1}^{2^m} \frac{k-1}{2^m} P^* [\Lambda \cap M_{k;m}; x] = \\ &= \lim_{m \rightarrow \infty} \sum_{k=1}^{2^m} \frac{k-1}{2^m} \int_{\Lambda} P^* [d\omega; x] P^* [M_{k;m}; x | \hat{F}_{t_0}^*] = \end{aligned}$$

$$\begin{aligned}
&= \lim_{m \rightarrow \infty} \sum_{k=1}^{2^m} \frac{k-1}{2^m} \int_{\Lambda} P^* [d\omega; x] P^* [M'_{k;m}; x_{t_0}^* (\omega)] = \\
&= \int_{\Lambda} P^* [d\omega; x] \left\{ \lim_{m \rightarrow \infty} \sum_{k=1}^{2^m} \frac{k-1}{2^m} P^* [M'_{k;m}; x_{t_0}^* (\omega)] \right\} = \\
&= \int_{\Lambda} P^* [d\omega; x] \int_{\Omega^*} P^* [d\omega_1; x_{t_0}^* (\omega)] y_{t_0} (\omega_1). \quad (1.217)
\end{aligned}$$

This ends the proof.

Finally, let  $y_{[C]}(\omega_1)$  and  $y(\omega)$  be two  $\omega$ -functions, satisfying

$$a) \quad 0 \leq y_{[C]}(\omega_1) \leq 1; \quad (1.218)$$

$$b) \quad y(\omega) = y_{[C]}(T[C](\omega)). \quad (1.219)$$

Using similar arguments as in the proof of lemma 1.35, we can prove:

Lemma 1.36

- 1) If  $y(\omega)$  is measurable with respect to  $F_{[C]}^*$ , then  $y_{[C]}(\omega_1)$  is measurable with respect to  $F^*$ .
- 2) If  $y_{[C]}(\omega)$  is measurable with respect to  $F^*$ , then  $y(\omega)$  is measurable with respect to  $F_{[C]}^*$ .
- 3) If  $\Lambda \in \hat{F}_{[C]}$ , if  $y(\omega)$  is measurable with respect to  $F_{[C]}^*$  and if  $\{\Omega^*; F^*; P^*\}$  is strongly Markovian, then

$$\int_{\Lambda} P^* [d\omega; x] y(\omega) = \int_{\Lambda} P^* [d\omega; x] \int_{\Omega^*} P^* [d\omega_1; x^*(\omega; [C])] y_{[C]}(\omega_1). \quad (1.220)$$

4. Stationary Markov processes and random losses

In this section our discussions are based on the following assumption:

Assumption 3

For each  $x \in X^*$  we have

$$P^* [\Lambda_{0;x}] = 1. \quad (1.221)$$



If  $\omega$  represents a realization of the basic stochastic process, let  $k_T(\omega)$  denote the costs incurred during the period  $[0, T)$ .

We shall show that, under conditions to be mentioned below, the limit

$$\lim_{T \rightarrow \infty} \frac{k_T(\omega)}{T} \quad (1.222)$$

almost surely exists.

Let us define the random variables  $\{\underline{x}_{t_0}^*; x; j; j=1, 2, \dots\}$  by

$$\underline{x}_{t_0}^*; x; j = \underline{x}_{jt_0}^*; x \quad (1.223)$$

If the functions  $\{p_{t_0}^j(B; x); j=1, 2, \dots\}$  are defined by

$$p_{t_0}^j(B; x) \stackrel{\text{def}}{=} P^* [\Lambda_{jt_0}^*; B; x], \quad (1.224)$$

then

$$\text{Prob} \{ \underline{x}_{t_0}^*; x; j \in B \} = p_{t_0}^j(B; x). \quad (1.225)$$

Let us assume that for each  $j \geq 1$ ,  $x \in X^*$ ,  $\Lambda \in \hat{F}_{t_0}^*$  and  $K \in F_{t_0}^*$  the Markov property

$$P^* [\Lambda \cap K; x] = \int_{\Lambda} P^* [d\omega; x] P^* [T_{t_0}(K); x_1^*(\omega; t_0)] \quad (1.226)$$

is true.

This property implies for each  $j$ ,  $x$  and  $B \in G^*$

$$p_{t_0}^j(B; x) = \int_{X^*} p_{t_0}^1(dx_1; x) p_{t_0}^{j-1}(B; x_1); j=1, 2, \dots \quad (1.227)$$

Since the functions  $p_{t_0}^j(B; x)$  are

- for each  $B \in G^*$  and for each  $j \geq 1$  measurable with respect to  $G^*$ ,
- for each  $x \in X^*$  and for each  $j \geq 1$  a probability measure defined on  $G^*$ ,

the relations (1.227) imply that the sequence of states  $\{\underline{x}_{t_0}^*; x; j; j=1, 2, \dots\}$  constitutes a stationary Markov process with a discrete time parameter (cf. [2] p.190 ff). So we have proved:

Lemma 1.37

Under (1.226) the sequence of states  $\{x_{t_0}^*; x; j; j=1,2,\dots\}$  constitutes a stationary Markov process with a discrete time parameter.

Let us make the following assumption:

There is a finite valued measure  $Q(C)$  of sets  $C \in G^*$  with  $Q(X^*) = 0$ , an integer  $k \geq 1$  and a positive  $\eta$ , such that for each  $x \in X^*$  (cf. [2], p. 192)

$$p_{t_0}^k(C; x) \leq 1 - \eta \text{ if } Q(C) \leq \eta. \quad (1.228)$$

This assumption is called the "Doeblin condition". The following lemma can be proved (cf. [2], p.214):

Lemma 1.38

Under (1.226) and (1.228) the function  $p_{t_0}^k(C; x)$ , given by

$$p_{t_0}^k(C; x) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n p_{t_0}^j(C; x), \quad (1.229)$$

defines for each  $x \in X^*$  a stationary absolute probability distribution.

For the meaning of the concepts: "ergodic sets", "cyclically moving subsets" etc., the reader is referred to [2]. In this book, however, we prefer the name "simple ergodic set" to the term "ergodic set". The latter can be mixed up with the set of all ergodic states.

In [2] on p.207 ff. the following lemma is proved:

Lemma 1.39

If the initial state  $x$  belongs to a simple ergodic set  $E_i$  and if the number of cyclically moving subsets of  $E_i$  is  $c_i$ , then, under (1.226) and (1.228), the sequences  $\{p_{t_0}^{nc_i+j}(C; x); n=1,2,\dots\}$  ( $j=1,2,\dots,c_i$ ) converge for  $n \rightarrow \infty$  exponentially fast and uniformly in  $C$  and  $x$  to a limit, denoted by  $p_{t_0}^{\infty c_i+j}(C; x)$ .

We now introduce an  $(M+1)$ -dimensional Cartesian space, which is the product space of  $X^*$  and the line  $\Gamma = (-\infty, +\infty)$ .

Let us consider the  $\omega$ -functions  $y_j(\omega; t_0)$ , given by

$$y_j(\omega; t_0) \stackrel{\text{def}}{=} (x_j(\omega; t_0), n_j(\omega; t_0; [A])); j=1, 2, \dots . \quad (1.230)$$

As we know the  $\omega$ -function  $n_j(\omega; t_0; [A])$  represents the number of entries in  $A$  during the period  $[(j-1)t_0, jt_0)$ .

The  $\omega$ -functions  $y_j(\omega; t_0)$  map  $\Omega^*$  into the product space  $X^* \times \Gamma$ .

Lemma 1.40

The  $\omega$ -functions  $\{y_j(\omega; t_0); j=1, 2, \dots\}$  are measurable with respect to  $F^*$ .

Proof:

Let  $L$  be a linear Borel field of sets in  $\Gamma$ . If  $U_1 \in G^*$  and if  $U_2 \in L$  we have for each  $j$

$$\begin{aligned} \{\omega \mid y_j(\omega; t_0) \in U_1 \times U_2\} &= \\ &= \{\omega \mid x_j(\omega; t_0) \in U_1\} \cap \{\omega \mid n_j(\omega; t_0; [A]) \in U_2\} . \end{aligned} \quad (1.231)$$

Let  $J_k$  be the class of  $(M+1)$ -dimensional Borel sets  $U$ , satisfying

$$\{\omega \mid y_k(\omega; t_0) \in U\} \in F^* . \quad (1.232)$$

So,  $J_k$  contains all  $(M+1)$ -dimensional intervals. In addition, we have

- a) if  $U \in J_k$ , then  $\bar{U} \in J_k$ ;
- b) if  $U^i \in J_k$  ( $i=1, 2, \dots$ ), then  $\bigcup_{i=1}^{\infty} U^i \in J_k$ .

Consequently,  $J_k$  is a  $\sigma$ -field that includes the  $(M+1)$ -dimensional intervals.

Hence,  $J_k$  is the class of  $(M+1)$ -dimensional Borel sets.

This ends the proof.

If  $U$  is an  $(M+1)$ -dimensional Borel set, let the  $\omega$ -set  $O_{k;U}$  be defined by

$$O_{k;U} \stackrel{\text{def}}{=} \{\omega \mid y_k(\omega; t_0) \in U\} . \quad (1.233)$$

Obviously,

$$O_{k;U} \in \Gamma_{(k-1)t_0} \quad (1.234)$$

We now define the set functions  $\{ 'p_{t_0}^k(U;x); k=1,2,\dots \}$  by

$$'p_{t_0}^1(U;x) \stackrel{\text{def}}{=} P^*[O_1;U;x] \quad (1.235)$$

$$'p_{t_0}^k(U;x) \stackrel{\text{def}}{=} \int_{X^*} p_{t_0}^{k-1}(dx_1;x) 'p_{t_0}^1(U;x_1); k=2,3,\dots \quad (1.236)$$

By means of (1.226) we can easily verify that

$$\begin{aligned} P^*[O_{k;U};x] &= \int_{\Omega^*} P^*[d\omega;x] P^*[O_1;U;x_{k-1}^*(\omega;t_0)] = \\ &= \int_{X^*} p_{t_0}^{k-1}(dx_1;x) P^*[O_1;U;x_{k-1}^*(\omega;t_0)] = \\ &= \int_{X^*} p_{t_0}^{k-1}(dx_1;x) 'p_{t_0}^1(U;x_1) = 'p_{t_0}^k(U;x). \end{aligned} \quad (1.237)$$

If  $a \in \Gamma$  and if  $y = (x,a)$ , let  $''p_{t_0}^k(U;y)$  be defined by

$$''p_{t_0}^k(U;y) \stackrel{\text{def}}{=} 'p_{t_0}^k(U;x). \quad (1.238)$$

We can easily verify, that for  $U_1 \in G^*$  we have

$$''p_{t_0}^k(U_1 \times \Gamma;y) = 'p_{t_0}^k(U_1 \times \Gamma;x) = p_{t_0}^k(U_1;x). \quad (1.239)$$

Consequently, (1.237) can be rewritten as follows:

$$''p_{t_0}^k(U;y) = \int_{X^*} ''p_{t_0}^{k-1}(dy_1;y) ''p_{t_0}^1(U;y_1). \quad (1.240)$$

It can easily be proved that

- a) for a given  $(M+1)$ -dimensional Borel set  $U$  the  $y$ -functions  $''p_{t_0}^k(U;y)$  are measurable with respect to the class of all  $(M+1)$ -dimensional Borel sets;

b) for a given  $y$  the set function  $"p_{t_0}^k(U; y)$  is a probability measure defined on the class of all  $(M+1)$ -dimensional Borel sets.

We now consider the stochastic variables  $\{y_{t_0}; x; k; k=1, 2, \dots\}$ , generated by the  $\omega$ -functions  $\{y_k(\omega; t_0); k=1, 2, \dots\}$  and the probability spaces  $\{\Omega^*; F^*; P^*\}$ .

Obviously, for each  $k \geq 1$  and  $x \in X^*$

$$\text{Prob } [y_{t_0}; x; k \in U] = P^* [O_k; U; x] = 'p_{t_0}^k(U; x). \quad (1.241)$$

The relations (1.241), (1.238) and (1.240) imply the following lemma:

Lemma 1.41

Under (1.226) the sequence of stochastic variables  $\{y_{t_0}; x; k; k=1, 2, \dots\}$  constitutes a stationary Markov process with a discrete time parameter.

Let us return to the Markov process  $\{x_{t_0}^*; x; k; k=1, 2, \dots\}$ . If  $x$  belongs to a simple ergodic set  $E_i$  and if  $c_i$  is the number of cyclically moving sets of  $E_i$ , then, according to lemma 1.39, the limits for  $n \rightarrow \infty$  of the sequences  $\{p_{t_0}^{nc_i+j}(U; x); n=1, 2, \dots\}$  converge to  $p_{t_0}^{\infty c_i+j}(U; x)$  exponentially fast and uniformly in  $U \in G^*$  and  $x \in E_i$ .

It follows from (1.239) and (1.240) that

$$\begin{aligned} \lim_{n \rightarrow \infty} "p_{t_0}^{nc_i+j}(U; y) &= \\ &= \lim_{n \rightarrow \infty} \int_{X^*} p_{t_0}^{nc_i+j-1}(dx_1; x) 'p_{t_0}^1(U; x_1) = \\ &= \int_{X^*} p_{t_0}^{\infty c_i+j-1}(dx_1; x) 'p_{t_0}^1(U; x_1). \end{aligned} \quad (1.242)$$

Consequently, the limit exists.

Now let  $"p_{t_0}^{\infty c_i+j}(U; y)$  be defined by

$$"p_{t_0}^{\infty c_i+j}(U; y) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} "p_{t_0}^{nc_i+j}(U; y). \quad (1.243)$$

By means of (1.240) we can easily verify that

$${}''p_{t_0}^{\infty c_i+j}(U;y) = \int_{X^*} {}''p_{t_0}^{\infty c_i+j-1}(dy_1;y) {}''p_{t_0}^1(U;y_1). \quad (1.244)$$

Lemma 1.42

Under (1.226) and (1.228) the sequences  $\{ {}''p_{t_0}^{nc_i+j}(U;y); n=1,2,\dots \}$  converge uniformly in  $y \in E_i \times \Gamma$  and  $U \in G^*(j=1,2,\dots,c_i)$ .

Proof:

Let the  $x$ -set  $B_{hr}^U$  be given by

$$B_{hr}^U \stackrel{\text{def}}{=} \{x \mid \frac{r-1}{2^h} < {}''p_{t_0}^1(U;x) \leq \frac{r}{2^h}\}. \quad (1.245)$$

It follows from (1.242), (1.240) and (1.245) that

$$| {}''p_{t_0}^{\infty c_i+j}(U;y) - \sum_{r=1}^{2^h} \frac{r}{2^h} {}''p_{t_0}^{\infty c_i+j-1}(B_{hr}^U;x) | \leq 2^{-h} \quad (1.246)$$

and

$$| {}''p_{t_0}^{nc_i+j}(U;y) - \sum_{r=1}^{2^h} \frac{r}{2^h} {}''p_{t_0}^{nc_i+j-1}(B_{hr}^U;x) | \leq 2^{-h}. \quad (1.247)$$

Consequently,

$$\begin{aligned} & | {}''p_{t_0}^{\infty c_i+j}(U;y) - {}''p_{t_0}^{nc_i+j}(U;y) | \leq \\ & \leq \sum_{r=1}^{2^h} \frac{r}{2^h} | {}''p_{t_0}^{\infty c_i+j-1}(B_{hr}^U;y) - {}''p_{t_0}^{nc_i+j-1}(B_{hr}^U;y) | + \\ & \quad + 2^{-h+1}. \end{aligned} \quad (1.248)$$

For each  $\eta > 0$  we can find an integer  $h_0$  such that for  $h \geq h_0$  we have  $2^{-h+1} < \frac{\eta}{2}$ . Since the sequences  $\{ {}''p_{t_0}^{nc_i+j-1}(B_{hr}^U;x); n=1,2,\dots \}$  converge uniformly in  $B_{hr}^U$  and  $x$ , an integer  $N_{ij}$  can be found such that for  $n \geq N_{ij}$

$$\sum_{r=1}^{2^h} \frac{r}{2^h} | {}''p_{t_0}^{\infty c_i+j-1}(B_{hr}^U;x) - {}''p_{t_0}^{nc_i+j-1}(B_{hr}^U;x) | \leq \frac{\eta}{2}. \quad (1.249)$$

Thus, for each  $\eta > 0$  an integer  $N_{ij}$  can be found such that uniformly in  $U$  and  $x$  we have for  $n \geq N_{ij}$

$$| {}^{\infty}p_{t_0}^{c_i+j}(U;x) - {}^{nc_i+j}p_{t_0}^{c_i+j}(U;x) | \leq \eta. \quad (1.250)$$

This ends the proof.

Lemma 1.43

If (1.226) and (1.228) hold, there is a finite valued measure  $Q^*(U)$  of  $(M+1)$ -dimensional Borel sets  $U$  with  $Q^*(X^* \times \Gamma) > 0$ , satisfying for each  $\eta > 0$ , each simple ergodic set  $E_i$ , each  $y \in E_i \times \Gamma$ , each  $j$  and some  $k_{ij} < \infty$

$${}^{k_{ij}c_i+j}p_{t_0}^{c_i+j}(U;y) \leq Q^*(U) + \eta. \quad (1.251)$$

Proof:

Suppose that the stochastic process  $\{x_{t_0}^*; x; k; k=1,2,\dots\}$  has  $m$  simple ergodic sets  $E_i$ . For each pair  $(i,j)$  the set functions  $\{ {}^{\infty}p_{t_0}^{c_i+j}(U;y); y \in E_i \times \Gamma \}$  are identical.

We now define  $Q^*(U)$  by

$$Q^*(U) \stackrel{\text{def}}{=} \sum_{i=1}^m \sum_{j=1}^{c_i} {}^{\infty}p_{t_0}^{c_i+j}(U;y_i), \quad (1.252)$$

where  $y_i$  is some point of  $E_i \times \Gamma$ .

The assertion is an immediate consequence of lemma 1.42.

This ends the proof.

The following lemma can be proved (cf. [2], p.207):

Lemma 1.44

If the stochastic process  $\{x_{t_0}^*; x; k; k=1,2,\dots\}$  has  $m$  simple ergodic sets  $E_i$  and if (1.226) and (1.228) hold, then for some  $\rho < 1$

$$1 - p_{t_0}^n \left( \bigcup_{i=1}^m E_i; x \right) \leq \text{const. } \rho^n; n=1,2,\dots \quad (1.253)$$

Now we shall prove that the stochastic process  $\{y_{t_0}; x; k; k=1, 2, \dots\}$  satisfies the Doeblin condition.

Lemma 1.45

If (1.226) and (1.228) hold, there is a finite valued measure  $Q^*(U)$  of  $(M+1)$ -dimensional Borel sets  $U$ , with  $Q^*(X^* \times \Gamma) > 0$ , an integer  $k \geq 1$  and an  $\eta' > 0$  such that for each  $y \in X^* \times \Gamma$

$${}^k p_{t_0}^k(U; y) \leq 1 - \eta', \text{ if } Q^*(U) \leq \eta'. \quad (1.254)$$

Proof:

Let the stochastic process  $\{x_{t_0}^*; x; k; k=1, 2, \dots\}$  have  $m$  simple ergodic sets  $E_i$  and let  $Q^*(U)$  be given by

$$Q^*(U) = \sum_{i=1}^m \sum_{j=1}^{c_i} {}^{\infty c_i + j} p_{t_0}^k(U; y_i), \quad (1.255)$$

where  $y_i$  is some point of  $E_i \times \Gamma$ .

Let  $k_1$  and  $k_2$  be two integers, such that for some positive  $\eta < \frac{1}{2}$ :

a) for each  $x$

$$1 - p_{t_0}^{k_1}(\bigcup_{i=1}^m E_i; x) \leq \eta; \quad (1.256)$$

b)  $k_2 = \max_{i,j} k_{ij} (c_i + 1)$  (cf. lemma 1.43). (1.257)

Obviously, by (1.251), (1.256) and (1.257) we have for each Borel set  $U$  and  $y$

$${}^{k_1+k_2} p_{t_0}^{k_1+k_2}(U; y) \leq \int_m p_{t_0}^{k_1}(dx_1; x) {}^{k_2} p_{t_0}^{k_2}(U; x_1) + \eta \leq Q^*(U) + 2\eta. \quad (1.258)$$

For sets  $U$ , with  $Q^*(U) \leq \frac{1}{2} - \eta$ , we find by means of (1.258)

$${}^{k_1+k_2} p_{t_0}^{k_1+k_2}(U; y) \leq \frac{1}{2} + \eta = 1 - (\frac{1}{2} - \eta). \quad (1.259)$$

Consequently, the triple  $(Q^*, k, \eta')$ , given by



$$Q^*(U) = \sum_{i=1}^m \sum_{j=1}^{c_i} p_{t_0}^{\infty c_i + j}(U; y), \quad (1.260)$$

$$k = k_1 + k_2 \quad (1.261)$$

$$\text{and } \eta' = \frac{1}{2} - \eta, \quad (1.262)$$

satisfies.

This ends the proof.

We now introduce for each  $x$  the assumption:

$$\sum_{n=1}^{\infty} \int_{X^*} p_{t_0}^*(dx_1; x) P^*[\Xi_{t_0}; [A]; n; x_1]^{<\infty} \quad (1.263)$$

If  $x$  is the initial state of the stochastic process  $\{\underline{x}_{t_0}; x; k; k=1, 2, \dots\}$ , the expected number of entries  $\bar{n}_{t_0}; x$  in  $A$  during a period of length  $t_0$  in the steady state is given by

$$\begin{aligned} \bar{n}_{t_0}; x &= \sum_{j=1}^{\infty} \int_{X^*} p_{t_0}^*(dx_1; x) j P^*[N_j; t_0; x_1] = \\ &= \sum_{j=1}^{\infty} \int_{X^*} p_{t_0}^*(dx_1; x) j \{ P^*[\Xi_{t_0}; [A]; j; x_1] + \\ &\quad - P^*[\Xi_{t_0}; [A]; j+1; x_1] \} = \\ &= \sum_{j=1}^{\infty} \int_{X^*} p_{t_0}^*(dx_1; x) P^*[\Xi_{t_0}; [A]; j; x_1]^{<\infty}. \end{aligned} \quad (1.264)$$

Let a function  $f(y)$  be real valued and measurable with respect to the class of all  $(M+1)$ -dimensional Borel sets and let the  $\omega$ -set  $F_I$  be defined by

$$F_I \stackrel{\text{def}}{=} \{ \omega \mid f(y(\omega; t_0)) \in I \}. \quad (1.265)$$

The following lemma can be proved (cf. [2], p.220):

Lemma 1.46

If (1.226) and (1.228) hold and if for each initial state  $x$

$$\int_{X^*} p_{t_0} (dx_1; x) \int_{-\infty}^{+\infty} |f| P^* [F_{df}; x_1] < \infty, \quad (1.266)$$

then for almost all  $\omega$  the limit

$$\lim_{r \rightarrow \infty} \frac{1}{r} \sum_{j=1}^r f(y_j(\omega; t_0)) \quad (1.267)$$

exists and is equal to

$$\int_{X^*} p_{t_0} (dx_1; x) \int_{-\infty}^{+\infty} f P^* [F_{df}; x_1] \quad (1.268)$$

if  $x$  belongs to a simple ergodic set of the stochastic process

$\{\underline{x}_{t_0}^*; x; j; j=1, 2, \dots\}$ .

It follows from (1.230) that  $n_j(\omega; t_0; [A])$  is a measurable function of  $y_j(\omega; t_0)$ . By lemma 1.46 and (1.264) we find:

Lemma 1.47

Under (1.226), (1.228) and (1.263), for almost all  $\omega$  the limit

$$\lim_{r \rightarrow \infty} \frac{1}{r} \sum_{j=1}^r n_j(\omega; t_0; [A]) \quad (1.269)$$

exists and is equal to

$$\sum_{j=1}^n \int_{X^*} p_{t_0} (dx_1; x) P^* [\underline{x}_{t_0}; [A]; j; x_1] \quad (1.270)$$

if  $x$  belongs to a simple ergodic set of the stochastic process

$\{\underline{x}_{t_0}^*; x; j; j=1, 2, \dots\}$ .

We now consider the sequence of  $\omega$ -functions  $\{k_j(\omega; t_0); j=1, 2, \dots\}$ . If (1.185) holds, the  $\omega$ -function  $k_j(\omega; t_0)$  represents the losses incurred in the period  $[(j-1)t_0, jt_0]$ .

Obviously, we have

$$\int_{X^*} p_{t_0} (dx_1; x) \int_{-\infty}^{+\infty} |k| P^* [K_{dk}; t_0; x_1] \leq t_0 \gamma_c + \bar{n}_{t_0; x} \gamma_d < \infty. \quad (1.271)$$

Using similar arguments as above, we can prove the following lemma:

Lemma 1.48

Under (1.226), (1.228) and (1.263), for almost all  $\omega$  the limit

$$\lim_{r \rightarrow \infty} \frac{1}{r} \sum_{j=1}^r k_j(\omega; t_0) \quad (1.272)$$

exists and is equal to

$$\int_{X^*} p_{t_0}(dx_1; x) \int_{-\infty}^{+\infty} kP^* [K_{dk; t_0}; x_1] \quad (1.273)$$

if  $x$  belongs to a simple ergodic set of the stochastic process

$$\{\underline{x}_{t_0}^*; x; j; j=1, 2, \dots\}.$$

As we know the  $(\omega, T)$ -function  $k_T(\omega)$  represents the losses incurred in the period  $[0, T)$ . We now prove:

Lemma 1.49

Under (1.226), (1.228) and (1.263), for almost all  $\omega$  the limit

$$\lim_{T \rightarrow \infty} \frac{k_T(\omega)}{T} \quad (1.274)$$

exists and is equal to

$$\frac{1}{t_0} \int_{X^*} p_{t_0}(dx_1; x) \int_{-\infty}^{+\infty} kP^* [K_{dk; t_0}; x_1] \quad (1.275)$$

if  $x$  belongs to a simple ergodic set of the stochastic process

$$\{\underline{x}_{t_0}^*; x; j; j=1, 2, \dots\}.$$

Proof:

It follows from lemma 1.47 that almost surely

$$\lim_{r \rightarrow \infty} \frac{n_r(\omega; t_0; [A])}{r} = 0. \quad (1.276)$$

Let the numbers  $n(T; t_0)$  be given by

$$n(T; t_0) = \left[ \frac{T}{t_0} \right]^-. \quad (1.277)$$

Obviously, we have almost surely

$$\begin{aligned} & \frac{\sum_{j=1}^{n(T;t_0)} k_j(\omega; t_0) - t_0 \gamma_c - n_{n(T;t_0)+1}(\omega; t_0; [A]) \gamma_d}{(n(T;t_0) + 1) \cdot t_0} \leq \\ & \leq \frac{k_T(\omega)}{T} \leq \frac{\sum_{j=1}^{n(T;t_0)} k_j(\omega; t_0) + t_0 \gamma_c + n_{n(T;t_0)+1}(\omega; t_0; [A]) \gamma_d}{n(T;t_0) \cdot t_0} \end{aligned} \quad (1.278)$$

and thus

$$\begin{aligned} & \frac{\frac{1}{n(T;t_0)} \sum_{j=1}^{n(T;t_0)} k_j(\omega; t_0) - \frac{t_0 \gamma_c}{n(T;t_0)} - \frac{n_{n(T;t_0)+1}(\omega; t_0; [A]) \gamma_d}{n(T;t_0)}}{t_0 \left(1 + \frac{1}{n(T;t_0)}\right)} \leq \\ & \leq \frac{k_T(\omega)}{T} \leq \\ & \leq \frac{\frac{1}{n(T;t_0)} \sum_{j=1}^{n(T;t_0)} k_j(\omega; t_0) + \frac{t_0 \gamma_c}{n(T;t_0)} + \frac{n_{n(T;t_0)+1}(\omega; t_0; [A]) \gamma_d}{n(T;t_0)}}{t_0} \end{aligned} \quad (1.279)$$

If  $T \rightarrow \infty$ , then the assertion is an immediate consequence of lemma 1.48 and the relations (1.276) and (1.279).

This ends the proof.

We shall now show that under certain conditions the limit

$$\lim_{T \rightarrow \infty} \frac{k_T(\omega)}{T} \quad (1.274)$$

can also be expressed in a different form.

To this end we consider the random variables  $\{x^*_{[C];x;j}; j=1,2,\dots\}$  again.

If the functions  $\{p_{[C]}^j(B;x); j=1,2,\dots\}$  are defined by

$$p_{[C]}^j(B;x) \stackrel{\text{def}}{=} P_{[C]}^* [T_{[C]}^{-j+1}(\Delta_{B; [C]}; x)] \quad (1.280)$$

and if for  $j=1,2,\dots$

$$P_{[C]}^* [T_{[C]}^{-j+1}(\Xi_{[0, \infty)}; [C]); x] = 1, \quad (1.281)$$

then

$$\text{Prob} \{ \underline{x}_{[C]}^*; x; j \in B \} = p_{[C]}^j(B;x). \quad (1.282)$$

Let us assume that for each  $j \geq 1$ ,  $x \in X^*$ ,  $\Lambda \in \hat{F}_{[C]}^*$  and  $K \in F_{[C]}^*$  the Markov property

$$P_{[C]}^* [\Lambda \cap K; x] = \int_{\Lambda} P_{[C]}^* [d\omega; x] P_{[C]}^* [T_{[C]}(K); x_1^*(\omega; t_0)] \quad (1.283)$$

is true.

Since

$$T_{[C]} [T_{[C]}^{-j+1}(\Delta_{B; [C]})] = T_{[C]}^{-j+2}(\Delta_{B; [C]}), \quad (1.284)$$

it follows from (1.280) and (1.283) that for  $j=2,3,\dots$

$$p_{[C]}^j(B;x) = \int_{X^*} p_{[C]}^1(dx_1; x) p_{[C]}^{j-1}(B;x_1). \quad (1.285)$$

Since the function  $p_{[C]}^j(B;x)$  is

- a) for each  $B \in G_{[C]}^*$  and for each  $j \geq 1$  measurable with respect to  $G^*$
- b) for each  $x \in X^*$  and for each  $j \geq 1$  a probability measure defined on  $G^*$ ,

the equations (1.285) imply that the sequence of states

$\{\underline{x}_{[C]}^*; x; j; j=1,2,\dots\}$  constitutes a stationary Markov process with a discrete time parameter.

#### Lemma 1.50

Under (1.283), the sequence of states  $\{\underline{x}_{[C]}^*; x; j; j=1,2,\dots\}$  constitutes a stationary Markov process with a discrete time parameter.

Let us make the following assumption:

There is a finite valued measure  $Q(B)$  of sets  $B \in G^*$  with  $Q(X^*) > 0$ , an integer  $k \geq 1$  and a positive  $\eta$ , such that for each  $x \in X^*$  (cf. [2], p. 192)

$$p_{[C]}^k(B; x) \leq 1 - \eta, \text{ if } Q(B) \leq \eta. \quad (1.286)$$

The following lemma can be proved (cf. [2], p.214):

Lemma 1.51

Under (1.283) and (1.286) the function  $p_{[C]}(B; x)$ , given by

$$p_{[C]}(B; x) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n p_{[C]}^j(B; x), \quad (1.287)$$

defines for each  $x \in X^*$  a stationary absolute probability distribution.

We now introduce an  $(M+1)$ -dimensional Cartesian space, which is the product space of  $X^*$  and the line  $\Gamma = [0, \infty)$ .

Let us consider the  $\omega$ -functions  $y_j(\omega; [C])$ , given by

$$y_j(\omega; [C]) \stackrel{\text{def}}{=} (x_j^*(\omega; [C]), t_j(\omega; [C])); j=1, 2, \dots \quad (1.288)$$

If for each  $x$

$$P_{[C]}^* [T^{-j+1}(\Xi_{[0, \infty)}; [C]); x] = 1, \quad (1.289)$$

the  $\omega$ -function  $t_j(\omega; [C])$  represents the length of the period between the  $(j-1)^{\text{st}}$  and the  $j^{\text{th}}$  entry in  $C$ .

The  $\omega$ -functions  $y_k(\omega; [C])$  ( $k=1, 2, \dots$ ) together with the probability spaces  $\{\Omega^*; F^*; P^*\}$  generate the stochastic variables  $\{y_{[C]}; x; k; k=1, 2, \dots\}$ . Using similar arguments as above, we can prove the following lemma:

Lemma 1.52

Under (1.283), (1.286) and (1.289) the stochastic process

$$\{y_{[C]}; x; k; k=1, 2, \dots\}$$

- a) is a stationary Markov process with a discrete time parameter;  
 b) satisfies the Doeblin condition.

Let us assume that for each  $x \in X^*$

$$0 < \int_{X^*} p[C](dx_1; x) \int_0^\infty t P^* [\Xi_{dt}; [C]; x_1] < \infty. \quad (1.290)$$

The following lemma can be proved (cf. lemma 1.48):

Lemma 1.53

Under (1.283), (1.286) and (1.290) for almost all  $\omega$  the limit

$$\lim_{r \rightarrow \infty} \frac{1}{r} \sum_{j=1}^r t_j(\omega; [C]) \quad (1.291)$$

exists and is equal to

$$\int_{X^*} p[C](dx_1; x) \int_0^\infty t P^* [\Xi_{dt}; [C]; x_1] \quad (1.292)$$

if  $x_1$  belongs to a simple ergodic set.

Lemma 1.54

We now consider the  $\omega$ -functions  $\{n_j(\omega; [C]; [A]); j=1, 2, \dots\}$ . The  $\omega$ -function  $n_j(\omega; [C]; [A])$  represents the number of entries in A during the period  $[\hat{t}_{j-1}(\omega; [C]), \hat{t}_j(\omega; [C])]$ .

Next we assume that

$$\sum_{j=1}^\infty \int_{X^*} p[C](dx_1; x) P^* [\Xi[C]; [A]; j; x_1] < \infty. \quad (1.293)$$

If  $x$  is the initial state of the stochastic process  $\{x[C]; x; j; j=1, 2, \dots\}$ , the expected number of entries  $\bar{n}[C]; x$  in A between two successive entries in C in the steady state is then given by

$$\begin{aligned} \bar{n}[C]; x &= \sum_{j=1}^\infty \int_{X^*} p[C](dx_1; x) j P^* [N_j; [C]; x_1] = \\ &= \sum_{j=1}^\infty \int_{X^*} p[C](dx_1; x) j \{P^* [\Xi[C]; [A]; j; x_1] + \end{aligned}$$

$$\begin{aligned}
& - P^* [ \Xi [C]; [A]; j+1; x_1 ] \} = \\
& = \sum_{j=1}^{\infty} \int_{X^*} P[C] (dx_1; x) P^* [ \Xi [C]; [A]; j; x_1 ] < \infty.
\end{aligned} \tag{1.294}$$

We can prove the following lemma (cf. lemma 1.47):

Lemma 1.55

Under (1.283), (1.286) and (1.293) for almost all  $\omega$  the limit

$$\lim_{r \rightarrow \infty} \frac{1}{r} \sum_{j=1}^r n_j(\omega; [C]; [A]) \tag{1.295}$$

exists and is equal to

$$\sum_{j=1}^n \int_{X^*} P[C] (dx_1; x) P^* [ \Xi [C]; [A]; j; x_1 ] \tag{1.296}$$

if  $x$  belongs to a simple ergodic set of the stochastic process  $\{x^*[C]; x; j; j=1,2,\dots\}$ .

We now consider the sequence of  $\omega$ -functions  $\{k_j(\omega; [C]); j=1,2,\dots\}$ . If (1.290) and (1.293) hold, the  $\omega$ -function  $k_j(\omega; [C])$  represents the losses incurred in the period  $[\hat{t}_{j-1}(\omega; [C]), \hat{t}_j(\omega; [C])]$ .

Obviously, we have

$$\begin{aligned}
& \int_{X^*} P[C] (dx_1; x) \int_{-\infty}^{+\infty} |k| P^* [K_{dk}; [C]; x_1] \leq \\
& \leq \gamma_c \int_{X^*} P[C] (dx_1; x) \int_0^{\infty} t P^* [ \Xi_{dt}; [C]; x_1 ] + \\
& \quad + \bar{n}[C]; x \gamma_d < \infty. \tag{1.297}
\end{aligned}$$

The following lemma can be proved (cf. lemma 1.48):

Lemma 1.56

Under (1.283), (1.286), (1.290) and (1.293) for almost all  $\omega$  the limit

$$\lim_{r \rightarrow \infty} \frac{1}{r} \sum_{j=1}^r k_j(\omega; [C]) \tag{1.298}$$



exists and is equal to

$$\int_{X^*} p[C](dx_1; x) \int_{-\infty}^{+\infty} kP^* [K_{dk}; [C]; x_1] \quad (1.299)$$

if  $x$  belongs to a simple ergodic set of the stochastic process  $\{x^*[C]; x; j; j=1, 2, \dots\}$ .

Finally, we prove:

Lemma 1.57

Under (1.283), (1.286), (1.290) and (1.293), for almost all  $\omega$  the limit

$$\lim_{T \rightarrow \infty} \frac{k_T(\omega)}{T} \quad (1.300)$$

exists and is equal to

$$\frac{\int_{X^*} p[C](dx_1; x) \int_{-\infty}^{+\infty} kP^* [K_{dk}; [C]; x_1]}{\int_{X^*} p[C](dx_1; x) \int_{-\infty}^{+\infty} tP^* [\Xi_{dt}; [C]; x_1]} \quad (1.301)$$

if  $x$  belongs to a simple ergodic set of the stochastic process  $\{x^*[C]; x; j; j=1, 2, \dots\}$ .

Proof:

It follows from lemmas 1.53 and 1.55 that almost surely

$$\lim_{r \rightarrow \infty} \frac{t_r(\omega; [C])}{r} = 0 \quad (1.302)$$

and

$$\lim_{r \rightarrow \infty} \frac{n_r(\omega; [C]; [A])}{r} = 0. \quad (1.303)$$

Let  $n(T; \omega)$  be the number of entries in  $C$  during the period  $[0, T)$ . Obviously, we have almost surely

$$\begin{aligned}
& \frac{\sum_{j=1}^{n(T;\omega)} k_j(\omega; [C]) - t_{n(T;\omega)+1}(\omega; [C]) \gamma_c - n_{n(T;\omega)+1}(\omega; [C]; [A]) \gamma_d}{\sum_{j=1}^{n(T;\omega)+1} t_j(\omega; [C])} \leq \\
& \leq \frac{k_T(\omega)}{T} \leq \\
& \leq \frac{\sum_{j=1}^{n(T;\omega)} k_j(\omega; [C]) + t_{n(T;\omega)+1}(\omega; [C]) \gamma_c + n_{n(T;\omega)+1}(\omega; [C]; [A]) \gamma_d}{\sum_{j=1}^{n(T;\omega)} t_j(\omega; [C])} \quad (1.304)
\end{aligned}$$

and thus

$$\begin{aligned}
& \frac{\frac{1}{n(T;\omega)} \sum_{j=1}^{n(T;\omega)} k_j(\omega; [C]) - \frac{t_{n(T;\omega)+1}(\omega; [C]) \gamma_c}{n(T;\omega)} - \frac{n_{n(T;\omega)+1}(\omega; [C]; [A]) \gamma_d}{n(T;\omega)}}{\frac{1}{n(T;\omega)} \sum_{j=1}^{n(T;\omega)} t_j(\omega; [C]) + \frac{t_{n(T;\omega)+1}(\omega; [C])}{n(T;\omega)}} \leq \\
& \leq \frac{k_T(\omega)}{T} \leq \\
& \leq \frac{\frac{1}{n(T;\omega)} \sum_{j=1}^{n(T;\omega)} k_j(\omega; [C]) + \frac{t_{n(T;\omega)+1}(\omega; [C]) \gamma_c}{n(T;\omega)} + \frac{n_{n(T;\omega)+1}(\omega; [C]; [A]) \gamma_d}{n(T;\omega)}}{\frac{1}{n(T;\omega)} \sum_{j=1}^{n(T;\omega)} t_j(\omega; [C])} \quad (1.305)
\end{aligned}$$

If  $T \rightarrow \infty$ , then the assertion is an immediate consequence of the lemmas 1.56 and 1.53 and the relations (1.302) and (1.303).

This ends the proof.

## CHAPTER 2

### The decision process

#### 1. The basic probability space

In this section we consider a family of stochastic processes  $\{S_x^0; x \in X\}$ . For the definition of these stochastic processes the reader is referred to chapter 1 (\* = 0, M=N).<sup>1)</sup>

Let  $\Lambda_0$  be an  $\omega^0$ -set with the following properties:

- 1) For each  $\omega^0 \in \bar{\Lambda}_0$ , the t-function  $x_t^0(\omega^0)$  is continuous from the right;
- 2) In each bounded time interval in  $[0, \infty)$  and for each  $\omega^0 \in \bar{\Lambda}_0$ , the t-function  $x_t^0(\omega^0)$  has only a finite number of discontinuities.

#### Assumption 1

For each  $x \in X$ , a set  $K_x \in H^0$  can be found such that

- a)  $\Lambda_0 \subset K_x$ ;
- b)  $P^0[K_x; x] = 0$ .

#### Assumption 2

If  $x(t)$  is any mapping of the time axis  $[0, \infty)$  into the state space  $X$ , one and only one point  $\omega^0$  can be found such that

$$x_t^0(\omega^0) = x(t), \quad (2.1)$$

These assumptions imply that the probability spaces  $\{\Omega^0; F^0; P^0\}$  have the properties of  $\{\Omega^*; F^*; P^*\}$ . These properties have been considered in chapter 1 of this part.

---

1) It is convenient to index the points  $\omega$  and to suppress the index 0 in  $X^0$  and  $G^0$ ; \* = 0 means read 0 where we wrote \* in chapter 1.

Moreover, we assume the following:

Assumption 3

For each  $x \in X$  we have (cf. p.1)

$$P^0 [\Lambda_{0;x}; x] = 1. \quad (2.2)$$

This assumption implies that the initial state of the stochastic process  $S_x^0$  is  $x$  with probability 1.

In part I a strategy  $z$  is given by means of a function  $z(B;x)$  of sets  $B \in G$  and points  $x \in X$  with the following properties:

- 1) For each  $x \in X$  the set function  $z(B;x)$  is a probability measure defined on  $G$ ;
- 2) For each  $B \in G$  the  $x$ -function  $z(B;x)$  is measurable with respect to  $G$ ;
- 3) A closed set  $A_z$  can be found such that (cf. p.9)
  - a)  $z(A_z;x) = 0$  and thus  $z(\bar{A}_z;x) = 1$  if  $x \in A_z$ ;
  - b)  $z(\{x\};x) = 1$  if  $x \in \bar{A}_z$  and if the set  $\{x\}$  only contains the single point  $x$ .
  - c) for each  $x \in X$  we have

$$P^0 [\bar{\Xi}_{[0,\infty)}; A_z; x] = 1 \quad (2.3)$$

$$\text{and} \quad \int_0^\infty t P^0 [\bar{\Xi}_{dt}; A_z; x] < \infty. \quad (2.4)$$

The application of a strategy  $z$  involves extra transitions. As soon as a state of  $A_z$ , say  $x_1$ , is assumed, an instantaneous transition will happen with transition probabilities  $z(B;x_1)$ .

In order to be able to describe the resulting random walk in  $X$ , the extra transitions have to be incorporated in the original stochastic process  $S_x^0$ . To this end we assume that, if the transition in question leads to a state  $x_2$ , the process goes on like a  $S_{x_2}^0$ -process. Note that by point 3a) of  $z(B;x)$  the state  $x_2$  almost surely belongs to  $\bar{A}_z$ .

Let us now consider the set functions

$$\{p^j(B;x;z); j=0,1,\dots\}, \quad (2.5)$$

defined by (cf.p.10)

$$p^0(B;x;z) \stackrel{\text{def}}{=} P^0[\Delta_{B;A_z};x], \quad (2.6)$$

$$p^1(B;x;z) \stackrel{\text{def}}{=} \begin{cases} \int_X z(dy;x)P^0[\Delta_{B;A_z};y], & \text{if } x \in A_z \\ \int_{A_z} p^0(dI_1;x;z) \int_X z(dy;I_1)P^0[\Delta_{B;A_z};y], & \text{if } x \in \bar{A}_z. \end{cases} \quad (2.7)$$

and

$$p^k(B;x;z) \stackrel{\text{def}}{=} \int_{A_z} p^{k-1}(dI_k;x;z)p^1(B;I_k;z); \quad k=2,3,\dots \quad (2.8)$$

recursively. 2)

The following lemma can easily be proved:

Lemma 2.1

The functions  $\{p^k(B;x;z); k=0,1,\dots\}$  satisfy:

- 1) For each  $x \in X$  the set function  $p^k(B;x;z)$  is a probability measure defined on  $G$  with

$$p^k(A_z;x;z) = 1; \quad (2.9)$$

- 2) For each set  $B \in G$ , the  $x$ -function  $p^k(B;x;z)$  is measurable with respect to  $G$ .

It follows from the construction of the set function  $p^k(B;x;z)$ , that it represents the probability distribution of the initial point  $I_{-k+1}$  of the  $(k+1)^{\text{st}}$  added transition.

We now consider a sequence of spaces  $\{\Omega^k; k=1,2,\dots\}$ . These spaces are isomorphic with  $\Omega^0$ . Consequently, there exist 1-1 point transformations

$$\omega^k = T_{kh}(\omega^h); \quad k,h=1,2,\dots \quad (\text{in } (2.10))$$

from  $\Omega^h$  onto  $\Omega^k$  satisfying:

- 
- 2) States in  $A_z$  are denoted by  $I, I_1, \dots$  etc.

$$a) \omega^k = T_{kk}(\omega^k), k=0,1,\dots; \quad (2.11)$$

$$b) \omega^k = T_{kh}(T_{hj}(\omega^j)); k,h,j=0,1,\dots \quad (2.12)$$

These point transformations induce set transformations, denoted by

$$K^k = T_{kh}(K^h); k,h=0,1,\dots \quad (2.13)$$

and defined by

$$T_{kh}(K^h) \stackrel{\text{def}}{=} \{\omega^k | \omega^k = T_{kh}(\omega^h); \omega^h \in K^h\} \quad (2.14)$$

The set transformation (2.13) has the following properties:

$$a) K^k = T_{kk}(K), k=0,1,\dots; \quad (2.15)$$

$$b) K^k = T_{kh}(T_{hj}(K^j)), \text{ if } K^k = T_{kj}(K^j) \\ k,h,j=0,1,2,\dots \quad (2.16)$$

Let the class  $F^k$  of sets  $K^k$  be defined by

$$F^k \stackrel{\text{def}}{=} \{K^k | K^k = T_{ko}(K^o); K^o \in F^o\} \quad (2.17)$$

Obviously, the class  $F^k$  is isomorphic with  $F^o$ . Consequently, the following lemma is true:

Lemma 2.2

The class  $F^k$  is a  $\sigma$ -field of  $\omega^k$ -sets.

In order to simplify the notation, from now on we drop the space-index  $k$  in the notation of the sets  $K^k$ . Corresponding sets in different spaces will be denoted by the same symbol.

Next we introduce the  $\omega^k$ -functions  $\{x_t^k(\omega^k); t \in [0, \infty)\}$ , defined by

$$x_t^k(\omega^k) = x_t^o(T_{ok}(\omega^k)); k=0,1,\dots \quad (2.18)$$

If  $B \in G$  and if  $\Lambda_{t;B}$  is defined by

$$\Lambda_{t;B} \stackrel{\text{def}}{=} \{\omega^0 \mid x_t^0(\omega^0) \in B\}, \quad (2.19)$$

then  $\Lambda_{t;B} \in F^0$  and thus

$$T_{k0}(\Lambda_{t;B}) = \{\omega^k \mid x_t^k(\omega^k) \in B\} \in F^k. \quad (2.20)$$

So we have proved:

Lemma 2.3

The  $\omega^k$ -functions  $\{x_t^k(\omega^k); t \in [0, \infty)\}$  are measurable with respect to  $F^k$  ( $k=1, 2, \dots$ ).

Finally, we introduce the set functions  $\{P^k [K; x; z]; k=1, 2, \dots\}$ , defined on  $F^k$  by

$$P^k [K; x; z] \stackrel{\text{def}}{=} \int_{A_z} p^{k-1}(dI_k; x; z) \int_X z(dy; I_k) P^0 [K; y]. \quad (2.21)$$

The proof of the following lemma is left to the reader:

Lemma 2.4

For each  $x$  and  $k$  the set function  $P^k [K; x; z]$  is a probability measure. For each set  $K$  the  $x$ -function  $P^k [K; x; z]$  is measurable with respect to  $G$ .

The  $\omega$ -functions  $\{x_t^k(\omega^k); t \in [0, \infty)\}$  together with the probability space  $\{\Omega^k; F^k; P^k\}$  generate the stochastic process  $S_x^k$  ( $k=0, 1, \dots$ ).

The initial state of this process is not  $x$  (cf. assumption 3), but obeys the probability distribution

$$\begin{aligned} q(B; x; z) &= P^k [\Lambda_{0;B}; x; z] = \\ &= \int_{A_z} p^{k-1}(dI_k; x; z) z(B; I_k). \end{aligned} \quad (2.22)$$

It can easily be verified that the set function  $q(B; x; z)$  represents the probability distribution of the state into which the system

is transferred by the  $k^{\text{th}}$  added transition. Note that apart from the initial distribution the process  $S_x^k$  does not depend on the strategy applied.

By (2.18) and (2.21) the set  $\Lambda_0 \in F^k$  and has the following properties:

- 1) For each  $\omega^k \in \bar{\Lambda}_0$ , the  $t$ -function  $x_t^k(\omega^k)$  is continuous from the right;
- 2) In each bounded time interval in  $[0, \infty)$  and for each  $\omega^k \in \bar{\Lambda}_0$ , the  $t$ -function  $x_t^k(\omega^k)$  has only a finite number of discontinuities;
- 3) For each  $x \in X$ , we have

$$P^k [\Lambda_0; x; z] = 0. \quad (2.23)$$

If  $x(t)$  is any mapping of the time axis  $[0, \infty)$  into the state space  $X$ , then it follows from assumption 2 and (2.18) that one and only one point  $\omega^k$  can be found such that

$$x_t^k(\omega^k) = x(t). \quad (2.24)$$

So we have proved the following lemma:

Lemma 2.5

The stochastic processes  $S_x^k$  ( $k=1, 2, \dots$ ) satisfy the assumptions 1 and 2 of the  $S_x^*$  process ( $*=k; M=N$ ) and have initial probability distributions.

Up to now the probability spaces  $\{\Omega^k; F^k; P^k\}$  have been considered separately. In the remainder of this section, however, we shall construct one single probability space  $\{\Omega; F; P\}$  which is in fact the Cartesian product of the probability spaces  $\{\Omega^k; F^k; P^k\}$ .

Let  $\Omega$  be the product space of the spaces  $\Omega^k$  ( $k=0, 1, \dots$ ) and let  $H$  be the smallest  $\sigma$ -field of sets  $K$  that contains the cylinder sets

$$K = K_0 \times K_1 \times \dots = \prod_{i=0}^{\infty} K_i, \quad (2.25)$$

where



a)  $K_i \in F^i$ ,  $i=0,1,\dots$ ;

b) only a finite number of  $\omega^i$ -sets  $K_i$  are different from  $\Omega^i$ .

The points of  $\Omega$  are denoted by  $\omega = (\omega^0, \omega^1, \dots)$ . We now consider the point transformation

$$\omega_1 = T_{(k)}(\omega), \quad (2.26)$$

defined by

$$\omega_1^j = \omega^{k+j}; \quad j=0,1,\dots \quad (2.27)$$

The definition of the point transformation  $T_{(k)}(\omega)$  implies:

Lemma 2.6

For each  $\omega \in \Omega$  one and only one point  $\omega_1 \in \Omega$  can be found such that

$$\omega_1 = T_{(k)}(\omega). \quad (2.28)$$

By means of the point transformation (2.26) we can define a set transformation  $'K = T_{(k)}(K)$ , given by

$$T_{(k)}(K) \stackrel{\text{def}}{=} \{\omega_1 \mid \omega_1 = T_{(k)}(\omega); \omega \in K\}. \quad (2.29)$$

Next we consider a set transformation of sets  $K \in H$ , denoted by

$$''K = T_{(k); \omega^0 \dots \omega^{k-1}}(K) \quad (2.30)$$

and defined by

$$T_{(k); \omega^0 \dots \omega^{k-1}}(K) \stackrel{\text{def}}{=} T_{(k)}(K \cap \{\omega^0\} \times \dots \times \{\omega^{k-1}\} \times \prod_{i=k}^{\infty} \Omega_i), \quad (2.31)$$

where  $\{\omega^j\}$  is the point set in  $\Omega^j$  containing the single point  $\omega^j$ .

We now prove the following lemma:

Lemma 2.7

For each  $\omega \in \Omega$  the set transformation  $T_{(k); \omega^0 \dots \omega^{k-1}}(K)$  induces a  $\sigma$ -homomorphism of  $H$  onto  $H$ .

Proof:

If  $K$  is a product set of the type

$$K = \prod_{j=0}^r K_j \times \prod_{j=r+1}^{\infty} \Omega^j, \quad (2.32)$$

with  $K_j \in F^j$  and  $r \geq k$ , then

$$T_{(k); \omega^0 \dots \omega^{k-1}}(K) = \begin{cases} \prod_{j=k}^r K_j \times \prod_{j=r+1}^{\infty} \Omega^j, & \text{if } \omega^j \in K_j; j \leq k-1 \\ \emptyset, & \text{if } \exists_{j \leq k-1} \omega^j \in \bar{K}_j. \end{cases} \quad (2.33)$$

Consequently,  $T_{(k); \omega^0 \dots \omega^{k-1}}(K) \in H$ .

From the definition of  $T_{(k); \omega^0 \dots \omega^{k-1}}(K)$  it follows that  $\omega_1 \in T_{(k); \omega^0 \dots \omega^{k-1}}(K)$  implies  $(\omega^0, \dots, \omega^{k-1}, \omega_1) \in K$  and conversely.

Hence,

$$T_{(k); \omega^0 \dots \omega^{k-1}}(\bar{K}) = \overline{T_{(k); \omega^0 \dots \omega^{k-1}}(K)} \quad (2.34)$$

and

$$T_{(k); \omega^0 \dots \omega^{k-1}}\left(\bigcup_{i=1}^{\infty} K^i\right) = \bigcup_{i=1}^{\infty} T_{(k); \omega^0 \dots \omega^{k-1}}(K^i). \quad (2.35)$$

Let  $J$  be the class of  $\omega$ -sets  $K \in H$  which satisfy

$$T_{(k); \omega^0 \dots \omega^{k-1}}(K) \in H.$$

Then, by (2.33) the class  $J$  contains the product sets (2.32). Because of (2.34) and (2.35)  $J$  is a  $\sigma$ -field and therefore,  $J = H$ .

Let  $J_1$  be the class of  $\omega_1$ -sets " $K$ ", defined by

$$J_1 \stackrel{\text{def}}{=} \{ "K \mid "K = T_{(k); \omega^0 \dots \omega^{k-1}}(K); K \in H \}. \quad (2.36)$$

We have just proved that  $J_1 \subset H$ .

On the other hand, if " $K \in H$ , then

$$"K = T_{(k); \omega^0 \dots \omega^{k-1}}\left(\prod_{i=0}^{k-1} \Omega_i \times "K\right) \quad (2.37)$$

and thus

$$K \in J_1.$$

Hence,

$$J_1 = H. \quad (2.38)$$

This proves the lemma completely.

If  $K_k \in F^k$ , let us introduce the  $\omega$ -functions  $P^k(K_k; \omega^0 \dots \omega^{k-1})$ , defined by (cf.p.10)

$$P^k [K_k; \omega^0 \dots \omega^{k-1}] \stackrel{\text{def}}{=} \int_X z(dy; x^{k-1}(\omega^{k-1}; A_z)) P^0 [K_k; y]. \quad (2.39)$$

The following lemma can easily be proved:

Lemma 2.8

For each  $K_k \in F^k$  the  $\omega$ -function  $P^k [K_k; \omega^0 \dots \omega^{k-1}]$  is measurable with respect to  $H$ . For each  $\omega$  the set function  $P^k [K_k; \omega^0 \dots \omega^{k-1}]$  is a probability measure, defined on  $F^k$ .

Next we prove:

Lemma 2.9

If  $K_k \in F^k$ , we have

$$P^k [K_k; x; z] = \int_{\Omega^0} P^0 [d\omega^0; x] \int_{\Omega^1} P^1 [d\omega^1; \omega^0] \dots \int_{\Omega^k} P^k [K_k; \omega^0 \dots \omega^{k-1}]; \quad k=1, 2, \dots \quad (2.40)$$

Proof:

This lemma is proved by induction.

If  $k=1$ , then according to (2.6), (2.21) and (2.39) we find

$$\begin{aligned} P^1 [K_1; x; z] &= \int_{A_z} P^0 (dI_1; x; z) \int_X z(dy; I_1) P^0 [K_1; y] = \\ &= \int_{\Omega^0} P^0 [d\omega^0; x] P^1 [K_1; \omega^0]. \end{aligned} \quad (2.41)$$

Thus, the assertion is true for  $k=1$ .

Let us now assume that the assertion is also true for  $k=n-1$  and let

$M_{j;m}$  be defined by

$$\begin{aligned} M_{j;m} &\stackrel{\text{def}}{=} \{ \omega^{n-1} \mid \frac{j-1}{2^m} < \int_X z(dy; x^{n-1}(\omega^{n-1}; A_Z)) P^O [K_n; y] \leq \frac{j}{2^m} \} = \\ &= \{ \omega^{n-1} \mid \frac{j-1}{2^m} < P^n [K_n; \omega^0 \dots \omega^{n-1}] \leq \frac{j}{2^m} \} . \end{aligned} \quad (2.42)$$

According to (2.21)

$$\begin{aligned} P^n [K_n; x; z] &= \int_{A_Z} P^{n-1}(dI_n; x; z) \int_X z(dy; I_n) P^O [K_n; y] = \\ &= \int_{\Omega^{n-1}} P^{n-1} [d\omega^{n-1}; x; z] \int_X z(dy; x^{n-1}(\omega^{n-1}; A_Z)) P^O [K_n; y] = \\ &= \lim_{m \rightarrow \infty} \sum_{j=1}^{2^m} \frac{j}{2^m} P^{n-1} [M_{j;m}; x; z] = \\ &= \lim_{m \rightarrow \infty} \sum_{j=1}^{2^m} \frac{j}{2^m} \int_{\Omega^0} P^O [d\omega^0; x] \dots \int_{\Omega^{n-1}} P^{n-1} [M_{j;m}; \omega^0 \dots \omega^{n-2}] = \\ &= \int_{\Omega^0} P^O [d\omega^0; x] \dots \left\{ \lim_{m \rightarrow \infty} \sum_{j=1}^{2^m} \frac{j}{2^m} P^{n-1} [M_{j;m}; \omega^0 \dots \omega^{n-2}] \right\} = \\ &= \int_{\Omega^0} P^O [d\omega^0; x] \dots \int_{\Omega^{n-1}} P^{n-1} [d\omega^{n-1}; \omega^0 \dots \omega^{n-2}] \cdot P^n [K_n; \omega^0 \dots \omega^{n-1}] . \end{aligned} \quad (2.43)$$

Hence the assertion is also true for  $k=n$ .

This ends the proof.

We now consider the cylinder set  $K \in H$ , given by

$$K = \prod_{i=0}^{\infty} K_i \quad (2.44)$$

with

- a)  $K_i \in F^i$ ;
- b) only a finite number of  $\omega^i$ -sets  $K_i$  different from  $\Omega^i$ .

For each cylinder set  $K$  of the type (2.44) we can define

- 1) a number  $m_K$  by

$$m_K \stackrel{\text{def}}{=} \inf \{i \mid \bigvee_{j \geq i+1} K_j = \Omega^j\}, \quad (2.45)$$

2) functions  $I_K(\omega^0 \dots \omega^{m_K})$  by

$$I_K(\omega^0 \dots \omega^{m_K}) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } (\omega^0 \dots \omega^{m_K}) \in \prod_{i=0}^{m_K} K_i. \\ 0, & \text{otherwise.} \end{cases} \quad (2.46)$$

Now we are in a position to define a set function  $P[K; x; z]$  on the class of all sets of the type (2.44).

Let  $P[K; x; z]$  be defined by

$$\begin{aligned} P[K; x; z] &= \\ &= \int_{\Omega^0} P^0[d\omega^0; x] \int_{\Omega^1} P^1[d\omega^1; \omega^0] \dots \int_{\Omega^{m_K}} P^{m_K}[d\omega^{m_K}; \omega^0 \dots \omega^{m_K-1}] \\ &\quad \cdot I_K(\omega^0 \dots \omega^{m_K}). \end{aligned} \quad (2.47)$$

It can easily be verified that the right hand side of (2.47) exists.

#### Lemma 2.10

a) The domain of definition of the set function  $P[K; x; z]$ , can be extended to  $H$ . For each  $x \in X$  the set function  $P[K; x; z]$  is a probability measure defined on  $H$ . For each  $K \in F$  the  $x$ -function  $P[K; x; z]$  is measurable with respect to  $G$ .

b) If  $K_k \in F^k$  and if  $K_k^C$  is given by

$$K_k^C = \prod_{j=0}^{k-1} \Omega^j \times K_k \times \prod_{j=k+1}^{\infty} \Omega^j, \quad (2.48)$$

then

$$P^k[K^k; x; z] = P[K_k^C; x; z]. \quad (2.49)$$

#### Proof:

Point a) has been proved by I. Tulcea (cf. [1] p.137).

We now consider point b). The proof runs as follows (cf. (2.40)):

$$\begin{aligned} P[K_k^C; x; z] &= \\ &= \int_{\Omega^0} P^0[d\omega^0; x] \dots \int_{\Omega^k} P^k[d\omega^k; \omega^0 \dots \omega^{k-1}] I_{K_k^C}(\omega^0 \dots \omega^k) = \end{aligned}$$

$$\begin{aligned}
&= \int_{\Omega^0} P^0 [d\omega^0; x] \dots \int_{\Omega^k} P^k [K_k; \omega^0 \dots \omega^{k-1}] = \\
&= P^k [K_k; x; z]. \quad (2.50)
\end{aligned}$$

So we have obtained a probability space  $\{\Omega; H; P\}$  with  $\{\Omega^i; F^i; P^i\}$  as factor probability spaces.

Since  $\omega^k$  is the  $k^{\text{th}}$  component of  $\omega$ , the  $\omega^k$ -functions  $\{x_t^k(\omega^k); t \in [0, \infty)\}$  can also be defined on  $\Omega$ . We define the  $\omega$ -functions  $\{x_t^k(\omega); t \in [0, \infty)\}$  by

$$x_t^k(\omega) \stackrel{\text{def}}{=} x_t^k(\omega^k). \quad (2.51)$$

It follows from (2.50) and (2.51), that the stochastic processes  $\{S_x^k; k=0, 1, \dots\}$  can also be defined by means of the  $\omega$ -functions  $\{x_t^k(\omega); t \in [0, \infty)\}$  and the probability spaces  $\{\Omega; H; P\}$ .

In the final part of this section we shall discuss some properties of the probability measure  $P [K; x; z]$ .

Lemma 2.11

If  $K \in H$ , the  $\omega$ -function

$$\int_X z(dy; x^{j-1}(\omega; A_z)) P [T_{(j)}; \omega^0 \dots \omega^{j-1}(K); y; z]; \quad j=1, 2, \dots \quad (2.52)$$

is measurable with respect to  $H$ .

Proof:

Let  $K$  be the product set

$$K = \prod_{h=0}^{\infty} K_h, \quad (2.53)$$

with  $K_h \in F^h$ .

Then, we have

$$\begin{aligned}
&\int_X z(dy; x^{j-1}(\omega; A_z)) P [T_{(j)}; \omega^0 \dots \omega^{j-1}(K); y; z] = \\
&= \begin{cases} \int_X z(dy; x^{j-1}(\omega; A_z)) P \left[ \prod_{h=j}^{\infty} K_h; y; z \right], & \text{if } \omega^i \in K_i; \\ & i=0, \dots, j-1. \\ \emptyset, & \text{otherwise.} \end{cases} \quad (2.54)
\end{aligned}$$

Hence,

$$\begin{aligned} \int_X z(dx; x^{j-1}(\omega; A_z)) P [T_{(j)}; \omega^0 \dots \omega^{j-1}(K); x; z] = \\ = \chi(\omega) \int_X z(dx; x^{j-1}(\omega; A_z)) P \left[ \prod_{i=j}^{\infty} K_i; x; z \right], \end{aligned} \quad (2.55)$$

where

$$\chi(\omega) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \omega^i \in K_i \text{ for } i=0, 1, \dots, j-1 \\ 0, & \text{otherwise.} \end{cases} \quad (2.56)$$

Thus, if  $K$  is of the form (2.53), the  $\omega$ -function (2.52) is measurable with respect to  $H$ .

Let  $J$  be the class of  $\omega$ -sets  $K \in H$  which satisfy the assertion. So  $J$  includes the product sets (2.53). It can easily be verified that  $J$  also contains:

- a) the complements of  $J$ -sets;
- b) the limit of any monotone sequence of  $J$ -sets.

Consequently,  $J=H$ .

This ends the proof.

Lemma 2.12

If  $K \in H$ , we have for each  $x \in X$  and  $j \geq 1$

$$\begin{aligned} P [K; x; z] &= \int_{\Omega} P [d\omega; x; z] \int_X z(dy; y^{j-1}(\omega; A_z)) \cdot \\ &\quad \cdot P [T_{(j)}; \omega^0 \dots \omega^{j-1}(K); y; z] = \\ &= \int_{\Omega} P [d\omega; x; z] P [T_{(j)}; \omega^0 \dots \omega^{j-1}(K); x_o^j(\omega); z]. \end{aligned} \quad (2.57)$$

Proof:

Let us consider the cylinder set  $K \in H$ , given by

$$K = \bigcap_{i=0}^{\infty} K_i \quad (2.58)$$

with

- a)  $K_i \in F^i$ ;
- b) only a finite number of sets  $K_i$  different from  $\Omega^i$ .

If  $m_K$  is defined by

$$m_K \stackrel{\text{def}}{=} \inf \{i \mid \bigvee_{h>i} K_h = \Omega^h\} \quad (2.59)$$

and if  $\chi(\omega)$  is defined by

$$\chi(\omega) = \begin{cases} 1, & \text{if } \omega^i \in K_i; \quad i=0,1,\dots,j-1 \\ 0, & \text{otherwise} \end{cases} \quad (2.60)$$

then for each  $j$  we have

$$\begin{aligned} P[K; x; z] &= \\ &= \int_{\Omega^0} P^0[d\omega^0; x] \dots \int_{\Omega^{j-1}} P^{j-1}[d\omega^{j-1}; \omega^0 \dots \omega^{j-2}] \cdot \chi(\omega) \cdot \\ &\quad \cdot \int_X z(dx_1; x^{j-1}(\omega^{j-1}; A_z)) \cdot P\left[\prod_{h=j}^{\infty} K_h; x_1; z\right] = \\ &= \int_{\Omega} P[d\omega; x; z] \int_X z(dx_1; x^{j-1}(\omega^{j-1}; A_z)) \cdot \\ &\quad \cdot P[T_{(j)}; \omega^0 \dots \omega^{j-1}(K); x_1; z]. \quad (2.61) \end{aligned}$$

From (2.61) we deduce that the product sets  $K$  satisfy the assertion. By using similar arguments as in the proof of lemma 2.11 we can complete the proof of this lemma.

We now consider the  $\omega$ -set  $M_0$ , defined by

$$M_0 \stackrel{\text{def}}{=} \bigcup_{j=0}^{\infty} \prod_{i=0}^{j-1} \Omega^i \times \Lambda_0 \times \prod_{i=j+1}^{\infty} \Omega^i. \quad (2.62)$$

Obviously, we have for each  $x \in X$

$$P[M_0; x; z] = 0. \quad (2.63)$$

By completing the measure, the domain of definition of  $P[K; x; z]$  is from now on extended to the  $\sigma$ -field  $F$ , the smallest  $\sigma$ -field including  $H$  and containing all subsets of  $M_0$ .



The proof of the following lemma is left to the reader (cf. lemma 1.1):

Lemma 2.13

For each set  $K \in F$ , the  $x$ -function  $P [K;x;z]$  is measurable with respect to  $G$ .

2. The probabilistic foundation of the decision process

In section 1 we have stipulated that the application of a strategy involves extra transitions. We assumed that an instantaneous transition with transition probabilities  $z(B;x_1)$  occurs if a state, say  $x_1$ , of a closed set  $A_z$  is reached. If such a transition leads to a state  $x_2$ , then the process goes on like a  $S_{x_2}^0$ -process. The resulting random walk is called the decision process and is denoted by  $S_{x;z}$  if  $x$  is the initial state.

Since the initial distribution of the  $S_x^k$ -process represents the probability distribution of the state into which the system is transferred by the  $k^{\text{th}}$  added transition (cf. p.60), the stochastic process  $S_x^k$  can be used for the description of that part of the decision process which will take place between the  $k^{\text{th}}$  and the  $(k+1)^{\text{st}}$  added transition. Hereafter this part is called the  $(k+1)^{\text{st}}$  stretch of the decision process.

In such a presentation the points  $\omega \in \Omega^k$  determine realizations of the  $(k+1)^{\text{st}}$  stretch. Hence the points  $\omega \in \Omega$  determine realizations of the whole decision process.

In this section we shall demonstrate that decision processes can also be defined by means of probability spaces  $\{\Omega;F;P\}$ .

Obviously, the successive states in  $A_z$ , reached by the system, can for almost all  $\omega$  be represented by  $\{x^j(\omega;A_z); j=0,1,\dots\}$ . The lengths  $\{t^j(\omega;A_z); j=0,1,\dots\}$  of the time intervals between the added transitions are defined and measurable with respect to  $F^j$  (cf. lemmas 1.5.1 and 1.5.2 with  $*=j$ ).

The sequence of  $\omega$ -functions  $\{x^j(\omega;A_z); j=0,1,\dots\}$  together with a probability space  $\{\Omega;F;P\}$  generate a sequence of stochastic variables,

denoted by  $\{\underline{I}_{j+1}; j=0,1,\dots\}$ .

We already know that the applied strategy  $z$  effects an extra transition in the random states  $\{\underline{I}_j; j=1,2,\dots\}$ .

Obviously, we have for  $j=1,2,\dots$  (cf. (2.6), (2.7) and (2.8))

$$\begin{aligned} \text{Prob } \{\underline{I}_{j+1} \in B \mid \underline{I}_1 = I_1, \dots, \underline{I}_j = I_j\} &= \\ &= \int_{\mathbf{X}} z(dy; I_j) P^0 [\Delta_{B; A_z}; y] = p^1(B; I_j; z) \end{aligned} \quad (2.64)$$

and

$$\text{Prob } \{\underline{I}_{j+1} \in B\} = P^j [\Delta_{B; A_z}; x; z] = p^j(B; x; z) \quad (2.65)$$

with

$$p^j(B; x; z) = \int_{A_z} p^{j-1}(dI_j; x; z) p^1(B; I_j; z). \quad (2.66)$$

The following theorem is an immediate consequence of lemma 2.1 and the equations (2.64) through (2.66) (cf. [2] p.190 ff.):

#### Theorem 1

The stochastic variables  $\{\underline{I}_k; k=1,2,\dots\}$  constitute a stationary Markov process with a discrete time parameter.

This stochastic process is called the decision process on  $A_z$ .

Henceforth our considerations are based on the following assumption:

#### Assumption 4

There is a finite valued measure  $Q(U)$  of sets  $U \in G$  with  $Q(X) > 0$ , an integer  $k \geq 1$  and a positive  $\eta$ , such that for each  $I \in A_z$  (cf. [2], p. 192)

$$p^k(U; I; z) \leq 1 - \eta, \text{ if } Q(U) \leq \eta. \quad (2.67)$$

We can now prove (cf. [2], p.214):

#### Lemma 2.14

Under (2.67), the function  $p(U; I_1; z)$ , given by

$$p(U; I_1; z) \stackrel{\text{def}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n p^j(U; I_1; z), \quad (2.68)$$

defines for each  $I_1 \in A_Z$  a stationary absolute probability distribution.

The sequence of  $\omega$ -functions  $\{t^j(\omega; A_Z); j=0,1,\dots\}$  together with the probability space  $\{\Omega; F; P\}$  also generate a sequence of stochastic variables. These stochastic variables are denoted by  $\{\underline{t}_j; j=0,1,2,\dots\}$ .

Using similar arguments as in the proof of lemma 1.47 we can prove:

Lemma 2.15

Under (2.4) and (2.67) the limit

$$\lim_{r \rightarrow \infty} \frac{1}{r} \sum_{j=1}^r t^j(\omega; A_Z) \quad (2.69)$$

exists for almost all  $\omega$  and is equal to

$$\int_{A_Z} p(dI; I_1; z) \int_X z(dy; I) \int_0^\infty tP^0 [\underline{\Xi}_{dt; A_Z}; y] \quad (2.70)$$

if the initial state  $I_1$  belongs to a simple ergodic set of the stochastic process  $\{\underline{I}_k; k=1,2,\dots\}$ .

Lemma 2.16

Under the assumptions 1,3 and 4 and by property 3<sup>a</sup>) of the function  $z(B;x)$ , we have for each  $I_1 \in A_Z$

$$\int_{A_Z} p(dI; I_1; z) \int_X z(dx; I) \int_0^\infty tP^0 [\underline{\Xi}_{dt; A_Z}; x] > 0. \quad (2.71)$$

Proof:

Obviously, we have by assumption 3 for each  $x$

$$P^0 [\underline{\Xi}_{[0, \frac{1}{n}]; A_Z}; x] = P^0 [\underline{\Xi}_{[0, \frac{1}{n}]; A_Z} \cap \Lambda_{O;x}; x]. \quad (2.72)$$

Let us consider the limit

$$\lim_{n \rightarrow \infty} P^0 [\underline{\Xi}_{[0, \frac{1}{n}]; A_Z}; x] = P^0 [\bigcap_{n=1}^\infty \underline{\Xi}_{[0, \frac{1}{n}]; A_Z} \cap \Lambda_{O;x}; x]. \quad (2.73)$$

If  $x \in \bar{A}_Z$ , by the definition of  $\Lambda_O$  we find

$$\bigcap_{n=1}^{\infty} \mathbb{E} \left[ \mathbb{E} \left[ 0, \frac{1}{n} \right]; A_z \cap \Lambda_{0;x} \subset \Lambda_0 \right]. \quad (2.74)$$

Hence, if  $x \in \bar{A}_z$ ,

$$\lim_{n \rightarrow \infty} P^0 \left[ \mathbb{E} \left[ 0, \frac{1}{n} \right]; A_z; x \right] = 0 \quad (2.75)$$

and thus

$$\lim_{n \rightarrow \infty} P^0 \left[ \mathbb{E} \left[ \frac{1}{n}, \infty \right]; A_z; x \right] = 1. \quad (2.76)$$

Therefore, if  $x \in \bar{A}_z$ ,

$$\int_0^{\infty} t P^0 \left[ \mathbb{E}_{dt}; A_z; x \right] > 0. \quad (2.77)$$

The assertion now is an immediate consequence of property 3<sup>a</sup>) of the function  $z(B;x)$ .

This ends the proof.

Now we are in a position to prove the following theorem:

Theorem 2

Under the assumptions 1,3 and 4 and by property 3<sup>a</sup>) of the function  $z(B;x)$ , the limit

$$\lim_{r \rightarrow \infty} \frac{1}{r} \sum_{j=1}^r t^j(\omega; A_z) \quad (2.78)$$

exists and is positive for almost all  $\omega$ . In particular, if the initial state  $I_1$  belongs to a simple ergodic set of the stochastic process  $\{I_k; k=1,2,\dots\}$ , the limit (2.78) is almost surely equal to

$$\int_{A_z} p(dI; I_1; z) \int_X z(dx; I) \int_0^{\infty} t P^0 \left[ \mathbb{E}_{dt}; A_z; x \right]. \quad (2.79)$$

Proof:

If the initial state  $I_1$  is an ergodic state, then the assertion is an immediate consequence of lemmas 2.15 and 2.16.

If the initial state is a transient state then for almost all  $\omega$  the system will stay outside all simple ergodic sets only a finite number of times in its transitions (cf. [2], p.207). Consequently, for almost all  $\omega$  an integer  $n(\omega)$  can be found such that for  $n \geq n(\omega)$  the states  $x^n(\omega; A_Z)$  are ergodic. Hence, for almost all  $\omega$

$$\begin{aligned} \lim_{r \rightarrow \infty} \frac{1}{r} \sum_{j=1}^r t^j(\omega; A_Z) &= \\ &= \int_{A_Z} p(dI; \hat{I}; z) \int_X z(dx; I) \int_0^\infty t P^0[\cdot; dt; A_Z; x] > 0, \end{aligned} \quad (2.80)$$

where  $\hat{I}$  stands for  $x^{n(\omega)}(\omega; A_Z)$ .

This completes the proof.

Let us now consider the  $\omega$ -functions  $\{u_{t;1}^k(\omega); t \in [0, \infty)\}$  and  $\{u_{t;2}^k(\omega); t \in [0, \infty)\}$  for  $k=0, 1, \dots$ , defined by

$$u_{t;1}^0(\omega) = u_{t;2}^0(\omega) \stackrel{\text{def}}{=} x_t^0(\omega), \quad (2.81)$$

$$u_{t;1}^k(\omega) \stackrel{\text{def}}{=} \begin{cases} u_{t;1}^{k-1}(\omega), & \text{if } t \leq \hat{t}_k(\omega; A_Z) \\ x_{t-\hat{t}_k}^k(\omega; A_Z)(\omega), & \text{if } t > \hat{t}_k(\omega; A_Z) \end{cases}; \quad k > 0 \quad (2.82)$$

and

$$u_{t;2}^k(\omega) \stackrel{\text{def}}{=} \begin{cases} u_{t;2}^{k-1}(\omega), & \text{if } t < \hat{t}_k(\omega; A_Z) \\ x_{t-\hat{t}_k}^k(\omega; A_Z)(\omega), & \text{if } t \geq \hat{t}_k(\omega; A_Z) \end{cases}; \quad k > 0 \quad (2.83)$$

where

$$\hat{t}_k(\omega; A_Z) \stackrel{\text{def}}{=} \sum_{j=0}^{k-1} t^j(\omega; A_Z). \quad (2.84)$$

Note that the  $t$ -functions  $u_{t;1}^k(\omega)$  and  $u_{t;2}^k(\omega)$  only differ if  $t = \hat{t}_j(\omega; A_Z)$  ( $j=1, 2, \dots, k$ ).

Lemma 2.17

The  $\omega$ -functions  $\{u_{t;1}^k(\omega); k=1,2,\dots; t \in [0,\infty)\}$  are measurable with respect to  $H$ .

Proof:

This lemma will be proved by induction. It follows from (2.81), that the assertion is true for  $k=0$ .

Let us assume the assertion to be true for  $k=n-1$  and let us consider the sequence of  $\omega$ -functions  $\{u_{m;t}^n(\omega); m=1,2,\dots\}$ , defined for  $k=1,\dots,2^m$  by

$$u_{m;t}^n(\omega) \stackrel{\text{def}}{=} \begin{cases} x_{t-\frac{(k-1)t}{2^m}}^n(\omega), & \text{if } \frac{(k-1)t}{2^m} \leq \hat{t}_n(\omega; A_Z) < \frac{kt}{2^m} \\ u_{t;1}^{n-1}(\omega), & \text{if } \hat{t}_n(\omega; A_Z) \geq t. \end{cases} \quad (2.85)$$

Let the  $\omega$ -functions  $\{\chi_k(\omega); k=0,\dots,2^m\}$  be defined by

$$\chi_0(\omega) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \hat{t}_n(\omega; A_Z) \geq t \text{ or } \omega^n \in \Lambda_0 \\ 0, & \text{otherwise} \end{cases}, \quad (2.86)$$

and

$$\chi_k(\omega) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } \frac{(k-1)t}{2^m} \leq \hat{t}_n(\omega; A_Z) < \frac{kt}{2^m} \text{ and } \omega^n \in \bar{\Lambda}_0 \\ 0, & \text{otherwise} \end{cases}, \quad (2.87)$$

where  $\omega^n$  is the  $n^{\text{th}}$  component of  $\omega$ .

It can easily be verified that the  $\omega$ -functions  $\{\chi_k(\omega); k=0,\dots,2^m\}$  are measurable with respect to  $H$ .

Consequently, the  $\omega$ -functions

$$u_{m;t}^n(\omega) = \chi_0(\omega)u_{t;1}^{n-1}(\omega) + \sum_{k=1}^{2^m} \chi_k(\omega)x_{t-\frac{(k-1)t}{2^m}}^n(\omega); m=1,2,\dots \quad (2.88)$$

are measurable with respect to  $H$ .

It can easily be verified that for each  $\omega$  the sequence  $\{u_{m;t}^n(\omega); m=1,2,\dots\}$  converges to a limit, which is by consequence measurable with respect to  $H$ . Since for  $\omega \in \prod_{j=0}^{n-1} \Omega^j \times \bar{\Lambda}_0 \times \prod_{j=n+1}^{\infty} \Omega^j$  this limit is equal to  $u_{t;1}^n(\omega)$ , the latter is measurable with respect to  $H$ .

By a similar reasoning, we can prove:

Lemma 2.18

The  $\omega$ -functions  $\{u_{t;2}^k(\omega); k=0,1,\dots; t \in [0,\infty)\}$  are measurable with respect to  $H$ .

For each fixed  $k$  the  $\omega$ -functions  $\{u_{t;1}^k(\omega); t \in [0,\infty)\}$  together with the probability space  $\{\Omega; F; P\}$  generate a stochastic process

$$\{u_{t;x;1}^k; t \in [0,\infty)\}. \quad (2.89)$$

Let the stochastic variables  $\{\hat{t}_{-j}; j=1,2,\dots\}$  be defined by

$$\hat{t}_{-j} = \sum_{k=0}^{j-1} t_{-k}. \quad (2.90)$$

The stochastic process (2.89) describes the state of the system in  $X$  if after the  $k^{\text{th}}$  extra transition no more extra transitions are added and if at  $\{\hat{t}_{-j}; j=1,2,\dots,k\}$  only the initial point of the corresponding extra transition is recorded.

Similarly, we find that for each  $k$  the  $\omega$ -functions  $\{u_{t;2}^k(\omega); t \in [0,\infty)\}$  together with the probability space  $\{\Omega; F; P\}$  generate a stochastic process

$$\{u_{t;x;2}^k; t \in [0,\infty)\}. \quad (2.91)$$

The stochastic processes (2.89) and (2.91) are identical with the exception of the random points of time  $\{\hat{t}_{-j}; j=1,2,\dots,k\}$ . In (2.91) the state after the effectuation of the extra transition is presented at  $\hat{t}_{-j}$ .

In order to evade difficulties in determining the state at  $\hat{t}_{-j}$  we introduce the product space  $X'$  of two  $N$ -dimensional Cartesian spaces  $X_1$  and  $X_2$ . The  $\sigma$ -fields of all  $2N$ -dimensional Borelsets in  $X'$  is denoted by  $G'$ , while the corresponding  $\sigma$ -fields in  $X_1$  and  $X_2$  are called  $G_1$  and  $G_2$  respectively. Note that the spaces  $X$ ,  $X_1$  and  $X_2$  are isomorphic.

Next let for each  $k$  and  $t \in [0,\infty)$  the  $\omega$ -functions  $u_{t;1}^k(\omega)$  and  $u_{t;2}^k(\omega)$  map  $\Omega$  into  $X_1$  and  $X_2$  respectively.

From now on states are represented by points  $x' \in X'$ .

If  $x' \in A_Z \times X_2$ , the  $x_1$ -component of the "state"  $x'$  determines the initial point of the extra transition, while the  $x_2$ -component describes the state just after the effectuation of the transition.

If  $x' \in \bar{A}_Z \times X_2$ , then the  $x_1$  and the  $x_2$ -components of  $x'$  are equal.

Obviously, the  $\omega$ -functions  $\{u_t^k(\omega); k=0,1,\dots; t \in [0,\infty)\}$ , defined

by

$$u_t^k(\omega) \stackrel{\text{def}}{=} (u_{t;1}^k(\omega); u_{t;2}^k(\omega)), \quad (2.92)$$

map  $\Omega$  into  $X'$ .

The proofs of the following lemmas are obvious.

Lemma 2.19

The  $\omega$ -functions  $\{u_t^k(\omega); k=0,1,\dots; t \in [0,\infty)\}$  are measurable with respect to  $F$ .

Lemma 2.20

The  $\omega$ -functions  $\{u_t^k(\omega); k=0,1,\dots; t \in [0,\infty)\}$  have the following properties:

- a) For each  $\omega \in \bar{M}_0$  the  $t$ -function  $u_{t;2}^k(\omega)$  is continuous from the right;
- b) For each  $\omega \in \bar{M}_0$  the  $t$ -function  $u_t^k(\omega)$  has only a finite number of discontinuities in each finite time interval.

We now consider the  $\omega$ -functions  $x_{t;1}(\omega)$  and  $x_{t;2}(\omega)$  for  $k=1,2,\dots$ , defined by

$$x_{t;1}(\omega) \stackrel{\text{def}}{=} u_{t;1}^{k-1}(\omega), \text{ if } t \leq \hat{t}_k(\omega; A_Z) \text{ and}$$

$$\lim_{h \rightarrow \infty} \hat{t}_h(\omega; A_Z) = \infty. \quad (2.93)$$

$$x_{t;2}(\omega) \stackrel{\text{def}}{=} u_{t;2}^{k-1}(\omega), \text{ if } t < \hat{t}_k(\omega; A_Z) \text{ and}$$

$$\lim_{h \rightarrow \infty} \hat{t}_h(\omega; A_Z) = \infty; \quad (2.94)$$

$$x_{t;1}(\omega) = x_{t;2}(\omega) = x_t^0(\omega), \text{ if } \lim_{h \rightarrow \infty} \hat{t}_h(\omega; A_Z) < \infty. \quad (2.95)$$



Note that the  $\omega$ -functions  $x_{t;1}(\omega)$  and  $x_{t;2}(\omega)$  only differ at the points of time  $\{\hat{t}_k(\omega; A_z); k=1,2,\dots\}$ .

The following lemmas can easily be verified:

Lemma 2.21

The  $\omega$ -functions  $\{x_{t;1}(\omega); t \in [0, \infty)\}$  and  $\{x_{t;2}(\omega); t \in [0, \infty)\}$  are measurable with respect to  $H$ .

Lemma 2.22

The  $\omega$ -functions  $\{x_t(\omega); t \in [0, \infty)\}$ , defined by

$$x_t(\omega) \stackrel{\text{def}}{=} (x_{t;1}(\omega), x_{t;2}(\omega)), \quad (2.96)$$

map  $\Omega$  into  $X'$  and are measurable with respect to  $F$ .

Lemma 2.23

The  $\omega$ -functions  $\{x_t(\omega); t \in [0, \infty)\}$  have the following properties:

- a) For each  $\omega \in \bar{M}_0$  the  $t$ -function  $x_{t;2}(\omega)$  is continuous from the right;
- b) For each  $\omega \in \bar{M}_0$  the  $t$ -function  $x_t(\omega)$  has only a finite number of discontinuities in each finite time interval.

The  $\omega$ -functions  $\{x_t(\omega); t \in [0, \infty)\}$  together with a probability space  $\{\Omega; F; P\}$  generate a stochastic process

$$\{\underline{x}_{t;x}; t \in [0, \infty)\} \quad (2.97)$$

in  $X'$ .

Since

- a) by theorem 2 we have almost surely

$$\lim_{h \rightarrow \infty} \hat{t}_h(\omega; A_z) = \infty; \quad (2.98)$$

- b) the stochastic processes  $\{\underline{u}_{t;x;1}^k; t \in [0, \infty)\}$  and  $\{\underline{u}_{t;x;2}^k; t \in [0, \infty)\}$  describe the evolution in the state of the system if only  $k$  extra transitions are added,

it follows from (2.93), (2.94) and (2.95) that the stochastic process  $\{\underline{x}_{t;x}; t \in [0, \infty)\}$  describes the whole decision process in  $X'$ .

If  $\underline{x}_{t;x} \in A_z \times X_2$ , the  $x_1$ -component of the state  $\underline{x}_{t;x}$  determines the initial point of the extra transition, while the  $x_2$ -component describes the state just after the effectuation of this transition. The two components are equal if  $\underline{x}_{t;x} \in \bar{A}_z \times X_2$ .

Note that by virtue of assumption 3 the  $x_1$ -component of the initial state  $\underline{x}_{0;x}$  is equal to  $x$  with probability 1.

If  $x \in \bar{A}_z$ , then the  $x_2$ -component of the initial state  $\underline{x}_{0;x}$  also is equal to  $x$  with probability 1.

The  $x_2$ -component of  $\underline{x}_{0;x}$  obeys the probability distribution  $z(B;x)$  if  $x \in A_z$ . Consequently, for  $x \in A_z$  the decision process has an initial distribution.

So we have proved:

Lemma 2.24

Under assumptions 1,3 and 4, the decision process in  $X'$  can be defined by means of a stochastic process.

3. Properties of the decision process

In this section we shall show that, notwithstanding the decision process does not satisfy assumption 1 (cf. chapter 1 of this part), the assertions stated in lemmas 1.5.1 through 1.9 can still be proved.

As we noted at the end of section 2 the decision process  $\{\underline{x}_{t;x}; t \in [0, \infty)\}$  has an initial probability distribution if  $x_1 \in A_z$ . In the coming discussion we shall demonstrate that decision processes with given initial  $x'$ -states can also be defined.

For that purpose we have to define set functions  $\{P[\underline{K}; x'; z]; x' \in X'\}$ . Properties of these set functions are investigated at the end of this section.

Let us start with introducing the  $\omega$ -functions  $\{v_t(\omega); t \in [0, \infty)\}$ , defined by

$$v_t(\omega) \stackrel{\text{def}}{=} (x_{t;2}(\omega); x_{t;2}(\omega)). \quad (2.99)$$

The assertion stated in the following lemma can easily be proved.

Lemma 2.25

The  $\omega$ -functions  $\{v_t(\omega); t \in [0, \infty)\}$  have the following properties:

- a) For each  $t \in [0, \infty)$  the  $\omega$ -function  $v_t(\omega)$  is measurable with respect to  $F$ ;
- b) For each  $\omega \in \bar{M}_0$  the  $t$ -function  $v_t(\omega)$  is continuous from the right;
- c) For each  $\omega \in \bar{M}_0$  the  $t$ -function  $v_t(\omega)$  has only a finite number of discontinuities in each finite interval.

Next we consider the  $\omega$ -functions  $\{x_j(\omega; A_Z \times X_2); j=1, 2, \dots\}$ , defined by

$$x_j(\omega; A_Z \times X_2) \stackrel{\text{def}}{=} (x^{j-1}(\omega; A_Z), x_0^j(\omega)). \quad (2.100)$$

We easily verify:

Lemma 2.26

The  $\omega$ -functions  $\{x_j(\omega; A_Z \times X_2); j=1, 2, \dots\}$  are measurable with respect to  $F$ .

It follows from (2.99) and (2.100) that

$$x_t(\omega) = \begin{cases} x_j(\omega; A_Z \times X_2), & \text{if } t = \hat{t}_j(\omega; A_Z) \\ v_t(\omega), & \text{if } t \neq \hat{t}_j(\omega; A_Z); j=1, 2, \dots \end{cases} \quad (2.101)$$

If  $C$  is a closed set in  $X'$ , the  $\omega$ -function  $t(\omega; C)$  represents the moment that the system is for the first time in  $C$ .

In other words (cf. (1.47))

$$t(\omega; C) \stackrel{\text{def}}{=} \begin{cases} \inf \{t | x_t(\omega) \in C\}, & \text{if } x_t(\omega) \in C \text{ for some} \\ \infty, & \text{otherwise.} \end{cases} \quad \text{finite } t \quad (2.102)$$

If  $'t(\omega; C)$  and  $''t(\omega; C)$  are defined by

$$'t(\omega; C) \stackrel{\text{def}}{=} \begin{cases} \hat{t}_j(\omega; A_Z), & \text{if } x_k(\omega; A_Z \times X_2) \in \bar{C} \quad (k=1, \dots, j-1) \\ & \text{and } x_j(\omega; A_Z \times X_2) \in C \\ \infty, & \text{otherwise} \end{cases} \quad (2.103)$$

and

$${}''t(\omega; C) \stackrel{\text{def}}{=} \begin{cases} \inf \{t | v_t(\omega) \in C\}, & \text{if } v_t(\omega) \in C \text{ for some} \\ \infty, & \text{otherwise} \end{cases} \quad \text{finite } t \quad (2.104)$$

respectively, then we obviously have

$$t(\omega; C) = \min({}'t(\omega; C), {}''t(\omega; C)). \quad (2.105)$$

Since the  $\omega$ -functions  $\hat{t}_j(\omega; A_Z)$  and  $x_k(\omega; A_Z \times X_2)$  are measurable with respect to  $F$ , it follows from (2.103) that the  $\omega$ -function  $'t(\omega; C)$  is measurable with respect to  $F$ . Moreover, lemmas 2.25 and 1.5.1 imply that the  $\omega$ -function  ${}''t(\omega; C)$  is measurable with respect to  $F$ . ( $M_0 = \Lambda_0^*$ ) Consequently, we find:

Lemma 2.27.1

If  $C$  is a closed set in  $X'$ , then the  $\omega$ -function  $t(\omega; C)$ , defined by (2.102), is measurable with respect to  $F$ .

We now consider the  $\omega$ -function  $x(\omega; C)$ , defined by (cf. (1.51))

$$x(\omega; C) \stackrel{\text{def}}{=} \begin{cases} x_{t(\omega; C)}(\omega), & \text{if } t(\omega; C) < \infty \\ x_0(\omega), & \text{if } t(\omega; C) = \infty. \end{cases} \quad (2.106)$$

By (2.103) and (2.104)

$$x(\omega; C) = \begin{cases} x_j(\omega; A_Z \times X_2), & \text{if } \hat{t}_{j-1}(\omega; A_Z) < t(\omega; C) = \hat{t}_j(\omega; A_Z) < \infty \\ v(\omega; C), & \text{if } t(\omega; C) = {}''t(\omega; C) \end{cases}, \quad (2.107)$$

where

$$v(\omega; C) \stackrel{\text{def}}{=} \begin{cases} v_{{}''t(\omega; C)}(\omega), & \text{if } {}''t(\omega; C) < \infty \\ x_0(\omega), & \text{if } {}''t(\omega; C) = \infty. \end{cases} \quad (2.108)$$

Using similar arguments as in the proof of lemma 1.5.2 we can prove that the  $\omega$ -function  $v(\omega; C)$  is measurable with respect to  $F$ .

Obviously, we have:

Lemma 2.27.2

If  $C$  is a closed set in  $X'$ , then the  $\omega$ -function  $x(\omega; C)$ , defined by (2.106), is measurable with respect to  $F$ .

If  $B$  is a  $2N$ -dimensional Borelset, if  $C$  is a  $2N$ -dimensional closed set and if  $I$  is an interval in  $[0, \infty)$ , the  $\omega$ -sets  $\Lambda_{t;B;z}$ ,  $\Xi_{I;C;z}$  and  $\Delta_{B;C;z}$  are defined by

$$\Lambda_{t;B;z} \stackrel{\text{def}}{=} \{\omega \mid x_t(\omega) \in B\}, \quad (2.109)$$

$$\Xi_{I;C;z} \stackrel{\text{def}}{=} \{\omega \mid t(\omega;C) \in I\}, \quad (2.110)$$

and

$$\Delta_{B;C;z} \stackrel{\text{def}}{=} \{\omega \mid x(\omega;C) \in B\} \quad (2.111)$$

respectively.

We now assume the set  $C$  to be chosen in such a way that for each  $x_1 \in X_1$  we have

$$P[\Xi_{[0,\infty);C;z;x_1;z}] = 1. \quad (2.112)$$

Since each combination of a measurable  $\omega$ -function and the probability space  $\{\Omega;F;P\}$  generates a stochastic variable, the  $\omega$ -functions  $t(\omega;C)$  and  $x(\omega;C)$  lead us to the stochastic variables  $\underline{t}_{C;x_1}$  and  $\underline{x}_{C;x_1}$ . The probability distributions of these stochastic variables are given by

$$\text{Prob}\{\underline{t}_{C;x_1} \in I\} \stackrel{\text{def}}{=} P[\Xi_{I;C;z;x_1;z}] \quad (2.113)$$

and

$$\text{Prob}\{\underline{x}_{C;x_1} \in B\} \stackrel{\text{def}}{=} P[\Delta_{B;C;z;x_1;z}] \quad (2.114)$$

respectively.

The stochastic variable  $\underline{t}_{C;x_1}$  represents the length of the time period preceding the moment at which the system first is in  $C$ , while  $\underline{x}_{C;x_1}$  denotes the state at the end of this period if (2.112) is true.

Summarizing:

Lemma 2.27

If the assumptions 1,2,3 and 4 and condition (2.112) are satisfied, the probability distribution of the length  $\underline{t}_{C;x_1}$  of the period preceding the moment at which the system first is in  $C$  and that of the state  $\underline{x}_{C;x_1}$  at that point of time are defined. They are given by (2.113) and (2.114) respectively.

Let  $B$  be a closed set in  $X'$  and let us define a family of  $\omega$ -functions  $\{x_t(\omega; B); t \in [0, \infty)\}$  by

$$x_t(\omega; B) \stackrel{\text{def}}{=} \begin{cases} x_{t(\omega; B)+t}(\omega), & \text{if } t(\omega; B) < \infty \\ x_t(\omega), & \text{if } t(\omega; B) = \infty. \end{cases} \quad (2.115)$$

Lemma 2.28

The  $\omega$ -functions  $\{x_t(\omega; B); t \in [0, \infty)\}$  are measurable with respect to  $F$ .

Proof.

We first consider the  $\omega$ -functions  $\{v_t(\omega; B); t \in [0, \infty)\}$ , defined by

$$v_t(\omega; B) \stackrel{\text{def}}{=} \begin{cases} v_{t(\omega; B)+t}(\omega), & \text{if } t(\omega; B) < \infty \\ x_t(\omega), & \text{if } t(\omega; B) = \infty. \end{cases} \quad (2.116)$$

It follows from lemma 2.25 and 1.6 ( $\Lambda_0^{**} = M_0$ ) that the  $\omega$ -function  $v_t(\omega; B)$  is measurable with respect to  $F$ .

We obviously have

$$x_t(\omega; B) = \begin{cases} v_t(\omega; B), & \text{if } t(\omega; B) = \infty \text{ or if for each } k \\ & t(\omega; B) + t \neq \hat{t}_k(\omega; A_Z) \\ x_k(\omega; A_Z \times X_2), & \text{if } t(\omega; B) + t = \hat{t}_k(\omega; A_Z) > \\ & > \hat{t}_{k-1}(\omega; A_Z). \end{cases} \quad (2.117)$$

We now define the  $\omega$ -functions  $\{\chi_k(\omega); k=0, 1, \dots\}$  by

$$\chi_0(\omega) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } t(\omega; B) = \infty \text{ or if for each } k \\ & t(\omega; B) + t \neq \hat{t}_k(\omega; A_Z) \\ 0, & \text{otherwise} \end{cases} \quad (2.118)$$

and

$$\chi_k(\omega) \stackrel{\text{def}}{=} \begin{cases} 1, & \text{if } t(\omega; B) + t = \hat{t}_k(\omega; A_Z) > \hat{t}_{k-1}(\omega; A_Z) \\ 0, & \text{otherwise.} \end{cases} \quad (2.119)$$

Next we consider the sequence

$$\{\chi_0(\omega) v_t(\omega; B) + \sum_{k=1}^n \chi_k(\omega) x_k(\omega; A_Z \times X_2); n=1, 2, \dots\}. \quad (2.120)$$

Since the  $\omega$ -functions  $\chi_k(\omega)$ ,  $v_t(\omega;B)$  and  $x_k(\omega;A_Z \times X_2)$  are measurable with respect to  $F$ , all elements of (2.120) are measurable.

Consequently, the limit  $x_t(\omega;B)$  of this sequence is measurable with respect to  $F$ .

This ends the proof.

If  $B$  and  $C$  are closed sets, let us introduce the  $\omega$ -functions  $t(\omega;B;C)$  and  $x(\omega;B;C)$ , defined by

$$t(\omega;B;C) \stackrel{\text{def}}{=} \begin{cases} \inf \{t | x_t(\omega;B) \in C\}, & \text{if } x_t(\omega;B) \in C \text{ for some} \\ \infty, & \text{otherwise} \end{cases} \quad \text{finite } t \quad (2.121)$$

and

$$x(\omega;B;C) \stackrel{\text{def}}{=} \begin{cases} x_{t(\omega;B;C)}(\omega;B), & \text{if } t(\omega;B;C) < \infty \\ x_0(\omega;B), & \text{if } t(\omega;B;C) = \infty \end{cases} \quad (2.122)$$

respectively.

Lemma 2.29

If  $B$  and  $C$  are closed sets in  $X'$ , the  $\omega$ -functions  $t(\omega;B;C)$  and  $x(\omega;B;C)$ , defined by (2.121) and (2.122), are measurable with respect to  $F$ .

Proof:

The  $t$ -function  $x_t(\omega;B)$  has the same properties as the function  $x_t(\omega)$ . Therefore, lemma 2.29 is a direct consequence of lemmas 2.27.1 and 2.27.2.

This ends the proof.

If  $C$  is a closed set in  $X'$  and if  $\omega$  is a realization of a stochastic process  $S_{z;x}$ , let  $t(\omega;[C])$  be the moment that the system enters into  $C$  for the first time.

Repeating the arguments used in the proof of lemma 1.8.1, we can prove the following lemma:

Lemma 2.30.1

The  $\omega$ -function  $t(\omega;[C])$  is measurable with respect to  $F$ .

Let us introduce the  $\omega$ -function  $x(\omega; [C])$ , defined by

$$x(\omega; [C]) \stackrel{\text{def}}{=} \begin{cases} x_{t(\omega; [C])}(\omega), & \text{if } t(\omega; [C]) < \infty \\ x_0(\omega), & \text{if } t(\omega; [C]) = \infty \end{cases} \quad (2.123)$$

Note that by this definition the state at the end of the period  $[0, t(\omega; [C])]$  is given by  $x(\omega; [C])$  unless  $t(\omega; [C]) = \infty$ .

Lemma 2.30.2

The  $\omega$ -function  $x(\omega; [C])$  is measurable with respect to  $F$ .

Proof:

We first consider the function  $v(\omega; [C])$ , defined by

$$v(\omega; [C]) \stackrel{\text{def}}{=} \begin{cases} v_{t(\omega; [C])}(\omega), & \text{if } t(\omega; [C]) < \infty \\ x_0(\omega), & \text{if } t(\omega; [C]) = \infty \end{cases} \quad (2.124)$$

By using similar arguments as in the proof of lemma 1.82 we can prove that the  $\omega$ -function  $v(\omega; [C])$  is measurable with respect to  $F$ .

Obviously, we have

$$x(\omega; [C]) = \begin{cases} v(\omega; [C]), & \text{if } t(\omega; [C]) \neq \hat{t}_k(\omega; A_Z) \text{ for each } k \\ x_k(\omega; A_Z \times X_2), & \text{if } t(\omega; [C]) = \hat{t}_k(\omega; A_Z) > \\ & > \hat{t}_{k-1}(\omega; A_Z). \end{cases} \quad (2.125)$$

The proof is immediate.

Let us introduce the  $\omega$ -sets  $\Xi_I; [C]; z$  and  $\Delta_B; [C]; z$ , defined by

$$\Xi_I; [C]; z \stackrel{\text{def}}{=} \{\omega \mid t(\omega; [C]) \in I\} \quad (2.126)$$

and

$$\Delta_B; [C]; z \stackrel{\text{def}}{=} \{\omega \mid x(\omega; [C]) \in B\} \quad (2.127)$$

respectively.

We now assume the closed set  $C$  to be chosen in such a way that for each  $x_1 \in X_1$



$$P \left[ \Xi_{[0, \infty)}; [C]; z; x_1; z \right] = 1. \quad (2.128)$$

The  $\omega$ -functions  $t(\omega; [C])$  and  $x(\omega; [C])$  together with the probability space  $\{\Omega; F; P\}$  generate the stochastic variables  $\underline{t}[C]; x_1$  and  $\underline{x}[C]; x_1$ ; the corresponding probability distributions are given by

$$\text{Prob} \{ \underline{t}[C]; x_1 \in I \} \stackrel{\text{def}}{=} P \left[ \Xi_I; [C]; z; x_1; z \right] \quad (2.129)$$

and

$$\text{Prob} \{ \underline{x}[C]; x_1 \in B \} \stackrel{\text{def}}{=} P \left[ \Delta_B; [C]; z; x_1; z \right] \quad (2.130)$$

respectively.

The stochastic variable  $\underline{t}[C]; x_1$  represents the length of the time period preceding the first entry in  $C$ , while  $\underline{x}[C]; x_1$  denotes the state at the end of that period if (2.128) is true.

Summarizing:

Lemma 2.30

If the assumptions 1,2,3 and 4 and condition (2.128) are satisfied, the probability distribution of the length  $\underline{t}[C]; x_1$  of the period preceding the first entry in  $C$  and that of the state  $\underline{x}[C]; x_1$  at that point of time are defined. They are given by (2.129) and (2.130) respectively.

We now consider the  $\omega$ -functions

$\{x_t(\omega; [C]); t \in [0, \infty)\}$ , defined by

$$x_t(\omega; [C]) \stackrel{\text{def}}{=} \begin{cases} x_{t(\omega; [C]) + t}(\omega), & \text{if } t(\omega; [C]) < \infty \\ x_t(\omega), & \text{if } t(\omega; [C]) = \infty \end{cases} \quad (2.131)$$

Repeating the arguments used in the proof of lemma 2.28, we can prove:

Lemma 2.31

The  $\omega$ -functions  $\{x_t(\omega; [C]); t \in [0, \infty)\}$  are measurable with respect to  $F$ .

We shall now demonstrate that decision processes with a given initial  $x'$ -state can also be defined. To this end we introduce the  $\sigma$ -field  $\hat{H}_0$ , the smallest  $\sigma$ -field of  $\omega$ -sets with respect to which the  $\omega$ -function  $x_0(\omega)$  is measurable.

Lemma 2.32

A conditional probability measure  $P[K; x_1; z | \hat{H}_0]$  can be defined on  $F$ .

Proof:

We first consider a set  $K$  of the following type:

$$K = K_0 \times 'K \quad (2.132)$$

with  $K_0 \in F^0$  and  $'K \in F$ . 3)

We can easily verify that the  $\omega$ -function  $\mu(K; \omega)$ , defined by

$$\mu(K; \omega) \stackrel{\text{def}}{=} \begin{cases} P[K; x_{0;2}(\omega); z], & \text{if } x_{0;1}(\omega) = x_{0;2}(\omega) \in \bar{A}_z \\ P[K_0^c; x_{0;1}(\omega); z] P['K; x_{0;2}(\omega); z], & \text{otherwise} \end{cases} \quad (2.133)$$

with  $K_0^c = K_0 \times \Omega$ , 3)

is measurable with respect to  $\hat{H}_0$ .

Since for  $B_1 \in G_1$  and  $B_2 \in G_2$  (cf. lemma 2.12)

$$\begin{aligned} \text{a) } P[K \cap \Lambda_{0; B_1 \times B_2; z}; x_1; z] &= \\ &= P[K \cap \Lambda_{0; B_1 \times B_2; z} \cap \Lambda_{0; A_z \times X_2; z}; x_1; z] + \\ &+ P[K \cap \Lambda_{0; B_1 \times B_2; z} \cap \Lambda_{0; \bar{A}_z \times X_2; z}; x_1; z]; \end{aligned} \quad (2.134)$$

---

3) Really,  $F$  is a  $\sigma$ -field of sets in  $\Omega = \prod_{j=0}^{\infty} \Omega^j$  and not in  $'\Omega = \prod_{j=1}^{\infty} \Omega^j$ . But, since the spaces  $\Omega$  and  $'\Omega$  are isomorphic, isomorphic  $\sigma$ -fields, denoted with the same symbol, do not cause confusion.

$$\begin{aligned}
\text{b) } & P [K \cap \Lambda_{0; B_1 \times B_2; z} \cap \Lambda_{0; A_z \times X_2; z; x_1; z}] = \\
& = P [K \cap \Lambda_{0; (B_1 \cap A_z) \times B_2; z; x_1; z}] = \\
& = \int_{\Lambda_{0; (A_z \cap B_1) \times X_2; z}} P [d\omega; x_1; z] \int_{B_2} z(dx_2; x_1^0(\omega; A_z)) P [T_1; \omega^0(K); x_2; z] = \\
& = \int_{\Lambda_{0; (A_z \cap B_1) \times X_2; z}} P [d\omega; x_1; z] \int_{B_2} z(dx_2; x_1) P [T_1; \omega^0(K); x_2; z] = \\
& = \int_{\Lambda_{0; (A_z \cap B_1) \times X_2; z} \cap K_o^c} P [d\omega; x_1; z] \int_{B_2} z(dx_2; x_1) P ['K; x_2; z] = \\
& = P [\Lambda_{0; (A_z \cap B_1) \times X_2; z} \cap K_o^c; x_1; z] \int_{B_2} z(dx_2; x_1) P ['K; x_2; z] = \\
& = \int_{\Lambda_{0; (A_z \cap B_1) \times B_2; z}} P [d\omega; x_1; z] P [K_o^c; x_{o;1}(\omega); z] P ['K; x_{o;2}(\omega); z]; \quad (2.135)
\end{aligned}$$

$$\begin{aligned}
\text{c) } & P [K \cap \Lambda_{0; B_1 \times B_2; z} \cap \Lambda_{0; \bar{A}_z \times X_2; z; x_1; z}] = \\
& = \int_{\Lambda_{0; (\bar{A}_z \cap B_1) \times B_2; z}} P [d\omega; x_1; z] P [K; x_{o;2}(\omega); z], \quad (2.136)
\end{aligned}$$

we find

$$\begin{aligned}
& P [K \cap \Lambda_{0; B_1 \times B_2; z; x_1; z}] = \\
& = \int_{\Lambda_{0; (A_z \cap B_1) \times B_2; z}} P [d\omega; x_1; z] P [K_o^c; x_{o;1}(\omega); z] P ['K; x_{o;2}(\omega); z] + \\
& + \int_{\Lambda_{0; (\bar{A}_z \cap B_1) \times B_2; z}} P [d\omega; x_1; z] P [K; x_{o;2}(\omega); z] = \int_{\Lambda_{0; B_1 \times B_2; z}} P [d\omega; x_1; z] \mu(K; \omega). \quad (2.137)
\end{aligned}$$

The class of all finite unions of disjoint product sets  $B_1 \times B_2$ , with  $B_1 \in G_1$  and  $B_2 \in G_2$ , is a field of  $x'$ -sets. If  $\bigcup_{i=1}^n B_{1i} \times B_{2i}$  is such a union, then we obviously have

$$\begin{aligned} P \left[ K \cap \Lambda_{\circ; \bigcup_{i=1}^n B_{1i} \times B_{2i}; z; x_1; z} \right] &= \\ &= \sum_{i=1}^n \int_{\Lambda_{\circ; B_{1i} \times B_{2i}; z}} P [d\omega; x_1; z] \mu(K; \omega) = \\ &= \int_{\Lambda_{\circ; \bigcup_{i=1}^n B_{1i} \times B_{2i}; z}} P [d\omega; x_1; z] \mu(K, \omega). \end{aligned} \quad (2.138)$$

Now let  $J_K$  be the class of sets  $B \in G'$  with the following property:

$$P \left[ K \cap \Lambda_{\circ; B; z; x_1; z} \right] = \int_{\Lambda_{\circ; B; z}} P [d\omega; x_1; z] \mu(K; \omega). \quad (2.139)$$

By (2.138) the class  $J_K$  includes the field of all finite unions of disjoint product sets. Moreover, we can easily prove that  $J_K$  contains all limits of monotone sequences of  $J_K$ -sets. Consequently,  $J_K$  includes the  $\sigma$ -field  $G'$ . Hence  $J_K = G'$ .

Since the  $\sigma$ -field  $\hat{H}_\circ$  consists of sets of the form  $\Lambda_{\circ; B; z}$  with  $B \in G'$ , we have now proved that

$$P \left[ K \cap \Lambda; x_1; z \right] = \int_{\Lambda} P [d\omega; x_1; z] \mu(K; \omega) \quad (2.140)$$

if  $K$  satisfies (2.132) and  $\Lambda \in \hat{H}_\circ$ .

The set function  $\mu(K; \omega)$  is for each  $\omega$  a product probability measure, defined on the class of all product sets of the type (2.132), and therefore the domain of definition can be extended to  $H$  and  $F$  uniquely.

Henceforth the set function  $\mu(K; \omega)$  is defined on  $F$ .

Now let  $J$  be the class of  $\omega$ -sets  $K \in F$  with the following properties:

- a) the  $\omega$ -function  $\mu(K; \omega)$  is measurable with respect to  $\hat{H}_0$ ;  
 b) for each  $\Lambda \in \hat{H}_0$

$$P [K \cap \Lambda; x_1; z] = \int_{\Lambda} P [d\omega; x_1; z] \mu(K; \omega). \quad (2.141)$$

It can easily be verified that

- a)  $\Omega \in J$ ;  
 b) if  $K \in J$ , then  $\bar{K} \in J$ ;  
 c) if  $K_i \in J$  and if  $K_i \subset K_{i+1} \dots (i=1, 2, \dots)$ , then  $\bigcup_{i=1}^{\infty} K_i \in J$ .

Consequently,  $J$  is a  $\sigma$ -field.

According to (2.140) the class  $J$  includes product sets. Thus,  $J \supset H$ .

It follows from (2.133) that  $K = \prod_{j=0}^{k-1} \Omega^j \times \Lambda_0 \times \prod_{j=k+1}^{\infty} \Omega^j$  satisfies for each  $\omega \in \Omega$

$$\mu(K; \omega) = 0. \quad (2.142)$$

Therefore, for each  $\omega \in \Omega$  (cf. (2.62))

$$\mu(M_0; \omega) = 0. \quad (2.143)$$

This implies that all subsets of  $M_0$  belong to  $J$ . Hence,  $J = F$ .

Finally, let us define  $P [K; x_1; z | \hat{H}_0]$  by

$$P [K; x_1; z | \hat{H}_0] = \mu(K; \omega). \quad (2.144)$$

The proof is complete.

The following properties of  $P [K; x_1; z | \hat{H}_0]$  can easily be proved and are stated for later reference:

- 1) for each  $\omega \in \Omega$  and  $K \in F$

$$P [K; x_1; z | \hat{H}_0] = P [K; x_{0;2}(\omega); z], \text{ if } x_{0;1}(\omega) = x_{0;2}(\omega) \in \bar{A}_z; \quad (2.145)$$

2) for each  $\omega \in \Omega$  and  $x' \in X'$

$$P \left[ \Lambda_{0; \{x_1\} \times \{x_2\}; z; x_1; z \mid \hat{H}_0 \right] = \begin{cases} 1, & \text{if } x_0(\omega) = (x_1, x_2) \\ 0, & \text{if } x_0(\omega) \neq (x_1, x_2) \end{cases} \quad (2.146)$$

where the product set  $\{x_1\} \times \{x_2\}$  is the point set containing the single point  $x' = (x_1, x_2)$ .

Let us define a family of probability measures  $\{P[K; x'; z] ; x' \in X'\}$ , with  $F$  as domain of definition, by

$$P[K; x'; z] \stackrel{\text{def}}{=} P[K; x_1; z \mid \hat{H}_0], \quad (2.147)$$

where  $\omega$  satisfies  $x_0(\omega) = x' = (x_1, x_2)$ .

The decision process  $\{x_t; x'; z; t \in [0, \infty)\}$  with initial state  $x'$  is now defined by means of the  $\omega$ -functions  $\{x_t(\omega); t \in [0, \infty)\}$  and the probability space  $\{\Omega; F; P\}$ , where  $P$  is given by (2.147).

Let  $H_z$  be the smallest  $\sigma$ -field of  $\omega$ -sets with respect to which the  $\omega$ -functions  $\{x_t(\omega); t \in [0, \infty)\}$  are measurable.

Presently, we need the following result:

Lemma 2.33

If  $K \in H_z$ , then

$$P[K; x_1; z \mid \hat{H}_0] = \begin{cases} P[K; x_{0;2}(\omega); z], & \text{if } x_{0;1}(\omega) = x_{0;2}(\omega) \in \bar{A}_z \\ P[T_{(1)}; \omega^0(K); x_{0;2}(\omega); z], & \text{if } x_{0;1}(\omega) \in A_z. \end{cases} \quad (2.148)$$

Proof:

Let us first consider a set  $K$ , given by

$$K = \Lambda_{t; B_1 \times B_2; z}, \quad (2.149)$$

with  $B_1 \in G_1$  and  $B_2 \in G_2$ .

Obviously, we have

$$\Lambda_{t;B_1 \times B_2; z} \cap \Lambda_{0;A_z \times X_2; z} = \begin{cases} \Lambda_{0;A_z} \times \Lambda_{t;B_1 \times B_2; z}, & \text{if } t > 0 \\ \Lambda_{0;A_z} \cap B_1 \times \Lambda_{0;B_2 \times X_2; z}, & \text{if } t=0 \end{cases}$$

(2.150)

where  $\Lambda_{t;B_1 \times B_2; z}$  and  $\Lambda_{0;B_2 \times X_2; z}$  are

a) sets in  $\Omega = \prod_{j=1}^{\infty} \Omega^j$ ;

b) isomorphic with  $\Lambda_{t;B_1 \times B_2; z}$  and  $\Lambda_{0;B_2 \times X_2; z}$  respectively.

Thus, if  $t > 0$ ,

$$\begin{aligned} \mu(\Lambda_{t;B_1 \times B_2; z}; \omega) &= \mu(\Lambda_{t;B_1 \times B_2; z} \cap \Lambda_{0;A_z \times X_2; z}; \omega) + \\ &\quad + \mu(\Lambda_{t;B_1 \times B_2; z} \cap \Lambda_{0;\bar{A}_z \times X_2; z}; \omega) = \\ &= \mu(\Lambda_{0;A_z} \times \Lambda_{t;B_1 \times B_2; z}; \omega) + \mu(\Lambda_{t;B_1 \times B_2; z} \cap \Lambda_{0;\bar{A}_z \times X_2; z}; \omega). \end{aligned}$$

(2.151)

By (2.133) and (2.145) we find, if  $t > 0$ ,

$$\begin{aligned} \mu(\Lambda_{t;B_1 \times B_2; z}; \omega) &= \\ &= \begin{cases} P[\Lambda_{t;B_1 \times B_2; z}; x_{0;2}(\omega); z], & \text{if } x_{0;1}(\omega) \in \bar{A}_z \\ P[\Lambda_{0;A_z}^c; x_{0;1}(\omega); z] P[\Lambda_{t;B_1 \times B_2; z}; x_{0;2}(\omega); z], & \\ & \text{if } x_{0;1}(\omega) \in A_z \end{cases} \end{aligned}$$

(2.152)

and consequently,

$$\mu(\Lambda_{t;B_1 \times B_2; z}; \omega) = P[\Lambda_{t;B_1 \times B_2; z}; x_{0;2}(\omega); z]. \quad (2.153)$$

However, if  $t=0$ , we find by (2.150)

$$\begin{aligned}
\mu(\Lambda_{t; B_1 \times B_2; z; \omega}) &= \mu(\Lambda_{0; A_z \cap B_1 \times B_2; z; \omega}) + \\
&\quad + \mu(\Lambda_{0; \bar{A}_z \cap B_1 \times B_2; z; \omega}) = \\
&= \mu(\Lambda_{0; A_z \cap B_1 \times B_2; z; \omega}) + \mu(\Lambda_{0; \bar{A}_z \cap B_1 \times B_2; z; \omega}).
\end{aligned} \tag{2.154}$$

By (2.133) and (2.145)

$$\begin{aligned}
\mu(\Lambda_{0; B_1 \times B_2; z; \omega}) &= \\
&= \begin{cases} P[\Lambda_{0; B_1 \times B_2; z; x_{0;2}(\omega); z}], & \text{if } x_{0;1}(\omega) \in \bar{A}_z \\ P[\Lambda_{0; A_z \cap B_1; x_{0;1}(\omega); z}] \cdot P[\Lambda_{0; B_2 \times X_2; z; x_{0;2}(\omega); z}], & \\ & \text{if } x_{0;1}(\omega) \in A_z \end{cases}
\end{aligned} \tag{2.155}$$

and consequently,

$$\begin{aligned}
\mu(\Lambda_{0; B_1 \times B_2; z; \omega}) &= \begin{cases} P[\Lambda_{0; B_1 \times B_2; z; x_{0;2}(\omega); z}], & \text{if } x_{0;1}(\omega) \in \bar{A}_z \\ 0, & \text{if } x_{0;1}(\omega) \in \bar{B}_1 \cap A_z \\ P[\Lambda_{0; B_2 \times X_2; z; x_{0;2}(\omega); z}], & \\ & \text{if } x_{0;1}(\omega) \in B_1 \cap A_z. \end{cases}
\end{aligned} \tag{2.156}$$

From (2.153) and (2.156) it follows that for any  $t$

$$\begin{aligned}
\mu(\Lambda_{t; B_1 \times B_2; z; \omega}) &= \begin{cases} P[\Lambda_{t; B_1 \times B_2; z; x_{0;1}(\omega); z}], & \text{if } x_{0;1}(\omega) \in \bar{A}_z \\ P[T_{(1); \omega^0}(\Lambda_{t; B_1 \times B_2; z; x_{0;2}(\omega); z)], & \\ & \text{if } x_{0;1}(\omega) \in A_z. \end{cases}
\end{aligned} \tag{2.157}$$

Hence, the  $\omega$ -sets of the type (2.149) satisfy the assertion.



Let  $J$  be the class of sets  $K$  with the following properties:

- a)  $J \subset H_z$ ;
- b) the sets  $K$  satisfy (2.148).

The following points can easily be verified:

- a)  $\Lambda_{t; B_1 \times B_2; z} \in J$ ;
- b) if  $K \in J$ , then  $\bar{K} \in J$ ;
- c) if  $K_i \in J$  and if  $K_i \subset K_{i+1} \dots$  ( $i=1, 2, \dots$ ), then  $\bigcup_{i=1}^{\infty} K_i \in J$ .

Consequently,  $J$  is a  $\sigma$ -field that includes the sets  $\Lambda_{t; B_1 \times B_2; z}$ . Hence,  $J = H_z$ .

Lemma 2.34

If  $K \in H$ , we have for each  $x' \in X'$  and  $j \geq 1$

$$\begin{aligned} P [K; x'; z] &= \int_{\Omega} P [d\omega; x'; z] \int_X z(dy; x^{j-1}(\omega; A_z)) \cdot \\ &\quad \cdot P [T_{(j)}; \omega^0 \dots \omega^{j-1}(K); y; z] = \\ &= \int_{\Omega} P [d\omega; x'; z] P [T_{(j)}; \omega^0 \dots \omega^{j-1}(K); x_0^j(\omega); z] . \end{aligned} \tag{2.158}$$

Proof:

The proof of this lemma is similar to that of lemma 2.12.

We now make the following assumption:

Assumption 5

The basic probability space  $\{\Omega^0; F^0; P^0\}$  is strongly Markovian.

In the final part of this section we shall prove two additional properties of the set function  $P [K; x'; z]$ . In virtue of these properties the decision process is a stationary strong Markov process, as we shall see in section 4.

Let us consider the  $\omega$ -functions  $\{\hat{x}_t(\omega; t_0); t \in [0, \infty)\}$ , defined by

$$\hat{x}_t(\omega; t_0) \stackrel{\text{def}}{=} \begin{cases} x_t(\omega), & \text{if } t \leq t_0 \\ x_{t_0}(\omega), & \text{if } t > t_0. \end{cases} \quad (2.159)$$

The following lemma can easily be proved:

Lemma 2.35

The  $\omega$ -functions  $\{\hat{x}_t(\omega; t_0); t \in [0, \infty)\}$  are measurable with respect to  $F$ .

Let the class of  $\omega$ -sets  $\hat{H}_{t_0}$  be the smallest  $\sigma$ -field with respect to which the  $\omega$ -functions  $\{\hat{x}_t(\omega; t_0); t \in [0, \infty)\}$  are measurable.

Note that  $\hat{H}_{t_0}$  also represents the smallest  $\sigma$ -field with respect to which the  $\omega$ -functions  $\{x_t(\omega); t \leq t_0\}$  are measurable.

Next we consider the  $\omega$ -sets  $\{\Xi_{j;t;z}^1; j=1,2,\dots\}$  and  $\{\Xi_{j;t;z}^2; j=0,1,\dots\}$ , defined by

$$\Xi_{j;t;z}^1 \stackrel{\text{def}}{=} \{\omega | \hat{t}_j(\omega; A_z) = t; \hat{t}_k(\omega; A_z) \neq t \text{ if } k \neq j\} \quad (2.160)$$

$$\Xi_{j;t;z}^2 \stackrel{\text{def}}{=} \{\omega | \hat{t}_j(\omega; A_z) < t < \hat{t}_{j+1}(\omega; A_z)\} \quad (2.161)$$

with  $\hat{t}_0(\omega; A_z) = 0$ .

Obviously, we have

$$\Xi_{j;t;z}^i \in H; i=1,2. \quad (2.162)$$

Lemma 2.36

If  $\Lambda \in \hat{H}_{t_0}$ , the  $\omega$ -sets

$$\Lambda \cap \Xi_{0;t_0;z}^2 \quad (2.163)$$

and

$$T_{(j); \omega^0 \dots \omega^{j-1}}(\Lambda \cap \Xi_{j;t_0;z}^2); j=1,2,\dots \quad (2.164)$$

are cylinder sets of the respective forms

$$(\Lambda \cap \Xi_{(t_0, \infty); A_z}) \times \prod_{h=1}^{\infty} \Omega^h \quad (2.165)$$

and

$$(\Lambda^j \cap \Xi_{(t_j, \infty)}; A_z) \times \prod_{h=1}^{\infty} \Omega^h; \quad j=1, 2, \dots, \quad (2.166)$$

where  $\Lambda^j \in \hat{F}_{t_j}^0$  (cf. p.38 with  $*=0$ ) and  $t_j^0 = t_j - \hat{t}_j(\omega; A_z)$ .

Proof:

Obviously, we have

$$\{\omega \mid \hat{x}_t(\omega; t_0) \in B\} = \begin{cases} \Lambda_{t; B; z}, & \text{if } t \leq t_0 \\ \Lambda_{t_0; B; z}, & \text{if } t > t_0 \end{cases}. \quad (2.167)$$

We first consider the case

$$\Lambda = \Lambda_{t; B_1 \times B_2; z}. \quad (2.168)$$

It can easily be verified that for  $t \leq t_0$ , we find

$$\text{a) } \Lambda_{t; B_1 \times B_2; z} \cap \Xi_{t_0; t_0; z}^2 = (\Lambda_{t; B_1 \cap B_2} \cap \Xi_{(t_0, \infty)}; A_z) \times \prod_{j=1}^{\infty} \Omega^j \quad (2.169)$$

Hence for  $t \leq t_0$  the set  $\Lambda_{t; B_1 \times B_2; z}$  satisfies the assertion;

$$\begin{aligned} \text{b) } T_{(j); \omega^0 \dots \omega^{j-1}}(\Lambda_{t; B_1 \times B_2; z} \cap \Xi_{j; t_0; z}^2) &= \\ = T_{(j); \omega^0 \dots \omega^{j-1}}(\Lambda_{t; B_1 \times B_2; z}) \cap T_{(j); \omega^0 \dots \omega^{j-1}}(\Xi_{j; t_0; z}^2) &= \end{aligned}$$

$$= \begin{cases} \emptyset, & \text{if } t < \hat{t}_j(\omega; A_z) \text{ and } \omega \in \Lambda_{t; \overline{B_1 \times B_2}; z} \text{ or if } t = \hat{t}_j(\omega; A_z) \text{ and} \\ & \omega \in \Lambda_{t; \overline{B_1 \times X_2}; z} \\ \Xi_{(t_0 - \hat{t}_j(\omega; A_z), \infty)}; A_z \times \prod_{j=1}^{\infty} \Omega^j, & \text{if } t < \hat{t}_j(\omega; A_z) \text{ and} \\ & \omega \in \Lambda_{t; B_1 \times B_2; z} \\ (\Lambda_{t_0; B_2} \cap \Xi_{(t_0 - \hat{t}_j(\omega; A_z), \infty)}; A_z) \times \prod_{j=1}^{\infty} \Omega^j, & \text{if } t = \hat{t}_j(\omega; A_z) \text{ and} \\ & \omega \in \Lambda_{t; B_1 \times X_2; z} \\ (\Lambda_{t - \hat{t}_j(\omega; A_z); B_1 \cap B_2} \cap \Xi_{(t_0 - \hat{t}_j(\omega; A_z), \infty)}; A_z) \times \prod_{j=1}^{\infty} \Omega^j, & \\ & \text{if } t_0 \geq t > \hat{t}_j(\omega; A_z). \end{cases} \quad (2.170)$$

Since the  $\omega$ -function  $\hat{t}_j(\omega; A_z)$  only depends on the components  $(\omega^0, \dots, \omega^{j-1})$ , it follows from (2.170) that sets of the type (2.168) with  $t \leq t_0$  satisfy the assertion.

Now let  $J$  be the class of  $\omega$ -sets  $\Lambda$  with the following properties:

- a)  $\Lambda \in \hat{H}_{t_0}$ ;
- b) the  $\omega$ -sets  $\Lambda$  satisfy the assertion.

The following points can easily be verified:

- a)  $\Lambda_{t; B_1 \times B_2; z} \in J$  if  $t \leq t_0$ ;
- b) if  $K \in J$ , then  $\bar{K} \in J$ ;
- c) if  $K_i \in J$  and if  $K_i \subset K_{i+1} \dots$  ( $i=1, 2, \dots$ ) then  $\bigcup_{i=1}^{\infty} K_i \in J$ .

Consequently,  $J$  is a  $\sigma$ -field that includes the sets  $\Lambda_{t; B_1 \times B_2; z}$  with  $t \leq t_0$ . Hence  $J = \hat{H}_{t_0}$ .

This ends the proof.

#### Lemma 2.37

If  $\Lambda \in \mathcal{F}_{t_1}^0$  and if  $B \in G'$ , then, under the assumptions 1 through 5, for each  $s \in [0, \infty)$  and  $x_1 \in X_1$  we have

$$\begin{aligned} & \int_{\Lambda \cap \Xi_{(t_1, \infty); A_z}} P^0 [d\omega^0; x_1] \int_{X_2} z(du; x^0(\omega^0; A_z)) P [T_{(1)}; \omega^0(\Lambda_{s+t_1}; B; z); u; z] \\ &= \int_{\Lambda \cap \Xi_{(t_1, \infty); A_z}} P^0 [d\omega^0; x_1] P [\Lambda_{s; B; z}; x_{t_1}^0(\omega^0); z]. \end{aligned} \quad (2.171)$$

#### Proof:

Let us consider the functions  $y_{t_1}(\omega_1^0)$  and  $y(\omega^0)$ , defined by (cf. (1.205) and (1.206))

$$y_{t_1}(\omega_1^0) \stackrel{\text{def}}{=} \int_{X_2} z(du; x^0(\omega_1^0; A_z)) P [T_{(1)}; \omega_1^0(\Lambda_{s; B; z}); u; z] \quad (2.172)$$

and (cf. (1.98))

$$y(\omega^0) = y_{t_1}(T_{t_1}(\omega^0)). \quad (2.173)$$

By lemmas 1.35 and 2.12

$$\begin{aligned}
& \int_{\Lambda \cap \Xi(t_1, \infty); A_Z} P^O [d\omega^O; x_1] y(\omega^O) = \\
& = \int_{\Lambda \cap \Xi(t_1, \infty); A_Z} P^O [d\omega^O; x_1] \int_{\Omega^O} P^O [d\omega_1^O; x_{t_1}^O(\omega^O)] y_{t_1}(\omega_1^O) = \\
& = \int_{\Lambda \cap \Xi(t_1, \infty); A_Z} P^O [d\omega^O; x_1] P [\Lambda_{S; B; Z}; x_{t_1}^O(\omega^O); z]. \quad (2.174)
\end{aligned}$$

If  $\omega^O \in \Xi(t_1, \infty); A_Z$  and if  $\omega_1^O = T_{t_1}(\omega^O)$ , then we can easily verify that

$$x^O(\omega^O; A_Z) = x^O(\omega_1^O; A_Z). \quad (2.175)$$

Since for  $\omega^O \in \Xi(t_1, \infty); A_Z$  and  $\omega_1^O = T_{t_1}(\omega^O)$  holds

$$t(\omega^O; A_Z) = t(\omega_1^O; A_Z) + t_1, \quad (2.176)$$

we find

$$T(1); \omega_1^O(\Lambda_{S; B; Z}) = T(1); \omega^O(\Lambda_{S+t_1; B; Z}). \quad (2.177)$$

Consequently, by (2.172), (2.173), (2.175) and (2.177),

$$\begin{aligned}
y(\omega^O) = \int_{X_2} z(du; x^O(\omega^O; A_Z)) P [T(1); \omega^O(\Lambda_{S+t_1; B; Z}); \\
; u; z]. \quad (2.178)
\end{aligned}$$

Hence, by (2.174) and (2.178)

$$\begin{aligned}
& \int_{\Lambda \cap \Xi(t_1, \infty); A_Z} P^O [d\omega^O; x_1] \int_{X_2} z(du; x^O(\omega^O; A_Z)) \\
& \quad P [T(1); \omega^O(\Lambda_{S+t_1; B; Z}); u; z] = \\
& \int_{\Lambda \cap \Xi(t_1, \infty); A_Z} P^O [d\omega^O; x_1] P [\Lambda_{S; B; Z}; x_{t_1}^O(\omega^O); z]. \quad (2.179)
\end{aligned}$$

This ends the proof.

Lemma 2.38

If  $\Lambda \in \hat{H}_{t_0}^1$ , under the assumptions 1 through 5, for each  $s \in [0, \infty)$ ,  $x \in X'$  and  $B \in G'$  we have

$$\begin{aligned} P[\Lambda \cap \Lambda_{s+t_0; B; z; x'; z}] &= \\ &= \int_{\Omega} P[d\omega; x'; z] P[\Lambda_{s; B; z; x_{t_0}(\omega); z}]. \end{aligned} \quad (2.180)$$

Proof:

It can easily be verified that for each  $x' \in X'$  and  $t_0 > 0$  we have (cf. (2.76), (2.160) and (2.161))

$$P[\bar{\cup}_{j=1}^{\infty} \bar{\equiv}^1_{j; t_0; z} \cup \bar{\cup}_{j=0}^{\infty} \bar{\equiv}^2_{j; t_0; z; x'; z}] = 1. \quad (2.181)$$

So that, by lemma 2.34,

$$\begin{aligned} P[\Lambda \cap \Lambda_{s+t_0; B; z; x'; z}] &= \\ &= P[\Lambda \cap (\bar{\cup}_{j=1}^{\infty} \bar{\equiv}^1_{j; t_0; z} \cap \bar{\cup}_{j=0}^{\infty} \bar{\equiv}^2_{j; t_0; z}) \cap \Lambda_{s+t_0; B; z; x'; z}] = \\ &= P[\Lambda \cap \bar{\equiv}^2_{0; t_0; z} \cap \Lambda_{s+t_0; B; z; x'; z}] + \\ &+ \sum_{j=1}^{\infty} \sum_{i=1}^2 \int_{\Omega} P[d\omega; x'; z] \int_{X_2} z(du; x^{j-1}(\omega; A_z)) \cdot \\ &\cdot P[T_{(j); \omega_0 \dots \omega_{j-1}}(\Lambda \cap \bar{\equiv}^i_{j; t_0; z} \cap \Lambda_{s+t_0; B; z; x'; z}); u; z]. \end{aligned} \quad (2.182)$$

We first consider the term  $P[\Lambda \cap \bar{\equiv}^2_{0; t_0; z} \cap \Lambda_{s+t_0; B; z; x'; z}]$  of the right hand side of (2.182).

By lemmas 2.36 and 2.37 we find

$$\begin{aligned}
& P [\Lambda \cap \overset{2}{\Xi}_{0;t_0;z} \cap \Lambda_{s+t_0;B;z}; x'; z] = \\
& = P [((\Lambda^0 \cap \Xi_{(t_0, \infty)}; A_z) \times \prod_{h=1}^{\infty} \Omega^h) \cap \Lambda_{s+t_0;B;z}; x'; z] = \\
& = \int_{\Lambda^0 \cap \Xi_{(t_0, \infty)}; A_z} P^0 [d\omega^0; x_1] \int_{X_2} z(du; x^0(\omega^0; A_z)) \cdot \\
& \quad \cdot P [T_{(1)}; \omega^0(\Lambda_{s+t_0;B;z}); u; z] = \\
& = \int_{\Lambda^0 \cap \Xi_{(t_0, \infty)}; A_z} P^0 [d\omega^0; x_1] P [\Lambda_{s;B;z}; x_{t_0}^0(\omega^0); z] = \\
& = \int_{\Lambda \cap \overset{2}{\Xi}_{0;t_0;z}} P [d\omega; x'; z] P [\Lambda_{s;B;z}; x_{t_0}(\omega); z]. \quad (2.183)
\end{aligned}$$

Since for  $\omega \in \overset{2}{\Xi}_{0;t_0;z}$  holds  $x_{t_0;1}(\omega) = x_{t_0;2}(\omega) \in \bar{A}_z$ , we find by means of (2.145) and (2.147)

$$P [\Lambda_{s;B;z}; x_{t_0;2}(\omega); z] = P [\Lambda_{s;B;z}; x_{t_0}(\omega); z]. \quad (2.184)$$

Hence, by (2.183) and (2.184)

$$\begin{aligned}
P [\Lambda \cap \overset{2}{\Xi}_{0;t_0;z} \cap \Lambda_{s+t_0;B;z}; x'; z] & = \int_{\Lambda \cap \overset{2}{\Xi}_{0;t_0;z}} P [d\omega; x'; z] \cdot \\
& \quad \cdot P [\Lambda_{s;B;z}; x_{t_0}(\omega); z]. \quad (2.185)
\end{aligned}$$

Next we consider the term

$$\begin{aligned}
& \int_{\Omega} P [d\omega; x'; z] \int_{X_2} z(du; x^{j-1}(\omega; A_z)) \cdot \\
& \quad \cdot P [T_{(j)}; \omega^0 \dots \omega^{j-1}(\Lambda \cap \overset{2}{\Xi}_{j;t_0;z} \cap \Lambda_{s+t_0;B;z}); u; z] \quad (2.186)
\end{aligned}$$

of the right hand side of (2.182).

By means of lemmas 2.34 and 2.36 the expression (2.186) can be rewritten in

$$\begin{aligned}
& \int_{\Omega} P [d\omega; x'; z] \int_{X_2} z(du; x^{j-1}(\omega; A_z)) \cdot \\
& \cdot \int_{\Lambda^j \cap \equiv (t_0^j; \infty); A_z} P^0 [d\omega_1^0; u] \int_{X_2} z(dv; x^0(\omega_1^0; A_z)) P [T_{(1)}; \omega_1^0(\Lambda_{s+t_0^j}; B; z); \\
& \quad ; v; z] \cdot \quad (2.187)
\end{aligned}$$

By means of lemma 2.37, we find for (2.187)

$$\begin{aligned}
& \int_{\Omega} P [d\omega; x'; z] \int_{X_2} z(du; x^{j-1}(\omega; A_z)) \cdot \\
& \cdot \int_{\Lambda^j \cap \equiv (t_0^j; \infty); A_z} P^0 [d\omega_1^0; u] \cdot P [\Lambda_{s; B; z}; x_{t_0^j}^0(\omega_1^0); z] = \\
& = \int_{\Lambda \cap \equiv \begin{smallmatrix} 2 \\ j; t_0; z \end{smallmatrix}} P [d\omega; x'; z] P [\Lambda_{s; B; z}; x_{t_0}(\omega); z] \cdot \quad (2.188)
\end{aligned}$$

Since, by (2.145) and (2.147),

$$P [\Lambda_{s; B; z}; x_{t_0}(\omega); z] = P [\Lambda_{s; B; z}; x_{t_0}^0(\omega); z] \quad (2.189)$$

for each  $\omega \in \equiv \begin{smallmatrix} 2 \\ j; t_0; z \end{smallmatrix}$ , (2.186) becomes

$$\int_{\Lambda \cap \equiv \begin{smallmatrix} 2 \\ j; t_0; z \end{smallmatrix}} P [d\omega; x'; z] P [\Lambda_{s; B; z}; x_{t_0}(\omega); z] \cdot \quad (2.190)$$

Thus,

$$\begin{aligned}
& \int_{\Omega} P [d\omega; x'; z] \int_{X_2} z(du; x^{j-1}(\omega; A_z)) \cdot \\
& \quad \cdot P [T_{(j)}; \omega^0 \dots \omega^{j-1}(\Lambda \cap \equiv \begin{smallmatrix} 2 \\ j; t_0; z \end{smallmatrix} \cap \Lambda_{s+t_0}; B; z); u; z] = \\
& = \int_{\Lambda \cap \equiv \begin{smallmatrix} 2 \\ j; t_0; z \end{smallmatrix}} P [d\omega; x'; z] P [\Lambda_{s; B; z}; x_{t_0}(\omega); z] \cdot \quad (2.191)
\end{aligned}$$



We now consider the term

$$\int_{\Omega} P [d\omega; x'; z] \int_{x_2} z(du; x^{j-1}(\omega; A_z)) \cdot \\ \cdot P [T_{(j)}; \omega^0 \dots \omega^{j-1} (\Lambda \cap \Xi_{j; t_0; z}^1 \cap \Lambda_{s+t_0; B; z}) ; \\ ; u; z] . \quad (2.192)$$

Since for each  $\omega \in \Xi_{j; t_0; z}^1$  we have  $\hat{t}_j(\omega; A_z) = t_0$ , (2.192) becomes

$$\int_{\Lambda \cap \Xi_{j; t_0; z}^1} P [d\omega; x'; z] P [T_{(j)}; \omega^0 \dots \omega^{j-1} (\Lambda_{s+t_0; B; z}); x_0^j(\omega^j); z] . \quad (2.193)$$

According to assumption 2 to each  $\omega \in \Omega$  corresponds one and only one point  $\omega_1 = (\omega_1^0, \omega_1^1, \dots) \in \Omega$ , given by

$$x_t^0(\omega_1) = x_{t+t_0-\hat{t}_{j-1}}^{j-1}(\omega; A_z) \quad (2.194)$$

and

$$x_t^k(\omega_1) = x_t^{k+j-1}(\omega); \quad k=1, 2, \dots . \quad (2.195)$$

We obviously have

$$x_0^1(\omega_1) = x_0^j(\omega) \quad (2.196)$$

and

$$T_{(j)}; \omega^0 \dots \omega^{j-1} (\Lambda_{s+t_0; B; z}) = T_{(1)}; \omega_1^0 (\Lambda_{s+t_0-\hat{t}_j(\omega; A_z); B; z}) . \quad (2.197)$$

Thus, if  $\hat{t}_j(\omega; A_z) = t_0$ ,

$$P [T_{(j)}; \omega^0 \dots \omega^{j-1} (\Lambda_{s+t_0; B; z}); x_0^j(\omega^j); z] = \\ = P [T_{(1)}; \omega_1^0 (\Lambda_s; B; z); x_0^1(\omega_1^1); z] . \quad (2.198)$$

By (2.147) and (2.148)

$$P [T_{(j)}; \omega^0 \dots \omega^{j-1} (\Lambda_{s+t_0; B; z}); x_0^j(\omega^j); z] =$$

$$\begin{aligned}
&= P \left[ T_{(1)}; \omega_1^0 (\Lambda_{s;B;z}) ; x_0^1(\omega_1^1); z \right] = P \left[ \Lambda_{s;B;z}; x_0(\omega_1); z \right] = \\
&= P \left[ \Lambda_{s;B;z}; x_{t_0}(\omega); z \right]. \quad (2.199)
\end{aligned}$$

Consequently, the expression (2.193) turns out to be equal to

$$\int_{\Lambda \cap \Xi_{j;t_0}^1} P [d\omega; x'; z] P \left[ \Lambda_{s;B;z}; x_{t_0}(\omega); z \right]. \quad (2.200)$$

Hence, by (2.192), (2.193) and (2.200)

$$\begin{aligned}
&\int_{\Omega} P [d\omega; x'; z] \int_{X_2} z(du; x^{j-1}(\omega; A_z)) \cdot \\
&\quad \cdot P \left[ T_{(j)}; \omega^0 \dots \omega^{j-1} (\Lambda \cap \Xi_{j;t_0}^1; z \cap \Lambda_{s+t_0;B;z}); \right. \\
&\quad \left. ; u; z \right] = \\
&= \int_{\Lambda \cap \Xi_{j;t_0}^1} P [d\omega; x'; z] P \left[ \Lambda_{s;B;z}; x_{t_0}(\omega); z \right]. \quad (2.201)
\end{aligned}$$

Finally, it follows from (2.185), (2.191) and (2.201) that

$$\begin{aligned}
P \left[ \Lambda \cap \Lambda_{s+t_0;B;z}; x'; z \right] &= \int_{\Lambda \cap \Xi_{0;t_0}^2} P [d\omega; x'; z] P \left[ \Lambda_{s;B;z}; x_{t_0}(\omega); z \right] + \\
&+ \sum_{i=1}^2 \sum_{j=1}^{\infty} \int_{\Lambda \cap \Xi_{j;t_0}^i} P [d\omega; x'; z] P \left[ \Lambda_{s;B;z}; x_{t_0}(\omega); z \right] = \\
&= \int_{\Lambda} P [d\omega; x'; z] P \left[ \Lambda_{s;B;z}; x_{t_0}(\omega); z \right]. \quad (2.202)
\end{aligned}$$

This ends the proof.

Let  $C$  be a closed set in  $X'$ , satisfying for each  $x' \in X'$

$$P \left[ \Xi_{[0,\infty)}; [C]; z; x'; z \right] = 1. \quad (2.203)$$

We now introduce the functions  $\{\hat{x}_t(\omega; [C]); t \in [0, \infty)\}$ , defined by

$$\hat{x}_t(\omega; [C]) \stackrel{\text{def}}{=} \begin{cases} x_t(\omega), & \text{if } t \leq t(\omega; [C]) \\ x(\omega; [C]), & \text{if } t \geq t(\omega; [C]). \end{cases} \quad (2.204)$$

The following lemma can easily be proved:

Lemma 2.39

The  $\omega$ -functions  $\{\hat{x}_t(\omega; [C]); t \in [0, \infty)\}$  are measurable with respect to  $F$ .

Let the class of  $\omega$ -sets  $\hat{H}_{[C]}$  be the smallest  $\sigma$ -field with respect to which the  $\omega$ -functions  $\{\hat{x}_t(\omega; [C]); t \in [0, \infty)\}$  are measurable.

If  $C$  is a closed set in  $X'$ , let the  $x_1$ -set  $\hat{C}$  be defined by

$$\hat{C} \stackrel{\text{def}}{=} \{x_1 \mid (x_1, x_1) \in C\}. \quad (2.205)$$

The proof of the following lemma is left to the reader.

Lemma 2.40

If  $C \in G'$ , then  $\hat{C} \in G_1$ .

Let us consider the  $\omega$ -sets  $\{\hat{\equiv}_j^1; [C]; z; j=1, 2, \dots\}$  and  $\{\hat{\equiv}_j^2; [C]; z; j=0, 1, \dots\}$ , defined by

$$\hat{\equiv}_j^1; [C]; z \stackrel{\text{def}}{=} \{\omega \mid t(\omega; [C]) = \hat{t}_j(\omega; A_z); \\ \hat{t}_k(\omega; A_z) \neq t(\omega; [C]), k \neq j\} \quad (2.206)$$

and

$$\hat{\equiv}_j^2; [C]; z \stackrel{\text{def}}{=} \{\omega \mid \hat{t}_j(\omega; A_z) < t(\omega; [C]) < \hat{t}_{j+1}(\omega; A_z)\} \quad (2.207)$$

with  $\hat{t}_0(\omega; A_z) = 0$ .

Obviously, we have

$$\hat{\equiv}_j^i; [C]; z \in H; i=1, 2 \quad (2.208)$$

Finally, the  $\omega^0$ -set  $\hat{\equiv}(t_{[C]}, \infty); A_z$ , defined by

$$\equiv (t_{[\hat{C}]}^{\infty}; A_z \stackrel{\text{def}}{=} \{\omega^0 \mid t(\omega^0; A_z) > t(\omega^0; [\hat{C}])\} \quad (2.209)$$

will be used in the coming discussion.

Clearly, (cf. p. 38)

$$\equiv (t_{[\hat{C}]}^{\infty}; A_z \in \hat{F}^0[\hat{C}]. \quad (2.210)$$

Lemma 2.41

If  $\Lambda \in \hat{H}[\hat{C}]$ , the  $\omega$ -sets

$$\Lambda \cap \equiv_0^2; [\hat{C}]; z \quad (2.111)$$

and

$$T_{(j); \omega^0 \dots \omega^{j-1}} (\Lambda \cap \equiv_j^2; [\hat{C}]; z); j=1, 2, \dots \quad (2.112)$$

are cylinder sets of the respective forms

$$(\Lambda^0 \cap \equiv (t_{[\hat{C}]}^{\infty}; A_z) \times \prod_{h=1}^{\infty} \Omega^h \quad (2.213)$$

and

$$(\Lambda^j \cap \equiv (t_{[\hat{C}]}^{\infty}; A_z) \times \prod_{h=1}^{\infty} \Omega^h; j=1, 2, \dots, \quad (2.214)$$

where  $\Lambda^j \in \hat{F}^0[\hat{C}]$ .

Proof:

The proof of this lemma is similar to that of lemma 2.36.

Let us introduce the  $\omega$ -set  $\Lambda_{s; B; [\hat{C}]; z}$ , defined by (cf. (2.131))

$$\Lambda_{s; B; [\hat{C}]; z} \stackrel{\text{def}}{=} \{\omega \mid x_s(\omega; [\hat{C}]) \in B\}. \quad (2.215)$$

Lemma 2.42

If  $\Lambda \in \hat{F}^0[\hat{C}]$  and if  $B \in G'$ , then, under the assumptions 1 through 5 and (2.203), for each  $s \in [0, \infty)$  and  $x_1 \in X_1$  we have

$$\begin{aligned}
& \int_{\Lambda \cap \Xi(t[\hat{C}], \infty); A_z} P^O [d\omega^O; x_1] \int_{X_2} z(du; x^O(\omega; A_z)) P [T_1; \omega^O(\Lambda_{s; B}; [C]); u; z] = \\
& = \int_{\Lambda \cap \Xi(t[\hat{C}], \infty); A_z} P^O [d\omega^O; x_1] P [\Lambda_{s; B; z}; x^O(\omega^O; [\hat{C}]); z] . \quad (2.216)
\end{aligned}$$

Proof:

The proof of this lemma is similar to that of lemma 2.37.

Lemma 2.43

If  $\Lambda \in \hat{H}[C]$ , under the assumptions 1 through 5 and (2.203), for each  $s \in [0, \infty)$ ,  $x' \in X'$  and  $B \in G'$  we have

$$\begin{aligned}
& P[\Lambda \cap \Lambda_{s; B; [C]}; z; x'; z] = \\
& = \int_{\Lambda} P [d\omega; x'; z] P [\Lambda_{s; B; z}; x(\omega; [C]); z] . \quad (2.217)
\end{aligned}$$

Proof:

The proof is similar to that of lemma 2.38.

#### 4. A new foundation of the decision process

In this section we shall give a formulation of the decision process which is similar to that of the fundamental stochastic process in chapter 1. Next we shall show that these two stochastic processes have nearly equal properties.

Let the class  $H_z$  be the smallest  $\sigma$ -field of  $\omega$ -sets with respect to which the  $\omega$ -functions  $\{x_t(\omega); t \in [0, \infty)\}$  are measurable.

We now introduce the  $\omega$ -set  $\bar{M}_{0; z}$ , the smallest set with the following properties:

- 1) for each  $\omega \in \bar{M}_{0; z}$ , the  $t$ -function  $x_{t; 2}(\omega)$  is continuous from the right;
- 2) in each bounded time interval in  $[0, \infty)$  and for each  $\omega \in \bar{M}_{0; z}$  the  $t$ -function  $x_t(\omega)$  has only a finite number of discontinuities.

Since  $M_{0;z} \subset M_0$  we have for each  $x_1$

$$P [M_{0;z}; x_1; z] = 0. \quad (2.218)$$

Let the class  $F_z$  be the smallest  $\sigma$ -field of  $\omega$ -sets with the following properties:

- 1)  $F_z \supset H_z$ ;
- 2)  $F_z$  contains all subsets of  $M_{0;z}$ ;
- 3) the  $\omega$ -functions  $t(\omega; C)$ ,  $t(\omega; [C])$ ,  $x(\omega; C)$  and  $x(\omega; [C])$  are measurable with respect to  $F_z$  if  $C$  is any closed set in  $X'$ .

We now consider

- 1) a space  $\Omega^z$  with points  $\omega^z$ ;
- 2) a family of  $\omega^z$ -functions  $\{x_t^z(\omega^z); t \in [0, \infty)\}$ , defined on  $\Omega^z$ , such that
  - a) for each  $t \in [0, \infty)$  the  $\omega^z$ -function  $x_t^z(\omega^z)$  maps  $\Omega^z$  into  $X'$ ;
  - b) if  $x'(t)$  is any mapping of the time axis into the state space  $X'$ , one and only one point  $\omega^z$  can be found such that

$$x_t^z(\omega^z) = x'(t); \quad t \in [0, \infty). \quad (2.219)$$

Consequently, a 1-1 correspondence exists between realizations of the decision process and points  $\omega^z \in \Omega$ .

Similar to the  $\omega$ -functions  $t(\omega; C)$ ,  $t(\omega; [C])$ ,  $x^*(\omega; C)$  and  $x^*(\omega; [C])$  in chapter 1 of this part, we can define  $\omega^z$ -functions  $t^z(\omega^z; C)$ ,  $t^z(\omega^z; [C])$ ,  $x^z(\omega^z; C)$  and  $x^z(\omega^z; [C])$ .<sup>4)</sup>

Since each point  $\omega \in \Omega$  corresponds to one and only one realization  $\{x'(t); t \in [0, \infty)\}$  of the decision process, (2.219) also defines a point transformation

$$\omega^z = T_z(\omega) \quad (2.220)$$

from  $\Omega$  onto  $\Omega^z$ .

If  $\omega^z = T_z(\omega)$ , then

---

4) In order to save confusion the  $\omega^z$ -functions  $t(\omega^z; C)$  and  $t(\omega^z; [C])$  have been indexed.

$$x_t(\omega) = x_t^Z(T_Z(\omega)), \quad (2.221)$$

$$t(\omega; C) = t^Z(T_Z(\omega); C), \quad (2.222)$$

$$t(\omega; [C]) = t^Z(T_Z(\omega); [C]), \quad (2.223)$$

$$x(\omega; C) = x^Z(T_Z(\omega); C), \quad (2.224)$$

$$\text{and } x(\omega; [C]) = x^Z(T_Z(\omega); [C]). \quad (2.225)$$

Let  $\Lambda_0^Z$  be the smallest  $\omega^Z$ -set with the following properties:

- 1) for each  $\omega^Z \in \bar{\Lambda}_0^Z$ , the t-function  $x_{t;2}^Z(\omega^Z)$  is continuous from the right;
- 2) in each bounded time interval in  $[0, \infty)$  and for each  $\omega^Z \in \bar{\Lambda}_0^Z$  the t-function  $x_t^Z(\omega^Z)$  has only a finite number of discontinuities.

Next we define the  $\omega^Z$ -sets  $\Lambda_{t;B}^Z, \Xi_{I;C}^Z, \Xi_{I;[C]}^Z, \Delta_{B;C}^Z, \Delta_{B;[C]}^Z$  and  $\Lambda_{s;B;[C]}^Z$  by

$$\Lambda_{t;B}^Z \stackrel{\text{def}}{=} \{ \omega^Z \mid x_t^Z(\omega^Z) \in B \}, \quad (2.226)$$

$$\Xi_{I;C}^Z \stackrel{\text{def}}{=} \{ \omega^Z \mid t^Z(\omega^Z; C) \in I \}, \quad (2.227)$$

$$\Xi_{I;[C]}^Z \stackrel{\text{def}}{=} \{ \omega^Z \mid t^Z(\omega^Z; [C]) \in I \}, \quad (2.228)$$

$$\Delta_{B;C}^Z \stackrel{\text{def}}{=} \{ \omega^Z \mid x^Z(\omega^Z; C) \in B \} \quad (2.229)$$

$$\Delta_{B;[C]}^Z \stackrel{\text{def}}{=} \{ \omega^Z \mid x^Z(\omega^Z; [C]) \in B \} \quad (2.230)$$

and

$$\Lambda_{s;B;[C]}^Z \stackrel{\text{def}}{=} \{ \omega^Z \mid x_s^Z(\omega^Z; [C]) \in B \}. \quad (2.231)$$

Obviously, if we define the set transformations

$$K = T_Z^{-1}(K_1) \quad (2.232)$$

and

$$K_1 = T_Z(K) \quad (2.333)$$

by

$$T_Z^{-1}(K_1) \stackrel{\text{def}}{=} \{\omega \mid \omega^Z = T_Z(\omega); \omega^Z \in K_1\} \quad (2.234)$$

and

$$T_Z(K) \stackrel{\text{def}}{=} \{\omega^Z \mid \omega^Z = T_Z(\omega); \omega \in K\} \quad (2.235)$$

respectively,

then

$$\Lambda_O^Z = T_Z(M_{O;z}), \quad (2.236)$$

$$M_{O;z} = T_Z^{-1}(\Lambda_O^Z), \quad (2.237)$$

$$\Lambda_{t;B}^Z = T_Z(\Lambda_{t;B;z}), \quad (2.238)$$

$$\Lambda_{t;B;z} = T_Z^{-1}(\Lambda_{t;B}^Z), \quad (2.239)$$

$$\Lambda_{s;B;[C]}^Z = T_Z(\Lambda_{s;B;[C];z}), \quad (2.240)$$

and

$$\Lambda_{s;B;[C];z} = T_Z^{-1}(\Lambda_{s;B;[C]}^Z). \quad (2.241)$$

Let the class  $H^Z$  be the smallest  $\sigma$ -field of  $\omega^Z$ -sets with respect to which the  $\omega^Z$ -functions  $\{x_t^Z(\omega); t \in [0, \infty)\}$  are measurable.

We now introduce  $F^Z$ , the smallest  $\sigma$ -field with the following properties:

- 1)  $F^Z \supset H^Z$ ;
- 2)  $F^Z$  contains all subsets of  $\Lambda_O^Z$ ;
- 3) the  $\omega$ -functions  $t^Z(\omega^Z; C)$ ,  $t^Z(\omega^Z; [C])$ ,  $x^Z(\omega^Z; C)$  and  $x^Z(\omega^Z; [C])$  are measurable with respect to  $F^Z$  if  $C$  is any closed set in  $X'$ .

The following lemma can easily be proved:

Lemma 2.44

The set transformation  $K = T_Z^{-1}(K_1)$  generates an isomorphism of  $F^Z$  with  $F_Z$ .

Now we are in a position to define probability measures on  $F^Z$ .

These set functions,

$$\{P^Z [K_1; x_1] ; x_1 \in X_1\}, \quad (2.242)$$



are defined on  $F^Z$  by

$$P^Z [K_1; x_1] \stackrel{\text{def}}{=} P [T_Z^{-1}(K_1); x_1; \bar{z}] . \quad (2.243)$$

Hence, the  $\omega^Z$ -functions  $\{x_t^Z(\omega^Z); t \in [0, \infty)\}$  and the probability space  $\{\Omega^Z; F^Z; P^Z[\cdot; x_1]\}$  provide us with an alternative description of the decision process in  $X'$ .

Decision processes, defined in this way, are denoted by

$$S_{x_1}^Z \equiv \{x_t^Z; x_1; t \in [0, \infty)\} . \quad (2.244)$$

We already know that, if a decision process is described by means of a set function  $P [K; x_1; \bar{z}]$ , the  $x_2$ -component of the initial state obeys an initial distribution. In section 3 we found a set function  $P [K; x'; \bar{z}]$  that describes the decision process in case the initial  $x_2$ -state has also been given.

If on  $F^Z$  the set functions

$$\{P^Z [K_1; x'] ; x' \in X'\} \quad (2.245)$$

are defined by

$$P^Z [K_1; x'] \stackrel{\text{def}}{=} P [T_Z^{-1}(K_1); x'; \bar{z}] , \quad (2.246)$$

then the  $\omega^Z$ -functions  $\{x_t^Z(\omega^Z); t \in [0, \infty)\}$  together with the probability space  $\{\Omega^Z; F^Z; P^Z[\cdot; x']\}$  generate the decision process with initial state  $x'$ .

Decision processes, defined in this way, are denoted by

$$S_{x'}^Z \equiv \{x_t^Z; x'; t \in [0, \infty)\} . \quad (2.247)$$

Finally, let us compare the fundamental stochastic processes  $\{S_x^*; x \in X^*\}$ , described in chapter 1, with the decision processes  $\{S_x^Z; x' \in X'\}$ .

It follows from lemma 2.23 and (2.221) that the decision processes  $\{S_x^Z; x' \in X'\}$  do not satisfy assumption 1 completely. (Cf. chapter 1,

p.2 ,\*=z and M=2N). 5)

In the points  $\{\hat{t}_j(\omega^z; A_z); j=1,2,\dots\}$  almost all t-functions  $\{x_t^z(\omega^z); \omega^z \in \Omega^z\}$  are not continuous from the right. Therefore the proofs of lemmas 1.1 through 1.9 do not apply to decision processes. However, in this chapter (lemma 2.22 ff.) we have demonstrated that the assertions stated in lemmas 1.1 through 1.9 remain true for these processes.

By the choice of  $\Omega^z$  the decision processes  $\{S_x^z; x' \in X'\}$  satisfy assumption 2. (Cf. chapter 1, p.17 , \*=z and M=2N). According to (2.146), (2.147), (2.238) and (2.246) assumption 3 (cf. chapter 1, p.40 , \*=z) is also fulfilled. This implies that the results obtained in chapter 1 of this part also apply to decision processes.

#### 5. Stationary strong Markovian decision processes

In this section we shall show, that if the basic probability space  $\{\Omega^0; F^0; P^0\}$  is strongly Markovian the decision processes  $\{S_x^z; x' \in X'\}$  are stationary strong Markov processes.

It follows from lemma 2.38, (2.226) and (2.246) that for each pair of non-negative values  $(t_0, s)$ ,  $B \in G'$ ,  $x' \in X'$  and  $\Lambda \in \hat{H}_{t_0}^z$  (cf. chapter 1, p.37 , \*=z)

$$P^z [\Lambda_{t_0+s}^z; B; x'] = \int_{\Lambda} P^z [d\omega^z; x'] P^z [\Lambda_s^z; B; x_{t_0}^z(\omega^z)] . \quad (2.248)$$

#### Lemma 2.45.1

If  $\Lambda \in \hat{H}_{t_0}^z$ ,  $t_0 \in [0, \infty)$ ,  $x' \in X'$  and  $B \in G'$  then, under the assumptions 1 through 5, we have for each  $K \in H_{t_0}^z$  (cf. chapter 1, p. 37, \*=z)

$$P^z [K \cap \Lambda; x'] = \int_{\Lambda} P^z [d\omega^z; x'] P^z [T_{t_0}(K); x_{t_0}^z(\omega^z)] . \quad (2.249)$$

---

5) \*=0 means: "read 0 where we wrote \*"

Proof:

Let  $J$  be the class of  $\omega^Z$ -sets  $K$  with the following properties:

- a)  $K \in \hat{H}_{t_0}^Z$ ;
- b) the sets  $K$  satisfy (2.249).

Obviously, by (2.248)

$$\Lambda_{t_0+s;B}^Z \in J; \quad s \geq 0. \quad (2.250)$$

We can easily verify that

- a)  $\Omega^Z \in J$ ;
- b) if  $K \in J$ , then  $\bar{K} \in J$ ;
- c) if  $K_i \in J$  ( $i=1,2,\dots$ ) and if  $K_i \subset K_{i+1} \subset \dots$ , then  $\bigcup_{i=1}^{\infty} K_i \in J$ .

Consequently,  $J$  is a  $\sigma$ -field that contains the sets  $\Lambda_{t_0+s;B}^Z$  with  $s \geq 0$ . Hence,  $J = \hat{H}_{t_0}^Z$ .

This ends the proof.

It follows from lemma 2.43, (2.231) and (2.246) that for each  $s \in [0, \infty)$ ,  $B \in G'$ ,  $x' \in X'$ ,  $\Lambda \in \hat{H}_{[C]}^Z$  (cf. chapter 1 p. 36,  $**=z$ ) and closed set  $C$  in  $X'$ , satisfying

$$P^Z [\equiv_{[0,\infty)}^Z; [C]; x'] = 1, \quad (2.251)$$

we have

$$\begin{aligned} P^Z [\Lambda_{s;B}^Z; [C] \cap \Lambda; x'] &= \\ &= \int_{\Lambda} P^Z [d\omega^Z; x'] P^Z [\Lambda_{s;B}^Z(\omega^Z); [C]] . \end{aligned} \quad (2.252)$$

Lemma 2.45.2

If  $\Lambda \in \hat{H}_{[C]}^Z$ ,  $x' \in X'$ ,  $B \in G'$  and  $C$  is a closed set in  $X'$ , then, under the assumptions 1 through 5 and (2.251), we have for  $K \in \hat{H}_{[C]}^Z$  (cf. chapter 1, p.36,  $**=z$ )

$$P^Z [K; x'] = \int_{\Lambda} P^Z [d\omega^Z; x'] P^Z [T_{[C]}^{*0}(K); x^Z(\omega^Z; [C])] . \quad (2.253)$$

Proof:

The proof is similar to that of lemma 2.45.1.

Lemma 2.45

Under the assumptions 1 through 5,

1) for each  $t_0 \in [0, \infty)$ ,  $K \in H_{t_0}^Z$  and  $x' \in X'$  the conditional probability measure  $P^Z [K; x' | \hat{H}_{t_0}^Z]$  can be defined by

$$P^Z [K; x' | \hat{H}_{t_0}^Z] = P^Z [T_{t_0}(K); x_{t_0}^Z(\omega^Z)] ; \quad (2.254)$$

2) for each  $x' \in X'$ , closed set  $C$  in  $X'$ , satisfying

$$P^Z [ \equiv_{[0, \infty)}^Z; [C]; x' ] = 1, \quad (2.255)$$

$K \in H_{[C]}^Z$ , the conditional probability measure  $P^Z [K; x' | \hat{H}_{[C]}^Z]$  can be defined by

$$P^Z [K; x' | \hat{H}_{[C]}^Z] = P^Z [T_{[C]}(K); x^Z(\omega^Z; [C])]. \quad (2.256)$$

Proof:

The assertions are immediate consequences of (2.249) and (2.253).

Finally, lemma 2.45 implies (cf. chapter 1 p.37):

Theorem 3

Under the assumptions 1 through 5, the decision processes  $\{S_x^Z; x' \in X'\}$  are stationary strong Markov processes.

REFERENCES

1. M. LOEVE, Probability theory. D. van Nostrand Cy., Princeton, New Jersey, 1963 (3rd. edition).
2. J.L. DOOB, Stochastic Processes. John Wiley & Sons, New York and Chapman & Hall Ltd., London, 1953.
3. R. BELLMAN, Dynamic Programming. Princeton University Press, Princeton, New Jersey, 1957.
4. R.A. HOWARD, Dynamic Programming and Markov Processes. The Technology Press of the Massachusetts Institute of Technology + John Wiley & Sons, New York, 1960.
5. E.B. DYNKIN, Die Grundlagen der Theorie der Markoffschen Prozesse. Springer Verlag, Berlin, 1961.
6. E.B. DYNKIN, Markov Processes, volume I. Springer Verlag, Berlin, 1965.
7. E.B. DYNKIN, Markov Processes, volume II. Springer Verlag, Berlin, 1965.

ERRATA AND ADDENDA  
(Part II)

page line

- 2 14 The largest "x-free" extension of  $H^*$  is the intersection of all  $\sigma$ -algebras obtained by completing  $H^*$  with respect to the measures  $P^*[\cdot; x]$ .
- 2 16 for  $\bar{\Lambda}_0^*$  be an  $\omega$ -set ...  
read  $\bar{\Lambda}_0^*$  be the smallest  $\omega$ -set ... .
- 2 19 for finite number of discontinuities.  
read finite number of jump discontinuities.
- 3 Lemmas 1.2.1, 1.2, 1.3, 1.4.1, 1.4.2, 1.4.3, 1.4 and 1.5.1 together are equivalent to the following statement:  
the moment that the system is for the first time in C is a  $F^*$ -measurable function if C is an open or closed subset of  $X^*$ .  
If C is open it follows that  

$$[t(\omega; C) < t] \cap \bar{\Lambda}_0^* = \bigcup_{\substack{r \text{ rational} \\ r < t}} [x_r^*(\omega) \in C] \cap \bar{\Lambda}_0^*$$
thus  $t(\omega; C)$  (for the definition see page 9) is  $F^*$ -measurable.  
If C is closed, define a sequence of  $\omega$ -functions  $\{t_k(\omega)\}_{k=1}^\infty$  on  $\bar{\Lambda}_0^*$  as follows:  $t_1(\omega)$  is the moment of first contact with C (cf. [1] page 580 and [6] page 105),  $t_{k+1}(\omega)$  is the moment of first contact after  $t_k(\omega)$  with C. The limit of these  $F^*$ -measurable functions is the restriction of  $t(\omega; C)$  to  $\bar{\Lambda}_0^*$  thus  $t(\omega; C)$  is  $F^*$ -measurable.
- 7 4 for the points of accumulation  $\{t_k^\alpha; \alpha=1,2,\dots\}$  of ...  
read the points of accumulation of ... .
- 7 7 for At least one of the points  $\{t_k^\alpha; \alpha=1,2,\dots\}$ ,  
say ... read At least one of these points, say ... .
- 7 9 for All points  $\{t_k^\alpha; \alpha=1,2,\dots\}$  are ...  
read All the accumulation points are ... .

page line

7 17 add From now on the accumulation points are denoted by  $\{t_k^\alpha; \alpha=1,2,\dots\}$ .

10 It is easy to see that if  $\Omega^*$  is restricted to  $\bar{\Lambda}_0^*$ , the corresponding stochastic process is right-continuous. A right-continuous process is strongly measurable, for strongly measurable processes it is known that  $x_{\tau(\omega)}(\omega)$  is measurable if  $\tau(\omega)$  is measurable. (Cf. [6] page 98 and [1] page 579.)

Each of the lemmas 1.5.2, 1.6, second part of 1.7, 1.8.2 and 1.9 is a direct consequence of this statement. As example we prove lemma 1.6.

$$\tau(\omega) \stackrel{\text{def.}}{=} \begin{cases} t(\omega;B) + t & \text{if } t(\omega;B) < \infty \\ t & \text{if } t(\omega;B) = \infty. \end{cases}$$

It follows that  $\tau(\omega)$  is  $(F^* \cap \bar{\Lambda}_0^*)$ -measurable and consequently  $x_{\tau(\omega)}(\omega) = x_{t(\omega;B)}^*(\omega;B)$  is  $(F^* \cap \bar{\Lambda}_0^*)$ -measurable on  $\bar{\Lambda}_0^*$ . This implies that  $x_t^*(\omega;B)$  is  $F^*$ -measurable on  $\Omega^*$ .

10 25 for  $\omega \in \Lambda_0^* \cup \Xi[0,\infty);C$  read  $\omega \in \Lambda_0^* \cup \overline{\Xi[0,\infty);C}$ .

13 7 for  $\chi_k(\omega)$  read  $\chi_{(m);k}(\omega)$

14,15 For an  $\omega$  with  $\begin{cases} x_t^*(\omega) \in C & \text{if } t \leq t_0 \\ x_t^*(\omega) \in B_n & \text{if } t > t_0 \text{ for some } n, \end{cases}$

it follows that  $t_\infty(\omega) = t_0$ . However we should expect  $t(\omega;[C]) = \infty$ . We define

$$t(\omega;[C]) = \inf \{t | x_t^*(\omega) \in C \text{ and } (\exists t_1 < t)(x_{t_1}^*(\omega) \in \bar{C})\}.$$

The following additions are necessary in the proof of lemma 1.8.1.

page line

page line

14 27 for  $t(\omega; [C]) \leq t_n(\omega)$   
read for  $\omega \in \overline{\Lambda}_0^*$   $t(\omega; [C]) \leq t_n(\omega)$ .

15 8 for It follows ... read If  $t(\omega; [C]) < \infty$   
it follows ... .

15 14 for  $t(\omega; [C]) = t(\omega; B_n) + t(\omega; B_n; C) + \delta$   
read  $t(\omega; [C]) \geq t(\omega; B_n) + t(\omega; B_n; C) - \delta$ .

15 16 for  $t(\omega; [C]) = t_\infty(\omega)$   
read for  $\omega \in \overline{\Lambda}_0^*$   $t(\omega; [C]) = t_\infty(\omega)$ .

16 9 for Since ... read Since for  $\omega \in \overline{\Lambda}_0^*$  ... .

16 12 for  $x^*(\omega; [C]) = x_{(n)}^*(\omega)$   
read  $x^*(\omega; [C]) = \lim_{n \rightarrow \infty} x_{(n)}^*(\omega)$ .

17 2 for  $t(\omega; [C])$  and  $x(\omega; [C])$   
read  $t(\omega; [C])$  and  $x^*(\omega; [C])$ .

17 14 for condition (1.86) ... read condition (1.91) ... .

20 22 for trnasformation ... read transformation ... .

21 18 for  $t(\omega''; [C]) > 0$  ... read  $0 < t(\omega''; [C]) < \infty$  and  
 $t(\omega''; \overline{C}) < t(\omega''; [C])$ .

23 1 for  $T_{[C]}^1(\omega) = T_{[C]}(\omega)$  ... read  $T_{[C]}^0(\omega) = \omega$ ,  
 $T_{[C]}^1(\omega) = T_{[C]}(\omega)$  ... .

23 18 omit If  $K \in F^*$ , then  $T_{[C]}^j(K) \in F^*$ .

24 7) for  $x_j(\omega; [C])$  read  $x_j^*(\omega; [C])$ .  
24 10)

25 7 for finite number of discontinuities ...  
read finite number of jump discontinuities ... .

31 8 for  $k(\omega; t(\omega; B))$  read  $k_{\text{cont}}(\omega; t(\omega; B))$ .

31 12 for  $k(\omega; t(\omega; [C]))$  read  $k_{\text{cont}}(\omega; t(\omega; [C]))$ .

31 22 for  $k(\omega; \frac{j}{n})$  read  $k_{\text{cont}}(\omega; \frac{j}{n})$ .

35 5 for  $P^*[T_{[C]}^{-j+1}(k_I; [C]); x]$   
read  $P^*[T_{[C]}^{-j+1}(K_I; [C]); x]$ .



page line

- 35 11 for  $k[C];x;j \dots$  read  $k[C];x;j \dots$
- 36 16 for Let  $\hat{y}(\omega)$  be a measurable ( $F^*$ ) and ...  
read Let  $\hat{y}(\omega)$  be a measurable ( $F_1^*$ ) and ...
- 36 17 for for some  $K \in F_1^*$  ... read for some  $K \in F^*$ .
- 36 19 for relative to  $F^*$ , ... read relative to  $F_1^*$ , ...
- 37 If we compare the definitions of a stationary strong Markov process given on this page with those given in [1] and [6], we will find:
- i) Assumption 3 (1.221) has to be added to the defining relations of a (regular) Markov process.
- ii) For (1.198) we find in [1]:  
for each  $t_0 \geq 0$ , for each  $K \in H^*$  and for each  $x \in X^*$   
 $P^*[T_{t_0}^{-1}K; x | \hat{H}_{t_0}^*] = P^*[K; x_{t_0}^*(\omega)] P^*[\cdot; x]$  almost sure.  
From  $T_{t_0}^{-1}H^* = H_{t_0}^*$  and  $T_{t_0} \circ T_{t_0}^{-1}K = K$  it follows that the two relations are equivalent.
- iii) It is not true that  $T[C] \circ T[C]^{-1}K = K$  for each  $K \in H^*$ . Consequently (1.200) is not equivalent to:  
for each  $K \in H^*$  and  $x \in X^*$   
 $P^*[T[C]^{-1}K; x | \hat{H}[C]] = P^*[K; x^*(\omega; [C])] P^*[\cdot; x]$  almost sure.  
However this relation must be used on page 53.
- iv)  $x^*(\omega; [C])$  is not  $\hat{H}[C]$ -measurable and therefore (1.200) is not correct.  $x^*(\omega; [C])$  is measurable on  $(\mathbb{E}[0, \infty); [C], \hat{H}[C] \cap \mathbb{E}[0, \infty); [C])$ . If we restrict  $\Omega^*$  to  $\mathbb{E}[0, \infty); [C] = [t(\omega; [C]) < \infty]$ , definitions like (1.84) and (1.94) are superfluous because:  
for each  $K \in H^*$  and each  $x \in X^*$  (3.1)  
 $P^*[T[C]^{-1}K; x | \hat{H}[C]] = P^*[K; x^*(\omega; [C])] (\mathbb{E}[0, \infty); [C], P^*[\cdot; x])$   
almost sure  
is equivalent to  
for each  $t \geq 0$  and each  $B \in G^*$   
 $P^*[[x_{t(\omega; [C])}^* \in B] | \hat{H}[C]] = P^*[[x_t^*(\omega) \in B] | x_{t(\omega; [C])}^*(\omega)]$   
 $(\mathbb{E}[0, \infty); [C], P^*[\cdot; x])$  almost sure.

page line

v) According to [1]  $t(\omega; [C])$  must be a Markov time.

$t(\omega; [C])$  is a Markov time if:

a) for each  $t \geq 0$

$x_{t(\omega; [C]) + t}^*$  is measurable;

b) for each  $t \geq 0$

$[t(\omega; [C]) \leq t] \in \hat{H}_t^*$ .

As a consequence of assumption 1 every  $t(\omega; [C])$

fulfils a).

However b) is not true for every  $t(\omega; [C])$ .

If it is assumed that  $t(\omega; [C])$  also fulfils b) and

$(\Omega^*, H^*)$  is restricted to  $(\bar{\Lambda}_0^*, H^* \cap \bar{\Lambda}_0^*)$ , it can be proved that  $\hat{H}_{[C]}^* \cap \bar{\Lambda}_0^* = \mathcal{B}(x_s^* \mid s \leq t(\omega; [C]))$ .<sup>1</sup>

The conclusion is that if for  $t(\omega; [C])$  b) is true (3.1)

is equivalent with the strong Markov property for  $t(\omega; [C])$

as defined in [1].

vi) The requirement that (3.1) is also true for a  $t(\omega; [C])$

not satisfying b) will strongly restrict the class of

admissible Markov processes. For an example the reader is

referred to [5] page 118. Defining  $C = \{\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots\} \cup$

$\cup \{0, 1, 2, \dots\}$   $t(\omega; [C])$  will equal the function  $\tau(\omega)$

defined there. On the measurable subspace

$(\bar{\Lambda}_0^* \cap [t(\omega; \bar{C}) < t(\omega; [C])], H^* \cap (\bar{\Lambda}_0^* \cap [t(\omega; \bar{C}) < t(\omega; [C])]))$

$t(\omega; [C])$  will satisfy b). Consequently, if

$P^*[[t(\omega; \bar{C}) < t(\omega; [C])]; x] = 1$  for each  $x \in X^*$  (3.2)

then (3.1) is equivalent with the strong Markov property

for  $t(\omega; [C])$  as defined in [1].

There are several conditions implying (3.2), e.g. (1.185).

vii) Summarizing a stationary strong Markov process might

be defined as a stochastic process with the properties:

<sup>1</sup>. For the definition of  $\mathcal{B}(x_s^* \mid s \leq t(\omega; [C]))$  see [1] page 580.

page line

1) it is a stationary Markov process according to [1]  
 2) for each  $t(\omega; [C])$  satisfying (1.199), (3.2) and a)  
 the strong Markov property as defined in [1] is  
 fulfilled.

38 22 for  $\Lambda \in \hat{F}_{t_0}$  ... read  $\Lambda \in \tilde{F}_{t_0}^*$  ... .

39 33 for  $\lim_{n \rightarrow \infty}$  ... read  $\lim_{m \rightarrow \infty}$  ... .

40 14 for  $\Lambda \in \hat{F}[C]$  ... read  $\Lambda \in \tilde{F}[C]^*$  ... .

40 16 for  $\int_{\Lambda} P^*[d\omega; x]$   $\int_{\Omega^*} P^*[d\omega_1; x^*(\omega; [C])] y[C](\omega_1)$ .  
read  $\int_{\Lambda} P^*[d\omega; x]$   $\int_{T[C]\Omega^*} P^*[d\omega_1; x^*(\omega; [C])] y[C](\omega_1)$ .

40 22 for  $P^*[\Lambda_0; x] = 1$  read  $P^*[\Lambda_0; \{x\}; x] = 1$ ,  
 where  $\{x\}$  denotes the set consisting of  $x$  only.

41 13 for that for each  $j \geq 1$ ,  $x \in X^*$ , ...  
read that for each  $x \in X^*$ , ... .

41 15 for  $\int_{\Lambda} P^*[d\omega; x] P^*[T_{t_0}(K); x_1^*(\omega; t_0)]$   
read  $\int_{\Lambda} P^*[d\omega; x] P^*[T_{t_0}(K); x_{t_0}^*(\omega)]$ .

42 5 for with  $Q(X^*) = 0$ , ... read with  $Q(X^*) > 0$ , ... .

43 1 define:  $x_j(\omega; t_0) = x_{jt_0}^*(\omega)$   $j = 1, 2, 3, \dots$ .  
 From the definition of  $y_j(\omega; t_0)$   $j = 1, 2, 3, \dots$  it follows  
 that  $\Omega^*$  is restricted to  $\bigcap_{j=1}^{\infty} [n_j(\omega; t_0; [A]) < \infty]$ .

43 4 add In order to avoid needless repetitions in the  
 argumentation we use the product space  $X^* \in \Gamma$  with  
 $\Gamma \equiv (-\infty, +\infty)$  instead of  $\Gamma \equiv \{0, 1, \dots\}$ .

44 1 for  $O_{k;U} \in F_{(k-1)t_0}$  read  $O_{k;U} \in F_{(k-1)t_0}^*$ .

44 3 for  $P^*[O_1; U; x]$  read  $P^*[O_1; U; x]$ .

44 5 for (1.226) ... read (1.226) if we take  $(k-1)t_0$  for  $t_0$  ... .

page line

44 7 for  $\int_{X^*} p_{t_0}^{k-1}(dx_1; x) P^*[O_1; U; \bar{x}_{k-1}^*(\omega; t_0)]$   
read  $\int_{X^*} v_{t_0}^{k-1}(dx_1; x) P^*[O_1; U; \bar{x}_1]$ .

44 14 for  $\int_{X^*} "p_{t_0}^{k-1}(dy_1; y) "p^1(U; y_1)$ .  
read  $\int_{X^* \times \Gamma} "p_{t_0}^{k-1}(dy_1; y) "p^1(U; y_1)$ .

46 2 for  $\int_{X^*} "p_{t_0}^{\infty c_i + j - 1}(dy_1; y) "p_{t_0}^1(U; y_1)$ .  
read  $\int_{X^* \times \Gamma} "p_{t_0}^{\infty c_i + j - 1}(dy_1; y) "p_{t_0}^1(U; y_1)$ .

46 15 for  $\sum_{r=1}^{2^h} \frac{r}{2^h} |p_{t_0}^{\infty c_i + j - 1}(B_{hr}^U; y) - p_{t_0}^{nc_i + j - 1}(B_{hr}^U; y)| \dots$   
read  $\sum_{r=1}^{2^h} \frac{r}{2^h} |p_{t_0}^{\infty c_i + j - 1}(B_{hr}^U; x) - p_{t_0}^{nc_i + j - 1}(B_{hr}^U; x)| \dots$ .

47 3 for  $|"p_{t_0}^{\infty c_i + j}(U; x) - "p_{t_0}^{nc_i + j}(U; x)| \leq \eta$ .  
read  $|"p_{t_0}^{\infty c_i + j}(U; y) - "p_{t_0}^{nc_i + j}(U; y)| \leq \eta$ .

49 18 for  $F_I \stackrel{\text{def.}}{=} \{\omega | f(y(\omega; t_0)) \in I\}$   
read  $F_I \stackrel{\text{def.}}{=} \{\omega | f(y_1(\omega; t_0)) \in I\}$ .

50 4) for is equal to ... read is almost surely equal to ... .  
 50 13

50 18 for represents the losses ... read represents almost surely the losses ... .

50 21 define  $K_{I; t_0} = \{\omega | k_1(\omega; t_0) \in I\}$ .

51 6 for is equal to ... read is almost surely equal to ... .

51 23 for  $n(T; t_0) = \left\lfloor \frac{T}{t_0} \right\rfloor$ . read  $n(T; t_0) = \left\lfloor \frac{T}{t_0} \right\rfloor$ .

page line

52 15 omit again.for 53-7 to 53-15read Assume that for  $t(\omega; [C])$  the following strong Markov property is true.

For each  $x \in X^*$ , for each  $\Lambda \in \hat{F}_{[C]}^*$  and each  $k \in F^*$

$$P^*[\Lambda \cap T_{[C]}^{-1} K; x] = \int_{\Lambda} P^*[d\omega; x] P^*_{[K; x_1^*](\omega; [C])}. \quad (1.283)$$
Then it follows that for  $j \geq 2$ 

$$\begin{aligned} p_{[C]}^j(B; x) &= P^*[T_{[C]}^{-j+1}(\Delta_B; [C]); x] \\ &= \int_{\Omega^*} P^*[d\omega; x] P^*[T_{[C]}^{j-2}(\Delta_B; [C]); x_1^*(\omega; [C])] \\ &= \int_{X^*} P^*[C](dx_1; x) p_{[C]}^{j-1}(B; x_1). \end{aligned} \quad (1.285)$$

54 12 From the definition of  $\Gamma$  it follows that  $\Omega^*$  is restricted to  $\bigcap_{j=1}^{\infty} [t_j(\omega; [C]) < \infty] = \{\omega \mid \lim_{T \rightarrow \infty} n(T; \omega) = \infty\}$  (cf. page 57 line 20).

From (1.281) it follows that for each  $x \in X^*$

$$P^*[\{\omega \mid \lim_{T \rightarrow \infty} n(T; \omega) = \infty\}; x] = 1.$$

55 4 remark  $E_I; [C] = \{\omega \mid t_1(\omega; [C]) \in I\}$ .

55 9 for is equal to ... read is almost surely equal to ... .

55 19 for  $\{x_{[C]}; x; j; j=1, 2, \dots\}$

read  $\{x_{[C]}^*; x; j; j=1, 2, \dots\}$ .

56 7 for is equal to ... read is almost surely equal to ... .

56 12 for If (1.290) and (1.293) hold, the  $\omega$ -function  $k_j(\omega; [C])$  represents ... read If (1.186) and (1.187) hold, the  $\omega$ -function  $k_j(\omega; [C])$  almost surely represents ... .

56 15 define  $K_I; [C] = \{\omega \mid k_1(\omega; [C]) \in I\}$ .

57 1) for is equal to ... read is almost surely equal to ... .

57 10

page line

- 58 9 for of the lemmas 1.56 and 1.53 ...  
read of the lemmas 1.53, 1.54 and 1.56 ... .
- 59 7 for Let  $\Lambda_0$  be an  $\omega^0$ -set ...  
read Let  $\Lambda_0$  be the smallest  $\omega^0$ -set ... .
- 59 10) for finite number of discontinuities  
64 9 read finite numbers of jump discontinuities.
- 60 3 for (cf. p.1) read (cf. p.40).
- 60 4 for  $P^0[\Lambda_0; x] = 1$  read  $P^0[\Lambda_0; \{x\}; x] = 1$ ,  
where  $\{x\}$  denotes the set containing the single  
point  $x$ .
- 61 24 for  $\omega^k = T_{kh}(\omega^h); k, h=1, 2, \dots$  (in (2.10))  
read  $\omega^k = T_{kh}(\omega^h); k, h=0, 1, 2, \dots$  (2.10).
- 62 9 for a)  $K^k = T_{kk}(K), k=0, 1, \dots;$   
read a)  $K^k = T_{kk}(K^k), k=0, 1, \dots;$ .
- 63 18 for the stochastic process  $S_x^k (k=0, 1, \dots)$   
read the stochastic process  $S_x^k (k=1, 2, \dots)$ .
- 65 18 for  $T_{(k)}(K \cap \{\omega^0\} \times \dots \times \{\omega^{k-1}\} \times \prod_{i=k}^{\infty} \Omega_i)$   
read  $T_{(k)}(K \cap \{\omega^0\} \times \dots \times \{\omega^{k-1}\} \times \prod_{i=k}^{\infty} \Omega_i)$ .
- 67 18 for  $\int P^k[K_k; \omega^0 \dots \omega^{k-1}]$  read  $P^k[K_k; \omega^0 \dots \omega^{k-1}]$ .
- 68 8 for  $\int_{\Omega^{n-1}} P^{n-1}[M_{j;m}; \omega^0 \dots \omega^{n-2}] = \dots$   
read  $P^{n-1}[M_{j;m}; \omega^0 \dots \omega^{n-2}] = \dots$ .
- 69 16 for  $K \in F \dots$  read  $K \in H \dots$ .
- 69 21 for  $P^k[K^k; x; z]$  read  $P^k[K_k; x; z]$ .

page line

70 1 for  $\int_{\Omega^k} P^k[K_k; \omega^0 \dots \omega^{k-1}] \dots$   
read  $P^k[K_k; \omega^0 \dots \omega^{k-1}] \dots$

72 omit from page 72 line 1 and line 2.

72 11 for  $\int_X z(dx_1; x^{j-1}(\omega^{j-1}; A_z)) \dots$   
read  $\int_X z(dx_1; x^{j-1}(\omega; A_z)) \dots$

72 19 for  $P[M^0; x; z] = 0$  read  $P[M_0; x; z] = 0$ .

74 7 for  $p^j(B; x; z)$   
read  $\int_{A_z} p^0(dx_1; x; z) p^j(B; x_1; z)$ .

75 10 for is equal to ... read is almost surely equal to ...

75 11 let  $\Xi_{I; A_z}$  be  $\{\omega \mid t(\omega^0; A_z) \in I\}$ .

75 20 for  $\Lambda_{0; x; \bar{x}}$  read  $\Lambda_{0; \{x\}; \bar{x}}$ .  
 22 }  
 76 1 }

78 9-25 for k read h.

79 24 for  $\sigma$ -fields ... read  $\sigma$ -field ...

80 20 for finite number of discontinuities ...  
 21 }  
 81 15 read finite number of jump discontinuities ...

86 20 for if  $t(\omega; B) + t = \hat{t}_k(\omega; A_z) > \dots$   
 27 read if  $\infty > t(\omega; B) + t =$   

$$= \hat{t}_k(\omega; A_z) \begin{cases} \geq 0 & k=1 \\ > \hat{t}_{k-1}(\omega; A_z) & k \geq 2. \end{cases}$$

page line

86 31 for  $\{\chi_0(\omega) v_t(\omega; B) + \dots$   
read  $\{(1 - \sum_{k=1}^n \chi_k(\omega)) v_t(\omega; B) + \dots .$

88 16 for  $v(\omega; [C]),$  if  $t(\omega; [C]) \neq \hat{t}_k(\omega; A_z) \dots$   
read  $v(\omega; [C]),$  if  $t(\omega; [C]) = \infty$  or if  
 $t(\omega; [C]) \neq \hat{t}_k(\omega; A_z) \dots .$

88 18 for  $x_k(\omega; A_z \times X_2),$  if  $t(\omega; [C]) = \hat{t}_k(\omega; A_z) \dots$   
read  $x_k(\omega; A_z \times X_2),$  if  $\infty > t(\omega; [C]) =$   
 $= \hat{t}_k(\omega; A_z) \begin{cases} \geq 0 & k=1 \\ > \hat{t}_{k-1}(\omega; A_z) & k \geq 2. \end{cases}$

102 6 for  $\int_{\Omega} P[d\omega; x'; z] P[\Lambda_{s; B; z}; x_{t_0}(\omega); z]$   
read  $\int_{\Lambda} P[d\omega; x'; z] P[\Lambda_{s; B; z}; x_{t_0}(\omega); z].$

102 13 for  $(\bigcup_{j=1}^{\infty} \Xi_j^1; t_0; z \cap \bigcup_{j=0}^{\infty} \Xi_j^2; t_0; z)$   
read  $(\bigcup_{j=1}^{\infty} \Xi_j^1; t_0; z \cup \bigcup_{j=0}^{\infty} \Xi_j^2; t_0; z).$

105 7 for to each  $\omega \in \Omega$  corresponds ...  
read to each  $\omega,$  satisfying  $\hat{t}_{j-1}(\omega; A_z) \leq t_0,$   
corresponds ... .

107 24 for H; read F;

109 24 } for finite number of discontinuities.  
111 11 } read finite number of jump discontinuities.

114 20 for  $P^Z[\Lambda_{t_0+s; B}^Z; x']$  read  $P^Z[\Lambda_{t_0+s; B}^Z \cap \Lambda; x'].$

115 12 for  $J = \hat{H}_{t_0}^Z$  read  $J = H_{t_0}^Z.$



page line

115 23 for  $K \in H^Z_{[C]}$  read  $K \in H^Z$ .

116 1 for  $P^Z[K; x'] = \int_{\Lambda} P^Z[d\omega^Z; x'] P^Z[T_{[C]}(K); x^Z(\omega^Z; [C])]$ .  
read  $P^Z[T_{[C]}^{-1}K \cap \Lambda; x'] = \int_{\Lambda} P^Z[d\omega^Z; x'] P^Z[K; x^Z(\omega^Z; [C])]$ .

116 9 to 116 13 cf. Errata and addenda for page 37.

LIST OF SYMBOLS

The numbers refer to the pages, where the symbol in question is used.

$B_n$	4	$H^Z$	112
$B_n^*$	4	$H^Z[C]$	115
$\hat{C}$	107	$H_{t_0}^Z$	114
$F$	72	$H^*$	1
$F_Z$	110	$H^*[C]$	36
$F^k$	61	$H_{t_0}^*$	36
$F^Z$	112	$H^0$	59
$F^{**}$	2	$\hat{H}_0$	90
$F_{t_0}^{**}$	18	$\hat{H}[C]$	107
$F^*[C]$	22	$\hat{H}_t$	98
$F^{**}[C]$	22	$\hat{H}^Z[C]$	115
$\hat{F}^{**}[C]$	38	$\hat{H}_{t_0}^Z$	114
$\hat{F}_{t_0}^{**}$	38	$\hat{H}^{**}[C]$	36
$\hat{F}_{t_0}^0$	99	$\hat{H}_{t_0}^{**}$	36
$G_1$	79	$I_K(\omega^0 \dots \omega^k)$	69
$G_2$	79	$\underline{I}_k$	61, 74
$G^*$	1	$K_{I;B}$	33
$G'$	79	$K_{I;t_0}$	33
$H$	64	$K_{I;[C]}$	33
$H_Z$	94		

$K_k^c$	69	$M_{I;B}$	3
$k(\omega; B)$	32	$N_{n;B}$	33
$k(\omega; [C])$	32	$N_{n;[C]}$	33
$k(\omega; t_0)$	32	$N_{n;t_0}$	33
$k_{\text{cont}}(\omega; B)$	31	$n(\omega; B; [A])$	28
$k_{\text{cont}}(\omega; [C])$	31	$n(\omega; [C]; [A])$	29
$k_{\text{cont}}(\omega; s)$	30	$n(\omega; t_0; [A])$	26
$k_{\text{disc}}(\omega; B)$	28	$n_j(\omega; [C]; [A])$	34, 55
$k_{\text{disc}}(\omega; [C])$	29	$n_j(\omega; t_0; [A])$	34
$k_{\text{disc}}(\omega; t_0)$	26	$\bar{n}_{B;x}$	33
$k_j(\omega; [C])$	34	$\bar{n}[C];x$	33
$k_j(\omega; t_0)$	34	$\bar{n}[C];x;j$	34
$\frac{k}{B;x}$	32	$\bar{n}_{t_0;x}$	33
$\frac{k}{[C];x}$	32	$\bar{n}_{t_0;x;j}$	34
$\frac{k}{[C];x;j}$	34	$\bar{\bar{n}}[C];x$	55
$\frac{k}{t_0;x}$	32	$\bar{\bar{n}}_{t_0;x}$	49
$\frac{k}{t_0;x;j}$	34	$O_k;U$	43
$k_T(\omega)$	41	$P[K;x;z]$	69
$M_0$	72	$P^0[K;x]$	59
$M_{0;z}$	109	$P^k[K;x;z]$	63

$P^k[K_k; \omega^0 \dots \omega^{k-1}]$	67	$T_{kh}$	61
$P^Z[K; x_1]$	113	$T_{k; \omega^0 \dots \omega^{k-1}}$	65
$P^Z[K; x']$	113	$T[C]$	20
$P^*[K; x]$	1	$T_{t_0}$	18
$P^*[K; x_1   F_1^*]$	36	$T_z$	110
$p(U; I; z)$	74	$T^j[C](\omega)$	22
$p[C](B; x)$	54	$t(\omega; C)$	9
$p_{t_0}(C; x)$	42	$t(\omega; B; C)$	13, 87
$p^j(B; x; z)$	60	$t(\omega; [C])$	14, 87
$p^j[C](B; x)$	53	$t_j(\omega; [C])$	23
$p_{t_0}^j(B; x)$	41	$\underline{t}_j$	75
$'p_{t_0}^k(U; x)$	44	$\underline{t}_{C; x}$	11
$"p_{t_0}^k(U; y)$	44	$\underline{t}_{C; x_1}$	85
$q(B; x; z)$	63	$\underline{t}[C]; x$	17
$S_{z; x}$	87	$\underline{t}[C]; x_1$	89
$S_x^k$	63	$\underline{t}[C]; x; j$	24
$S_{x_1}^Z$	113	$t^j(\omega; A_z)$	73
$S_x^*$	1	$t^Z(\omega^Z; C)$	110
$S_x^0$	59	$t^Z(\omega^Z; [C])$	110
$T(k)$	65	$\hat{t}_k(\omega; A_z)$	77

$\hat{t}_n(\omega; [A])$	26	$x(\omega; [C])$	88	88
$\hat{t}_j$	79	$x_j(\omega; A_z \times X_2)$		83
$\hat{t}[C]; x; n$	35	$x_j(\omega; t_0)$		43
$'t(\omega; C)$	83	$x_t(\omega; B)$		86
$''t(\omega; C)$	83	$x_t(\omega; [C])$		89
$u_t^k(\omega)$	80	$x_{t;1}(\omega)$		80
$u_{t;1}^k(\omega)$	77	$x_{t;2}(\omega)$		80
$u_{t;2}^k(\omega)$	77	$\underline{x}_{C;x_1}$		85
$\underline{u}_{t;x;1}^k$	79	$\underline{x}[C]; x_1$		89
$\underline{u}_{t;x;2}^k$	79	$\underline{x}_{t;x}$		81
$v(\omega; C)$	84	$\underline{x}_{t;x';z}$		94
$v(\omega; [C])$	88	$x_t^0(\omega^0)$		59
$v_t(\omega)$	82	$x_t^k(\omega)$		70
$v_t(\omega; B)$	86	$x^j(\omega; A_z)$		73
$X_1$	79	$x^z(\omega^z; C)$		110
$X_2$	79	$x^z(\omega^z; [C])$		110
$X'$	79	$x_t^z(\omega^z)$		110
$X^*$	1	$x^*(t)$		17
$x$	1	$x^*(\omega; B; C)$		13
$x(\omega; B; C)$	87	$x^*(\omega; z)$		10
$x(\omega; C)$	84	$x^*(\omega; [C])$		15

$x_j^*(\omega; [C])$	24	$y_{t_0; x; k}$	45
$x_t^*(\omega)$	1	$z(B; x)$	60
$x_t^*(\omega; B)$	12	$\gamma_c$	25
$x_t^*(\omega; [C])$	17	$\gamma_{cont}(x)$	25
$x_t^*(\omega; t_0)$	18	$\gamma_d$	25
$\underline{x}_{C; x}$	11	$\gamma_{disc}(x)$	25
$\underline{x}[C]; x$	17	$\Delta_{B; C}$	10
$\underline{x}[C]; x; j$	24, 52	$\Delta_{B; [C]}$	16
$\underline{x}_t; x_0$	1	$\Delta_{B; C; z}$	85
$\underline{x}_{t_0}; x; j$	41	$\Delta_{B; [C]; z}$	88
$x_t^{***}(\omega)$	29	$\Delta_{B; C}^z$	111
$x'$	80	$\Delta_{B; [C]}^z$	111
$\hat{x}_t(\omega; [C])$	107	$\Lambda_0$	59
$\hat{x}_t(\omega; t_0)$	98	$\Lambda_{I; B}$	4
$\hat{x}_t^*(\omega; [C])$	35	$\Lambda_{s; B; [C]; z}$	108
$\hat{x}_t^*(\omega; t_0)$	35	$\Lambda_{t; A}$	1, 63
$y$	44	$\Lambda_{t; B; z}$	85
$y_j(\omega; [C])$	54	$\Lambda_0^z$	111
$y_j(\omega; t_0)$	43	$\Lambda_{s; B; [C]}^z$	111
$\underline{y}[C]; x; k$	54	$\Lambda_{t; B}^z$	111

$\Lambda_0^*$	2	$\Omega^*$	1
$\mu(K; \omega)$	90	$\Omega^*[C]$	21
$E_{B; [A]; n}$	28	$\{\Omega; H; P\}$	70
$E[C]; [A]; n$	29	$\{\Omega^0; F^0; P^0\}$	59
$E_{I; C}$	9	$\{\Omega^k; F^k; P^k\}$	63
$E_{I; [C]}$	16	$\{\Omega^z; F^z; P^z[.; x_1]\}$	113
$E_{I; [C]; z}$	88	$\{\Omega^*, H^*, P^*\}$	2
$E_{t; B; z}$	85	$\omega$	1, 65
$E_{t_0; [A]; n}$	26	$\omega^0$	59
$E(t[\hat{C}], \infty); A_z$	108	$\omega^k$	61
$E_J^1; [C]; z$	107	$\omega^z$	110
$E_J^2; [C]; z$	107		
$E_J^1; t; z$	98		
$E_J^2; t; z$	98		
$E_{I; C}^z$	111		
$E_{I; [C]}^z$	111		
$\Omega$	64		
$\Omega^k$	61		
$\Omega^z$	110		

