



*Printed at the Mathematical Centre, 413 Kruislaan, Amsterdam.*

*The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O.).*

MATHEMATICAL CENTRE TRACTS 58

---

W. ALBERS

**ASYMPTOTIC EXPANSIONS  
AND THE DEFICIENCY CONCEPT  
IN STATISTICS**

Second edition

---

MATHEMATISCH CENTRUM

AMSTERDAM 1982

---

1980 Mathematics subject classification: 62G10, 62G20

---

ISBN 90 6196 102 5

First printing: 1974  
Second edition: 1982

## CONTENTS

1. INTRODUCTION	1
2. THE OPTIMAL RATIO OF SAMPLE SIZES FOR COMPARING TWO DISTRIBUTIONS	8
2.1. Introduction	8
2.2. Preliminaries	9
2.3. An expansion for the unconditional power	21
2.4. Optimal ratio and deficiencies	26
2.5. Comparison of sampling rules for Bernoulli experiments	29
3. ASYMPTOTIC EXPANSIONS FOR NONPARAMETRIC TESTS FOR THE ONE SAMPLE PROBLEM	31
3.1. Introduction	31
3.2. The basic expansion	32
3.3. Contiguous alternatives	40
3.4. Exact and approximate scores and contiguous location alternatives	47
3.5. The sign test	55
4. DEFICIENCIES BETWEEN VARIOUS TESTS FOR THE ONE SAMPLE PROBLEM	59
4.1. Introduction	59
4.2. Parametric tests	59
4.3. Permutation tests	70
4.4. Scale invariant tests	91
4.5. Randomized rank score tests	92
4.6. Deficiencies	102
5. DEFICIENCIES OF SOME RELATED ESTIMATORS	107
5.1. Introduction	107
5.2. Maximum likelihood estimators	107
5.3. Hodges-Lehmann estimators	113

6. FINITE SAMPLE COMPUTATIONS	115
6.1. Introduction	115
6.2. The normal scores and the Wilcoxon test against normal and logistic alternatives	118
6.3. The Wilcoxon test against Cauchy alternatives	121
6.4. The sign test against normal, logistic and double exponential alternatives	122
6.5. Deficiencies between tests for the normal case	128
TABLES	140
REFERENCES	143
INDEX	

## CHAPTER 1

## INTRODUCTION

As "deficiency" is the unifying topic of this study, we shall start by introducing this concept. Let there be given two statistical procedures A and B. If A is based on  $N$  observations, we need  $k_N$  observations for B to attain the same level of performance with both procedures.

Usually, A and B are compared by means of the ratio  $N/k_N$ . If it exists,  $\lim_{N \rightarrow \infty} N/k_N$  is called the *asymptotic relative efficiency* (ARE) of B with respect to A and denoted as  $e$ . Such efficiency computations are by now almost classical: as early as 1925 Fisher (1925) found  $e = 2/\pi$  in comparing the median and the mean for the estimation of normal location. It should always be kept in mind that the information contained in the single number  $e$  is of an asymptotic nature. As we are interested in finite, preferably even small sample sizes  $N$ , this information becomes more valuable according as the convergence of  $N/k_N$  towards  $e$  becomes faster. Hence, when  $e$  has been found, the natural next step is to investigate this rate of convergence, for example by looking at the behaviour of  $ek_N - N$  as  $N \rightarrow \infty$ . This may be done for all cases where  $0 < e < \infty$ , but we shall always restrict ourselves to the by far most interesting case where  $e = 1$ . For then we have a second, perhaps even stronger reason for further investigation: from the fact that  $e = 1$  we cannot even deduce which of the two procedures A and B is better. Hence a study of the difference  $k_N - N$  is not merely useful to get information about the rate of convergence of  $N/k_N$  towards 1, but here it may also reveal which of the two procedures is preferable. For  $e \neq 1$ ,  $ek_N - N$  only supplies some additional information, but for  $e = 1$  this number becomes of importance in its own right.

Although this difference  $k_N - N$  seems to be a very natural quantity to examine, historically the ratio  $N/k_N$  was preferred by almost all authors in view of its simpler behaviour. The first general investigation of  $k_N - N$  was carried out by Hodges and Lehmann (1970). They name  $k_N - N$  the *deficiency* of B with respect to A and denote it as  $d_N$ . If  $\lim_{N \rightarrow \infty} d_N$  exists, it is called the *asymptotic deficiency* of B with respect to A and denoted as  $d$ . At points where no confusion is likely, we shall simply call  $d$  the deficiency of B with respect to A.

Under the assumption  $e = 1$  we evaluate  $d_N$  and  $d$  in the following way. Denote the performance criteria for A and B as  $P_{A,N}$  and  $P_{B,N}$  respectively. If A and B are tests,  $P_{A,N}$  and  $P_{B,N}$  may be the powers of these tests, if A and B are point estimators,  $P_{A,N}$  and  $P_{B,N}$  may be expected squared errors, etc. By definition,  $d_N = k_N - N$  may, for each N, be found from

$$(1.1) \quad P_{A,N} = P_{B,k_N}.$$

In order to solve (1.1),  $k_N$  has to be treated as a continuous variable. This can be done in a satisfactory manner by defining  $P_{B,k_N}$  for non-integral  $k_N$  as

$$P_{B,k_N} = (1 - k_N + [k_N])P_{B,[k_N]} + (k_N - [k_N])P_{B,[k_N]+1}$$

(cf. Hodges and Lehmann (1970)).

Generally  $P_{A,N}$  and  $P_{B,N}$  are not known exactly and we have to use approximations. Here these are obtained by observing that  $P_{A,N}$  and  $P_{B,N}$  will typically satisfy asymptotic expansions of the form

$$(1.2) \quad P_{A,N} = \frac{c}{N^r} + \frac{a}{N^{r+s}} + o\left(\frac{1}{N^{r+s}}\right),$$

$$P_{B,N} = \frac{c}{N^r} + \frac{b}{N^{r+s}} + o\left(\frac{1}{N^{r+s}}\right),$$

for certain  $c$ ,  $a$  and  $b$  not depending on  $N$  and certain constants  $s > 0$  and  $r \neq 0$ . The leading term in both expansions is the same in view of the fact that  $e = 1$ . From (1.1) and (1.2) it now easily follows that

$$(1.3) \quad d_N = \frac{(b-a)}{rc} N^{(1-s)} + o(N^{(1-s)}).$$

Hence

$$(1.4) \quad d = \begin{cases} \pm \infty & , \quad 0 < s < 1, \\ \frac{(b-a)}{rc} & , \quad s = 1, \\ 0 & , \quad s > 1. \end{cases}$$

A useful property of deficiencies is the following: if a third procedure C



is given, for which the performance criterion  $P_{C,N}$  also has an expansion of the form (1.2), the deficiency  $d$  of  $C$  with respect to  $A$  satisfies  $d = d_1 + d_2$ , where  $d_1$  is the deficiency of  $C$  with respect to  $B$  and  $d_2$  is the deficiency of  $B$  with respect to  $A$ .

The situation where  $s = 1$  seems to be the most interesting one. Hodges and Lehmann (1970) demonstrate the use of deficiency in a number of simple examples for which this is the case. One of these problems is the following: consider a sample  $X_1, \dots, X_N$  from a distribution  $F$  with mean  $\xi$  and variance  $\sigma^2$ . Now  $\sigma^2$  can be estimated by  $M_N = N^{-1} \sum_{i=1}^N (X_i - \xi)^2$ , but also by  $M'_N = (N-1)^{-1} \sum_{i=1}^N (X_i - \bar{X})^2$ , with  $\bar{X} = N^{-1} \sum_{i=1}^N X_i$ , if we do not know  $\xi$  or do not dare to rely on its given value. Both estimators are unbiased and therefore we compare  $\sigma^2(M_N)$  and  $\sigma^2(M'_N)$

$$(1.5) \quad \sigma^2(M_N) = \sigma^4 \frac{\gamma}{N}, \quad \sigma^2(M'_N) = \sigma^4 \frac{\gamma(N-1)+2}{N(N-1)},$$

where  $\gamma+1 = \mu_4/\sigma^4$ , the standardized fourth central moment of  $F$ . Application of (1.2) and (1.3) to (1.4) shows that  $d = 2/\gamma$ . If  $F$  is normal,  $\gamma = 2$  and hence  $d = 1$ : the price of not knowing the mean is asymptotically one additional observation. Note that in the normal case not only  $d = 1$ , but also  $d_N \equiv 1$ ; in fact,  $M_N$  and  $M'_{N+1}$  are identically distributed.

The present thesis consists of a number of applications of the deficiency concept. Below we give for each of the problems considered an indication of the problem, of the results, and of the way in which these are obtained.

In chapter 2 the following problem is considered:  $X_1, \dots, X_m$  are independent random variables (r.v.'s), all having distribution  $P_\theta$ ,  $Y_1, \dots, Y_n$  are independent r.v.'s, all having distribution  $P_{\tilde{\theta}}$ , where  $\theta, \tilde{\theta} \in \Theta \subset \mathbb{R}^1$ . From Lehmann (1959) it follows that the test for  $\theta = \tilde{\theta}$  against  $\theta > \tilde{\theta}$  that rejects the hypothesis for large values of  $\sum_{i=1}^m X_i$ , conditionally given  $\sum_{i=1}^m X_i + \sum_{j=1}^n Y_j$ , is uniformly most powerful unbiased (UMPU) for this situation under suitable conditions. These conditions are satisfied for example in the case of the  $2 \times 2$  table, where  $\sum_{i=1}^m X_i$  and  $\sum_{j=1}^n Y_j$  are binomial r.v.'s.

Usually the test is performed with equal sample sizes  $m = n$ . If the criterion of optimality is the unconditional power of the test, this choice is known to be asymptotically optimal in the sense that the optimal value of  $\gamma = m/(m+n)$  satisfies  $\gamma_{\text{opt}} = \frac{1}{2} + o(1)$  as  $(m+n)$ , the total number of experiments, tends to infinity. Here we obtain the optimal value of  $\gamma$  to

$o((m+n)^{-1/2})$ . Attention is restricted to the case where  $|\theta - \tilde{\theta}| \rightarrow 0$  as  $(m+n) \rightarrow \infty$ , at such a rate that for a fixed size  $\alpha > 0$ ,  $\beta$ , the error of the second kind, remains bounded away from 0 and  $1-\alpha$ .

In order to compute deficiencies we need expansions like (1.2) for the power functions. We first expand the conditional distribution function (d.f.) of the test statistic, both under hypothesis and alternative. From this we obtain an expansion for the conditional power. By taking expectations we arrive at an expansion for the unconditional power. Finally, from this expansion  $\gamma_{\text{opt}}$  can be determined. Comparison of the expansion of the power for  $m = \gamma_{\text{opt}}(m+n)$  and  $m = \frac{1}{2}(m+n)$ , gives the deficiency  $d_N$  of the traditional choice  $m = n$  with respect to the optimal choice. The asymptotic deficiency  $d$  proves to be finite. For the special case where  $\alpha = \beta$  we even have  $d_N = O(N^{-1/2})$ , and hence  $d = 0$ . For other choices of  $\alpha$  and  $\beta$ ,  $d$  is positive, but usually rather small. For example, let  $\sum_{i=1}^m X_i$  and  $\sum_{j=1}^n Y_j$  be binomial r.v.'s with parameters  $(m, p_1)$  and  $(n, p_2)$  respectively, where  $p_1 = p_2$  under the hypothesis and  $p_1 > p_2 = p$ ,  $p_1 - p_2 = O((m+n)^{-1/2})$  under the alternative. Then  $d$  satisfies

$$d = \frac{(2p-1)^2}{36p(1-p)} (u_\alpha - u_\beta)^2.$$

Here  $u_\alpha = \Phi^{-1}(1-\alpha)$ ,  $u_\beta = \Phi^{-1}(1-\beta)$ , where  $\Phi^{-1}$  is the inverse of the standard normal d.f. and  $\alpha(\beta)$  is the error of the first (second) kind. For  $0.03 \leq p \leq 0.97$  this gives  $d \leq (u_\alpha - u_\beta)^2$ , for  $0.01 \leq p \leq 0.99$ ,  $d \leq 3(u_\alpha - u_\beta)^2$ . In most applications  $u_\alpha - u_\beta$  will satisfy  $|u_\alpha - u_\beta| \leq 1$ . Then we have that for  $p$  in the given intervals the price of not using the optimal choice but simply  $m = n$  is asymptotically at most 1 or 3 additional observations.

In chapters 3 and 4 we compare various tests for the one sample problem. In its most general form this problem can be formulated in the following way: given a sample  $X_1, \dots, X_N$  of independent identically distributed (i.i.d.) r.v.'s with common d.f.  $G$ , we have to test the hypothesis  $H_0$  that the distribution determined by  $G$  is symmetric about zero, i.e.  $G(x) + G(-x) = 1$  for all  $x$ , against the alternative  $H_1$  that it is not.  $H_0$  is called the hypothesis of symmetry. In this formulation,  $H_1$  is too large to construct tests, having optimal properties against all points of  $H_1$ . Therefore one usually restricts  $H_1$  to some class of interesting alternatives, after which optimal tests against this family are derived. The most common choice is

$\bar{H}_1 : G(x) = F(x-\theta), \theta > 0$ , the family of one-sided location alternatives for a fixed d.f.  $F$  that is symmetric about zero.

A well-known class of tests for the one sample problem is the class of linear one sample rank tests, for example Wilcoxon's signed rank test or the absolute normal scores test. Such tests are not only distribution-free, but also relatively easy to compute, a combination which in general is not achieved by other tests for the one sample problem, such as the test based on  $\bar{X} = N^{-1} \sum_{i=1}^N X_i$ , the one sample t-test or the permutation test based on  $\bar{X}$ . One would expect that the price for these desirable properties would be a loss of efficiency, but, at least asymptotically to first order, this is not the case if one considers contiguous location alternatives  $H_1^* : G(x) = F(x-\theta), \theta = O(N^{-1/2})$ . Hájek and Šidák (1967) prove that the asymptotically most powerful rank test against  $H_1^*$  has ARE  $e = 1$  with respect to the asymptotically most powerful test. For the normal case this means that the absolute normal scores test has  $e = 1$  with respect to the t-test and the  $\bar{X}$ -test. The restriction to  $\theta = O(N^{-1/2})$  is rather natural, as for such sequences of alternatives the power remains bounded away from 1.

In view of the above, it seems interesting to know deficiencies of asymptotically most powerful rank tests with respect to the other types of test, as was suggested by Hodges and Lehmann (1970). To this end we need asymptotic expansions as in (1.2) for the power functions of the tests involved. For linear rank tests for the one sample problem these expansions have been obtained by Albers, Bickel and van Zwet (1974); the two sample problem is dealt with by Bickel and van Zwet (1974). A review of asymptotic expansions in nonparametric statistics is given by Bickel (1974).

Chapter 3 is devoted to asymptotic expansions for one sample rank tests. It contains the results of Albers, Bickel and van Zwet (1974) and some extensions; an outline of the proofs is given but we omit a number of technical details for which the interested reader is referred to Albers, Bickel and van Zwet (1974). The idea is that the rank test statistic is a sum of independent random variables, conditionally under the vector  $Z$  of order statistics of  $|X_1|, \dots, |X_N|$ . Hence we can give Edgeworth expansions in this conditional situation.

An unconditional expansion for the distribution of the test statistic, and hence for the power of the test, follows by taking the expectation with respect to  $Z$  of the conditional expansion. The evaluation of this expecta-

tion is a highly technical matter.

In order to be able to justify the above mentioned Edgeworth expansions we have to exclude cases where the lattice character of the statistic is too pronounced. This occurs for example with the sign test. This test is therefore dealt with separately.

In chapter 4 similar expansions are derived for several other tests: parametric tests, permutation tests and the randomized rank score tests due to Bell and Doksum (1965). After this, deficiencies can be evaluated of the rank tests with respect to the other types of test. For example, the deficiency of the absolute normal scores test with respect to the t-test and the  $\bar{X}$ -test satisfies  $d_N = O(\log \log N)$ ; the asymptotic deficiency of the permutation test based on  $\bar{X}$  with respect to the t-test equals zero.

Chapter 5 is devoted to the application of the results of chapters 3 and 4 to estimation of location. Consider again the situation where  $X_1, \dots, X_N$  are i.i.d. r.v.'s from  $F(x-\theta)$ , in which  $F(-x) = 1-F(x)$  for all  $x$ . For some of the test statistics considered in chapters 3 and 4 there exists a well-known estimator of  $\theta$  which is closely related to this test statistic in the sense that, for all  $a$ ,

$$(1.5) \quad P_{-a}(T < 0) \leq P_0(S < a) \leq P_{-a}(T \leq 0),$$

where  $T$  is the test statistic,  $S$  is the estimator and  $P_\theta$  denotes probability under  $\theta$ . From (1.5) it is clear that the expansion for the d.f. of  $T$ , obtained in chapter 3 or 4, immediately leads to an expansion for the d.f. of  $S$ .

The above correspondence exists for example between the maximum likelihood estimators and the locally most powerful parametric tests of section 4.2 and between the estimators due to Hodges and Lehmann (1956) and the corresponding rank tests of chapter 3. The expansions thus obtained can be used for deficiency comparisons between these estimators. It appears that the deficiency between two estimators equals the deficiency between the corresponding tests for size  $\alpha = \frac{1}{2}$ .

By using certain generalizations of the Cramér-Rao bound - the so called Bhattacharyya bounds - we obtain a lower bound to order  $N^{-1}$  for the variance of an unbiased estimator. We conclude the chapter by evaluating

the deficiency of the estimators considered with respect to this lower bound.

Finally, in chapter 6 we give the results of a number of numerical investigations. These give an indication of the extent to which the asymptotic results obtained in chapters 3 and 4, are of value for small to moderate sample sizes.

In the first place we investigate the behaviour of the expansions for the power of the rank tests in chapter 3 as approximations of the finite sample power. For this purpose we have at our disposal a number of exact power results from literature. These are available for rather small sample sizes -5 to 20- for e.g. Wilcoxon's signed rank test and the absolute normal scores test against normal and logistic location alternatives. Comparison of these values with those resulting from our expansions shows that here the expansions perform very well for these sample sizes already. It also shows that they are much better than the usual normal approximations.

One should keep in mind that the optimistic conclusions above depend on the kind of test and alternative under consideration. If we have long-tailed distributions under the alternative, the situation becomes entirely different. For example in the case of Wilcoxon's signed rank test against Cauchy alternatives not only the normal approximation, but also our expansion leads to very bad results for the same range of sample sizes as above.

In section 4.6 we found approximations to  $o(1)$  for the deficiencies between various tests. Here we compare some of these asymptotic expressions to deficiency values that are obtained numerically. We consider the absolute normal scores test, the t-test and the  $\bar{X}$ -test against normal location alternatives for sample sizes 5-10, 20 and 50. The results thus obtained show a satisfactory agreement between the numerical and asymptotic values.

## CHAPTER 2

## THE OPTIMAL RATIO OF SAMPLE SIZES FOR COMPARING TWO DISTRIBUTIONS

## 2.1. INTRODUCTION

Let  $X_1, \dots, X_m, Y_1, \dots, Y_n$  be independent r.v.'s, the  $X_i$  all having distribution  $P$ , the  $Y_i$  all having distribution  $\tilde{P}$ . We assume that  $P$  and  $\tilde{P}$  belong to a family of distributions  $\mathcal{P}$ , characterized by a real parameter  $\theta : P, \tilde{P} \in \mathcal{P} = \{P_\theta\}, \theta \in \Theta \subset R^1$ .

DEFINITION 2.1.1.  $\Psi$  is the class of continuous functions  $\psi$  on  $R^1$  with  $\psi(0) = 1, 0 < \psi(t) < 1$  for  $t \neq 0$  and  $\sup\{\psi(t) : |t| > \pi\} < 1$ .

DEFINITION 2.1.2. For  $\psi \in \Psi, \mathcal{P}_{1\psi}$  is the class of all distributions  $P$  on  $\{0, 1, 2, \dots\}$  with characteristic functions  $\rho$  satisfying  $|\rho(t)| \leq \psi(t)$  for  $|t| \leq \pi$ . For  $\psi \in \Psi, \mathcal{P}_{2\psi}$  is the class of all distributions  $P$  that are absolutely continuous with respect to Lebesgue measure and satisfy  $|\rho(t)| \leq \psi(t)$  for all  $t$ .

Throughout this chapter we suppose that the family  $\mathcal{P}$  belongs to either  $\mathcal{P}_{1\psi}$  or  $\mathcal{P}_{2\psi}$  for a suitable choice of  $\psi$ .

Before continuing, we shall give an explanation of the definitions above. First we recall some results about lattice distributions (cf. Feller (1966)). A r.v.  $X$  has a lattice distribution  $P$  if there exist real numbers  $a$  and  $h$  with  $h > 0$  such that all values of  $X$  can be represented as  $x = a + vh$  for some integer  $v$ . If  $h_0$  is the largest number such that all values of  $X$  can be represented as  $a + vh_0$ , it is called the span of  $P$ . The characteristic function (c.f.)  $\rho$  of  $X$  is periodic with period  $2\pi/h_0$  and  $|\rho(t)| < 1$  for  $0 < t < 2\pi/h_0$ .

Now it is easy to see how we arrive at  $\mathcal{P}_{1\psi}$ . We are interested in families of distributions that are concentrated on a fixed lattice and that all have the same span. Then it is no loss of generality to assume this lattice to be  $\{0, 1, 2, \dots\}$  and this span to be 1. Note that it is equivalent to assume that the lattice is  $\{0, 1, 2, \dots\}$  and that  $|\rho(t)| < 1$  for  $0 < |t| \leq \pi$ . We arrive at  $\mathcal{P}_{1\psi}$  by using the above stronger version of the second assumption. In this way distributions with span larger than 1 and degenerate distributions are not only excluded, but the elements of  $\mathcal{P}_{1\psi}$  remain bounded away

from such distributions and we can formulate results that are uniformly true for all  $P \in \mathcal{P}_{1\psi}$ .

A similar explanation can be given for  $\mathcal{P}_{2\psi}$ . Here the condition on  $\rho$  serves to ensure that the elements of  $\mathcal{P}_{2\psi}$  remain bounded away from lattice distributions.

We continue our exposition by introducing

$$(2.1.1) \quad X = \sum_{i=1}^m X_i, Y = \sum_{j=1}^n Y_j, T = X+Y, N = m+n.$$

Let  $\theta$  and  $\tilde{\theta}$  be the parameter values corresponding to  $P$  and  $\tilde{P}$ , respectively. A possible way to compare  $P$  and  $\tilde{P}$  on the basis of  $X_1, \dots, X_m$  and  $Y_1, \dots, Y_n$  is to test  $H_0 : \theta = \tilde{\theta}$  against  $H_1 : \theta > \tilde{\theta}$  by rejecting  $H_0$  for large values of  $X$ , conditionally on  $T = t$ . This test is in many cases the uniformly most powerful unbiased (UMPU) test for  $H_0$  against  $H_1$ , as will be shown in section 2.2.

Usually the test is performed with equal sample sizes  $m = n = \frac{1}{2}N$ . In this chapter it is investigated which choice of  $m/N$  is optimal, given  $N$ , as  $N \rightarrow \infty$ . Here the criterion of optimality is the unconditional power of the test. We restrict attention to the Pitman-case: a sequence of alternatives is chosen which converges to the hypothesis at such a rate that for fixed level of significance  $\alpha$  the power remains bounded away from  $\alpha$  and 1. Then it is shown that  $m = n$  is optimal to first order only, but that the deficiency of this choice with respect to the optimal choice is finite.

In section 2.2 conditions are given under which the test is UMPU. Also in this section an expansion is derived for the conditional distribution of  $X$  given  $T = t$ . In section 2.3 the unconditional power is obtained, which enables us to find an expression for the optimal ratio  $m/N$  and for the deficiency of the traditional choice  $m = \frac{1}{2}N$ . This is done in section 2.4. Finally, section 2.5 contains another application of the results in section 2.3.

## 2.2. PRELIMINARIES

First we shall show that the test considered in the previous section is UMPU for the case where  $P$  is an exponential family with monotone likelihood ratio.

LEMMA 2.2.1. Let  $\mathcal{P} = \{P_\theta : \theta \in \Theta\}$  be a family of distributions on  $\mathbb{R}^1$  having densities  $f_\theta(x) = c(\theta) h(x) \exp(Q(\theta)x)$  with respect to a fixed  $\sigma$ -finite measure  $\mu$ . Suppose that  $\Theta$  is an interval in  $\mathbb{R}^1$  and that  $Q$  is continuous and increasing on  $\Theta$ . Let  $X_1, \dots, X_m, Y_1, \dots, Y_n$  be independent r.v.'s, the  $X_i$  all having distribution  $P_\theta$ , the  $Y_i$  all having distribution  $P_{\tilde{\theta}}$ . Finally, let  $X, Y, T$  and  $N$  be defined as in (2.1.1). Then the test for  $\theta = \tilde{\theta}$  against  $\theta > \tilde{\theta}$  that rejects the hypothesis for large values of  $X$ , conditionally on  $T = t$ , is UMPU.

PROOF. The joint density of  $X$  and  $Y$  with respect to an appropriately chosen measure  $\nu$  on  $\mathbb{R}^2$  is

$$f_\theta(x, y) = c^m(\theta) c^n(\tilde{\theta}) \exp[(Q(\theta) - Q(\tilde{\theta}))x + Q(\tilde{\theta})t],$$

where  $t = x + y$ . As  $\Theta \times \Theta$  is a rectangle and  $Q$  is continuous and increasing,  $\{(Q(\theta) - Q(\tilde{\theta}), Q(\tilde{\theta})) : \theta, \tilde{\theta} \in \Theta\}$  is a quadrangle. In view of theorem 4.4.3 of Lehmann (1959) this shows that the test under consideration is UMPU for  $H_0 : Q(\theta) = Q(\tilde{\theta})$  against  $H_1 : Q(\theta) > Q(\tilde{\theta})$ . Hence, by the monotonicity of  $Q$ , the desired result follows.  $\square$

REMARK. In the case where  $Q$  is decreasing, the test is of course UMPU for  $\theta = \tilde{\theta}$  against  $\theta < \tilde{\theta}$ .

The conditions of this lemma are often satisfied for families  $\mathcal{P} \subset \mathcal{P}_{1\psi}$  or  $\mathcal{P} \subset \mathcal{P}_{2\psi}$ . In the first case  $\mu$  can be taken as counting measure, in the second case as Lebesgue measure. We consider the following examples, where  $\eta$  is a positive constant,

EXAMPLE 2.2.1.  $\mathcal{P} = \{P_p\}$  with  $P_p\{1\} = 1 - P_p\{0\} = p$ ,  $\eta \leq p \leq 1 - \eta$ . Hence  $X$  and  $Y$  are binomial r.v.'s. The c.f.  $\rho_p$  of  $P_p$  satisfies  $|\rho_p(t)| = |1 - p + pe^{it}| = \{1 - 2p(1-p)(1 - \cos t)\}^{1/2} \leq \{1 - 2\eta(1-\eta)(1 - \cos t)\}^{1/2}$ . This shows that  $\mathcal{P} \subset \mathcal{P}_{1\psi}$ , for a suitable choice of  $\psi$ . Furthermore,  $Q(p) = \log\{p/(1-p)\}$ , which increases.

EXAMPLE 2.2.2. The family of Poisson distributions  $\mathcal{P} = \{P_\lambda\}$  with  $P_\lambda\{k\} = e^{-\lambda} \lambda^k / (k!)$ ,  $k = 0, 1, 2, \dots$  and  $\lambda \geq \eta$ . Again, for a suitable  $\psi$ ,  $\mathcal{P} \subset \mathcal{P}_{1\psi}$  and  $Q(\lambda) = \log \lambda$  is increasing.



EXAMPLE 2.2.3. The family of geometric distributions  $\mathcal{P} = \{P_p\}$  with  $P_p\{k\} = p(1-p)^{(k-1)}$ ,  $k = 1, 2, \dots$  and  $0 < p \leq 1-n$ . Again, for a suitable  $\psi$ ,  $\mathcal{P} \subset \mathcal{P}_{1\psi}$ , but now  $Q(p) = \log(1-p)$  decreases in  $p$  (cf. the remark following lemma 2.2.1).

EXAMPLE 2.2.4. The family of exponential distributions  $\mathcal{P} = \{P_\lambda\}$  where  $P_\lambda$  is determined by the density  $f_\lambda(x) = \lambda e^{-\lambda x}$ ,  $x > 0$  and  $0 < \lambda \leq \eta$ . Now  $\mathcal{P} \subset \mathcal{P}_{2\psi}$ , for a suitable choice of  $\psi$ , and  $Q(\lambda) = -\lambda$  decreases.

The examples above can all be placed in the framework of a  $2 \times 2$ -table. In the first example we compare two Bernoulli experiments in the following way: the first experiment, which has probability of success  $p_1$ , is performed  $m$  times and the second experiment, which has probability of success  $p_2$ , is performed  $n$  times. The hypothesis  $p_1 = p_2$  is tested against the alternative  $p_1 > p_2$  on the basis of  $X$ , the number of successes obtained with the first experiment, conditionally on  $X+Y$ , the total number of successes. It is well-known that the conditional distribution of  $X$  under the hypothesis is hypergeometric. The second example can be looked at as a limiting case of the first one for small  $p$ -values, as the binomial distribution with parameters  $(N, \frac{\lambda}{N})$  tends to the Poisson distribution with parameter  $\lambda$ . The conditional distribution of  $X$  under the hypothesis is binomial for this case.

In the third example we also compare two Bernoulli experiments, but now the first (second) experiment is performed until  $m(n)$  successes - or, equivalently, failures - have occurred. The above hypothesis is tested here on the basis of the number of trials with the first experiment, conditionally on the total number of trials performed. Finally, the fourth example is the continuous analogue of the third example.

We now return to the general case, where  $\mathcal{P} \subset \mathcal{P}_{1\psi}$  or  $\mathcal{P}_{2\psi}$ . In order to find the optimal ratio  $m/N$ , we must compute the unconditional power of the test, which is the expectation with respect to  $T$  of the conditional power. For the evaluation of this conditional power, we have to know the conditional distribution of  $X$ , given  $T = t$ , under  $H_0$ , as well as under  $H_1$ . First this distribution is found for  $\mathcal{P} \subset \mathcal{P}_{1\psi}$ ; the case  $\mathcal{P} \subset \mathcal{P}_{2\psi}$  follows by analogy. As we shall restrict attention to local alternatives, the following lemma is more general than needed here since it holds for general  $\theta$  and  $\tilde{\theta}$ . It is given in the present form as it may be of some interest of its own.

LEMMA 2.2.2. Suppose that  $X_1, \dots, X_m, Y_1, \dots, Y_n$  are independent r.v.'s, the  $X_i$  with distribution  $P_\theta$ , the  $Y_i$  with distribution  $P_\theta^*$ , where  $P_\theta, P_\theta^* \in \mathcal{P} \subset \mathcal{P}_{1\psi}$ , for some  $\psi \in \Psi$ . Let  $X, Y, T$  and  $N$  be defined as in (2.1.1); let  $\mu, \sigma^2, \mu_k$  ( $\tilde{\mu}, \tilde{\sigma}^2, \tilde{\mu}_k$ ) be the expectation, variance and  $k$ -th central moment of  $X_1$  ( $Y_1$ ),  $k = 3, 4, \dots$ . Moreover, assume that positive constants  $b, B$  and  $V$  exist such that  $\sigma^2 \geq b, \tilde{\sigma}^2 \geq b, Ee^{vX_1} \leq B$  and  $Ee^{vY_1} \leq B$  for  $|v| \leq V$ . Then for all non-negative integers  $m, n$  and  $t$  such that  $|t - m\mu - n\tilde{\mu}| \leq cN^{2/3}/\log N$  and  $\varepsilon \leq m/N \leq 1 - \varepsilon$  for some positive constants  $c$  and  $\varepsilon$ , we have for each non-negative integer  $l$  the following expansion

$$(2.2.1) \quad P(X \leq l \mid T=t) = \phi(y_{1+\frac{1}{2}}) + \phi(y_{1+\frac{1}{2}})(a_0 + a_1 y_{1+\frac{1}{2}} + a_2 (y_{1+\frac{1}{2}}^2 + 2)) + R,$$

where

$$|R| \leq A[N^{-1} + N^{-4}(t - m\mu - n\tilde{\mu})^6].$$

Here

$$(2.2.2) \quad y_1 = \frac{(m\sigma^2 + n\tilde{\sigma}^2)^{1/2}}{mn\sigma^2\tilde{\sigma}^2} \left(1 - \frac{n\tilde{\sigma}^2 m\mu + m\sigma^2(t - n\tilde{\mu})}{m\sigma^2 + n\tilde{\sigma}^2}\right),$$

$$a_0 = \frac{1}{2}(m\sigma^2 + n\tilde{\sigma}^2)^{-1/2} \left[ \frac{\mu_3}{\sigma^2} \left(\frac{n\tilde{\sigma}^2}{m\sigma^2}\right)^{1/2} - \frac{\tilde{\mu}_3}{\tilde{\sigma}^2} \left(\frac{m\sigma^2}{n\tilde{\sigma}^2}\right)^{1/2} \right] +$$

$$- \frac{1}{2}(t - m\mu - n\tilde{\mu})^2 (m\sigma^2 + n\tilde{\sigma}^2)^{-5/2} \left[ \mu_3 \left(\frac{mn\tilde{\sigma}^2}{\sigma^2}\right)^{1/2} - \tilde{\mu}_3 \left(\frac{mn\sigma^2}{\tilde{\sigma}^2}\right)^{1/2} \right],$$

$$a_1 = - \frac{1}{2}(t - m\mu - n\tilde{\mu}) (m\sigma^2 + n\tilde{\sigma}^2)^{-2} \left[ \frac{\mu_3 n\tilde{\sigma}^2}{\sigma^2} + \frac{\tilde{\mu}_3 m\sigma^2}{\tilde{\sigma}^2} \right],$$

$$a_2 = - \frac{1}{6}(m\sigma^2 + n\tilde{\sigma}^2)^{-3/2} \left[ \mu_3 \frac{n^{3/2}\tilde{\sigma}^3}{m^{1/2}\sigma^3} - \tilde{\mu}_3 \frac{m^{3/2}\sigma^3}{n^{1/2}\tilde{\sigma}^3} \right],$$

and  $A$  depends on  $P_\theta, P_\theta^*$  in  $\mathcal{P}_{1\psi}$ ,  $m, n, t$  and  $l$  only through  $b, B, V, c, \varepsilon$  and  $\psi$ .

PROOF. We have

$$(2.2.3) \quad P(X \leq l \mid T=t) = \left[ \sum_{k=0}^l P(X=k)P(Y=t-k) \right] / \left[ \sum_{k=0}^t P(X=k)P(Y=t-k) \right].$$

The procedure is as follows: we give an expansion for  $P(X=k)$  for the central  $k$ -values. From this expansion we immediately derive a similar expansion for  $P(Y=t-k)$ , which leads to an expansion for  $P(X=k)P(Y=t-k)$ , for central  $k$ -values. After showing that the  $P(X=k)P(Y=t-k)$  for  $k$  in the tail of the distribution can be neglected, the sums in (2.2.3) can be evaluated.

First we give an expansion for  $P(X=k)$ . Let  $\rho_Z$  be the characteristic function of a r.v.  $Z$ . As  $\rho_X(u) = \mathbb{E}e^{iuX} = \sum_{k=0}^{\infty} P(X=k)e^{iuk}$ , we have

$$(2.2.4) \quad P(X=k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \rho_X(u) e^{-iuk} du, \quad k = 0, 1, 2, \dots$$

From the definition of  $P_{1\psi}$  it follows that for each  $c_1 \in (0, \pi)$  there exists  $\epsilon_1 \in (0, 1)$  such that  $|\rho_{X_1}(u)| \leq 1 - \epsilon_1$  for  $c_1 \leq |u| \leq \pi$ , uniformly for all  $P \in P_{1\psi}$ . This implies that

$$(2.2.5) \quad |\rho_X(u)| \leq (1 - \epsilon_1)^m, \text{ for } c_1 \leq |u| \leq \pi.$$

Using (2.2.5) and the fact that  $\epsilon \leq m/N \leq 1 - \epsilon$ , (2.2.4) becomes

$$(2.2.6) \quad P(X=k) = \frac{1}{2\pi m^{1/2} \sigma} \int_{-c_1 m^{1/2} \sigma}^{c_1 m^{1/2} \sigma} \rho_{X-m\mu} \left( \frac{u}{m^{1/2} \sigma} \right) e^{\frac{-iu(k-m\mu)}{m^{1/2} \sigma}} du + O(e^{-\tilde{C}N}),$$

for some  $\tilde{C} > 0$ .

Let  $w = u + iv$  be a complex number. From the fact that  $\mathbb{E}e^{vX} \leq B$  for  $|v| \leq V$  it follows that  $\rho_{X_1}(w)$  is analytic for  $|v| < V$ . Also, in analogy to Feller (1966) one may derive a bound for the error in approximating  $\rho_{X_1}(w)$  in the region  $|v| < V$  by finitely many terms of its power series. For  $M = 1, 2, \dots$  define

$$p_M(w) = e^{iw} - 1 - \frac{iw}{1!} - \dots - \frac{(iw)^{M-1}}{(M-1)!}.$$

From

$$|p_1(w)| = \left| \int_0^w e^{iz} dz \right| \leq \int_0^w e^{-\text{Im } z} |dz| \leq |w| \max(1, e^{-V}),$$

and

$$p_M(w) = i \int_0^w p_{M-1}(z) dz, \quad M = 2, 3, \dots,$$

it follows that

$$(2.2.7) \quad |p_M(w)| \leq \frac{|w|^M}{M!} \max(e^{-v}, 1), \quad M = 1, 2, \dots$$

Substitution of  $wX_1$  for  $w$  in (2.2.7) and taking expectations shows

$$(2.2.8) \quad \left| \rho_{X_1}(w) - 1 - \frac{iw}{1!} EX_1 - \dots - \frac{(iw)^{M-1}}{(M-1)!} EX_1^{M-1} \right| \leq \\ \leq \frac{|w|^M}{M!} E[|X_1|^M \max(1, e^{-vX_1})] = O\left(\frac{|w|^M}{M!}\right), \quad M = 1, 2, \dots,$$

for  $|v| < V$ , as  $Ee^{vX_1} \leq B$  for  $|v| \leq V$ . As an application of (2.2.8) we have, since  $\sigma^2 \geq b > 0$ ,

$$\rho_{X_1-\mu}\left(\frac{w}{m^{1/2}\sigma}\right) = 1 - \frac{w^2}{2m} - \frac{i\mu_3 w^3}{6m^{3/2}\sigma^3} + O\left(\frac{|w|^4}{m^2}\right), \text{ for } |v| < m^{1/2}\sigma V.$$

For some positive constant  $c_2$ , which we may choose  $< V$ ,  $\rho_{X_1-\mu}$  satisfies  $\operatorname{Re}\{\rho_{X_1-\mu}(w/(m^{1/2}\sigma))\} > \frac{1}{2}$  for  $|w| \leq c_2 m^{1/2}\sigma$ . Hence, for these  $w$ , we may expand  $\log \rho_{X_1-\mu}(w/(m^{1/2}\sigma))$  to obtain

$$\log \rho_{X_1-\mu}\left(\frac{w}{m^{1/2}\sigma}\right) = -\frac{w^2}{2m} - \frac{i\mu_3 w^3}{6m^{3/2}\sigma^3} + O\left(\frac{|w|^4}{m^2}\right).$$

Consequently

$$\log \rho_{X-m\mu}\left(\frac{w}{m^{1/2}\sigma}\right) = -\frac{w^2}{2} - \frac{i\mu_3 w^3}{6m^{1/2}\sigma^3} + O\left(\frac{|w|^4}{m}\right),$$

for  $|w| \leq c_2 m^{1/2}\sigma$ . Furthermore, there exists a positive constant  $c_3$ , which we may choose  $\leq c_2$ , such that for  $|w| \leq c_3 m^{1/2}\sigma$

$$(2.2.9) \quad \left| \log \rho_{X-m\mu}\left(\frac{w}{m^{1/2}\sigma}\right) + \frac{w^2}{2} \right| \leq \frac{|w|^2}{4}.$$

Hence, for these values of  $w$

$$(2.2.10) \quad \rho_{X-m\mu}\left(\frac{w}{m^{1/2}\sigma}\right) = e^{-w^2/2} \left( 1 - \frac{i\mu_3 w^3}{6m^{1/2}\sigma^3} + O([\mathbb{N}^{-1} + \mathbb{N}^{-1}|w|^6]e^{|w|^2/4}) \right).$$

Using (2.2.9) and (2.2.10) we can now deal with the integral in (2.2.6).

Suppose  $|k-m\mu| \leq CN^{2/3}/\log N$  for some positive constant  $C$  and choose  $c_1 < c_3$ . Then the rectangle with vertices  $\pm c_1 m^{1/2}\sigma$  and  $\pm c_1 m^{1/2}\sigma - i\frac{(k-m\mu)}{m^{1/2}\sigma}$

is contained in  $|w| \leq c_3 m^{1/2} \sigma$  for  $N$  sufficiently large. By complex integration of  $\rho_{X-m\mu} (w/m^{1/2}\sigma) \exp\{-iw(k-m\mu)/(m^{1/2}\sigma)\}$  over this rectangle, we arrive at

$$(2.2.11) \quad P(X=k) = \frac{1}{2\pi m^{1/2}\sigma} \int_{(-c_1 m^{1/2}\sigma - i\frac{k-m\mu}{m^{1/2}\sigma})}^{(c_1 m^{1/2}\sigma - i\frac{k-m\mu}{m^{1/2}\sigma})} \rho_{X-m\mu} \left( \frac{w}{m^{1/2}\sigma} \right) e^{-iw\frac{(k-m\mu)}{m^{1/2}\sigma}} dw + \\ + O\left( \left| \frac{k-m\mu}{m^{1/2}\sigma} \right| \sup_{0 \leq v \leq 1} \left| \rho_{X-m\mu} \left( \pm c_1 - i v \frac{(k-m\mu)}{m\sigma} \right) \right| + e^{-\tilde{C}N} \right).$$

Application of (2.2.10) to the integral in (2.2.11) and of (2.2.9) to the remainder in (2.2.11) leads to

$$(2.2.12) \quad P(X=k) = \frac{e^{-\frac{1}{2} \frac{(k-m\mu)^2}{m\sigma^2}}}{2\pi m^{1/2}\sigma} \int_{-c_1 m^{1/2}\sigma}^{c_1 m^{1/2}\sigma} e^{-\frac{1}{2} u^2} \left[ 1 - \frac{i\mu_3 (u - i \frac{(k-m\mu)}{m^{1/2}\sigma})^3}{6m^{1/2}\sigma^3} + \right. \\ \left. + O\left( [N^{-1} + N^{-1} |u - i \frac{(k-m\mu)}{m^{1/2}\sigma}|^6] e^{\frac{1}{4} (u^2 + \frac{(k-m\mu)^2}{m\sigma^2})} \right) \right] du + \\ + O\left( \left| \frac{k-m\mu}{m^{1/2}\sigma} \right| e^{-\frac{1}{4} c_1^2 m\sigma^2 + \frac{3}{4} \frac{(k-m\mu)^2}{m\sigma^2}} + e^{-\tilde{C}N} \right).$$

Upon evaluation of the integral in (2.2.12), where we use the fact that  $|k-m\mu| \leq CN^{2/3}/\log N$ , we finally arrive at

$$(2.2.13) \quad P(X=k) = \frac{e^{-\frac{1}{2} \frac{(k-m\mu)^2}{m\sigma^2}}}{(2\pi m\sigma^2)^{1/2}} \left[ 1 - \frac{\mu_3}{6m^{1/2}\sigma^3} \left\{ 3 \left( \frac{k-m\mu}{m^{1/2}\sigma} \right) - \left( \frac{k-m\mu}{m^{1/2}\sigma} \right)^3 \right\} + \right. \\ \left. + O(N^{-1} + N^{-4} |k-m\mu|^6) \right],$$

where the remainder only depends on  $P_\theta$  through  $b, B, V$  and  $\psi$ .

Since  $|t-m\mu-\tilde{n}\mu| \leq cN^{2/3}/\log N$ ,  $|k-m\mu| \leq CN^{2/3}/\log N$  implies  $|t-k-\tilde{n}\mu| \leq (C+c)N^{2/3}/\log N$ . Hence, for  $|k-m\mu| \leq CN^{2/3}/\log N$  we also have

$$(2.2.14) \quad P(Y=t-k) = \frac{e^{-\frac{1}{2} \frac{(t-k-n\tilde{\mu})^2}{n\tilde{\sigma}^2}}}{(2\pi n\tilde{\sigma}^2)^{1/2}} \left[ 1 - \frac{\tilde{\mu}_3}{6n^{1/2}\tilde{\sigma}^3} \left\{ 3\left(\frac{t-k-n\tilde{\mu}}{n^{1/2}\tilde{\sigma}}\right) - \left(\frac{t-k-n\tilde{\mu}}{n^{1/2}\tilde{\sigma}}\right)^3 \right\} + \right. \\ \left. + O(N^{-1} + N^{-4} |t-k-n\tilde{\mu}|^6) \right].$$

As we shall have to sum over  $k$ , the expression, obtained for  $P(X=k)$ .  $P(Y=t-k)$  from (2.2.13) and (2.2.14), has to be replaced by an integral over the interval  $(k - \frac{1}{2}, k + \frac{1}{2})$ . First note that

$$(2.2.15) \quad k - m\tilde{\mu} = \left[ \frac{mn\tilde{\sigma}^2\tilde{\sigma}^2}{m\tilde{\sigma}^2 + n\tilde{\sigma}^2} \right]^{1/2} y_k + \frac{m\tilde{\sigma}^2}{m\tilde{\sigma}^2 + n\tilde{\sigma}^2} (t - m\mu - n\tilde{\mu}),$$

$$(2.2.16) \quad t - k - n\tilde{\mu} = - \left[ \frac{mn\tilde{\sigma}^2\tilde{\sigma}^2}{m\tilde{\sigma}^2 + n\tilde{\sigma}^2} \right]^{1/2} y_k + \frac{n\tilde{\sigma}^2}{m\tilde{\sigma}^2 + n\tilde{\sigma}^2} (t - m\mu - n\tilde{\mu}),$$

where

$$(2.2.17) \quad y_k = \left[ \frac{m\tilde{\sigma}^2 + n\tilde{\sigma}^2}{nm\tilde{\sigma}^2\tilde{\sigma}^2} \right]^{1/2} \left[ k - \frac{n\tilde{\sigma}^2}{m\tilde{\sigma}^2 + n\tilde{\sigma}^2} m\mu - \frac{m\tilde{\sigma}^2}{m\tilde{\sigma}^2 + n\tilde{\sigma}^2} (t - n\tilde{\mu}) \right].$$

As a consequence we have

$$(2.2.18) \quad \frac{(k - m\mu)^2}{m\tilde{\sigma}^2} + \frac{(t - k - n\tilde{\mu})^2}{n\tilde{\sigma}^2} = y_k^2 + \frac{(t - m\mu - n\tilde{\mu})^2}{m\tilde{\sigma}^2 + n\tilde{\sigma}^2}.$$

From (2.2.15) it follows that  $|k - m\mu| \leq cN^{2/3}/\log N$  and  $|t - m\mu - n\tilde{\mu}| \leq cN^{2/3}/\log N$  imply  $|y_k| \leq \bar{c}N^{1/6}/\log N$ , for some  $\bar{c} > 0$ . Hence, if  $y_x$  is defined by (2.2.17) with  $k$  replaced by  $x$

$$(2.2.19) \quad \int_{k-1/2}^{k+1/2} e^{-\frac{1}{2} y_x^2} dx = e^{-\frac{1}{2} y_k^2} \int_{k-1/2}^{k+1/2} \left[ 1 - \left( \frac{m\tilde{\sigma}^2 + n\tilde{\sigma}^2}{nm\tilde{\sigma}^2\tilde{\sigma}^2} \right)^{1/2} (x-k) y_k + \right. \\ \left. + O\left( \frac{m\tilde{\sigma}^2 + n\tilde{\sigma}^2}{nm\tilde{\sigma}^2\tilde{\sigma}^2} (x-k)^2 y_k^2 \right) \right] \left[ 1 + O\left( \frac{m\tilde{\sigma}^2 + n\tilde{\sigma}^2}{nm\tilde{\sigma}^2\tilde{\sigma}^2} (x-k)^2 \right) \right] dx = \\ = e^{-\frac{1}{2} y_k^2} \{ 1 + O(N^{-1} + N^{-1} y_k^2) \}.$$

In the same way it can be shown that the lower order terms can be replaced by appropriate integrals. For example

$$\begin{aligned}
(2.2.20) \quad & \int_{k-1/2}^{k+1/2} \left(\frac{x-m\mu}{m^{1/2}\sigma}\right)^3 e^{-\frac{1}{2}y_x^2} dx = \\
& = \left[ \left(\frac{k-m\mu}{m^{1/2}\sigma}\right)^3 + O\left(\frac{k-m\mu}{m^{3/2}\sigma^3}\right) \right] \int_{k-1/2}^{k+1/2} e^{-\frac{1}{2}y_x^2} dx = \\
& = e^{-\frac{1}{2}y_k^2} \left[ \left(\frac{k-m\mu}{m^{1/2}\sigma}\right)^3 + O(N^{-1/2} + N^{-1/2} y_k^4 + N^{-5/2} |t-m\mu-n\tilde{\mu}|^4) \right].
\end{aligned}$$

From (2.2.18), (2.2.19) and expressions similar to (2.2.20) it follows that every term in the product of the expressions in (2.2.13) and (2.2.14) can be replaced by the corresponding integral. Upon doing this and using (2.2.15) we arrive at

$$\begin{aligned}
(2.2.21) \quad & P(X=k)P(Y=t-k) = \exp\left[-\frac{(t-m\mu-n\tilde{\mu})^2}{2(m\sigma^2+n\tilde{\sigma}^2)}\right] / [2\pi(m\sigma^2+n\tilde{\sigma}^2)]^{1/2} \left[ g(y_k) + \right. \\
& \quad \left. + O\left([N^{-1} + N^{-1} y_k^6 + N^{-4} (t-m\mu-n\tilde{\mu})^6]\right) \int_{y_{k-1/2}}^{y_{k+1/2}} e^{-\frac{1}{2}y^2} dy \right],
\end{aligned}$$

for  $|k-m\mu| \leq CN^{2/3}/\log N$ , with

$$\begin{aligned}
(2.2.22) \quad & g(y_k) = \\
& = \int_{y_{k-1/2}}^{y_{k+1/2}} e^{-\frac{1}{2}y^2} \left( 1 - \frac{\mu_3}{6m^{1/2}\sigma^3} \left\{ 3\left[\left(\frac{n\tilde{\sigma}^2}{m\sigma^2+n\tilde{\sigma}^2}\right)^{1/2}y + \frac{m^{1/2}\sigma}{m\sigma^2+n\tilde{\sigma}^2}(t-m\mu-n\tilde{\mu})\right] + \right. \right. \\
& \quad \left. \left. - \left[\left(\frac{n\tilde{\sigma}^2}{m\sigma^2+n\tilde{\sigma}^2}\right)^{1/2}y + \frac{m^{1/2}\sigma}{m\sigma^2+n\tilde{\sigma}^2}(t-m\mu-n\tilde{\mu})\right]^3 \right\} + \right. \\
& \quad \left. + \frac{\tilde{\mu}_3}{6n^{1/2}\tilde{\sigma}^3} \left\{ 3\left[\left(\frac{m\sigma^2}{m\sigma^2+n\tilde{\sigma}^2}\right)^{1/2}y - \frac{n^{1/2}\tilde{\sigma}}{m\sigma^2+n\tilde{\sigma}^2}(t-m\mu-n\tilde{\mu})\right] + \right. \right. \\
& \quad \left. \left. - \left[\left(\frac{m\sigma^2}{m\sigma^2+n\tilde{\sigma}^2}\right)^{1/2}y - \frac{n^{1/2}\tilde{\sigma}}{m\sigma^2+n\tilde{\sigma}^2}(t-m\mu-n\tilde{\mu})\right]^3 \right\} \right) dy.
\end{aligned}$$

With (2.2.21) and (2.2.22) we have found an approximation for the central  $k$ -values; it remains to show that the sums over  $|k-m\mu| > CN^{2/3}/\log N$  of both  $P(X=k)P(Y=t-k)$  and  $g(y_k)$  are sufficiently small.

First  $g(y_k)$  is considered. Choose  $C > 2c$ , then  $|k-m\mu| > CN^{2/3}/\log N$  and  $|t-m\mu-n\tilde{\mu}| \leq cN^{2/3}/\log N$  imply in view of (2.2.15) that  $|y_k| > \tilde{c}N^{1/6}/\log N$ , for some constant  $\tilde{c} > 0$ . Hence

$$(2.2.23) \quad \sum_{|k-m\mu| > CN^{2/3}/\log N} |g(y_k)| \leq \sum_{|y_k| > \tilde{c}N^{1/6}/\log N} |g(y_k)| = \\ = O\left(\int_{|y| > \tilde{c}N^{1/6}/\log N} (1+|y|^3)e^{-\frac{1}{2}y^2} dy\right) = O(N^{-1}).$$

For  $P(X=k)P(Y=t-k)$  we proceed in the following way. According to (2.2.21) to (2.2.23) we have

$$(2.2.24) \quad \sum_{m\mu-CN^{2/3}/\log N}^{m\mu+CN^{2/3}/\log N} P(X=k)P(Y=t-k) = \exp\left[-\frac{(t-m\mu-n\tilde{\mu})^2}{2(m\sigma^2+n\tilde{\sigma}^2)}\right] / [2\pi(m\sigma^2+n\tilde{\sigma}^2)]^{1/2} \\ [1 + O(N^{-1}+N^{-4}(t-m\mu-n\tilde{\mu})^6)].$$

As  $X+Y$  is a sum of independent r.v.'s, an expansion similar to (2.2.14) holds for  $P(X+Y=t)$  for  $|t-m\mu-n\tilde{\mu}| \leq cN^{2/3}/\log N$ . In particular,  $P(X+Y=t)$  equals the right side of (2.2.24). This implies that

$$(2.2.25) \quad \sum_{|k-m\mu| > CN^{2/3}/\log N} P(X=k)P(Y=t-k) = \\ = O([N^{-3/2}+N^{-9/2}(t-m\mu-n\tilde{\mu})^6] \exp[-\frac{1}{2} \frac{(t-m\mu-n\tilde{\mu})^2}{(m\sigma^2+n\tilde{\sigma}^2)}]).$$

The sums in (2.2.3) can now be approximated. For any  $l$  we have

$$(2.2.26) \quad \sum_{k=0}^l P(X=k)P(Y=t-k) = \exp\left[-\frac{(t-m\mu-n\tilde{\mu})^2}{2(m\sigma^2+n\tilde{\sigma}^2)}\right] / [2\pi(m\sigma^2+n\tilde{\sigma}^2)]^{1/2} \left\{ \sum_{k=0}^l g(y_k) + \right. \\ \left. + O(N^{-1}+N^{-4}(t-m\mu-n\tilde{\mu})^6) \right\},$$

where we use (2.2.21) for  $|k-m\mu| \leq CN^{2/3}/\log N$  and (2.2.24) and (2.2.25) for  $|k-m\mu| > CN^{2/3}/\log N$ . From (2.2.26) it immediately follows that



$$(2.2.27) \quad P(X \leq 1 | T=t) = \left[ \sum_{k=0}^1 g(y_k) + O(N^{-1} + N^{-4}(t - m\mu - n\tilde{\mu})^6) \right] / \left[ \sum_{k=0}^t g(y_k) + O(N^{-1} + N^{-4}(t - m\mu - n\tilde{\mu})^6) \right].$$

It remains to insert (2.2.22) in (2.2.27). We can write  $g(y_k)$  as

$$g(y_k) = \int_{y_{k-1/2}}^{y_{k+1/2}} e^{-\frac{1}{2}y^2} (1 + b - a_0 y - a_1 y^2 - a_2 y^3) dy,$$

where  $b$ ,  $a_0$ ,  $a_1$  and  $a_2$  are all  $O(N^{-1/2} + N^{-3/2}(t - m\mu - n\tilde{\mu})^2)$  and  $a_0$ ,  $a_1$  and  $a_2$  are given in (2.2.2). This implies

$$\sum_{k=0}^1 g(y_k) = \phi(y_{1+1/2})(1 + b - a_1) + \phi(y_{1+1/2})[a_0 + a_1 y_{1+1/2} + a_2 (y_{1+1/2}^2 + 2)].$$

In particular,  $\sum_{k=0}^t g(y_k) = 1 + b - a_1 + O(N^{-1})$ . Substitution of these results in (2.2.27) leads to the desired expressions (2.2.1) and (2.2.2). Finally the uniformity of the  $O$ -symbol in  $m$ ,  $l$  and  $t$  is evident from the method of proof.  $\square$

**COROLLARY 2.2.1.** *Under the conditions of lemma 2.2.2 we have for each  $\delta \in (0, 1]$  an expansion to  $O(N^{-1} + N^{-4}(t - m\mu - n\tilde{\mu})^6)$  for  $P(X < 1 | T=t) + \delta P(X=1 | T=t)$  if we replace  $y_{1+1/2}$  by  $y_{1+\delta-1/2}$  in (2.2.1).*

**PROOF.** The result follows immediately from lemma 2.2.2 by noting that  $P(X < 1 | T=t) + \delta P(X=1 | T=t) = (1-\delta)P(X < 1 - 1 | T=t) + \delta P(X \leq 1 | T=t)$  and that  $(1-\delta)\phi(y_{1-1/2}) + \delta\phi(y_{1+1/2}) = \phi(y_{1+\delta-1/2}) + O(N^{-1})$ .  $\square$

In the second case, where  $P \in \mathcal{P}_{2\psi}$ , a similar expansion holds.

**LEMMA 2.2.3.** *Let  $\rho_{X_1}(\rho_{Y_1})$  be the characteristic function of  $X_1(Y_1)$  and suppose that  $\int |\rho_{X_1}(u)|^v du \leq B$ ,  $\int |\rho_{Y_1}(u)|^v du \leq B$  for some  $v \geq 1$ . Let  $P_\theta \in \mathcal{P}_{2\psi}$  for some  $\psi \in \Psi$ , let  $t$  and  $l$  be real numbers and suppose that the remaining conditions of lemma 2.2.2 continue to hold.*

*Then the conclusion in (2.2.1) continues to hold, if  $y_{1+1/2}$  is replaced by  $y_1$ .*

PROOF. As  $X_1$  and  $Y_1$  have densities  $f_{X_1}(x)$  and  $f_{Y_1}(y)$ , the conditional density of  $X = \sum_{i=1}^m X_i$  given  $T = t$  is given by

$$f_{X|T=t}(x) = \begin{cases} \frac{f_X(x) \cdot f_Y(t-x)}{f_T(t)} & \text{if } f_T(t) > 0 \\ 0 & \text{otherwise.} \end{cases}$$

Hence  $P(X \leq 1 | T=t) = \int_{-\infty}^1 f_X(x) f_Y(t-x) dx / \int_{-\infty}^{\infty} f_X(x) f_Y(t-x) dx$ . For  $m \geq \nu$ ,  $|\rho_X| = |\rho_{X_1}|^m \leq |\rho_{X_1}|^\nu$  and therefore  $\int_{-\infty}^{\infty} |\rho_X(u)| du < B$ . Under this condition, the Fourier inversion theorem yields

$$(2.2.28) \quad f_X(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \rho_X(u) e^{-iux} du = \\ = \frac{1}{2\pi m^{1/2}\sigma} \int_{-c_1 m^{1/2}\sigma}^{c_1 m^{1/2}\sigma} \rho_{(X-m\mu)}\left(\frac{u}{m^{1/2}\sigma}\right) e^{-iu\left(\frac{x-m\mu}{m^{1/2}\sigma}\right)} du + \\ + \frac{1}{2\pi} \int_{|u| > c_1} \rho_X(u) e^{-iux} dx.$$

From the definition of  $\mathcal{P}_{2\psi}$  it follows that for each  $c_1 > 0$  there exists  $\varepsilon_1 \in (0, 1)$  such that  $|\rho_{X_1}(u)| \leq 1 - \varepsilon_1$  for  $|u| \geq c_1$ , uniformly for all  $P \in \mathcal{P}_{2\psi}$ . This implies that

$$|\rho_X(u)| = |\rho_{X_1}(u)|^m \leq (1 - \varepsilon_1)^{m-\nu} |\rho_{X_1}(u)|^\nu,$$

for  $|u| > c_1$ . As  $|\rho_{X_1}(u)|^\nu$  is summable, it follows that the second integral in (2.2.28) is exponentially small.

Since (2.2.10) still holds in the present case, the first integral in (2.2.28) can be handled in the same way as in the proof of lemma 2.2.2. Hence, for  $|x - m\mu| \leq CN^{2/3}/\log N$ ,  $f_X(x)$  can be approximated by the right side of (2.2.14), upon changing  $k$  into  $x$ . The remaining part of the proof is analogous to and simpler than the proof of lemma 2.2.2.  $\square$

## 2.3. AN EXPANSION FOR THE UNCONDITIONAL POWER

The conditional power of the test of  $\theta = \tilde{\theta}$  against  $\theta > \tilde{\theta}$  (or  $\theta < \tilde{\theta}$ ) that rejects the hypothesis for large values of  $X$ , given  $T = t$ , is denoted as  $\pi_{t\alpha}$ . Corollary 2.2.1 and lemma 2.2.3 enable us to find an expansion for  $\pi_{t\alpha}$  for those  $t$  that satisfy  $|t - m\mu - n\tilde{\mu}| \leq cN^{2/3}/\log N$ . This will suffice to find an expansion for the unconditional power, which is denoted as  $\pi_\alpha$ , under the assumption of local alternatives. In the following lemma we give such an expansion.

LEMMA 2.3.1. Let  $P_\theta, P_{\tilde{\theta}} \in \mathcal{P}$ , where  $\mathcal{P} \subset \mathcal{P}_{1\psi}$  or  $\mathcal{P} \subset \mathcal{P}_{2\psi}$  for some  $\psi \in \Psi$ . Suppose there exist positive constants  $\epsilon, b, C, V$  and  $0 < \alpha < 1$  such that  $\epsilon \leq m/N \leq 1 - \epsilon$ ,  $Ee^{vX_1} \leq C$ ,  $Ee^{vY_1} \leq C$  for  $|v| \leq V$ ,  $\sigma^2, \tilde{\sigma}^2 \geq b$ ,  $|\mu - \tilde{\mu}| \leq CN^{-1/2}$ ,  $|\sigma^2 - \tilde{\sigma}^2| \leq CN^{-1/2}$  and  $|\mu_3 - \tilde{\mu}_3| \leq CN^{-1/2}$ . If  $\mathcal{P} \subset \mathcal{P}_{2\psi}$  then also assume that  $\int |\rho_{X_1}|^v \leq C$ ,  $\int |\rho_{Y_1}|^v \leq C$  for some  $v \geq 1$ .

Then the unconditional power  $\pi_\alpha$  satisfies

$$(2.3.1) \quad |\pi_\alpha - g(u_\alpha)| \leq AN^{-1},$$

where

$$(2.3.2) \quad g(u_\alpha) = 1 - \Phi\left[u_\alpha - \left(\frac{mn}{N}\right)^{1/2} \left(\frac{\mu - \tilde{\mu}}{\sigma}\right)\right] - \Phi\left[u_\alpha - \left(\frac{mn}{N}\right)^{1/2} \left(\frac{\mu - \tilde{\mu}}{\tilde{\sigma}}\right)\right] \\ \left[ u_\alpha \left\{ \frac{\tilde{\mu}_3}{6\tilde{\sigma}^3} \left(\frac{\mu - \tilde{\mu}}{\tilde{\sigma}}\right) \left(\frac{n+N}{N}\right) - \frac{n}{2N} \left(\frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2}\right) \right\} + \right. \\ \left. + \left(\frac{mn}{N}\right)^{1/2} \left(\frac{\mu - \tilde{\mu}}{\tilde{\sigma}}\right) \left\{ \frac{n}{2N} \left(\frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2}\right) - \frac{\tilde{\mu}_3}{6\tilde{\sigma}^3} \left(\frac{\mu - \tilde{\mu}}{\tilde{\sigma}}\right) \left(\frac{n-m}{N}\right) \right\} \right],$$

and where  $A$  depends on  $P_\theta, P_{\tilde{\theta}}$  in  $\mathcal{P}_{1\psi}$  or  $\mathcal{P}_{2\psi}$ ,  $m$  and  $n$  only through  $\epsilon, b, C, V$  and  $\psi$ .

PROOF. We give the proof for the case where  $\mathcal{P} \subset \mathcal{P}_{1\psi}$ . The proof for the other case is similar but requires some minor notational changes which are left to the reader. As long as nothing is said to the contrary, we restrict attention to those  $t$  that satisfy  $|t - m\mu - n\tilde{\mu}| \leq cN^{2/3}/\log N$ , for some  $c > 0$ . According to corollary 2.2.1 we have under  $H_1$

$$(2.3.3) \quad P_1(X < 1 | T=t) + \delta P_1(X=1 | T=t) = \phi(y_{1+\delta-1/2}) + \phi(y_{1+\delta-1/2})[a_0 + a_1 y_{1+\delta-1/2} + a_2 (y_{1+\delta-1/2}^2 + 2)] + O(N^{-1} + N^{-4}(t - m\mu - n\tilde{\mu})^6),$$

with  $y_1$ ,  $a_0$ ,  $a_1$  and  $a_2$  as defined in (2.2.2). Under  $H_0$  we have

$$(2.3.4) \quad P_0(X < 1 | T=t) + \delta P_0(X=1 | T=t) = \phi(\tilde{y}_{1+\delta-1/2}) + \phi(\tilde{y}_{1+\delta-1/2})[\tilde{a}_0 + \tilde{a}_1 \tilde{y}_{1+\delta-1/2} + \tilde{a}_2 (\tilde{y}_{1+\delta-1/2}^2 + 2)] + O(N^{-1} + N^{-4}(t - m\mu - n\tilde{\mu})^6),$$

where  $\tilde{y}_1$ ,  $\tilde{a}_0$ ,  $\tilde{a}_1$  and  $\tilde{a}_2$  are derived from  $y_1$ ,  $a_0$ ,  $a_1$  and  $a_2$  by replacing  $\mu$ ,  $\sigma^2$  and  $\mu_3$  by  $\tilde{\mu}$ ,  $\tilde{\sigma}^2$  and  $\tilde{\mu}_3$  everywhere.

Let  $1_{t\alpha}$  and  $\delta_{t\alpha}$  be such that  $P_0(X < 1_{t\alpha} | T=t) + \delta_{t\alpha} P_0(X=1_{t\alpha} | T=t) = 1 - \alpha = \Phi(u_\alpha)$ . From (2.3.4) and the fact that  $u_\alpha = O(1)$  it follows that

$$(2.3.5) \quad \tilde{y}_{1_{t\alpha} + \delta_{t\alpha} - 1/2} = u_\alpha - [\tilde{a}_0 + \tilde{a}_1 u_\alpha + \tilde{a}_2 (u_\alpha^2 + 2)] + O(N^{-1} + N^{-4}(t - m\mu - n\tilde{\mu})^6),$$

as  $\tilde{a}_0$ ,  $\tilde{a}_1$  and  $\tilde{a}_2$  are  $O(N^{-1/2} + N^{-3/2}(t - m\mu - n\tilde{\mu})^2)$ . In order to find  $\pi_{t\alpha}$ , we must know  $y_{1_{t\alpha} + \delta_{t\alpha} - 1/2}$ . To this end, note that

$$\frac{n\tilde{\sigma}^2 m\mu + m\sigma^2 (t - n\tilde{\mu})}{m\sigma^2 + n\tilde{\sigma}^2} = (t - n\tilde{\mu}) - \frac{n\tilde{\sigma}^2}{m\sigma^2 + n\tilde{\sigma}^2} (t - m\mu - n\tilde{\mu}),$$

and hence

$$\begin{aligned} \frac{n\tilde{\sigma}^2 m\mu + m\sigma^2 (t - n\tilde{\mu})}{m\sigma^2 + n\tilde{\sigma}^2} - \frac{n\tilde{\sigma}^2 m\tilde{\mu} + m\tilde{\sigma}^2 (t - n\tilde{\mu})}{m\tilde{\sigma}^2 + n\tilde{\sigma}^2} &= \frac{mn}{N} (\mu - \tilde{\mu}) + \\ &+ \left[ \frac{mn}{N^2} \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2} \right) + O(N^{-1}) \right] (t - m\mu - n\tilde{\mu}). \end{aligned}$$

Combining this result and (2.3.5) we get

$$(2.3.6) \quad \begin{aligned} y_{1_{t\alpha} + \delta_{t\alpha} - 1/2} &= [\tilde{y}_{1_{t\alpha} + \delta_{t\alpha} - 1/2} - \left( \frac{N}{mn\tilde{\sigma}^2} \right)^{1/2} \left( \frac{mn}{N} (\mu - \tilde{\mu}) + \right. \\ &+ \left. \left\{ \frac{mn}{N^2} \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2} \right) + O(N^{-1}) \right\} (t - m\mu - n\tilde{\mu}) \right)] \left( \frac{N\sigma^2}{m\sigma^2 + n\tilde{\sigma}^2} \right)^{-1/2} = \\ &= \left[ u_\alpha - \left( \frac{mn}{N} \right)^{1/2} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) \right] \left[ 1 - \frac{n}{2N} \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2} \right) \right] - \left( \frac{mn}{N} \right)^{1/2} \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^3} \right) \frac{(t - m\mu - n\tilde{\mu})}{N} + \\ &- [\tilde{a}_0 + \tilde{a}_1 u_\alpha + \tilde{a}_2 (u_\alpha^2 + 2)] + O(N^{-1} + N^{-4}(t - m\mu - n\tilde{\mu})^6), \end{aligned}$$

as  $|\mu - \tilde{\mu}| = O(N^{-1/2})$  and  $|\sigma^2 - \tilde{\sigma}^2| = O(N^{-1/2})$ . From (2.3.3) it now follows that

$$(2.3.7) \quad \begin{aligned} 1 - \pi_{t\alpha} &= \Phi\left\{ \left[ u_\alpha - \left( \frac{m\tilde{\mu}}{N} \right)^{1/2} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) \right] \left[ 1 - \frac{n}{2N} \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2} \right) \right] \right\} + \\ &+ \Phi\left\{ \left[ u_\alpha - \left( \frac{m\tilde{\mu}}{N} \right)^{1/2} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) \right] \left[ - \left( \frac{m\tilde{\mu}}{N} \right)^{1/2} \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^3} \right) \frac{(t - m\mu - n\tilde{\mu})}{N} + \right. \right. \\ &+ a_0 + a_1 \left[ u_\alpha - \left( \frac{m\tilde{\mu}}{N} \right)^{1/2} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) \right] + a_2 \left\{ \left[ u_\alpha - \left( \frac{m\tilde{\mu}}{N} \right)^{1/2} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) \right]^2 + 2 \right\} + \\ &\left. \left. - \tilde{a}_0 - \tilde{a}_1 u_\alpha - \tilde{a}_2 (u_\alpha^2 + 2) \right] \right\} + O(N^{-1} + N^{-4} (t - m\mu - n\tilde{\mu})^6). \end{aligned}$$

From (2.2.2) it follows that  $|a_0 - \tilde{a}_0| = O(N^{-1} + N^{-2} (t - m\mu - n\tilde{\mu})^2)$ ,  $|a_2 - \tilde{a}_2| = O(N^{-1})$ ,

$$a_1 - \tilde{a}_1 = \frac{m}{2N} \frac{\tilde{\mu}_3}{\tilde{\sigma}^3} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) + O(N^{-1} + N^{-3/2} (t - m\mu - n\tilde{\mu})).$$

Inserting these results and the expressions for  $a_1$  and  $a_2$  from (2.2.2), we get

$$(2.3.8) \quad \begin{aligned} \pi_{t\alpha} &= g(u_\alpha) + \Phi\left\{ \left[ u_\alpha - \left( \frac{m\tilde{\mu}}{N} \right)^{1/2} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) \right] \left( \frac{m\tilde{\mu}}{N} \right)^{1/2} \frac{(t - m\mu - n\tilde{\mu})}{N\tilde{\sigma}} \left[ \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2} \right) + \right. \right. \\ &\left. \left. - \frac{1}{2} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) \frac{\tilde{\mu}_3}{\tilde{\sigma}^3} \right] \right\} + O(N^{-1} + N^{-4} (t - m\mu - n\tilde{\mu})^6), \end{aligned}$$

where  $g(u_\alpha)$  is defined by (2.3.2).

The expansion in (2.3.8) only holds for  $|t - m\mu - n\tilde{\mu}| < cN^{2/3}/\log N$ , but this suffices to find  $\pi_\alpha = E\pi_{T\alpha}$ . Let  $S$  be a set satisfying

$$S \subset \{t : |t - m\mu - n\tilde{\mu}| < cN^{2/3}/\log N\}$$

and let  $I_S(t)$  be the indicator function of  $S$ . Then

$$(2.3.9) \quad \begin{aligned} \pi_\alpha - g(u_\alpha) &= E(\pi_{T\alpha} - g(u_\alpha)) = E[(T - m\mu - n\tilde{\mu}) I_S(T)] O(N^{-1}) + \\ &+ O(N^{-1} + N^{-4} E[(T - m\mu - n\tilde{\mu})^6 I_S(T)]) + O(E[|\pi_{T\alpha} - g(u_\alpha)| I_{(S^c)}(T)]). \end{aligned}$$

As  $ET = m\mu + n\tilde{\mu}$ ,  $E[(T - m\mu - n\tilde{\mu}) I_S(T)] = -E[(T - m\mu - n\tilde{\mu}) I_{(S^c)}(T)]$ . Because  $Ee^{vX_1} \leq B$ ,  $Ee^{vY_1} \leq B$  for  $|v| \leq v$  and  $T = \sum_{i=1}^m X_i + \sum_{j=1}^n Y_j$ , we have  $E|T - m\mu - n\tilde{\mu}|^r = O(N^{r/2})$  for all positive real  $r$ , by the Marcinkiewitz-Zygmund-

Chung-inequality (cf. Chung (1951)). In particular,  $E(T - m\mu - n\tilde{\mu})^6 = O(N^3)$ . Furthermore,  $\pi_{T\alpha}$  and  $g(u_\alpha)$  are both bounded. In view of these facts, (2.3.9) simplifies to

$$(2.3.10) \quad |\pi_\alpha - g(u_\alpha)| = O(N^{-1} + E\{[|T - m\mu - n\tilde{\mu}| + 1]^r I_{(S^c)}(T)\}).$$

Application of Hölder's inequality shows that for all  $r, s > 1$  with  $1/r + 1/s = 1$

$$(2.3.11) \quad E\{[|T - m\mu - n\tilde{\mu}| + 1]^r I_{(S^c)}(T)\} \leq \{E[|T - m\mu - n\tilde{\mu}| + 1]^r\}^{1/r} \{E[I_{(S^c)}(T)]^s\}^{1/s} = \\ = O(N^{1/2} [P(T \in S^c)]^{1/s}).$$

From Chebyshev's inequality we have, if we choose  $S = \{t: |t - m\mu - n\tilde{\mu}| \leq N^{1/2+\tau}\}$ , for some  $0 < \tau < 1/6$

$$(2.3.12) \quad P(T \in S^c) = O(E|T - m\mu - n\tilde{\mu}|^p P_N^{-p(1/2+\tau)}) = O(N^{-p\tau}),$$

for all  $p > 0$ . Now (2.3.1) follows from (2.3.10), (2.3.11) and (2.3.12). The uniformity in  $P_\theta$ ,  $P_{\tilde{\theta}}$  and  $m$  follows again from the method of proof.  $\square$

For exponential families with monotone likelihood ratio the expression for  $g(u_\alpha)$  in (2.3.2) can be simplified by using the result in the next lemma.

LEMMA 2.3.2. Let  $\mathcal{P} = \{P_\theta : \theta \in \Theta \subset \mathbb{R}^1\}$  be a family of distributions on  $\mathbb{R}^1$  having densities  $f_\theta(x) = \exp(Q(\theta)x)/c(\theta)$  with respect to a  $\sigma$ -finite measure  $\nu$ . Let  $\mu$ ,  $\sigma^2$  and  $\mu_3$  ( $\tilde{\mu}$ ,  $\tilde{\sigma}^2$  and  $\tilde{\mu}_3$ ) be the expectation, variance and third central moment of a r.v. with distribution  $P_\theta$  ( $P_{\tilde{\theta}}$ ). Let  $\theta, \tilde{\theta} \in \Theta$  be such that  $|Q(\theta) - Q(\tilde{\theta})| \leq CN^{-1/2}$  and  $0 < \tilde{c} \leq \int \exp\{(Q(\theta) + \epsilon)x\} d\nu \leq \tilde{C} < \infty$ ,  $0 < \tilde{c} \leq \int \exp\{(Q(\theta) - \epsilon)x\} d\nu \leq \tilde{C} < \infty$ , for constants  $\tilde{c}$ ,  $C$ ,  $\tilde{C}$  and  $\epsilon > 0$ . Then we have, uniformly for fixed  $\tilde{c}$ ,  $C$ ,  $\tilde{C}$  and  $\epsilon$

$$(2.3.13) \quad (\mu - \tilde{\mu})\mu_3 - (\sigma^2 - \tilde{\sigma}^2)\sigma^2 = O(N^{-1}).$$

PROOF. Let  $\tau = Q(\theta)$  and define  $d(\tau) = \int e^{\tau x} d\nu(x)$ . Then  $\int x^k f_\theta(x) d\nu(x) = d^{(k)}(\tau)/d(\tau)$ ,  $k=1,2,\dots$ , and therefore the expression on the left side of (2.3.13) may be expressed in terms of the values of the functions  $d^{(k)}$  and

$d$  at the points  $Q(\theta)$  and  $Q(\tilde{\theta})$ , for  $k = 1, 2, 3$ . Under the conditions above on  $\theta$ ,  $d^{(k)}$  and  $d$  are uniformly bounded away from zero and infinity at the point  $Q(\theta)$ . As  $|Q(\theta) - Q(\tilde{\theta})| \leq CN^{-1/2}$ , this will also hold at the point  $Q(\tilde{\theta})$  for  $N$  sufficiently large. Furthermore,

$$d^{(k)}(Q(\tilde{\theta})) = d^{(k)}(Q(\theta)) + \{Q(\tilde{\theta}) - Q(\theta)\}d^{(k+1)}(Q(\theta)) + O(N^{-1}).$$

Inserting this in  $(\mu - \tilde{\mu})\mu_3 - (\sigma^2 - \tilde{\sigma}^2)\sigma^2$  leads to the desired result.  $\square$

If (2.3.13) holds,  $g(u_\alpha)$  simplifies to

$$(2.3.14) \quad g(u_\alpha) = 1 - \Phi\left[u_\alpha - \left(\frac{mn}{N}\right)^{1/2} \left(\frac{\mu - \tilde{\mu}}{\sigma}\right)\right] - \phi\left[u_\alpha - \left(\frac{mn}{N}\right)^{1/2} \left(\frac{\mu - \tilde{\mu}}{\sigma}\right)\right] \\ \left[ \frac{\tilde{\mu}_3}{6\tilde{\sigma}^3} \left(\frac{\mu - \tilde{\mu}}{\sigma}\right) \left(\frac{N-2n}{N}\right) u_\alpha + \left(\frac{mn}{N}\right)^{1/2} \left(\frac{\mu - \tilde{\mu}}{\sigma}\right)^2 \frac{\tilde{\mu}_3}{6\tilde{\sigma}^3} \left(\frac{n+N}{N}\right) \right].$$

We now apply the results of this section to the four examples given in section 2.2. Here lemma's 2.2.1 and 2.3.2 clearly apply. Hence  $\pi_\alpha$  is the power of the UMPU test for these cases and under the conditions of lemma 2.3.1 it may be approximated by (2.3.14). Let  $\eta_1$  and  $\eta_2$  be positive constants.

EXAMPLE 2.3.1.  $P(X_1=1) = 1 - P(X_1=0) = p_1$ ,  $P(Y_1=1) = 1 - P(Y_1=0) = p_2 = p$ , where  $\eta_2 \leq p_2 < p_1 \leq 1 - \eta_2$  and  $p_1 = p_2 + bN^{-1/2}$ ,  $b = O(1)$ . Then  $\mu = p_1$ ,  $\sigma^2 = p_1(1-p_1)$ ,  $\mu_3 = (1-2p_1)p_1(1-p_1)$  and therefore  $\mu - \tilde{\mu} = bN^{-1/2}$ ,  $\sigma^2 - \tilde{\sigma}^2 = (1-2p)bN^{-1/2} + O(N^{-1})$ ,  $\mu_3 - \tilde{\mu}_3 = O(N^{-1/2})$ . Furthermore,  $Ee^{vX_1} = (1-p_1) + p_1 e^v$  for all  $v$ . It follows that we may apply lemma 2.3.1. Hence

$$(2.3.15) \quad \pi_\alpha = 1 - \Phi\left(u_\alpha - \left[\frac{mn}{p(1-p)}\right]^{1/2} \frac{b}{N}\right) - \phi\left(u_\alpha - \left[\frac{mn}{p(1-p)}\right]^{1/2} \frac{b}{N}\right) \\ \left[ \frac{(1-2p)(N-2n)}{6p(1-p)N^{3/2}} bu_\alpha + \left[\frac{mn}{p(1-p)}\right]^{1/2} \frac{(1-2p)(n+N)}{6p(1-p)N^{5/2}} b^2 \right] + O(N^{-1}).$$

EXAMPLE 2.3.2.  $X_1(Y_1)$  has a Poisson distribution with parameter  $\lambda_1(\lambda_2)$ , where  $\eta_2 \leq \lambda_2 < \lambda_1 \leq \eta_1$ ,  $\lambda_2 = \lambda$ ,  $\lambda_1 = \lambda + bN^{-1/2}$ ,  $b = O(1)$ . Then  $\mu = \sigma^2 = \mu_3 = \lambda_1$ , and therefore  $\mu - \tilde{\mu} = \sigma^2 - \tilde{\sigma}^2 = \mu_3 - \tilde{\mu}_3 = bN^{-1/2}$ . Furthermore,  $Ee^{vX_1} = e^{-\lambda_1 + \lambda_1 e^v}$  for all  $v$ . Hence

$$(2.3.16) \quad \pi_\alpha = 1 - \phi\left(u_\alpha - \left[\frac{mn}{\lambda}\right]^{1/2} \frac{b}{N}\right) - \phi\left(u_\alpha - \left[\frac{mn}{\lambda}\right]^{1/2} \frac{b}{N}\right) \left[ \frac{(N-2n)}{6\lambda N^{3/2}} bu_\alpha + \right. \\ \left. + \left[\frac{mn}{\lambda}\right]^{1/2} \frac{(n+N)}{6\lambda N^{5/2}} b^2 \right] + O(N^{-1}).$$

EXAMPLE 2.3.3.  $X_1(Y_1)$  has a geometric distribution with parameter  $p_1(p_2)$ , with  $n_2 \leq p_1 < p_2 \leq 1-n_2$ ,  $p_2 = p$ ,  $p_1 = p + bN^{-1/2}$ ,  $b = O(1)$ . Then  $\mu = \frac{1}{p_1}$ ,  $\sigma^2 = \frac{(1-p_1)}{p_1^2}$ ,  $\mu_3 = \frac{(1-p_1)(2-p_1)}{p_1^3}$  and therefore  $\mu - \tilde{\mu} = -\frac{bN^{-1/2}}{p^2} + O(N^{-1})$ ,  $\sigma^2 - \tilde{\sigma}^2 = -\frac{(2-p)}{p^3} bN^{-1/2} + O(N^{-1})$ ,  $\mu_3 - \tilde{\mu}_3 = O(N^{-1/2})$ . Furthermore,  $Ee^{vX_1} = p_1 e^v / (1 - [1-p_1]e^v)$ , for  $v < \log\left(\frac{1}{1-p_1}\right)$ . Hence

$$(2.3.17) \quad \pi_\alpha = 1 - \phi\left(u_\alpha + \left[\frac{mn}{1-p}\right]^{1/2} \frac{b}{pN}\right) - \phi\left(u_\alpha + \left[\frac{mn}{1-p}\right]^{1/2} \frac{b}{pN}\right) \left[ \frac{(p-2)(N-2n)}{6p(1-p)N^{3/2}} bu_\alpha + \right. \\ \left. + \left[\frac{mn}{1-p}\right]^{1/2} \frac{(2-p)(n+N)}{6p^2(1-p)N^{5/2}} b^2 \right] + O(N^{-1}).$$

EXAMPLE 2.3.4.  $X_1(Y_1)$  has an exponential distribution with parameter  $\lambda_1(\lambda_2)$ , with  $n_2 \leq \lambda_1 < \lambda_2 \leq n_1$ ,  $\lambda_2 = \lambda$ ,  $\lambda_1 = \lambda + bN^{-1/2}$ ,  $b = O(1)$ . Then  $\mu = \frac{1}{\lambda_1}$ ,  $\sigma^2 = \frac{1}{\lambda_1^2}$ ,  $\mu_3 = \frac{2}{\lambda_1^3}$ , and therefore  $\mu - \tilde{\mu} = -\frac{b}{\lambda^2} N^{-1/2} + O(N^{-1})$ ,  $\sigma^2 - \tilde{\sigma}^2 = -\frac{2b}{\lambda} N^{-1/2} + O(N^{-1})$ ,  $\mu_3 - \tilde{\mu}_3 = O(N^{-1/2})$ . Furthermore,  $Ee^{vX_1} = \lambda / (\lambda - v)$ , for  $v < \lambda$ , and  $|\rho_{X_1}(u)| = \left(1 + \frac{u^2}{\lambda^2}\right)^{-1/2}$ . Hence

$$(2.3.18) \quad \pi_\alpha = 1 - \phi\left(u_\alpha + \frac{(mn)^{1/2} b}{\lambda N}\right) - \phi\left(u_\alpha + \frac{(mn)^{1/2} b}{\lambda N}\right) \left[ -\frac{(N-2N)}{3\lambda N^{3/2}} bu_\alpha + \right. \\ \left. + \frac{(mn)^{1/2} (N+n)}{3\lambda^2 N^{5/2}} b^2 \right] + O(N^{-1}).$$

#### 2.4. OPTIMAL RATIO AND DEFICIENCIES

In this section we solve the main problem of this chapter: which choice of  $\gamma = m/N$  is optimal as  $N \rightarrow \infty$ ? From (2.3.2) it is evident that  $\gamma = \frac{1}{2} + O(N^{-1/2})$ . In order to find the second order term we set  $\gamma = \frac{1}{2} + fN^{-1/2}$ , with  $f = O(1)$ . For a given  $f$ , denote the power at level  $\alpha$  by  $\pi_{\alpha, f}$  and the expression in (2.3.2) by  $g(u_\alpha, f)$ . Under the conditions of lemma 2.3.1 we have



$$(2.4.1) \quad \pi_{\alpha, f} - \pi_{\alpha, 0} = g(u_{\alpha}, f) - g(u_{\alpha}, 0) + O(N^{-3/2}),$$

as changes in  $f$  cause changes of  $O(N^{-3/2})$  in  $\pi_{\alpha, f} - g(u_{\alpha}, f)$ . From (2.3.2) and (2.4.1) it follows that

$$(2.4.2) \quad \begin{aligned} \pi_{\alpha, f} - \pi_{\alpha, 0} &= N^{-1} \phi \left[ u_{\alpha} - \frac{N^{1/2}(\mu - \tilde{\mu})}{2\tilde{\sigma}} \right] \left[ -f^2 \frac{N^{1/2}(\mu - \tilde{\mu})}{\tilde{\sigma}} + \right. \\ &+ f u_{\alpha} \left\{ \frac{\tilde{\mu}_3}{6\tilde{\sigma}^3} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) N^{1/2} - \frac{1}{2} \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2} \right) N^{1/2} \right\} + \\ &\left. + f \frac{N^{1/2}(\mu - \tilde{\mu})}{\tilde{\sigma}} \left\{ \frac{N^{1/2}}{4} \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2} \right) - \frac{\tilde{\mu}_3 N^{1/2}}{6\tilde{\sigma}^3} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) \right\} \right] + O(N^{-3/2}). \end{aligned}$$

Obviously, (2.4.2) reaches its maximum for

$$(2.4.3) \quad \begin{aligned} f_0 &= \left[ u_{\alpha} \left\{ \frac{\tilde{\mu}_3}{6\tilde{\sigma}^3} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) - \frac{1}{2} \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2} \right) \right\} + N^{1/2} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) \left\{ \frac{1}{4} \left( \frac{\sigma^2 - \tilde{\sigma}^2}{\tilde{\sigma}^2} \right) - \frac{\tilde{\mu}_3}{6\tilde{\sigma}^3} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) \right\} \right] / \\ &[2 \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right)] + O(N^{-1/2}). \end{aligned}$$

Consequently,

$$(2.4.4) \quad \pi_{\alpha, f_0} - \pi_{\alpha, 0} = N^{-1/2} \phi \left[ u_{\alpha} - \frac{N^{1/2}(\mu - \tilde{\mu})}{2\tilde{\sigma}} \right] \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right) f_0^2 + O(N^{-3/2}).$$

Define  $b_N^* = \frac{N^{1/2}}{2} \left( \frac{\mu - \tilde{\mu}}{\tilde{\sigma}} \right)$ . Note that  $b_N^*$  is bounded. As  $\alpha$  is a constant, with  $0 < \alpha < 1$ , it follows that

$$(2.4.5) \quad \pi_{\alpha, f_0} = 1 - \phi \left[ u_{\alpha} - b_N^* + A(u_{\alpha}, b_N^*) N^{-1/2} - 2b_N^{*2} f_0^2 N^{-1} + O(N^{-3/2}) \right],$$

$$\pi_{\alpha, 0} = 1 - \phi \left[ u_{\alpha} - b_N^* + A(u_{\alpha}, b_N^*) N^{-1/2} + O(N^{-3/2}) \right],$$

where  $|A(u_{\alpha}, b_N^*)| \leq C$  for some  $C > 0$ . Application of (1.2) to (2.4.5) shows that the deficiency  $d_N$  of the choice  $m = n$  with respect to the optimal choice  $m/N = \frac{1}{2} + f_0 N^{-1/2}$ , satisfies

$$(2.4.6) \quad d_N = (2f_0)^2 + O(N^{-1/2}).$$

Hence the asymptotic deficiency  $d$  equals  $(2f_0)^2$ . This is finite, which may seem surprising in view of the fact that the difference between the optimal choice  $\gamma_0$  and  $\frac{1}{2}$  is  $O(N^{-1/2})$  rather than  $O(N^{-1})$ . This suggests the

asymptotic deficiency to be infinite. That this is not the case is explained as follows: the leading term in  $g(u_\alpha, f)$  is symmetric around  $\gamma = \frac{1}{2}$ , in fact  $\pi_\alpha = 1 - \Phi(u_\alpha - [N\gamma(1-\gamma)]^{1/2}(\frac{\mu - \tilde{\mu}}{\tilde{\sigma}})) + O(N^{-1/2})$ , and  $\gamma = \frac{1}{2} + O(N^{-1/2})$  implies  $\gamma(1-\gamma) = \frac{1}{4} + O(N^{-1})$ .

We finally apply these results to our examples. In the first place, (2.3.13) holds in all examples considered, and hence (2.4.3) simplifies to

$$(2.4.7) \quad f_0 = -\frac{\tilde{\mu}_3}{24\tilde{\sigma}^3} [4u_\alpha - N^{1/2}(\frac{\mu - \tilde{\mu}}{\tilde{\sigma}})] + O(N^{-1/2}).$$

The expression in (2.4.7) becomes more transparent if we eliminate  $N^{1/2}(\mu - \tilde{\mu})/\tilde{\sigma}$  by using its relation to the power of the test. Denote  $1 - \pi_\alpha$ , the error of the second kind, as  $\beta$ . From  $\beta = \Phi(u_\alpha - \frac{N^{1/2}(\mu - \tilde{\mu})}{2(\frac{\mu - \tilde{\mu}}{\tilde{\sigma}})}) + O(N^{-1/2})$  it then follows that

$$\frac{N^{1/2}}{2} (\frac{\mu - \tilde{\mu}}{\tilde{\sigma}}) = (u_\alpha + u_\beta) + O(N^{-1/2}).$$

Hence

$$(2.4.8) \quad f_0 = -\frac{\tilde{\mu}_3}{12\tilde{\sigma}^3} (u_\alpha - u_\beta) + O(N^{-1/2}),$$

$$(2.4.9) \quad d_N = \frac{\tilde{\mu}_3^2}{36\tilde{\sigma}^6} (u_\alpha - u_\beta)^2 + O(N^{-1/2}).$$

In most applications  $|u_\alpha - u_\beta| < 1$ , which makes  $(\frac{\tilde{\mu}_3}{6\tilde{\sigma}^3})^2$  a reasonable upper bound for  $d$ . In the special case where the errors of the first and second kind are equal,  $f_0 = O(N^{-1/2})$  and the choice of  $m = n$  is therefore optimal to  $O(N^{-1})$ . We conclude this section by making some remarks on each of the examples of section 2.3 separately.

EXAMPLE 2.4.1. For example 2.3.1, (2.4.8) specializes to

$$f_0 = \frac{(2p-1)}{12[p(1-p)]^{1/2}} (u_\alpha - u_\beta).$$

Remembering that  $m/N = \gamma_0 = \frac{1}{2} + f_0 N^{-1/2}$ , we may conclude that, if the error of the first kind has to be smaller (larger) than the error of the second kind, one should perform more (less) often the experiment whose probability of success differs most from  $\frac{1}{2}$ . Furthermore,  $(2p-1)^2/(36p(1-p)) =$

$= (2p-1)^2/[9(1-(2p-1)^2)]$ . Hence, if  $|u_\alpha - u_\beta| < 1$ , then  $d \leq D$  for  $|p-1/2| \leq \frac{1}{2}[9D/(9D+1)]^{1/2} \approx \frac{1}{2} - \frac{1}{36D}$ . This gives for example  $d \leq 1$  for  $0.03 \leq p \leq 0.97$  and  $d \leq 3$  for  $0.01 \leq p \leq 0.99$ .

EXAMPLE 2.4.2. For example 2.3.2 we find  $f_0 = -(u_\alpha - u_\beta)/(12\sqrt{\lambda})$ . If one wants to have  $\alpha < \beta$ , then one should perform the experiment with the smallest parameter more often. Furthermore,  $d \leq D$  if  $\lambda \leq \frac{D}{36}$  and  $|u_\alpha - u_\beta| < 1$ .

EXAMPLE 2.4.3. For example 2.3.3  $f_0 = -(2-p)(u_\alpha - u_\beta)/[12(1-p)]^{1/2}$ .

EXAMPLE 2.4.4. For example 2.3.4  $f_0 = -(u_\alpha - u_\beta)/6$ .

## 2.5. COMPARISON OF SAMPLING RULES FOR BERNOULLI EXPERIMENTS

In the previous section the results of section 2.3 were used to solve the main problem of this chapter. Here we briefly discuss another application of these results.

Consider two Bernoulli experiments, with probability of success  $p_1$  and  $p_2$ , respectively. We shall give an asymptotic comparison of the performance of two sampling rules in testing the hypothesis  $p_1 = p_2$  against the alternative  $p_2 = p$ ,  $p_1 = p + \Delta$ ,  $\Delta > 0$ . The first of these sampling rules is the "Vector-at-a-Time" (VT) rule, which simply states that  $m$  and  $n$ , the numbers of trials with both experiments, are equal. (cf. Sobel and Weiss (1970)). From example 2.3.1 it follows that for this rule the power  $\pi_\alpha$  of the UMPU test satisfies

$$(2.5.1) \quad \pi_\alpha = 1 - \Phi\left\{u_\alpha - \Delta\left(\frac{n}{2p(1-p)}\right)^{1/2}\left(1 - \frac{(1-2p)\Delta}{4p(1-p)}\right)\right\} + O(n^{-1}),$$

for  $\Delta = O(n^{-1/2})$  and  $\varepsilon \leq p_2 < p_1 \leq 1 - \varepsilon$  for some constant  $\varepsilon > 0$ .

The second sampling rule we consider is the "Play-the-Winner" (PW) rule, which prescribes that one continues with the same experiment after each success and that one switches to the opposite experiment after each failure. As soon as  $r$  failures have occurred with both experiments, sampling is terminated. In this situation there also exists an UMPU test for

$H_0 : p_1 = p_2$  against  $H_1 : p_2 = p$ ,  $p_1 = p + \Delta$ ,  $\Delta > 0$ . An approximation to the power  $\pi_\alpha^*$  of this test is supplied by example 2.3.3, if we interchange  $p$  and  $1-p$  in (2.3.17). We find for  $\Delta = O(r^{-1/2})$  and  $\varepsilon \leq p_2 < p_1 \leq 1 - \varepsilon$

$$(2.5.2) \quad \pi_{\alpha}^* = 1 - \Phi\left\{u_{\alpha} - \frac{\Delta}{(1-p)}\left(\frac{r}{2p}\right)^{1/2}\left(1 - \frac{(1+p)\Delta}{4p(1-p)}\right)\right\} + O(r^{-1}).$$

In order to make the PW rule and the VT rule comparable, it seems reasonable to choose  $r$ , for each  $n$ , in such a way that the powers of the two tests under consideration are equal. From (2.5.1) and (2.5.2) it follows that  $|\pi_{\alpha} - \pi_{\alpha}^*| = O(n^{-1})$  for

$$(2.5.3) \quad r = n(1-p)\left(1 + \frac{3}{2(1-p)} \Delta\right) + O(1).$$

Now there are various criteria according to which we can compare the two sampling rules. For example, we may prefer the rule that has the lowest expected number of trials on the poorer experiment, i.e. the experiment with the smallest probability of success. Another criterion is the expected number of failures that could have been avoided by using the better experiment throughout. Finally, a third criterion is the expected total number of trials.

We consider the first criterion. For the VT rule the expected number of trials on the poorer experiment obviously equals  $n$ . From (2.5.3) we obtain that for the corresponding PW rule this expectation is

$$n\left(1 + \frac{3}{2(1-p)} \Delta\right) + O(1).$$

Hence the PW rule requires in expectation  $3n\Delta/\{2(1-p)\}$  additional observations on the poorer experiment.

As concerns the other criteria mentioned above, we note that the second criterion is equivalent to the first, whereas the third criterion will certainly not prefer the PW rule if the first criterion prefers the VT rule. Hence, according to all criteria, the PW rule is asymptotically worse than the VT rule for the problem of this section, uniformly in  $p$ . Apparently the fact that the PW rule has a tendency to use the better experiment more often, is outweighed by the fact that the negative binomial distribution has a larger skewness than the binomial distribution.

## CHAPTER 3

## ASYMPTOTIC EXPANSIONS FOR NONPARAMETRIC TESTS FOR THE ONE SAMPLE PROBLEM

## 3.1. INTRODUCTION

In this chapter we shall give asymptotic expansions for the distribution functions of one sample linear rank statistics and also for the power functions of the corresponding tests. These have been derived by Albers, Bickel and van Zwet (1974); the present chapter contains the results of this paper and two extensions. We only sketch the proofs.

Our starting point in establishing the above expansions will be the so called Edgeworth expansions (cf. Cramér (1946), Feller (1966)). For the distribution function (d.f.)  $R(x)$  of any r.v.  $X$  with mean 0 and variance 1 we can give a formal Edgeworth expansion  $\tilde{R}(x)$  in powers of  $N^{-\frac{1}{2}}$ . For example, to  $O(N^{-1})$  this looks like

$$(3.1.1) \quad \tilde{R}(x) = \Phi(x) - \phi(x) \left[ \frac{N^{-\frac{1}{2}} \kappa_3}{6} (x^2 - 1) + \frac{N^{-1} \kappa_4}{24} (x^3 - 3x) + \frac{N^{-1} \kappa_3^2}{72} (x^5 - 10x^3 + 15x) \right],$$

where  $\kappa_3$  and  $\kappa_4$  are the third and fourth cumulant of  $X$ , multiplied by  $N^{\frac{1}{2}}$  and  $N$ , respectively, and  $\Phi$  and  $\phi$  denote the d.f. and the density of the standard normal distribution.

Such expansions have been used for rank tests before, for example by Hodges and Fix (1955), Fellingham and Stoker (1964), Sundrum (1954), Witting (1960) and Rogers (1971). These authors, however, restrict attention to the special case of the one or two sample Wilcoxon test. Furthermore, with the exception of Rogers, they do not bother to show that (3.1.1) is a valid expansion, but merely recommend it as an approximation on purely numerical grounds. Rogers gives an Edgeworth expansion  $\tilde{R}(x)$  to  $O(N^{-1})$  for the two sample Wilcoxon distribution  $R(x)$  under the hypothesis, and proves that  $\sup_x |R(x) - \tilde{R}(x)| = o(N^{-1})$ .

Here we shall justify expansions to  $O(N^{-1})$ , not only under the hypothesis, but also under contiguous alternatives, for quite general test scores. In section 3.2. we give a basic expansion for the d.f. of a linear rank

statistic, without any assumptions at all about the alternative. In the next section we restrict attention to contiguous alternatives and obtain a more explicit form for this expansion. This expression still involves sums of functions of moments of order statistics. These are replaced by appropriate integrals in section 3.4 under the assumption of smooth scores and contiguous location alternatives. Finally, in section 3.5, we consider the sign test as a separate case, since it cannot be handled by the general methods of this chapter, because of its pronounced lattice character.

### 3.2. THE BASIC EXPANSION

Let  $X_1, \dots, X_N$  be i.i.d. r.v.'s with common d.f.  $G$  and density  $g$ , and let  $0 < Z_1 < Z_2 < \dots < Z_N$  denote the order statistics of the absolute values of  $X_1, \dots, X_N$ . If  $|X_{R_j}| = Z_j$ , define

$$(3.2.1) \quad v_j = \begin{cases} 1 & \text{if } X_{R_j} > 0, \\ 0 & \text{otherwise.} \end{cases}$$

We introduce a vector of scores  $a = (a_1, \dots, a_N)$  and define the statistic

$$(3.2.2) \quad T = \sum a_j v_j.$$

Throughout this section  $\sum$  always means  $\sum_{j=1}^N$ , unless stated otherwise. We shall be concerned with obtaining an asymptotic expansion for the distribution of  $T$  as  $N \rightarrow \infty$ .

Our notation strongly suggests that we are considering a fixed underlying d.f.  $G$  and perhaps also a fixed infinite sequence of scores as  $N \rightarrow \infty$ . However, this is merely a matter of notational convenience and our main concern will in fact be the case where the d.f. depends on  $N$  and the scores form a triangular array  $a_{j:N}$ ,  $j = 1, \dots, N$ ,  $N = 1, 2, \dots$ .

The r.v.  $T$  is of course the general linear rank statistic for testing the hypothesis that  $g$  is symmetric about zero. Under this hypothesis,  $V_1, \dots, V_N$  are i.i.d. with  $P(V_j=1) = 1/2$ . For general  $G$ ,  $V_1, \dots, V_N$  are not independent. However, one easily verifies that conditionally on  $Z = (Z_1, \dots, Z_N)$ , the r.v.'s  $V_1, \dots, V_N$  are independent with

$$(3.2.3) \quad P_j = P(V_j=1|Z) = \frac{g(Z_j)}{g(Z_j)+g(-Z_j)}.$$

As independence allows us to obtain expansions of Edgeworth type, we shall carry out the following program to arrive at an expansion for the distribu-

tion of  $T$ . First we obtain an Edgeworth expansion for the distribution of  $\sum a_j W_j$ , where  $W_1, \dots, W_N$  are independent with  $p_j = P(W_j=1) = 1 - P(W_j=0)$ . Having done this, we substitute the random vector  $P = (P_1, \dots, P_N)$  defined in (3.2.3) for  $p = (p_1, \dots, p_N)$  in this expansion. The expected value of the resulting expression will then give us an expansion for the distribution of  $T$ .

In carrying out the first part of this program, we shall indicate any dependence on  $p = (p_1, \dots, p_N)$  in our notation. Consider the r.v.

$$(3.2.4) \quad \frac{\sum a_j (W_j - p_j)}{\tau(p)},$$

where

$$(3.2.5) \quad \tau^2(p) = \sum p_j (1-p_j) a_j^2$$

denotes the variance of  $\sum a_j W_j$ . Obviously (3.2.4) has expectation 0 and variance 1. Let  $R(x, p)$  and  $\rho(t, p)$  denote the d.f. and the characteristic function (c.f.) of (3.2.4) respectively. Denote the Edgeworth expansion to  $O(N^{-1})$  for  $R(x, p)$  as  $\tilde{R}(x, p)$  (cf. (3.1.1)). Let  $\tilde{r}(x, p)$  be  $\frac{\partial}{\partial x} \tilde{R}(x, p)$  and  $\tilde{\rho}(t, p) = \int_{-\infty}^{\infty} \exp(itx) \tilde{r}(x, p) dx$ , the Fourier transform of  $\tilde{r}$ .

To justify a formal Edgeworth expansion  $\tilde{R}$ , i.e. to show that  $|R - \tilde{R}|$  is indeed  $o(N^{-1})$ , one usually invokes the following result (Feller (1966)).

LEMMA 3.2.1. *Let  $R$  be a d.f. with vanishing expectation and c.f.  $\rho$ . Suppose that  $R - \tilde{R}$  vanishes at  $\pm\infty$  and that  $\tilde{R}$  has a derivative  $\tilde{r}$  such that  $|\tilde{r}| \leq m$ . Finally, suppose that  $\tilde{r}$  has a continuously differentiable Fourier transform  $\tilde{\rho}$  such that  $\tilde{\rho}(0) = 1$  and  $\tilde{\rho}'(0) = 0$ . Then, for all  $x$  and  $T > 0$ ,*

$$(3.2.6) \quad |R(x) - \tilde{R}(x)| \leq \frac{1}{\pi} \int_{-T}^T \left| \frac{\rho(t) - \tilde{\rho}(t)}{t} \right| dt + \frac{24m}{\pi T}.$$

PROOF. See Feller (1966).

To prove that  $|R - \tilde{R}| = o(N^{-1})$ , it suffices to show that e.g. for  $T = b \cdot N^{3/2}$ , the integral in (3.2.6) is  $o(N^{-1})$ . For the case we are considering, this may be done largely in the standard manner (Feller (1966)), by splitting the integral into several parts. Require that for some positive constants  $C$  and  $c$ ,  $\sum a_j^4 \leq CN$  and  $\tau^2(p) = \sum p_j (1-p_j) a_j^2 \geq cN$ . As this implies

that  $(\max_j a_j)/\tau(p) \leq (Cc^{-2})^{1/4} N^{-1/4}$ , we can expand

$$\rho(t,p) = \exp \left[ \sum \log \rho_{W_j - p_j} (a_j t / \tau(p)) \right]$$

around  $t = 0$  for  $|t| \leq c_1 N^{1/4}$ , for some positive  $c_1$ , depending on  $C$  and  $c$ . Comparison of this expansion with  $\tilde{\rho}(t,p)$  shows that  $|\rho(t,p) - \tilde{\rho}(t,p)|$  is sufficiently small on this interval to ensure that

$$\int_{|t| \leq c_1 N^{1/4}} \left| \frac{\rho(t,p) - \tilde{\rho}(t,p)}{t} \right| dt = O(N^{-5/4}).$$

On  $c_1 N^{1/4} \leq |t| \leq b_1 N^{3/2}$  we cannot expand  $\rho$  anymore, but as  $\rho$  and  $\tilde{\rho}$  are both small here, we simply use

$$\int \left| \frac{\rho(t,p) - \tilde{\rho}(t,p)}{t} \right| dt \leq \int \left| \frac{\rho(t,p)}{t} \right| dt + \int \left| \frac{\tilde{\rho}(t,p)}{t} \right| dt.$$

The last integral on the right is shown to be sufficiently small for  $|t| \geq \log(N+1)$  without any difficulties. As it can be shown that  $|\rho(t,p)| \leq \exp[-\frac{1}{2}t^2 + Ct^4/(96c^2N)]$ , we also have that the following integral is sufficiently small

$$\int_{c_1 N^{1/4} \leq |t| \leq b_1 N^{1/2}} \left| \frac{\rho(t,p)}{t} \right| dt,$$

for some positive constant  $b_1$ . Hence it remains to estimate

$$(3.2.7) \quad \int_{b_1 N^{1/2} \leq |t| \leq bN^{3/2}} \left| \frac{\rho(t,p)}{t} \right| dt.$$

Here one usually makes what Feller calls the extravagantly luxurious assumption that the c.f.'s of all summands are uniformly bounded away from 1 in absolute value outside every neighbourhood of 0. Obviously, this condition is not satisfied in our case where the summands  $a_j W_j$  are lattice r.v.'s. Weaker sufficient conditions of this type are known, but all seem to imply at the very least that the sum itself is non-lattice. In our case this would exclude for instance both the sign test and the Wilcoxon test. On the other hand, it is clear that one has to exclude cases where the sum (3.2.4) can only assume relatively few different values. As  $\tilde{R}$  is continuous, one cannot allow  $R$  to have jumps of  $O(N^{-1})$  or larger. Thus the sign test, where jumps of order  $N^{-1/2}$  occur, will certainly have to be excluded.



However, it is exactly the simple lattice character of this statistic that makes it easily amenable to other methods of expansion (see section 3.5). For the Wilcoxon statistic on the other hand, all jumps are  $O(N^{-3/2})$  and the assumptions we shall make will not rule out this case.

For  $\rho(t,p)$  we have

$$(3.2.8) \quad |\rho(t,p)| = \prod_{j=1}^N \left\{ 1 - 2p_j(1-p_j) \left( 1 - \cos \frac{a_j t}{\tau(p)} \right) \right\}^{1/2}.$$

This is exponentially small if for a positive fraction of indices  $j$  the following two conditions are simultaneously satisfied:  $\epsilon \leq p_j \leq 1-\epsilon$  and  $a_j t / \tau(p)$  differs at least  $\eta$  from the nearest multiple of  $2\pi$ , for some  $0 < \epsilon < 1$ ,  $\eta > 0$ . For our purpose this must hold for all  $t$  with  $b_1 N^{1/2} \leq |t| \leq b_2 N^{3/2}$ . If this is the case, then obviously (3.2.7) is sufficiently small, and the Edgeworth expansion is justified.

We summarize this result in the following theorem, where the two conditions above are replaced by one weaker, but less intuitive condition.

**THEOREM 3.2.1.** *Suppose that positive numbers  $c$ ,  $C$ ,  $\delta$  and  $\epsilon$  exist such that  $\sum a_j^4 \leq CN$ ,  $\sum a_j^2 p_j(1-p_j) \geq cN$  and  $\gamma(\epsilon, \zeta, p) = \lambda\{x | \exists_j |x - a_j| < \zeta\}$ ,  $\epsilon \leq p_j \leq 1-\epsilon\} \geq \delta N \zeta$  for some  $\zeta \geq N^{-3/2} \log N$ , where  $\lambda$  is Lebesgue measure. Then*

$$(3.2.9) \quad \sup_x |R(x,p) - \tilde{R}(x,p)| \leq A.N^{-5/4},$$

where  $A$  depends on  $N$ ,  $a$  and  $p$  only through  $c$ ,  $C$ ,  $\delta$  and  $\epsilon$ .

**PROOF.** For a formal and detailed proof see Albers, Bickel and van Zwet (1974).  $\square$

**REMARK.** If we require  $\sum |a_j|^5 \leq CN$  instead of  $\sum a_j^4 \leq CN$ , we get  $A.N^{-3/2}$  instead of  $A.N^{-5/4}$  in (3.2.9). This is the "natural" order of the remainder.

Before we replace  $p$  by the random vector  $P = (P_1, \dots, P_N)$  defined in (3.2.3) and compute the unconditional distribution of  $T$  by taking the expected value, two modifications must be performed.

In the first place, we have to change the standardization of  $\sum a_j W_j$  into one

that does not involve  $p$ . As before, let  $W_1, \dots, W_N$  be i.i.d. with  $P(W_j=1) = 1 - P(W_j=0) = p_j$ , let  $\tilde{p} = (\tilde{p}_1, \dots, \tilde{p}_N)$  be a vector with  $0 \leq \tilde{p}_j \leq 1$  for all  $j$ , and consider

$$(3.2.10) \quad R^*(x, p, \tilde{p}) = P\left(\frac{\sum_j a_j (W_j - \tilde{p}_j)}{\tau(\tilde{p})} \leq x\right).$$

Here  $\tau^2(\tilde{p}) = \sum_j \tilde{p}_j (1 - \tilde{p}_j) a_j^2$  in accordance with (3.2.5). From the fact that

$$R^*(x, p, \tilde{p}) = R\left(\left[x - \frac{\sum_j (p_j - \tilde{p}_j) a_j}{\tau(\tilde{p})}\right] \frac{\tau(\tilde{p})}{\tau(p)}, p\right),$$

we can immediately derive an expansion for  $R^*$  by means of (3.2.9).

The second modification is that everywhere in this expansion we expand  $\tau(\tilde{p})/\tau(p)$  in powers of  $[\tau^2(p) - \tau^2(\tilde{p})]/\tau^2(\tilde{p})$ ; the reasons for this will become clear in the sequel. The result of these steps is the following lemma.

LEMMA 3.2.2. *If the conditions of theorem 3.2.1 are satisfied and in addition  $\sum_j \tilde{p}_j (1 - \tilde{p}_j) a_j^2 \geq cN$ , we have*

$$(3.2.11) \quad \sup_x |R^*(x, p, \tilde{p}) - \tilde{R}^*(x, p, \tilde{p})| \leq \\ \leq A\{N^{-5/4} + N^{-3/2} \sum_j (p_j - \tilde{p}_j)^2 |a_j|^3 + N^{-3} |\tau^2(p) - \tau^2(\tilde{p})|^3\},$$

where  $A > 0$  depends on  $N, a, p, \tilde{p}$  only through  $c, C, \delta$  and  $\epsilon$  and where

$$(3.2.12) \quad \tilde{R}^*(x, p, \tilde{p}) = \tilde{R}(x-u, \tilde{p}) - \phi(x-u) \left\{ \frac{1}{2} \frac{\tau^2(p) - \tau^2(\tilde{p})}{\tau^2(\tilde{p})} (x-u) + \right. \\ + \frac{1}{6} \frac{\sum_j (p_j - \tilde{p}_j) (1 - 6\tilde{p}_j + 6\tilde{p}_j^2) a_j^3}{\tau^3(\tilde{p})} [(x-u)^2 - 1] + \\ + \frac{1}{8} \left( \frac{\tau^2(p) - \tau^2(\tilde{p})}{\tau^2(\tilde{p})} \right)^2 [(x-u)^3 - 3(x-u)] + \\ \left. + \frac{\kappa_3(\tilde{p})}{12N^{3/2}} \frac{\tau^2(p) - \tau^2(\tilde{p})}{\tau^2(\tilde{p})} [(x-u)^4 - 6(x-u)^2 + 3] \right\},$$

with  $u = [\sum_j (p_j - \tilde{p}_j) a_j] / \tau(\tilde{p})$ .

PROOF. See Albers, Bickel and van Zwet (1974).

We shall now replace  $p$  by  $P = (P_1, \dots, P_N)$  in  $\tilde{R}^*(x, p, \tilde{p})$  and take expectations. Define the vector  $\pi = (\pi_1, \dots, \pi_N)$  by

$$(3.2.13) \quad \pi_j = EP_j, \quad j = 1, \dots, N.$$

It will play the role of  $\tilde{p}$ . Then the following theorem can be formulated.

**THEOREM 3.2.2.** *Let  $X_1, \dots, X_N$  be i.i.d. with common d.f.  $G$  and density  $g$ , and let  $T$ ,  $P$  and  $\pi$  be defined by (3.2.2), (3.2.3) and (3.2.13). Suppose that positive numbers  $c$ ,  $C$ ,  $\delta$ ,  $\delta'$  and  $\epsilon$  exist with  $\delta' < \min(\delta/2, c^2 C^{-1})$  and such that*

$$(3.2.14) \quad \sum a_j^2 \geq cN, \quad \sum a_j^4 \leq CN,$$

$$(3.2.15) \quad \gamma(\zeta) = \lambda\{x \mid \exists_j |x - a_j| < \zeta\} \geq \delta N \zeta \text{ for some } \zeta \geq N^{-3/2} \log N,$$

$$(3.2.16) \quad P\{\epsilon \leq \frac{g(X_1)}{g(X_1) + g(-X_1)} \leq 1 - \epsilon\} \geq 1 - \delta'.$$

Then there exists  $A > 0$  depending on  $N$ ,  $a$  and  $G$  only through  $c$ ,  $C$ ,  $\delta$ ,  $\delta'$  and  $\epsilon$  and such that

$$(3.2.17) \quad \sup_x \left| P\left(\frac{\sum_{j=1}^T a_j \pi_j}{\tau(\pi)}\right) - E\tilde{R}^*(x, P, \pi) \right| \leq \\ \leq A \left\{ N^{-5/4} + N^{-3/4} \left[ \sum \{E(P_j - \pi_j)^2\}^{5/2} \right]^{2/5} + N^{-3/2} \left[ \sum \{E|P_j - \pi_j|^3\}^{2/3} \right]^{3/2} \right\}.$$

PROOF. Here we only sketch the proof. For a complete proof see Albers, Bickel and van Zwet (1974). Using (3.2.16) one can show that for  $\delta'' \in (\delta', \min(\delta/2, c^2 C^{-1}))$

$$P(E) \leq e^{-2N(\delta'' - \delta')^2},$$

where  $E = \{P \mid \epsilon \leq P_j \leq 1 - \epsilon \text{ for less than } (1 - \delta'')N \text{ indices } j\}$ . On  $E^c$ ,  $\sum a_j^2 \geq cN$  implies both  $\sum a_j^2 P_j (1 - P_j) \geq c^* N$  and  $\sum a_j^2 \pi_j (1 - \pi_j) \geq c^* N$ , for some  $c^*$  depending only on  $c$ ,  $C$ ,  $\delta''$ ,  $\delta'$  and  $\epsilon$  and satisfying  $0 < c^* < c$ . Also  $\gamma(\zeta) \geq \delta N \zeta$  implies  $\gamma(\epsilon, \zeta, P) \geq (\delta - 2\delta'')N \zeta$  on  $E^c$ , where  $\delta - 2\delta''$  is positive by

assumption. Hence, under the conditions of this theorem,  $a$ ,  $P$  and  $\pi$  satisfy on  $E^c$  the conditions for  $a$ ,  $p$  and  $\tilde{p}$  in lemma 3.2.2, if  $c$ ,  $C$ ,  $\delta$  and  $\epsilon$  are replaced there by positive numbers  $c^*$ ,  $C$ ,  $\delta - 2\delta''$  and  $\epsilon$ , depending only on  $c$ ,  $C$ ,  $\delta$ ,  $\delta'$  and  $\epsilon$ . In dealing with the set  $E$  it will suffice to note that  $\tilde{R}^*(x, P, \pi)$  is bounded. Of course,  $R^*(x, P, \pi)$ , being a probability, is also bounded.

As  $P([\tau - \sum a_j \pi_j] / \tau(\pi) \leq x) = ER^*(x, P, \pi)$ , the left side of (3.2.17) is bounded above by

$$(3.2.18) \quad E \sup_x |R^*(x, P, \pi) - \tilde{R}^*(x, P, \pi)|.$$

Applying lemma 3.2.2 on  $E^c$  and using the boundedness of  $|R^*(x, P, \pi) - \tilde{R}^*(x, P, \pi)|$  together with  $P(E) = O(N^{-5/4})$  we find that (3.2.18) is

$$O(N^{-5/4} + N^{-3/2} \sum [E(P_j - \pi_j)^2 |a_j|^3 + N^{-3} E|\tau^2(P) - \tau^2(\pi)|^3]),$$

where the order symbol is uniform for fixed  $c$ ,  $C$ ,  $\delta$ ,  $\delta'$  and  $\epsilon$ . This expression can be shown to be of the order of the right side in (3.2.17).  $\square$

We note that the boundedness of  $\tilde{R}^*(x, P, \pi)$  on  $E$  plays an important role in the above proof. Because  $\tau(P)$  may be arbitrarily small on  $E$ , this explains why we had to remove  $\tau(p)$  from the denominator of the expansion in lemma 3.2.2 by expanding  $\tau(\tilde{p})/\tau(p)$  in powers of  $[\tau^2(p) - \tau^2(\tilde{p})]/\tau^2(\tilde{p})$ .

Although theorem 3.2.2 is formally stated as a result for a fixed, but arbitrary value of  $N$ , it is of course meaningless for fixed  $N$  because we do not investigate the way in which  $A$  depends on  $c$ ,  $C$ ,  $\delta$ ,  $\delta'$  and  $\epsilon$ . In fact the theorem is a purely asymptotic result. Let us for a moment indicate dependence on  $N$  by a superscript. Thus for  $N = 1, 2, \dots$ , consider the distribution of the statistic  $T^{(N)}$  based on a vector of scores  $a^{(N)} = (a_1^{(N)}, \dots, a_N^{(N)})$  when the underlying d.f. is  $G^{(N)}$ . Fix positive values of  $c$ ,  $C$ ,  $\delta$ ,  $\delta'$  and  $\epsilon$  with  $\delta' < \min(\delta/2, c^2 C^{-1})$ . The theorem asserts that if for each  $N$ ,  $a^{(N)}$  and  $G^{(N)}$  satisfy (3.2.14) - (3.2.16) for these fixed  $c$ ,  $C$ ,  $\delta$ ,  $\delta'$  and  $\epsilon$ , then the error of the approximation  $ER^*(x, P^{(N)}, \pi^{(N)})$  is

$$O(N^{-5/4} + N^{-3/4} \sum [E(P_j^{(N)} - \pi_j^{(N)})^2]^{5/2} ]^{2/5} + N^{-3/2} \sum [E|P_j^{(N)} - \pi_j^{(N)}|^3]^{2/3} ]^{3/2}),$$

as  $N \rightarrow \infty$ . Moreover, the order of the remainder is uniform for all such sequences  $a^{(N)}, G^{(N)}, N = 1, 2, \dots$ .

Assumption (3.2.15) may need some clarification. It is clear from the sketch of the proof of theorem 3.2.1 that the role of the  $\gamma(\epsilon, \zeta, p)$  and  $\gamma(\zeta)$  conditions in theorem 3.2.1 and theorem 3.2.2 respectively, is to ensure that the  $a_j$  do not cluster too much around too few points. Assumption (3.2.15) is certainly satisfied if for some  $k \geq \delta N/2$ , indices  $j_1, j_2, \dots, j_k$  exist such that  $a_{j_{i+1}} - a_{j_i} \geq 2N^{-3/2} \log N$  for  $i = 1, \dots, k$ . Under condition (3.2.14) this will typically be the case. Consider for instance the important case where  $a_j = EJ(U_{j:N})$ , where  $U_{1:N} < U_{2:N} < \dots < U_{N:N}$  are order statistics from the uniform distribution on  $(0, 1)$  and  $J$  is a continuously differentiable, nonconstant function on  $(0, 1)$  with  $\int J^4 < \infty$ . Here both (3.2.14) and (3.2.15) are satisfied for all  $N$  with fixed  $c, C$  and  $\delta$ . The same is true if  $a_j = J(j/(N+1))$ , provided that  $J$  is monotone near 0 and 1.

For a large class of underlying d.f.'s  $G$ , the right side of (3.2.17) is uniformly  $o(N^{-1})$ . Still theorem 3.2.2 does not yet provide an explicit expansion to order  $N^{-1}$  for the distribution of  $T$  since we are still left with the task of computing the expected value of  $\tilde{R}^*(x, P, \pi)$ . This is of course a trivial matter under the hypothesis that  $g$  is symmetric about zero and, more generally, in the case where, for some  $\eta > 0$ ,  $g(x)/g(-x) = \eta$  for all  $x > 0$ . In this case  $P_j = \eta(1+\eta)^{-1}$  with probability 1 for all  $j$  and an expansion for the distribution of  $T$  is already contained in theorem 3.2.1. For fixed alternatives in general, however, the computation of  $E\tilde{R}^*(x, P, \pi)$  presents a formidable problem that we shall not attempt to solve here. It would seem that what is needed, is an expansion for the distribution of a linear combination of functions of order statistics.

In the remaining part of this chapter we shall restrict attention to sequences of alternatives that are contiguous to the hypothesis. Heuristically the situation is now as follows. Since  $g(x)/(g(x)+g(-x)) = \frac{1}{2} + O(N^{-1/2})$ ,  $P_j - \frac{1}{2}$  and  $\pi_j - \frac{1}{2}$  will be  $O(N^{-1/2})$ , whereas  $P_j - \pi_j$  will be  $O(N^{-1})$  instead of  $O(N^{-1/2})$  as before. In the first place this allows us to simplify  $E\tilde{R}^*(x, P, \pi)$  considerably as a number of terms may now be relegated to the remainder and functions of  $\pi_j$  may be expanded about the point  $\pi_j = \frac{1}{2}$ . Much more important, however, is the fact that  $U^* = \tau^{-1}(\pi) \sum (P_j - \pi_j) a_j$  will now be  $O(N^{-1/2})$  and that we may therefore expand  $\tilde{R}^*(x, P, \pi)$  in powers of  $U^*$ . This means that we shall be dealing with low moments of linear combinations of

functions of order statistics rather than with their distributions. We need hardly point out that a heuristic argument like this can be entirely misleading and that the actual order of the remainder in our expansion will of course have to be investigated.

Define

$$(3.2.19) \quad \tilde{K}(x) = \phi(x) + \phi(x) \left\{ \frac{\sum a_j^2 E(2P_j - 1)^2 - 4\sigma^2(\sum a_j P_j)}{2\sum a_j^2} x + \right. \\ \left. + \frac{\sum a_j^3 (2\pi_j - 1)}{3(\sum a_j^2)^{3/2}} (x^2 - 1) + \frac{\sum a_j^4}{12(\sum a_j^2)^2} (x^2 - 3x) \right\},$$

where  $\sigma^2(Z)$  denotes the variance of a r.v.  $Z$ . Carrying out the type of computation outlined above we arrive at the following simplified version of theorem 3.2.2.

THEOREM 3.2.3. *Theorem 3.2.2 continues to hold if (3.2.17) is replaced by*

$$(3.2.20) \quad \sup_x \left| P\left(\frac{2T - \sum a_j}{(\sum a_j^2)^{1/2}} \leq x\right) - \tilde{K}\left(x - \frac{\sum a_j (2\pi_j - 1)}{(\sum a_j^2)^{1/2}}\right) \right| \leq \\ \leq A \left\{ N^{-5/4} + \sum \{E(2P_j - 1)^4\}^{5/4} + N^{-3/4} \left[ \sum \{E|P_j - \pi_j|^3\}^{4/9} \right]^{9/4} \right\}.$$

PROOF. See Albers, Bickel and van Zwet (1974).

Theorem 3.2.3 provides the basic expansion for the distribution of  $T$  under contiguous alternatives. In section 3.3 we shall be concerned with a further simplification of this expansion and a precise evaluation of the order of the remainder term.

### 3.3. CONTIGUOUS ALTERNATIVES

We first consider the case of contiguous location alternatives. Let  $F$  be a d.f. with a density  $f$  that is positive on  $R^1$ , symmetric about zero and four times differentiable with derivatives  $f^{(i)}$ ,  $i = 1, \dots, 4$ . Define functions

$$(3.3.1) \quad \psi_i = \frac{f^{(i)}}{f}, \quad i = 1, \dots, 4,$$

and suppose that positive numbers  $\epsilon$  and  $C$  exist such that for

$$(3.3.2) \quad m_1 = 6, m_2 = 3, m_3 = \frac{4}{3}, m_4 = 1,$$

$$\sup \left\{ \int_{-\infty}^{\infty} |\psi_i(x+y)|^{m_i} f(x) dx : |y| \leq \epsilon \right\} \leq C, \quad i = 1, \dots, 4.$$

Let  $X_1, X_2, \dots, X_N$  be i.i.d. with common d.f.  $G(x) = F(x-\theta)$  where

$$(3.3.3) \quad 0 \leq \theta \leq CN^{-1/2}$$

for some positive  $C$ . Note that (3.3.2) and (3.3.3) together imply continuity. Let  $0 < Z_1 < \dots < Z_N$  again denote the order statistics of  $|X_1|, \dots, |X_N|$ , and let  $T$  be defined by (3.2.2). Probabilities, expected values and variances under  $G$  will be denoted by  $P_\theta, E_\theta$  and  $\sigma_\theta^2$ ; under  $F$  they will be indicated as  $P_0, E_0$  and  $\sigma_0^2$ . Define

$$(3.3.4) \quad K_\theta(x) = \Phi(x) + \phi(x) \left\{ \frac{\sum_j^4 a_j}{12(\sum_j a_j^2)^2} (x^3 - 3x) - \theta \frac{\sum_j^3 E_0 \psi_1(Z_j)}{3(\sum_j a_j^2)^{3/2}} (x^2 - 1) + \right.$$

$$\left. + \frac{\theta^2}{2\sum_j a_j^2} [\sum_j^2 E_0 \psi_1^2(Z_j) - \sigma_0^2(\sum_j \psi_1(Z_j))]x + \right.$$

$$\left. + \frac{\theta^3}{6(\sum_j a_j^2)^{1/2}} \sum_j^1 E_0 [3\psi_1^3(Z_j) - 6\psi_1(Z_j)\psi_2(Z_j) + \psi_3(Z_j)] \right\},$$

$$(3.3.5) \quad \eta = -\theta \frac{\sum_j^1 E_0 \psi_1(Z_j)}{(\sum_j a_j^2)^{1/2}}.$$

We shall show that  $K_\theta(x-\eta)$  is an expansion to order  $N^{-1}$  for the d.f. of  $(2T - \sum_j a_j) / (\sum_j a_j^2)^{1/2}$ . The expansion will be established in theorem 3.3.1 and an evaluation of the order of the remainder will be given in theorem 3.3.2.

Let  $\pi(\theta)$  denote the power of the one-sided level  $\alpha$  test based on  $T$  for the hypothesis of symmetry against the alternative  $G(x) = F(x-\theta)$ . Suppose that for some  $\epsilon > 0$

$$(3.3.6) \quad \epsilon \leq \alpha \leq 1-\epsilon.$$

We prove that an expansion for  $\pi(\theta)$  is given by

$$(3.3.7) \quad \tilde{\pi}(\theta) = 1 - K_{\theta}(u_{\alpha} - \eta) + \phi(u_{\alpha} - \eta) \frac{\sum a_j^4}{12(\sum a_j^2)^2} (u_{\alpha}^3 - 3u_{\alpha}),$$

where  $u_{\alpha} = \Phi^{-1}(1-\alpha)$  denotes the upper  $\alpha$ -point of the standard normal distribution.

**THEOREM 3.3.1.** *Suppose that positive numbers  $c, C, \delta$  and  $\epsilon$  exist such that (3.2.14), (3.2.15), (3.3.2) and (3.3.3) are satisfied. Then there exists  $A > 0$  depending on  $N, a, F$  and  $\theta$  only through  $c, C, \delta$  and  $\epsilon$  and such that*

$$(3.3.8) \quad \sup_x |P_{\theta}\left(\frac{2T - \sum a_j}{(\sum a_j^2)^{1/2}} \leq x\right) - K_{\theta}(x - \eta)| \leq A\{N^{-5/4} + N^{-3/4}\theta^3[\sum\{E_0|\psi_1(Z_j) - E_0\psi_1(Z_j)|^3\}^{4/9}]^{9/4}\},$$

$$(3.3.9) \quad |\eta| \leq A,$$

$$(3.3.10) \quad \theta \frac{|\sum a_j^3 E_0 \psi_1(Z_j)|}{(\sum a_j^2)^{3/2}} \leq AN^{-1}, \quad \theta^2 \frac{\sum a_j^2 E_0 \psi_1^2(Z_j)}{\sum a_j^2} \leq AN^{-1},$$

$$\frac{\theta^3}{(\sum a_j^2)^{1/2}} |\sum a_j E_0 [3\psi_1^3(Z_j) - 6\psi_1(Z_j)\psi_2(Z_j) + \psi_3(Z_j)]| \leq AN^{-1}.$$

If, in addition, (3.3.6) is satisfied there exists  $A' > 0$  depending on  $N, a, F, \theta$  and  $\alpha$  only through  $c, C, \delta$  and  $\epsilon$  and such that

$$(3.3.11) \quad |\pi(\theta) - \tilde{\pi}(\theta)| \leq A'\{N^{-5/4} + N^{-3/4}\theta^3[\sum\{E_0|\psi_1(Z_j) - E_0\psi_1(Z_j)|^3\}^{4/9}]^{9/4}\}.$$

**PROOF.** The first step is to show that the conditions of the present theorem imply the conditions of theorem 3.2.3. This comes down to the verification of (3.2.16), which is easily done by applying (3.3.2) and (3.3.3). Hence theorem 3.2.3 holds and we must show that (3.3.8) is implied by (3.2.20). This is achieved by Taylor expansion with respect to  $\theta$ , which is a highly technical and laborious procedure; the interested reader is referred to appendix 1 of Albers, Bickel and van Zwet (1974). The main problem is that not only  $P_j = f(Z_j - \theta) / [f(Z_j - \theta) + f(Z_j + \theta)]$ , but also its distribution depends on  $\theta$  because  $Z_j$  is the  $j$ -th absolute order statistic of a sample from  $F(x - \theta)$ .

Once the expansion for  $\tilde{K}(x - [\sum a_j(2\pi_j - 1)] / [(\sum a_j^2)^{1/2}])$  has been established,



it remains to show that  $|\tilde{K}(x - [\sum a_j(2\pi_j - 1)] / [(\sum a_j^2)^{1/2}]) - K_\theta(x - \eta)|$  and the remainder in (3.2.20) are of the order of the remainder in (3.3.8). Then (3.3.8) is proved.

As concerns (3.3.9) and (3.3.10) we note that both are immediate consequences of the results of appendix 1 in Albers, Bickel and van Zwet (1974). The one-sided level  $\alpha$  test based on  $T$  rejects the hypothesis if  $(2T - \sum a_j) / (\sum a_j^2)^{1/2} \geq \xi_\alpha$ , with possible randomization if equality occurs. Taking  $\theta = 0$  in (3.3.8) we find that

$$1 - \Phi(\xi_\alpha) - \phi(\xi_\alpha) \frac{\sum a_j^4}{12(\sum a_j^2)^2} (\xi_\alpha^3 - 3\xi_\alpha) = \alpha + O(N^{-5/4}),$$

and hence because of (3.2.14) and (3.3.6)

$$(3.3.12) \quad \xi_\alpha = u_\alpha - \frac{\sum a_j^4}{12(\sum a_j^2)^2} (u_\alpha^3 - 3u_\alpha) + O(N^{-5/4}).$$

The power of this test against the alternative  $F(x - \theta)$  is

$$(3.3.13) \quad \begin{aligned} \pi(\theta) = & 1 - K_\theta(\xi_\alpha - \eta) + \\ & + O(N^{-5/4} + N^{-3/4} \theta^3 [\sum \{E_0 |\psi_1(Z_j) - E_0 \psi_1(Z_j)|^3\}^{4/9}]^{9/4}). \end{aligned}$$

In (3.3.13) we expand  $K_\theta(\xi_\alpha - \eta)$  around  $u_\alpha - \eta$ . Noting that  $|\xi_\alpha - u_\alpha| = O(N^{-1})$  and using (3.2.14) and (3.3.10) we find (3.3.11).  $\square$

REMARK. As we shall see in section 3.4,  $\sigma_0^2(\sum a_j \psi_1(Z_j))$  is typically  $O(N)$  and therefore our expansion is of the form  $K_\theta(x) = \Phi(x) + N^{-1}A(x) + o(N^{-1})$ , for a certain bounded  $A(x)$ . There is no term of the order  $N^{-1/2}$  because of a certain symmetry in the situation.

For  $i = 1, 2, 3$ , define functions  $\Psi_i$  on  $(0, 1)$  by

$$(3.3.14) \quad \Psi_i(t) = \psi_i(F^{-1}(\frac{1+t}{2})) = \frac{f^{(i)}(F^{-1}(\frac{1+t}{2}))}{f(F^{-1}(\frac{1+t}{2}))}.$$

THEOREM 3.3.2. Suppose that positive numbers  $C$  and  $\delta$  exist such that (3.3.3) is satisfied and that  $|\Psi_i'(t)| \leq C(t(1-t))^{-4/3+\delta}$  for all  $0 < t < 1$ . Then there exists  $A'' > 0$  depending on  $N$ ,  $F$  and  $\theta$  only through  $C$  and  $\delta$  and such that

$$N^{-3/4} \theta^3 \left[ \sum \{E_0 |\psi_1(Z_j) - E_0 \psi_1(Z_j)|^3\}^{4/9} \right]^{9/4} \leq A'' N^{-5/4}.$$

For the highly technical proof of this result the reader is referred to appendix 2 of Albers, Bickel and van Zwet (1974).

The more general case where  $X_1, X_2, \dots, X_N$  are i.i.d. r.v.'s with common d.f.  $G(x) = F(x, \theta)$  can be dealt with analogously; the computations become even more laborious but no new techniques are needed. For this reason we only give the results and omit the proofs.

Suppose that (3.3.3) holds and that

$$f_{ij}(x, \theta) = \frac{\partial^{i+j+1}}{\partial x^{i+1} \partial \theta^j} F(x, \theta)$$

exists for  $i+j \leq 4$ ,  $i \geq -1$ ,  $j \geq 0$ . We shall simply write  $f(x, \theta)$  for the density  $f_{00}(x, \theta)$  and assume that  $f(x, 0) = f(-x, 0)$  for all  $x$ . Next we define

$$\tilde{\psi}_{ij} = \frac{f_{ij}}{f},$$

(3.3.15) for  $i+j \leq 4$ ,  $i \geq -1$ ,  $j \geq 0$ .

$$\psi_{ij}(x) = \tilde{\psi}_{ij}(x, 0),$$

Assume that positive numbers  $\varepsilon$  and  $C$  exist such that for

$$(3.3.16) \quad m_1 = 6, m_2 = 3, m_3 = \frac{4}{3}, m_4 = 1, i \geq -1, j \geq 0, 1 \leq i+j \leq 4, \\ \sup \left\{ \int_{-\infty}^{\infty} |\tilde{\psi}_{ij}(x, y)|^{m_{i+j}} [1 + \tilde{\psi}_{-11}^6(x, y)] f(x, 0) dx : |y| \leq \varepsilon \right\} \leq C.$$

Under the additional assumption

$$\int_{-\infty}^{\infty} f_{0i}(x, \theta) dx = 0, i = 1, 2, 0 \leq \theta \leq CN^{-1/2},$$

it may be shown that (3.3.3) and (3.3.16) together imply contiguity.

Define

$$\begin{aligned}
 (3.3.17) \quad \bar{K}_\theta(x) = & \phi(x) + \phi(x) \left\{ \frac{\sum a_j^4}{12(\sum a_j^2)^2} (x^3 - 3x) + \theta \frac{\sum a_j^3 E_0 [\psi_{01}(z_j) - \psi_{01}(-z_j)]}{6(\sum a_j^2)^{3/2}} (x^2 - 1) \right. \\
 & + \theta^2 \frac{\sum a_j^2 E_0 [\psi_{01}(z_j) - \psi_{01}(-z_j)]^2 - \sigma_0^2 (\sum a_j [\psi_{01}(z_j) - \psi_{01}(-z_j)])}{8\sum a_j^2} x + \\
 & - \frac{\theta^4}{2\sum a_j^2} (\sum a_j E_0 \tilde{p}_{02}(z_j))^2 x - \frac{\theta^2}{(\sum a_j^2)^{1/2}} \sum a_j E_0 \tilde{p}_{02}(z_j) + \\
 & \left. - \frac{\theta^3}{3(\sum a_j^2)^{1/2}} \sum a_j E_0 \tilde{p}_{03}(z_j) \right\},
 \end{aligned}$$

where

$$\begin{aligned}
 (3.3.18) \quad \tilde{p}_{02}(x) = & \frac{1}{4} [\psi_{02}(x) - \psi_{02}(-x) - \psi_{01}^2(x) + \psi_{01}^2(-x) + \\
 & - (\psi_{-11}(x) - \psi_{-11}(-x)) (\psi_{11}(x) + \psi_{11}(-x) - \psi_{10}(x) [\psi_{01}(x) - \psi_{01}(-x)])],
 \end{aligned}$$

$$\begin{aligned}
 (3.3.19) \quad \tilde{p}_{03}(x) = & \frac{1}{4} \{ \psi_{03}(x) - \psi_{03}(-x) - 3\psi_{01}(x)\psi_{02}(x) + 3\psi_{01}(-x)\psi_{02}(-x) + \\
 & + \frac{3}{2} (\psi_{01}(x) + \psi_{01}(-x)) (\psi_{01}^2(x) - \psi_{01}^2(-x)) + \\
 & - \frac{3}{2} (\psi_{11}(x) + \psi_{11}(-x) - \psi_{10}(x) [\psi_{01}(x) - \psi_{01}(-x)]) (\psi_{-12}(x) - \psi_{-12}(-x)) \} + \\
 & + 3\{ \psi_{12}(x) + \psi_{12}(-x) - 3\psi_{01}(x)\psi_{11}(x) - 3\psi_{01}(-x)\psi_{11}(-x) - \psi_{11}(x)\psi_{01}(-x) + \\
 & - \psi_{01}(x)\psi_{11}(-x) - \psi_{10}(x) (\psi_{02}(x) - \psi_{02}(-x)) \} + \\
 & + 3\psi_{10}(x) (\psi_{01}^2(x) - \psi_{01}^2(-x)) \} \xi_{01}(x) + 3\{ \psi_{21}(x) - \psi_{21}(-x) + \\
 & - 3\psi_{10}(x) (\psi_{11}(x) + \psi_{11}(-x)) - \psi_{20}(x) (\psi_{01}(x) - \psi_{01}(-x)) \} + \\
 & + 3\psi_{10}^2(x) (\psi_{01}(x) - \psi_{01}(-x)) \} \xi_{01}^2(x).
 \end{aligned}$$

Here  $\xi_{01}(x) = -\frac{1}{2} [\psi_{-11}(x) - \psi_{-11}(-x)]$ . Finally define

$$(3.3.20) \quad \bar{n} = \frac{\theta}{2} \frac{\sum a_j E_0 [\psi_{01}(z_j) - \psi_{01}(-z_j)]}{(\sum a_j^2)^{1/2}},$$

$$(3.3.21) \quad \bar{\pi}(\theta) = 1 - \bar{K}_\theta(u_\alpha - \bar{\eta}) + \phi(u_\alpha - \bar{\eta}) \frac{\sum a_j^4}{12(\sum a_j^2)^2} (u_\alpha^3 - 3u_\alpha).$$

Now we have in analogy to theorem 3.3.1

THEOREM 3.3.3. Suppose that positive numbers  $c, C, \delta$  and  $\epsilon$  exist such that (3.2.14), (3.2.15), (3.3.3) and (3.3.16) are satisfied. Then there exists  $A > 0$  depending on  $N, a, F$  and  $\theta$  only through  $c, C, \delta$  and  $\epsilon$  and such that

$$(3.3.22) \quad \sup_x |P_\theta\left(\frac{2T - \sum a_j}{(\sum a_j^2)^{1/2}} \leq x\right) - \bar{K}_\theta(x - \bar{\eta})| \leq$$

$$\leq A\{N^{-5/4} + N^{-3/4}\theta^3[\sum\{E_0|\psi_{01}(Z_j) - E_0\psi_{01}(Z_j)|^3\}^{4/9}]^{9/4} +$$

$$+ N^{-3/4}\theta^3[\sum\{E_0|\psi_{01}(-Z_j) - E_0\psi_{01}(-Z_j)|^3\}^{4/9}]^{9/4} +$$

$$+ N^{-1/4}\theta^3\sigma_0(\sum a_j[\psi_{01}(Z_j) - \psi_{01}(-Z_j)])\}.$$

If, in addition, (3.3.6) is satisfied, there exists  $A' > 0$  depending on  $N, a, F, \theta$  and  $\alpha$  only through  $c, C, \delta$  and  $\epsilon$  and such that

$$(3.3.23) \quad |\pi(\theta) - \bar{\pi}(\theta)| \leq A'\{N^{-5/4} + N^{-3/4}\theta^3[\sum\{E_0|\psi_{01}(Z_j) - E_0\psi_{01}(Z_j)|^3\}^{4/9}]^{9/4} +$$

$$+ N^{-3/4}\theta^3[\sum\{E_0|\psi_{01}(-Z_j) - E_0\psi_{01}(-Z_j)|^3\}^{4/9}]^{9/4} +$$

$$+ N^{-1/4}\theta^3\sigma_0(\sum a_j[\psi_{01}(Z_j) - \psi_{01}(-Z_j)])\}.$$

We conclude this section with some remarks on the relation between the general and the location case. In the first place, condition (3.3.16) is a straightforward generalization of condition (3.3.2) except for the  $\tilde{\psi}_{-11}^6$ -term. In the location case,  $|\tilde{\psi}_{-11}(x, \theta)| = |\frac{\partial}{\partial \theta} F(x - \theta) / \frac{\partial}{\partial x} F(x - \theta)| = 1$ , which explains why it does not occur in (3.3.2).

In the location case  $f(x, \theta)$  is not only symmetric in  $x$  about  $x = \theta$ , but also in  $\theta$  about  $\theta = x$ . Then  $\psi_{01}(x) = -\psi_{01}(-x) = -\psi_1(x)$ ,  $\tilde{p}_{02}(x) \equiv 0$ ,  $\xi_{01}(x) \equiv 0$  and  $\tilde{p}_{03}(x) = -\frac{1}{2}[3\psi_1^3(x) - 6\psi_1(x)\psi_2(x) + \psi_3(x)]$ . Inserting these results in (3.3.17) and (3.3.20) we again obtain (3.3.4) and (3.3.5). In view of these facts, the main difference between  $K_\theta(x)$  and  $\bar{K}_\theta(x)$  is the presence of the terms

$$-\frac{\theta^2}{(\sum a_j^2)^{1/2}} \sum a_j E_0 \tilde{p}_{02}(Z_j) - \frac{\theta^4}{2 \sum a_j^2} (\sum a_j E_0 \tilde{p}_{02}(Z_j))^2_x$$

in (3.3.17). The first of these terms is in general not  $O(N^{-1})$  but only  $O(N^{-1/2})$ .

The first three terms on the right side of (3.3.22) and (3.3.23) are again generalizations of the remainder in (3.3.8) and (3.3.11). The last term in (3.3.22) and (3.3.23), however, is new. It is due to the fact that  $\tilde{p}_{02}(x) \equiv 0$  does not hold in general. As  $\sigma_0^2(\sum a_j [\psi_{01}(Z_j) - \psi_{01}(-Z_j)])$  is typically  $O(N)$ , this remainder term is of the right order, i.e.  $O(N^{-5/4})$ . To be more precise: with the aid of theorem 3.3.2 it can be shown to be  $O(N^{-7/6})$ . If in this theorem  $|\Psi_1'(t)| = O((t(1-t))^{-5/4+\delta})$  instead of  $O((t(1-t))^{-4/3+\delta})$ , we get  $O(N^{-5/4})$ .

This concludes our treatment of the general case; in the next section we again restrict attention to contiguous location alternatives.

#### 3.4. EXACT AND APPROXIMATE SCORES AND CONTIGUOUS LOCATION ALTERNATIVES

The expansions given in section 3.3 for contiguous location alternatives can be simplified further if we make certain assumptions about the scores  $a_j$ . Consider a continuous function  $J$  on  $(0,1)$  and let  $U_{1:N} < U_{2:N} < \dots < U_{N:N}$  denote order statistics of a sample of size  $N$  from the uniform distribution on  $(0,1)$ . For  $N = 1, 2, \dots$  we define the exact scores generated by  $J$  by

$$(3.4.1) \quad a_j = a_{jN} = EJ(U_{j:N}), \quad j = 1, 2, \dots, N,$$

and the approximate scores generated by  $J$  by

$$(3.4.2) \quad a_j = a_{jN} = J\left(\frac{j}{N+1}\right), \quad j = 1, \dots, N.$$

For almost all well known linear rank tests the scores are of one of these two types. The locally most powerful rank test against location alternatives of type  $F$  is based on exact scores generated by the function  $-\Psi_1$ , where  $\Psi_1$  is defined in (3.3.14).

So far, we have systematically kept the order of the remainder in our expansions down to  $O(N^{-5/4})$ . From this point on, however, we shall be content with a remainder that is  $O(N^{-1})$ , because otherwise we would have to impose

rather restrictive conditions. In the previous sections, we have also consistently stressed the fact that the remainder depends on  $a$  and  $F$  only through certain constants occurring in our conditions, thus in effect indicating classes of scores and distributions for which the expansion holds uniformly. As the number of these constants is becoming rather large, we prefer to formulate our results from here on for a fixed score function  $J$  and a fixed d.f.  $F$ . The reader can easily construct uniformity classes for himself by using the results of section 3.3 and tracing the development of appendix 2 of Albers, Bickel and van Zwet (1974).

DEFINITION 3.4.1.  $J$  is the class of functions  $J$  on  $(0,1)$  that are twice continuously differentiable and nonconstant on  $(0,1)$  and satisfy

$$(3.4.3) \quad \int_0^1 J^4(t) dt < \infty.$$

$$(3.4.4) \quad \limsup_{t \rightarrow 0,1} t(1-t) \left| \frac{J''(t)}{J'(t)} \right| < \frac{3}{2}.$$

$F$  is the class of d.f.'s  $F$  on  $R^1$  with positive densities  $f$  that are symmetric about zero, four times differentiable and such that for  $\psi_i = f^{(i)}/f$ ,  $\psi_i(t) = \psi_i(F^{-1}(\frac{1+t}{2}))$ ,  $m_1 = 6$ ,  $m_2 = 3$ ,  $m_3 = \frac{4}{3}$ ,  $m_4 = 1$ ,

$$(3.4.5) \quad \limsup_{y \rightarrow 0} \int_{-\infty}^{\infty} |\psi_i(x+y)|^{m_i} f(x) dx < \infty, \quad i = 1, \dots, 4,$$

$$(3.4.6) \quad \limsup_{t \rightarrow 0,1} t(1-t) \left| \frac{\psi_1''(t)}{\psi_1'(t)} \right| < \frac{3}{2}.$$

For  $J \in J$  and  $F \in F$ , let

$$(3.4.7) \quad \begin{aligned} \tilde{K}_\theta(x) = & \phi(x) + \phi(x) \left\{ \frac{N^{-1} \int_0^1 J^4(t) dt}{12 \left( \int_0^1 J^2(t) dt \right)^2} (x^3 - 3x) + \right. \\ & - N^{-1/2} \frac{\int_0^1 J^3(t) \psi_1(t) dt}{3 \left( \int_0^1 J^2(t) dt \right)^{3/2}} (x^2 - 1) + \frac{\theta^2}{2 \int_0^1 J^2(t) dt} \\ & \cdot \left[ \int_0^1 J^2(t) \psi_1^2(t) dt - \int_0^1 \int_0^1 J(s) \psi_1'(s) J(t) \psi_1'(t) (s \wedge t - st) ds dt \right] x + \\ & \left. + \frac{N^{1/2} \theta^3}{6 \left( \int_0^1 J^2(t) dt \right)^{1/2}} \int_0^1 J(t) [3\psi_1^3(t) - 6\psi_1(t)\psi_2(t) + \psi_3(t)] dt \right\}, \end{aligned}$$

$$(3.4.8) \quad K_{\theta,1}(x) = \tilde{K}_{\theta}(x) + \phi(x) \frac{N^{-1/2} \theta}{2(\int_0^1 J^2(t) dt)^{1/2}} \left\{ \frac{\int_0^1 J(t) \psi_1(t) dt}{\int_0^1 J^2(t) dt} \sum_{j=1}^N \sigma^2(J(U_{j:N})) + 2 \sum_{j=1}^N \text{cov}(J(U_{j:N}), \psi_1(U_{j:N})) \right\},$$

$$(3.4.9) \quad K_{\theta,2}(x) = \tilde{K}_{\theta}(x) + \phi(x) \frac{N^{-1/2} \theta}{2(\int_0^1 J^2(t) dt)^{1/2}} \left\{ \frac{\int_0^1 J(t) \psi_1(t) dt}{\int_0^1 J^2(t) dt} \int_{1/N}^{1-1/N} (J'(t))^2 t(1-t) dt + 2 \int_{1/N}^{1-1/N} J'(t) \psi_1'(t) t(1-t) dt \right\},$$

$$(3.4.10) \quad \tilde{\eta} = -N^{1/2} \theta \frac{\int_0^1 J(t) \psi_1(t) dt}{(\int_0^1 J^2(t) dt)^{1/2}},$$

$$(3.4.11) \quad \pi_i(\theta) = 1 - K_{\theta,i}(u_{\alpha}^{-\tilde{\eta}}) + \phi(u_{\alpha}^{-\tilde{\eta}}) N^{-1} \frac{\int_0^1 J^4(t) dt}{12(\int_0^1 J^2(t) dt)^2} (u_{\alpha}^3 - 3u_{\alpha}),$$

for  $i = 1, 2$ . Then, in the notation of section 3.3, we have for contiguous location alternatives and exact scores

**THEOREM 3.4.1.** *Let  $F \in \mathcal{F}$ ,  $J \in \mathcal{J}$ ,  $a_j = EJ(U_{j:N})$  for  $j = 1, \dots, N$ , and let  $0 \leq \theta \leq CN^{-1/2}$ ,  $\epsilon \leq \alpha \leq 1 - \epsilon$  for positive  $C$  and  $\epsilon$ . Then, for every fixed  $J, F, C$  and  $\epsilon$ , there exist positive numbers  $A, \delta_1, \delta_2, \dots$  such that  $\lim_{N \rightarrow \infty} \delta_N = 0$  and for every  $N$*

$$(3.4.12) \quad \sup_x \left| P_{\theta} \left( \frac{2T - \sum a_j}{(\sum a_j^2)^{1/2}} \leq x \right) - K_{\theta,1}(x - \tilde{\eta}) \right| \leq \delta_N N^{-1},$$

$$(3.4.13) \quad \sup_x \left| P_{\theta} \left( \frac{2T - \sum a_j}{(\sum a_j^2)^{1/2}} \leq x \right) - K_{\theta,2}(x - \tilde{\eta}) \right| \leq \delta_N N^{-1} + A \cdot N^{-3/2} \int_{1/N}^{1-1/N} |J'(t)| (|J'(t)| + |\psi_1'(t)|) t(1-t)^{1/2} dt,$$

$$(3.4.14) \quad |\pi(\theta) - \pi_1(\theta)| \leq \delta_N N^{-1},$$

$$(3.4.15) \quad |\pi(\theta) - \pi_2(\theta)| \leq \delta_N N^{-1} + \\ + AN^{-3/2} \int_{1/N}^{1-1/N} |J'(t)| (|J'(t)| + |\Psi_1'(t)|) (t(1-t))^{1/2} dt.$$

PROOF. For fixed  $J \in \mathcal{J}$ , positive constants  $c$ ,  $C$ , and  $\delta$  exist for which (3.2.14) and (3.2.15) hold for all  $N$  (cf. one of the remarks following the proof of theorem 3.2.2). Similarly, for fixed  $F \in \mathcal{F}$ , (3.3.2) is satisfied and it follows that the conclusions of theorem 3.3.1 hold with  $A$  and  $A'$  depending only on  $F$ ,  $J$ ,  $C$  and  $\epsilon$ . From appendix 2 of Albers, Bickel and van Zwet (1974) it is clear that (3.4.5) and (3.4.6) imply that the conclusion of theorem 3.3.2 holds with  $A''$  depending only on  $F$  and  $C$ . In this appendix it is also shown that

$$(3.4.16) \quad \frac{1}{N} \sum_{j=1}^N a_j^2 = \int_0^1 J^2(t) dt + o(1),$$

$$(3.4.17) \quad \frac{1}{N} \sum_{j=1}^N a_j^k E \Psi_1^{4-k}(U_{j:N}) = \int_0^1 J^k(t) \Psi_1^{4-k}(t) dt + o(1), \quad k = 1, \dots, 4,$$

$$(3.4.18) \quad \frac{1}{N} \sum_{j=1}^N a_j E \Psi_1(U_{j:N}) \Psi_2(U_{j:N}) = \int_0^1 J(t) \Psi_1(t) \Psi_2(t) dt + o(1),$$

$$(3.4.19) \quad \frac{1}{N} \sum_{j=1}^N a_j E \Psi_3(U_{j:N}) = \int_0^1 J(t) \Psi_3(t) dt + o(1),$$

$$(3.4.20) \quad \frac{1}{N} \sigma^2 \left( \sum_{j=1}^N a_j \Psi_1(U_{j:N}) \right) = \int_0^1 \int_0^1 J(s) J(t) \Psi_1'(s) \Psi_1'(t) [s \wedge t - st] ds dt + o(1),$$

and furthermore

$$(3.4.21) \quad \frac{N^{-1/2} \sum_{j=1}^N a_j E \Psi_1(U_{j:N})}{\left( \sum_{j=1}^N a_j^2 \right)^{1/2}} = \\ = \frac{\int_0^1 J(t) \Psi_1(t) dt}{\left( \int_0^1 J^2(t) dt \right)^{1/2}} - \frac{1}{N} \frac{\sum_{j=1}^N \text{covar}(J(U_{j:N}), \Psi_1(U_{j:N}))}{\left( \int_0^1 J^2(t) dt \right)^{1/2}} + \\ + \frac{1}{2N} \frac{\int_0^1 J(t) \Psi_1(t) dt}{\left( \int_0^1 J^2(t) dt \right)^{3/2}} \sum_{j=1}^N \sigma^2(J(U_{j:N})) +$$



$$\begin{aligned}
+ o(N^{-1}) &= \frac{\int_0^1 J(t)\Psi_1(t)dt}{(\int_0^1 J^2(t)dt)^{1/2}} - \frac{1}{N} \frac{\int_{1/N}^{1-1/N} J'(t)\Psi_1'(t)t(1-t)dt}{(\int_0^1 J^2(t)dt)^{1/2}} + \\
&+ \frac{1}{2N} \frac{\int_0^1 J(t)\Psi_1(t)dt}{(\int_0^1 J^2(t)dt)^{3/2}} \int_{1/N}^{1-1/N} (J'(t))^2 t(1-t)dt + \\
&+ o(N^{-1}) + O(N^{-3/2} \int_{1/N}^{1-1/N} |J'(t)|(|J'(t)| + |\Psi_1'(t)|)(t(1-t))^{1/2} dt).
\end{aligned}$$

From (3.4.16) to (3.4.20) it is clear that  $K_\theta(x)$  in (3.3.4) satisfies  $K_\theta(x) = \tilde{K}_\theta(x) + o(N^{-1})$ . Applying this to the expansions  $K_\theta(x-\eta)$  and  $\tilde{\pi}(\theta)$  in theorem 3.3.1 and expanding these functions of  $\eta$  around the point  $\eta = \tilde{\eta}$ , we get the desired results in view of (3.4.21).  $\square$

In general the expansions given in theorem 3.4.1 will not hold if the exact scores are replaced by approximate scores  $a_j = J(\frac{j}{N+1})$ , because  $\eta - \tilde{\eta}$  will then give rise to a different term of order  $N^{-1}$ . If  $J = -\Psi_1$ , however, it is clear from appendix 2 of Albers, Bickel and van Zwet (1974), that expansions (3.4.13) and (3.4.15) are valid for approximate as well as exact scores. Also for  $J = -\Psi_1$ , these expansions may be simplified because  $F \in \mathcal{F}$  implies that by partial integration (cf. lemma 4.2.1)

$$(3.4.22) \quad \int_0^1 \int_0^1 \Psi_1(s)\Psi_1'(s)\Psi_1(t)\Psi_1'(t)(s\wedge t - st)dsdt = \frac{1}{4} \int_0^1 \Psi_1^4(t)dt - \frac{1}{4} (\int_0^1 \Psi_1^2(t)dt)^2,$$

$$(3.4.23) \quad \int_0^1 \Psi_1(t)[6\Psi_1(t)\Psi_2(t) - \Psi_3(t)]dt = \frac{10}{3} \int_0^1 \Psi_1^4(t)dt + \int_0^1 \Psi_2^2(t)dt.$$

It follows that in this case  $\tilde{\eta}$ ,  $K_{\theta,2}(x-\tilde{\eta})$  and  $\pi_2(\theta)$  reduce to

$$(3.4.24) \quad \eta_1 = N^{1/2} \theta [E_0 \psi_1^2(X_1)]^{1/2},$$

$$(3.4.25) \quad L_\theta(x) = \Phi(x-\eta_1) + \frac{\phi(x-\eta_1)}{72N} \{ \eta_2 [6(x^3-3x) + 6\eta_1(x^2-1) - 3\eta_1^2 x - 5\eta_1^3] + \\ + 12\eta_3 \eta_1^3 + 9\eta_1^2(x-\eta_1) + 36 \frac{\int_{1/N}^{1-1/N} (\Psi_1'(t))^2 t(1-t)dt}{E_0 \psi_1^2(X_1)} \eta_1 \},$$

where

$$(3.4.26) \quad \eta_2 = E_0 \psi_1^4(X_1) / [E_0 \psi_1^2(X_1)]^2, \\ \eta_3 = E_0 \psi_2^2(X_1) / [E_0 \psi_1^2(X_1)]^2,$$

and

$$(3.4.27) \quad \pi^*(\theta) = 1 - \Phi(u_\alpha - \eta_1) + \frac{\eta_1 \phi(u_\alpha - \eta_1)}{72N} \{-6\eta_2(u_\alpha^2 - 1) + 3(\eta_2 - 3)\eta_1 u_\alpha + \\ + (5\eta_2 - 12\eta_3 + 9)\eta_1^2 - 36 \frac{\int_{1/N}^{1-1/N} (\Psi_1'(t))^2 t(1-t) dt}{E_0 \psi_1^2(X_1)}\}.$$

Finally we note that for  $F \in \mathcal{F}$ ,  $-\Psi_1$  cannot be constant on  $(0,1)$  because the density  $f(x) = \frac{1}{2} \lambda \exp(-\lambda|x|)$  of the double exponential distribution is not differentiable at zero. It follows that  $-\Psi_1 \in \mathcal{J}$  for every  $F \in \mathcal{F}$ . We have shown

**THEOREM 3.4.2.** *Let  $F \in \mathcal{F}$  and let either  $a_j = -E\Psi_1(U_{j:N})$  for  $j = 1, \dots, N$  or  $a_j = -\Psi_1(\frac{j}{N+1})$  for  $j = 1, 2, \dots, N$ . Suppose that  $0 \leq \theta \leq CN^{-1/2}$  and  $\varepsilon \leq \alpha \leq 1 - \varepsilon$  for positive  $C$  and  $\varepsilon$ . Then, for every fixed  $F$ ,  $C$  and  $\varepsilon$ , there exist positive numbers  $A$ ,  $\delta_1$ ,  $\delta_2$ , ... such that  $\lim_{N \rightarrow \infty} \delta_N = 0$  and for every  $N$*

$$(3.4.28) \quad \sup_x \left| P_\theta \left( \frac{2T - \sum_j a_j}{(\sum_j a_j)^{1/2}} \leq x \right) - L_\theta(x) \right| \leq \\ \leq \delta_N N^{-1} + AN^{-3/2} \int_{1/N}^{1-1/N} (\Psi_1'(t))^2 t(1-t)^{1/2} dt,$$

$$(3.4.29) \quad |\pi(\theta) - \pi^*(\theta)| \leq \delta_N N^{-1} + AN^{-3/2} \int_{1/N}^{1-1/N} (\Psi_1'(t))^2 t(1-t)^{1/2} dt.$$

At this point it may be useful to make some remarks concerning the assumptions in theorems 3.4.1 and 3.4.2. Conditions (3.4.4) and (3.4.6) ensure that  $J'$  and  $\Psi_1'$  do not oscillate to wildly near 0 and 1. They also limit the growth of these functions near 0 and 1, but in this respect conditions (3.4.3) and (3.4.5) for  $i = 1$  are typically much stronger. Together with (3.4.4) and (3.4.6) they imply that  $J'(t) = o((t(1-t))^{-5/4})$  and  $\Psi_1'(t) = o((t(1-t))^{-7/6})$  near 0 and 1.

For expansions (3.4.13), (3.4.15), (3.4.28) and (3.4.29) to be meaningful rather than just formally correct, even stronger growth conditions have to be imposed. Consider, for example, expansion (3.4.29) and suppose, as is typically the case, that  $\Psi_1'$  remains bounded near 0. If  $\Psi_1'(t) = o((1-t)^{-1})$  near 1, then the right side in (3.4.29) is  $o(N^{-1})$  and the expansion makes sense. However, if  $\Psi_1'(t)$  is of exact order  $(1-t)^{-1}$ , the expansion reduces to

$$\pi(\theta) = 1 - \Phi(u_\alpha - \eta_1) - \frac{\eta_1 \phi(u_\alpha - \eta_1) \int_0^{1-1/N} (\psi_1'(t))^2 t(1-t) dt}{2N \int_0^1 \psi_1^2(t) dt} + O(N^{-1}),$$

and if  $\psi_1'(t) \sim (1-t)^{-1-\delta}$  near 1 for some  $0 < \delta < \frac{1}{6}$ , all we have left in (3.4.29) is  $\pi(\theta) = 1 - \Phi(u_\alpha - \eta_1) + O(N^{-1+2\delta})$ .

We conclude this section with a few applications of theorems 3.4.1 and 3.4.2. The tedious computations will be omitted almost completely. First we consider Wilcoxon's signed rank test (W), which is based on the scores  $a_j = \frac{j}{N+1}$ . This is the locally most powerful rank test against logistic (L) location alternatives  $G(x) = F(x-\theta)$ , with  $F(x) = 1/(1+e^{-x})$  and  $\theta = O(N^{-1/2})$ . As in this case  $-\Psi_1(t) = t$  the conditions of theorem 3.4.2 are easily verified and we get

$$(3.4.30) \quad \pi_{W,L}(\theta) = 1 - \Phi(u_\alpha - \eta_1) - \frac{\phi(u_\alpha - \eta_1) \eta_1}{20N} \{3u_\alpha^2 + u_\alpha \eta_1 + 2 + \eta_1^2\} + o(N^{-1}),$$

where  $\eta_1 = (\frac{N}{3})^{1/2} \theta$ . The power of W against normal (N) alternatives  $G(x) = \Phi(x-\theta)$ ,  $\theta = O(N^{-1/2})$  may be found from theorem 3.4.1. We now have  $\Psi_1(t) = -\Phi^{-1}(\frac{1+t}{2})$  and therefore

$$\begin{aligned} \lim_{t \rightarrow 1} (1-t) \left| \frac{\Psi_1''(t)}{\Psi_1'(t)} \right| &= \lim_{t \rightarrow 1} (1-t) \frac{\phi^{-1}(\frac{1+t}{2})}{2\phi(\phi^{-1}(\frac{1+t}{2}))} = \\ &= \lim_{x \rightarrow \infty} \frac{x(1-\phi(x))}{\phi(x)} = 1 < \frac{3}{2}, \end{aligned}$$

as for positive x

$$\phi(x) \left( \frac{1}{x} - \frac{1}{3} \right) < 1 - \phi(x) < \frac{\phi(x)}{x}.$$

The remaining conditions of theorem 3.4.1 are verified easily and we may apply (3.4.15). The expression thus obtained can be simplified further by evaluating for  $k = 0, 1, 2, 3, 4$

$$\int_0^1 y^k [\phi^{-1}(\frac{1+y}{2})]^{4-k} dy.$$

This can be done by partial integration. In doing so, only two integrals occur that are not entirely trivial:  $\int_0^\infty \phi^2(x) \phi(x) dx$  and  $\int_0^\infty \phi^2(x) \phi^2(x) dx$ . For the first one we note

$$\begin{aligned}
\int_0^\infty \phi^2(x) \Phi(x) dx &= (2\pi)^{-3/2} \int_0^\infty \int_{-\infty}^x e^{-x^2 - \frac{1}{2}y^2} dy dx = \\
&= (2\pi)^{-3/2} \int_{-\pi/2}^{\pi/4} \int_0^\infty e^{-\frac{1}{2}r^2(2\cos^2\phi + \sin^2\phi)} r dr d\phi = \\
&= (2\pi)^{-3/2} \int_{-\pi/2}^{\pi/4} \frac{d\phi}{1 + \cos^2\phi} = (2\pi)^{-3/2} \int_{-\infty}^1 \frac{dt}{2+t^2} = \\
&= \frac{1}{4\sqrt{\pi}} \left( \frac{\arctan \frac{1}{2}\sqrt{2}}{\pi} + \frac{1}{2} \right).
\end{aligned}$$

Furthermore, for  $x > 0$ , we have

$$\begin{aligned}
\phi^2(x) &= \frac{1}{2\pi} \int_{-\infty}^x \int_{-\infty}^x e^{-\frac{1}{2}(y^2+z^2)} dy dz = \frac{1}{2\pi} \int_{-\pi/2}^{\pi/4} (1 - e^{-\frac{x^2}{2\cos^2\phi}}) d\phi + \\
&+ \frac{1}{2\pi} \int_{\pi/4}^{\pi} (1 - e^{-\frac{x^2}{2\sin^2\phi}}) d\phi + \frac{1}{2\pi} \int_{\pi}^{3\pi/2} d\phi = 1 - \frac{1}{\pi} \int_{\pi/4}^{\pi} e^{-\frac{x^2}{2\sin^2\phi}} d\phi.
\end{aligned}$$

Hence,

$$\begin{aligned}
\int_0^\infty \phi^2(x) \Phi^2(x) dx &= \int_0^\infty \phi^2(x) dx - \frac{1}{2\pi^2} \int_{\pi/4}^{\pi} \int_0^\infty e^{-\frac{x^2}{2}\left(2 + \frac{1}{\sin^2\phi}\right)} dx d\phi = \\
&= \frac{1}{4\sqrt{\pi}} - \frac{1}{2\pi\sqrt{\pi}} \int_{\pi/4}^{\pi} \left(2 + \frac{1}{\sin^2\phi}\right)^{-1/2} d\phi = \frac{1}{8\sqrt{\pi}}.
\end{aligned}$$

Application of these results leads to

$$\begin{aligned}
(3.4.31) \quad \pi_{W,N}(\theta) &= 1 - \Phi(u_\alpha - \tilde{\eta}) - \frac{\tilde{\eta}\phi(u_\alpha - \tilde{\eta})}{N} \left\{ u_\alpha^2 \left( \frac{6 \arctan \frac{1}{4}\sqrt{2}}{\pi} - \frac{9}{20} \right) + \right. \\
&+ u_\alpha \tilde{\eta} \left( \frac{49}{20} - \frac{2}{3}\sqrt{3} - \frac{12 \arctan \frac{1}{4}\sqrt{2}}{\pi} \right) + \left( \frac{44}{20} - \sqrt{2} - \frac{6 \arctan \frac{1}{4}\sqrt{2}}{\pi} \right) + \\
&+ \tilde{\eta}^2 \left( \frac{6 \arctan \frac{1}{4}\sqrt{2}}{\pi} + \frac{2}{3}\sqrt{3} - \frac{43}{20} + \frac{\pi}{9} \right) \left. \right\} + o(N^{-1}),
\end{aligned}$$

where  $\tilde{\eta} = \left(\frac{3N}{\pi}\right)^{1/2}\theta$ .

The second test we consider is the one sample normal scores test (NS) which is based on the scores  $a_j = E\Phi^{-1}\left(\frac{1+U_j}{2}\right)$ . Its power against the normal and logistic alternatives described above satisfies

$$(3.4.32) \quad \pi_{NS,N}(\theta) = 1 - \Phi(u_\alpha - \eta_1) - \frac{\eta_1 \phi(u_\alpha - \eta_1)}{4N} \{u_\alpha^2 - 1 + \\ + 2 \int_0^{\Phi^{-1}(1 - \frac{1}{2N})} \frac{(2\Phi(x) - 1)(1 - \Phi(x))}{\phi(x)} dx\} + o(N^{-1}),$$

where  $\eta_1 = N^{1/2}\theta$  and

$$(3.4.33) \quad \pi_{NS,L}(\theta) = 1 - \Phi(u_\alpha - \tilde{\eta}) - \frac{\tilde{\eta} \phi(u_\alpha - \tilde{\eta})}{N} \left\{ \frac{u_\alpha^2}{12} + u_\alpha \tilde{\eta} \left( -\frac{5}{12} + \frac{\pi}{6} \right) + \left( \frac{23}{12} - \sqrt{2} \right) + \right. \\ \left. \tilde{\eta}^2 \left( \arctan \frac{23\sqrt{2} + 1}{20} - \frac{\pi}{3} \right) - \frac{1}{2} \int_0^{\Phi^{-1}(1 - \frac{1}{2N})} \frac{(2\Phi(x) - 1)(1 - \Phi(x))}{\phi(x)} dx \right\} + o(N^{-1}),$$

where now  $\tilde{\eta} = \left(\frac{N}{\pi}\right)^{1/2}\theta$ . We note that theorem 3.4.2 ensures that (3.4.32) will also hold for van der Waerden's one sample test which is based on the approximate scores  $a_j = \Phi^{-1}[(N+1+j)/(2(N+1))]$ . As concerns the integral in (3.4.32) and (3.4.33), one can show that (cf. Bickel and van Zwet (1978), formulas (5.54) and (5.55))

$$(3.4.34) \quad \int_0^{\Phi^{-1}(1 - \frac{1}{2N})} \frac{(2\Phi(x) - 1)(1 - \Phi(x))}{\phi(x)} dx = \frac{1}{2} \log \log N + \frac{1}{2}\gamma + o(1),$$

where  $\gamma$  is Euler's constant

$$\gamma = \lim_{k \rightarrow \infty} \left\{ \sum_{j=1}^k \frac{1}{j} - \log k \right\} = 0.577216\dots$$

### 3.5. THE SIGN TEST

The method developed in this chapter cannot be used for the sign test, as was pointed out in section 3.2. The lattice character of the sign test is too pronounced, which is caused by the fact that all scores are equal for this test. However, it is exactly this equality of scores that yields another, very simple method of finding and justifying a power expansion for the sign test: it makes it possible to write  $T$  as

$$T = \sum_{X_j > 0} 1,$$

the number of positive elements in the sample. This has a binomial distribution, both under the hypothesis and under alternatives. With the aid of a well-known expansion for the binomial distribution function, this leads to the desired expansion for the power of the sign test.

Let  $X_1, X_2, \dots, X_N$  be i.i.d. r.v.'s with continuous d.f.  $F(x-\theta)$ , where  $F(-x) = 1 - F(x)$  for all  $x$ . Define

$$(3.5.1) \quad \tau = N^{1/2}(2F(\theta)-1).$$

Let  $0 < \alpha < 1$  and let  $\pi(\theta)$  be the power of the sign test with size  $\alpha$  for  $H_0 : \theta = 0$  against  $H_1 : \theta > 0$ . Define

$$(3.5.2) \quad \tilde{\pi}(\theta) = 1 - \Phi(u_\alpha - \tau) - \frac{\tau \phi(u_\alpha - \tau)}{12N} \{u_\alpha^2 + u_\alpha \tau - 3\tau^2 + 24\gamma_\alpha(1-\gamma_\alpha) - 3\}.$$

Here

$$(3.5.3) \quad \gamma_\alpha = \frac{(N+1)}{2} + \frac{N^{1/2}u_\alpha}{2} - \left[ \frac{(N+1)}{2} + \frac{N^{1/2}u_\alpha}{2} \right] + O(N^{-1/2}),$$

where  $[y]$  is the integer part of  $y$ .

LEMMA 3.5.1. *Suppose that  $|\tau| \leq C$  for some constant  $C > 0$ . Then there exists  $A > 0$  depending on  $N, \theta$  and  $F$  only through  $C$  such that*

$$|\pi(\theta) - \tilde{\pi}(\theta)| \leq AN^{-3/2}.$$

PROOF. Let  $Y$  have a binomial distribution with parameters  $N$  and  $p$ , where  $0 < p < 1$ . Define  $q = 1-p$ ,  $\sigma^2 = Npq$ ,  $u_k = (k-Np)/\sigma$  for  $k = 0, 1, \dots, N$ . If  $u_k = O(1)$ , the following expansion holds

$$(3.5.4) \quad P(Y \leq k) = \\ = \Phi\left\{u_{k+1/2} + \frac{p-q}{6\sigma}u_{k+1/2}^2\right\} + \frac{(5-14pq)u_{k+1/2}^3 + (-2+2pq)u_{k+1/2}}{72\sigma^2} \\ + O(N^{-3/2}).$$

Expansions of this type are given by Molenaar (1970).

Clearly  $T = \sum_{X_j > 0} 1$  has a binomial distribution with parameters  $N$  and  $p$ , with  $p = P_\theta(X_1 > 0) = P_0(X_1 > -\theta) = 1 - F(-\theta) = F(\theta) = \frac{1}{2}(1 + \tau N^{-1/2})$ . For this choice of  $p$ ,

$$\frac{p-q}{6\sigma} = \frac{\tau}{3N} + O(N^{-2}),$$

and hence (3.5.4) simplifies to

$$(3.5.5) \quad P_\theta(T \leq k) = \Phi\{u_{k+1/2} + \frac{\tau}{3N}(u_{k+1/2}^2 - 1) + \frac{1}{12N}(u_{k+1/2}^3 - u_{k+1/2})\} + O(N^{-3/2}).$$

For power computations, we need an expansion for  $P_\theta(T < k) + \gamma P_\theta(T = k)$ , for  $0 < \gamma \leq 1$ . Note that

$$P_\theta(T < k) + \gamma P_\theta(T = k) = \gamma P_\theta(T \leq k) + (1-\gamma)P_\theta(T \leq k-1)$$

and

$$\begin{aligned} & \gamma \Phi(u_{k+1/2}) + (1-\gamma)\Phi(u_{k-1/2}) = \\ & = \Phi(u_{k+\gamma-1/2}) + \frac{\gamma(1-\gamma)}{2\sigma^2} \phi'(u_{k+\gamma-1/2}) + O(N^{-3/2}) = \\ & = \Phi(u_{k(\gamma)}) + O(N^{-3/2}), \end{aligned}$$

with

$$(3.5.6) \quad u_{k(\gamma)} = \frac{k+\gamma - Np - \frac{1}{2}}{\sigma} \left(1 - \frac{\gamma(1-\gamma)}{2\sigma^2}\right).$$

Hence

$$(3.5.7) \quad \begin{aligned} P_\theta(T < k) + \gamma P_\theta(T = k) = \\ = \Phi\{u_{k(\gamma)} + \frac{\tau}{3N}(u_{k(\gamma)}^2 - 1) + \frac{(u_{k(\gamma)}^3 - u_{k(\gamma)})}{12N}\} + O(N^{-3/2}). \end{aligned}$$

Under  $H_0$   $\tau = 0$ . Let  $k_\alpha$  and  $\gamma_\alpha$  be such that  $P_0(T < k_\alpha) + \gamma_\alpha P_0(T = k_\alpha) = 1 - \alpha = \Phi(u_\alpha)$ , with  $k_\alpha$  an integer and  $0 < \gamma_\alpha \leq 1$ . As  $0 < \alpha < 1$ ,  $\alpha$  constant, we have  $u_\alpha = O(1)$  and we may use (3.5.7). This leads to

$$(3.5.8) \quad \frac{k_\alpha + \gamma_\alpha - 1/2N - 1/2}{1/2N^{1/2}} \left(1 - \frac{2\gamma_\alpha(1-\gamma_\alpha)}{N}\right) = u_\alpha - \frac{(u_\alpha^3 - u_\alpha)}{12N} + O(N^{-3/2}),$$

which shows that  $\gamma_\alpha$  satisfies (3.5.3). Under  $H_1$  we have  $\tau > 0$  and  $1 - \pi(\theta) = P_\theta(T < k_\alpha) + \gamma_\alpha P_\theta(T = k_\alpha)$ . Before applying (3.5.7) again we note that

$$(3.5.9) \quad \begin{aligned} & \frac{k_\alpha + \gamma_\alpha - 1/2N(1+\tau N^{-1/2}) - 1/2}{1/2N^{1/2}} \left(1 - \frac{2\gamma_\alpha(1-\gamma_\alpha)}{N}\right) \left(1 - \frac{\tau}{N}\right)^{-1/2} = \\ & = \left(\frac{k_\alpha + \gamma_\alpha - 1/2N - 1/2}{1/2N^{1/2}} - \tau\right) \left(1 - \frac{2\gamma_\alpha(1-\gamma_\alpha)}{N}\right) \left(1 + \frac{\tau}{2N} + O(N^{-2})\right) = \\ & = (u_\alpha - \tau) \left(1 + \frac{\tau}{2N}\right) + \frac{2\tau\gamma_\alpha(1-\gamma_\alpha)}{N} - \frac{(u_\alpha^3 - u_\alpha)}{12N} + O(N^{-3/2}). \end{aligned}$$

Inserting this result in (3.5.7), we get  $\pi(\theta) = \tilde{\pi}(\theta) + O(N^{-3/2})$ . The uniformity of the  $O$ -symbol is evident from the proof.  $\square$

COROLLARY 3.5.1. If  $|\tau| \leq C$  and  $F$  has a density  $f$  that is three times differentiable with  $\sup_{|x| \leq \epsilon} |f^{(3)}(x)| \leq c$ , for some constants  $\epsilon$ ,  $C$  and  $c > 0$ , we may replace  $\tilde{\pi}(\theta)$  in Lemma 3.5.1 by

$$\begin{aligned} \tilde{\pi}(\theta) = & 1 - \phi(u_\alpha - 2N^{1/2}\theta f(0)) - \frac{\theta f(0)}{6\sqrt{N}} \phi(u_\alpha - 2N^{1/2}\theta f(0)) \{u_\alpha^2 + \\ & + 2N^{1/2}\theta f(0)u_\alpha - 2N\theta^2 \left(\frac{f''(0)}{f(0)} + 6(f(0))^2\right) + 24\gamma_\alpha(1-\gamma_\alpha) - 3\}. \end{aligned}$$

The constant  $A$  depends on  $N$ ,  $\theta$  and  $F$  only through  $\epsilon$ ,  $C$  and  $c$  for this choice of  $\tilde{\pi}(\theta)$ .

PROOF. Immediate.  $\square$

The conditions of the lemma and its corollary are satisfied for e.g. the normal or the logistic d.f.; the double-exponential d.f. satisfies the conditions of the lemma, but not of its corollary.



## CHAPTER 4

## DEFICIENCIES BETWEEN VARIOUS TESTS FOR THE ONE SAMPLE PROBLEM

## 4.1. INTRODUCTION

One of the results of the previous chapter is an expansion to  $o(N^{-1})$  for the power of the locally most powerful (LMP) rank test for the one sample problem. In this chapter we obtain similar expansions for various other types of test that are in some sense optimal for this one sample problem. Using the expansions thus obtained, we can evaluate deficiencies between any pair of the tests involved. Such evaluations make sense since the LMP rank test has asymptotic relative efficiency 1 with respect to all tests considered in this chapter.

In section 4.2 we investigate two parametric tests; in section 4.3 permutation tests are dealt with. Scale invariant tests are considered in section 4.4, the randomized rank score tests due to Bell and Doksum (1965) in section 4.5. Finally, the deficiency evaluations mentioned above, take place in section 4.6.

## 4.2. PARAMETRIC TESTS

Let  $X_1, X_2, \dots, X_N$  be i.i.d. r.v.'s with d.f.  $F(x-\theta)$ , where  $F$  is known and has a density  $f$  that is positive on  $R^1$  and symmetric about zero. For the testing problem  $H_0 : \theta = 0$  against  $H_1 : \theta > 0$  the Neyman-Pearson lemma asserts that the test that rejects  $H_0$  for large values of  $\prod_{j=1}^N [f(X_j - \theta_1)/f(X_j)]$  is most powerful against the alternative  $\theta = \theta_1$ . It follows that for every  $\theta > 0$ , the envelope power at  $\theta$  equals the power at  $\theta$  of the test that rejects  $H_0$  for large values of  $S = \sum_{j=1}^N S_j$ , where  $S_j = \frac{1}{\theta} \log [f(X_j - \theta)/f(X_j)]$ .

In general, a uniformly most powerful (UMP) test against  $H_1$  does not exist and no single test attains the envelope power for all  $\theta > 0$ . If this is the case, one may consider the use of the LMP test in the sense of Lehmann (1959), p. 342. A test  $\phi_0$  is LMP if, given any other test  $\phi$  with the same level, there exists  $\Delta > 0$  such that the powers satisfy  $\pi_{\phi_0}(\theta) \geq \pi_{\phi}(\theta)$  for all  $0 < \theta < \Delta$ . A LMP test of  $\theta = 0$  against  $\theta > 0$  exists and is defined by the fact that it maximizes  $\pi'(0)$  among all tests with the same level. It

follows that in the present case the LMP test is based on the teststatistic  $S^* = \sum_{j=1}^N S_j^*$ , with  $S_j^* = -f'(X_j)/f(X_j)$ .

In the sequel we shall derive power expansions to  $o(N^{-1})$  for the envelope power and for the test based on  $S^*$ . First we shall deal with the envelope power; the result for  $S^*$  is proved in a similar fashion.

Since  $S$  is a sum of i.i.d. r.v.'s, the obvious thing to do is to establish Edgeworth expansions to  $o(N^{-1})$  for the d.f. of  $S$ , both under  $F(x)$  and  $F(x-\theta)$ . Before we can evaluate the standardized cumulants required for such expansions, the following preliminaries are needed. Assume that  $f$  is five times continuously differentiable and define

$$(4.2.1) \quad \begin{aligned} \psi_i(x) &= \frac{d^i f(x)}{dx^i} / f(x), & i &= 1, 2, \dots, 5, \\ \zeta_i(x) &= \frac{d^i}{dx^i} (\log f(x)), & i &= 0, 1, \dots, 5, \end{aligned}$$

where  $\zeta_0 = \log f$ . The connection between the  $\psi_i$  and  $\zeta_i$  is as follows

$$(4.2.2) \quad \begin{aligned} \zeta_1 &= \psi_1, \quad \zeta_2 = \psi_2 - \psi_1^2, \quad \zeta_3 = \psi_3 - 3\psi_1\psi_2 + 2\psi_1^3, \\ \zeta_4 &= \psi_4 - 3\psi_2^2 - 4\psi_1\psi_3 + 12\psi_1^2\psi_2 - 6\psi_1^4, \\ \zeta_5 &= \psi_5 - 5\psi_1\psi_4 - 10\psi_2\psi_3 + 30\psi_1\psi_2^2 + 20\psi_1^2\psi_3 - 60\psi_1^3\psi_2 + 24\psi_1^5. \end{aligned}$$

Next we give two lemmas. For lemma 4.2.1, compare lemma a of Hájek and Šidák (1967), p.19; the second lemma is proved in Albers, Bickel and van Zwet (1974) and is an application of Taylor's formula with Cauchy's form of the remainder.

LEMMA 4.2.1. *Let  $h$  be a real differentiable function with derivative  $h'$ . If  $|h|$  and  $|h'|$  are both summable, we have  $\lim_{x \rightarrow \pm\infty} h(x) = 0$ .*

PROOF.  $\int_{-\infty}^{\infty} |h'(x)| dx < \infty$  and hence by the dominated convergence theorem  $\lim_{v \rightarrow \infty} \int_{c'_v}^{c_v} h'(x) dx = 0$  for any pair of sequences  $\{c_v\}$  and  $\{c'_v\}$  with  $c_v \rightarrow \infty$ ,  $c'_v \rightarrow \infty$  for  $v \rightarrow \infty$  and  $c'_v \leq c_v$  for all  $v$ . This implies that  $h(c_v) - h(c'_v) \rightarrow 0$  for  $v \rightarrow \infty$ , i.e.  $\lim_{x \rightarrow \infty} h(x)$  exists. Since  $h$  itself is also summable it follows that  $\lim_{x \rightarrow \infty} h(x) = 0$ . In the same way of course  $\lim_{x \rightarrow -\infty} h(x) = 0$ .  $\square$



LEMMA 4.2.3. Let  $\{\delta_N\}$  be a sequence of positive real numbers with  $\lim_{N \rightarrow \infty} \delta_N = 0$ . Let for some  $N$

$$(4.2.5) \quad 0 < \theta < \delta_N.$$

Suppose that positive constants  $\epsilon$  and  $C$  exist such that

$$(4.2.6) \quad \sup \left\{ \int_{-\infty}^{\infty} |\psi_i(x+y)|^{5/i} f(x) dx : |y| \leq \epsilon \right\} \leq C, \quad i = 1, \dots, 5.$$

Then there exists  $A > 0$  depending on  $N$ ,  $\theta$  and  $F$  only through  $\{\delta_N\}$ ,  $\epsilon$  and  $C$  and such that

$$(4.2.7) \quad \begin{aligned} & |E_0 S - \left\{ -\frac{N\theta}{2} E_0 \psi_1^2(X_1) + \frac{N\theta^3}{24} [E_0 \psi_2^2(X_1) - \frac{2}{3} E_0 \psi_1^4(X_1)] \right\}| \leq AN\theta^4, \\ & |\sigma_0^2(S) - \left\{ NE_0 \psi_1^2(X_1) + \frac{N\theta^2}{36} [5E_0 \psi_1^4(X_1) - 3E_0 \psi_2^2(X_1) - 9(E_0 \psi_1^2(X_1))^2] \right\}| \leq AN\theta^3, \\ & |\kappa_3^0 - \frac{\theta}{2(E_0 \psi_1^2(X_1))^{3/2}} [3(E_0 \psi_1^2(X_1))^2 - E_0 \psi_1^4(X_1)]| \leq A\theta^2, \\ & |\kappa_4^0 - \left\{ \frac{E_0 \psi_1^4(X_1)}{(E_0 \psi_1^2(X_1))^2} - 3 \right\}| \leq A\theta. \end{aligned}$$

PROOF. As  $E_0 S = NE_0 S_1$  we have to find  $E_0 S_1$ . Expansion around  $\theta = 0$  shows that

$$(4.2.8) \quad \begin{aligned} S_1 &= \frac{1}{\theta} [\zeta_0(X_1 - \theta) - \zeta_0(X_1)] = -\zeta_1(X_1) + \frac{\theta}{2} \zeta_2(X_1) + \\ & - \frac{\theta^2}{6} \zeta_3(X_1) + \frac{\theta^3}{24} \zeta_4(X_1) + \frac{1}{\theta} [\zeta_0(X_1 - \theta) - \sum_{j=0}^4 \zeta_j(X_1) \frac{(-\theta)^j}{j!}]. \end{aligned}$$

Application of lemma 4.2.2 with  $q(x, t) = \zeta_0(x - t)$  gives

$$\begin{aligned} E_0 \left| \frac{1}{\theta} [\zeta_0(X_1 - \theta) - \sum_{j=0}^4 \zeta_j(X_1) \frac{(-\theta)^j}{j!}] \right| &\leq \\ &\leq \frac{\theta^4}{5!} \sup \{ E_0 |\zeta_5(X_1 - v\theta)|; 0 \leq v \leq 1 \}. \end{aligned}$$

Conditions (4.2.5) and (4.2.6) ensure that  $E_0 |\psi_i^{5/i}(X_1 - \theta)|$  is bounded for

$i = 1, \dots, 5$  and for  $N$  sufficiently large. Hence by Hölder's inequality and the triangle inequality, we get from (4.2.2) that  $|E_0 \zeta_i^{5/i}(X_1 - \theta)|$  is also bounded. This shows that the expectation of the last term in (4.2.8) is  $O(\theta^4)$ .

The remaining terms in (4.2.8) are dealt with in the following way. First we note that  $E_0 \zeta_i(2i-1)(X_1) = 0$ ,  $i = 1, 2, 3$ , by virtue of the symmetry of  $f$ . From (4.2.6) we have  $\int_{-\infty}^{\infty} |f^{(i)}(x)| dx < \infty$ ,  $i = 0, 1, \dots, 5$ . Hence by lemma 4.2.1,  $\lim_{x \rightarrow \pm\infty} f^{(i-1)}(x) = 0$ ,  $i = 1, \dots, 5$ , where  $f^{(0)} = f$ , and thus

$E_0 \psi_i(X_1) = \int_{-\infty}^{\infty} f^{(i)}(x) dx = 0$ ,  $i = 1, \dots, 5$ . These two steps show that

$$E_0 S_1 = -\frac{\theta}{2} E_0 \psi_1^2(X_1) + \frac{\theta^3}{24} E_0 [-3\psi_2^2(X_1) - 4\psi_1(X_1)\psi_3(X_1) + 12\psi_1^2(X_1)\psi_2(X_1) + 6\psi_1^4(X_1)] + O(\theta^4).$$

By partial integration and another application of lemma 4.2.1 we observe that  $E_0 \psi_1^2(X_1)\psi_2(X_1) = 2/3 E_0 \psi_1^4(X_1)$  and  $E_0 \psi_1(X_1)\psi_3(X_1) = 2/3 E_0 \psi_1^4(X_1) - E_0 \psi_2^2(X_1)$ . With this the statement for  $E_0 S$  is proved.

The formula for  $\sigma_0^2(S)$  is proved analogously; the only point that needs some explanation is the expansion of  $S_1^2$ .

$$\begin{aligned} S_1^2 &= -\zeta_1(X_1)S_1 + \frac{\theta}{2}\zeta_2(X_1)S_1 - \frac{\theta^2}{6}\zeta_3(X_1)S_1 + \\ &+ \frac{1}{\theta}[\zeta_0(X_1 - \theta) - \sum_{j=0}^3 \zeta_j(X_1) \frac{(-\theta)^j}{j!}]S_1 = \\ &= \zeta_1^2(X_1) - \theta\zeta_1(X_1)\zeta_2(X_1) + \frac{\theta^2}{3}\zeta_1(X_1)\zeta_3(X_1) + \frac{\theta^2}{4}\zeta_2^2(X_1) + \\ &+ \frac{1}{\theta}[\zeta_0(X_1 - \theta) - \sum_{j=0}^3 \zeta_j(X_1) \frac{(-\theta)^j}{j!}] [-\zeta_1(X_1) + \frac{1}{\theta}\{\zeta_0(X_1 - \theta) - \zeta_0(X_1)\}] + \\ &+ \frac{1}{\theta}[\zeta_0(X_1 - \theta) - \sum_{j=0}^2 \zeta_j(X_1) \frac{(-\theta)^j}{j!}] \frac{\theta}{2}\zeta_2(X_1) + \\ &- \frac{1}{\theta}[\zeta_0(X_1 - \theta) - \sum_{j=0}^1 \zeta_j(X_1) \frac{(-\theta)^j}{j!}] \frac{\theta^2}{6}\zeta_3(X_1). \end{aligned}$$

By expanding  $S_1^2$  in this way it is possible to deal with the remainder terms without using more than (4.2.6).

As concerns  $\kappa_3^0$  and  $\kappa_4^0$ , we have by definition

$$\begin{aligned}\kappa_3^0 &= N^{1/2} \left[ \sum_{j=1}^N E_0(S_j - E_0 S_j)^3 \right] / \left[ \sum_{j=1}^N E_0(S_j - E_0 S_j)^2 \right]^{3/2} = \\ &= E_0(S_1 - E_0 S_1)^3 / \sigma_0^3(S_1), \\ \kappa_4^0 &= N \left[ \sum_{j=1}^N \{E_0(S_j - E_0 S_j)^4 - 3\sigma_0^4(S_j)\} \right] / \left[ \sum_{j=1}^N \sigma_0^2(S_j) \right]^2 = \\ &= E_0(S_1 - E_0 S_1)^4 / \sigma_0^4(S_1) - 3.\end{aligned}$$

The third and fourth central moment of  $S_1$  are found in the same way as  $E_0 S_1$  and  $\sigma_0^2(S_1)$ . We get  $E_0(S_1 - E_0 S_1)^3 = (\theta/2)[3\{E_0 \psi_1^2(X_1)\}^2 - E_0 \psi_1^4(X_1)] + O(\theta^2)$  and  $E_0(S_1 - E_0 S_1)^4 = E_0 \psi_1^4(X_1) + O(\theta)$ . From this the required expressions for  $\kappa_3^0$  and  $\kappa_4^0$  follow immediately.  $\square$

Let  $R_0$  and  $\rho_0$  be the d.f. and c.f., respectively, of  $(S - E_0 S) / \sigma_0(S)$  under  $F(x)$ . Let  $\tilde{R}_0$  be the Edgeworth expansion to  $o(N^{-1})$  for  $R_0$  (cf. (3.1.1)) and let  $\tilde{\rho}_0$  be the Fourier transform of  $\tilde{R}_0$ . Define

$$(4.2.9) \quad \tilde{R}_0^*(x) = \phi(x) + \phi(x) \left[ - \frac{\theta}{12N^{1/2} (E_0 \psi_1^2(X_1))^{3/2}} \{3(E_0 \psi_1^2(X_1))^2 + \right. \\ \left. - E_0 \psi_1^4(X_1)\} (x^2 - 1) - \frac{1}{24N} \left( \frac{E_0 \psi_1^4(X_1)}{(E_0 \psi_1^2(X_1))^2} - 3 \right) (x^3 - 3x) \right].$$

The following lemma gives an upper bound for  $|R_0 - \tilde{R}_0^*|$ .

LEMMA 4.2.4. Let  $\{\delta_N\}$  be a sequence of positive real numbers with  $\lim_{N \rightarrow \infty} \delta_N = 0$  and suppose that (4.2.5) and (4.2.6) hold. Suppose in addition that there exist positive constants  $c$  and  $\eta$  such that

$$(4.2.10) \quad \left| \frac{d}{dy} \psi_1(F^{-1}(y)) \right| \geq c, \quad y \in \tau \subset (0, 1),$$

where the interval  $\tau$  has length at least  $\eta$ . Then there exists  $A > 0$  depending on  $N$ ,  $\theta$  and  $F$  only through  $\{\delta_N\}$ ,  $\epsilon$ ,  $C$ ,  $c$  and  $\eta$  and such that

$$(4.2.11) \quad \sup_x |R_0(x) - \tilde{R}_0^*(x)| \leq A\{N^{-3/2} + N^{-1/2}\theta^2\}.$$

PROOF. Upon inserting the expressions for  $\kappa_3^0$  and  $\kappa_4^0$  from (4.2.7) in the Edgeworth expansion  $\tilde{R}_0$ , we obtain in view of (4.2.9) that

$$\sup_x |\tilde{R}_0(x) - \tilde{R}_0^*(x)| \leq A\{N^{-3/2} + N^{-1/2}\theta^2\}.$$

Hence it remains to show that  $\sup_x |R_0(x) - \tilde{R}_0(x)| \leq A\{N^{-3/2} + N^{-1/2}\theta^2\}$ . As the proof of this result is almost standard (cf. Feller (1966)), we do not give many details here. The emphasis will be on showing that, under condition (4.2.10), the distribution of  $S_1$  satisfies a strong non-lattice condition, i.e. that the modulus of its c.f. remains bounded away from 1 outside a neighbourhood of 0.

From lemma 3.2.1 it is clear that it is sufficient to prove that

$$\int_{-T}^T \left| \frac{\rho_0(t) - \tilde{\rho}_0(t)}{t} \right| dt \leq A\{N^{-3/2} + N^{-1/2}\theta^2\},$$

where  $T = bN^{1/2} \min(N, \theta^{-2})$ , for some  $b > 0$ . In analogy to the rank test case we bound the integral by

$$(4.2.12) \quad \int_{|t| \leq c_1 N^{1/2}} \left| \frac{\rho_0(t) - \tilde{\rho}_0(t)}{t} \right| dt + \int_{c_1 N^{1/2} \leq |t| \leq bN^{1/2} \min(N, \theta^{-2})} \left| \frac{\rho_0(t)}{t} \right| dt + \\ + \int_{c_1 N^{1/2} \leq |t| \leq bN^{1/2} \min(N, \theta^{-2})} \left| \frac{\tilde{\rho}_0(t)}{t} \right| dt,$$

and show that these three integrals are sufficiently small.

First consider the interval  $|t| \leq c_1 N^{1/2}$ . Let  $w_0(t)$  be the c.f. of  $S_1 - E_0 S_1$  under  $H_0$ , then  $\rho_0(t) = \exp[N \log w_0\{t/(N^{1/2}\sigma_0(S_1))\}]$ . From (4.2.10) it follows that  $E_0 \psi_1^2(X_1) \geq \bar{c}$ , for some  $\bar{c} > 0$ , depending on  $c$  and  $\eta$ . According to (4.2.7),  $\sigma_0^2(S_1) = E_0 \psi_1^2(X_1) + O(\theta^2)$  and hence  $\sigma_0^2(S_1) \geq \bar{c}/2$  for  $N$  sufficiently large. This proves that for some  $c_1 > 0$ ,  $\operatorname{Re}\{w_0\{t/(N^{1/2}\sigma_0(S_1))\}\} \geq \frac{1}{2}$  for  $|t| \leq c_1 N^{1/2}$ . Therefore  $\log w_0\{t/(N^{1/2}\sigma_0(S_1))\}$ , and hence  $\log \rho_0(t)$ , can be expanded around  $t = 0$ , for  $|t| \leq c_1 N^{1/2}$ . Doing so, we find that  $\rho_0(t) = \tilde{\rho}_0(t) + M(t)$ , where  $|M(t)| \leq N^{-3/2} |t|^5 Q(|t|) \exp(-t^2/4)$  for

$|t| \leq c_1 N^{1/2}$ , and  $Q$  is a polynomial with coefficients that depend only on  $\{\delta_N\}$ ,  $\epsilon$ ,  $C$ ,  $c$  and  $n$ . From this it is clear that the first integral in (4.2.12) is  $O(N^{-3/2})$ .

Just as in the sketch of the proof of theorem 3.2.1, one can show that

$$\int_{|t| \geq \log(N+1)} \left| \frac{\tilde{\rho}_0(t)}{t} \right| dt$$

is sufficiently small. Hence it remains to show that the second integral in (4.2.12) is  $O(N^{-3/2} + N^{-1/2}\theta^2)$ . Using the fact that  $\zeta_0$  is an even and  $\zeta_1$  is an odd function, we find

$$(4.2.13) \quad |w_0(t)| = \left| \int_0^\infty \left[ e^{it\zeta_1(x)} + e^{-it\zeta_1(x)} \right] e^{\frac{it}{\theta}[\zeta_0(x-\theta) - \zeta_0(x) + \theta\zeta_1(x)]} + \right. \\ \left. + e^{it\zeta_1(x)} e^{\frac{it}{\theta}[\zeta_0(x+\theta) - \zeta_0(x) - \theta\zeta_1(x)]} - e^{\frac{it}{\theta}[\zeta_0(x-\theta) - \zeta_0(x) + \theta\zeta_1(x)]} \right] f(x) dx \Big| \leq \\ \leq 2 \int_0^\infty |\cos t\psi_1(x)| f(x) dx + \int_0^\infty \left| e^{it\left[\frac{\zeta_0(x+\theta) - \zeta_0(x-\theta)}{\theta} - 2\zeta_1(x)\right]} - 1 \right| f(x) dx.$$

The last integral in (4.2.13) is less than or equal to

$$(4.2.14) \quad |t| \int_0^\infty \left| \frac{\zeta_0(x+\theta) - \zeta_0(x-\theta)}{\theta} - 2\zeta_1(x) \right| f(x) dx \leq \\ \leq \frac{|t|\theta^2}{12} E_0 |2\zeta_3(X_1 + v\theta)| \leq \theta^2 |t| \tilde{C},$$

for  $|v| \leq 1$  and some  $\tilde{C} > 0$ . The first inequality is a consequence of lemma 4.2.2, the second of (4.2.2) and (4.2.6).

Next we investigate the behaviour of  $\psi_1(F^{-1}(y))$  on  $\tau$ . This interval has the form  $\tau = (a, a + \tilde{\eta})$ , where  $0 < a < 1 - \tilde{\eta}$  and  $\eta \leq \tilde{\eta} < 1$ . Let  $\delta$  and  $\tilde{c}$  be constants with  $0 < \delta < \frac{\pi}{2}$  and  $\tilde{c} > 0$  and define, for a fixed  $|t| \geq \tilde{c}$ , the set  $\tilde{\tau}$  by

$$\tilde{\tau} = \{y | \exists \text{ integer } k, k\pi + \delta \leq t\psi_1(F^{-1}(y)) \leq (k+1)\pi - \delta\} \cap \tau.$$



Let  $\lambda$  be Lebesgue measure and denote  $\lambda(\tilde{\tau})$  as  $\tilde{\lambda}$ . We shall show that  $\tilde{\lambda} \geq \eta/2$  for  $\delta$  sufficiently small and depending only on  $c, \tilde{c}, C$  and  $\eta$ . Obviously, we have  $\lambda(\tau \cap \tilde{\tau}^c) = \tilde{\eta} - \tilde{\lambda}$  and therefore

$$(4.2.15) \quad \lambda(\{y | a + \frac{\tilde{\eta} - \tilde{\lambda}}{4} \leq y \leq a + \tilde{\eta} - \frac{\tilde{\eta} - \tilde{\lambda}}{4}\} \cap \tilde{\tau}^c) \geq \frac{\tilde{\eta} - \tilde{\lambda}}{2}.$$

Condition (4.2.10) implies that  $|\frac{d}{dy} t\psi_1(F^{-1}(y))| \geq |t|c$  on  $\tau$ . From the differentiability conditions on  $f$  and the assumption that  $f(x)$  is positive for all  $x$ , it follows that  $\frac{d}{dy} \psi_1(F^{-1}(y))$  is continuous on  $\tau$ . Hence (4.2.10) also implies that  $\psi_1(F^{-1}(y))$  is monotone on  $\tau$ . This gives, together with (4.2.15), that  $t\psi_1(F^{-1}(y))$  traverses at least  $[\frac{(\tilde{\eta} - \tilde{\lambda})c|t|}{4\delta}]$  intervals of the form  $(k\pi - \delta, k\pi + \delta)$  as  $y$  increases from  $a + (\tilde{\eta} - \tilde{\lambda})/4$  to  $a + \tilde{\eta} - (\tilde{\eta} - \tilde{\lambda})/4$ . Here  $[z]$  stands for the integer part of  $z$ . Because  $t\psi_1(F^{-1}(y))$  is monotone on  $\tau$

$$(4.2.16) \quad |t| |\psi_1(F^{-1}(a + \tilde{\eta} - \frac{\tilde{\eta} - \tilde{\lambda}}{4})) - \psi_1(F^{-1}(a + \frac{\tilde{\eta} - \tilde{\lambda}}{4}))| \geq \{[\frac{(\tilde{\eta} - \tilde{\lambda})c|t|}{4\delta}] - 1\}\pi.$$

This inequality leads to the desired lower bound for  $\tilde{\lambda}$ . Take  $\delta = \min \{nc\tilde{c}/16, c\pi\eta^2/(128C)\}$ . If  $(\tilde{\eta} - \tilde{\lambda})c\tilde{c} \leq 8\delta$ , then  $\tilde{\lambda} \geq \tilde{\eta} - 8\delta/(c\tilde{c}) \geq \tilde{\eta} - \eta/2 \geq \eta/2$ . If  $(\tilde{\eta} - \tilde{\lambda})c\tilde{c} \geq 8\delta$ , (4.2.16) implies that

$$|\psi_1(F^{-1}(a + \tilde{\eta} - \frac{\tilde{\eta} - \tilde{\lambda}}{4})) - \psi_1(F^{-1}(a + \frac{\tilde{\eta} - \tilde{\lambda}}{4}))| \geq \frac{(\tilde{\eta} - \tilde{\lambda})c\pi}{8\delta},$$

and hence

$$(4.2.17) \quad \int_0^1 |\psi_1(F^{-1}(y))| dy \geq \frac{(\tilde{\eta} - \tilde{\lambda})^2 c\pi}{32\delta}.$$

According to (4.2.6) the integral in (4.2.17) is at most  $C$  and therefore  $\tilde{\lambda} \geq \tilde{\eta} - \{32\delta C/(c\pi)\}^{1/2} \geq \eta/2$ .

In view of the above, there exists a positive constant  $\tilde{\delta}$ , depending only on  $c, \tilde{c}, C$  and  $\eta$  and such that, for all  $|t| \geq \tilde{c}$ , we have  $\lambda(\tilde{\tau}) = \tilde{\lambda} \geq \eta/2$  and on  $\tilde{\tau}$

$$|\cos\{t\psi_1(F^{-1}(y))\}| \leq 1 - \tilde{\delta}.$$

Hence, for  $|t| \geq \tilde{c}$ ,

$$(4.2.18) \quad 2 \int_0^\infty |\cos\{t\psi_1(x)\}| f(x) dx = \int_0^1 |\cos\{t\psi_1(F^{-1}(y))\}| dy \leq 1 - \frac{\tilde{\delta}\eta}{2}.$$

Combination of (4.2.14) and (4.2.18) leads to

$$|w_0\{t/(N^{1/2}\sigma_0(S_1))\}| \leq 1 - \frac{\tilde{\delta}\eta}{2} + \theta^2 |t|\tilde{c}/(N^{1/2}\sigma_0(S_1)),$$

for  $|t| \geq \tilde{c}N^{1/2}\sigma_0(S_1)$ . Now  $\sigma_0^2(S_1)$  is bounded above because of (4.2.6). Therefore,  $\tilde{c}\sigma_0(S_1) \leq c_1$  if  $\tilde{c}$  is chosen sufficiently small. On the other hand,  $\sigma_0^2(S_1) \geq \bar{c}/2$ . Together this gives

$$|w_0\{t/(N^{1/2}\sigma_0(S_1))\}| \leq 1 - \frac{\tilde{\delta}\eta}{2} + N^{-1/2}\theta^2 |t|\tilde{c}(\frac{\bar{c}}{2})^{-1/2},$$

for  $|t| \geq c_1N^{1/2}$ . Hence, there exists a constant  $b > 0$  such that for  $c_1N^{1/2} \leq |t| \leq bN^{1/2}\theta^{-2}$  we have  $|w_0\{t/(N^{1/2}\sigma_0(S_1))\}| \leq 1 - \tilde{\delta}\eta/4$ . But then  $|\rho_0(t)| \leq (1 - \tilde{\delta}\eta/4)^N$  and

$$\int_{c_1N^{1/2} \leq |t| \leq bN^{1/2}\theta^{-2}} \frac{\rho_0(t)}{t} |dt| \leq \frac{b}{c_1} N(1 - \frac{\tilde{\delta}\eta}{4})^N = O(N^{-3/2}). \quad \square$$

Let  $R_\theta$  be the d.f. of  $(S - E_\theta S)/\sigma_\theta(S)$  under  $F(x-\theta)$ . If  $X_1$  has d.f.  $F(x-\theta)$ , the r.v.  $Y_1 = X_1 - \theta$  clearly has d.f.  $F(x)$ . Furthermore,

$$S_1 = (1/\theta)\log\{f(X_1 - \theta)/f(X_1)\} = (1/(-\theta))\log\{f(Y_1 - (-\theta))/f(Y_1)\}.$$

Hence by changing  $\theta$  into  $(-\theta)$  in the right side of (4.2.9), we get an expansion  $\tilde{R}_\theta^*$  for  $R_\theta$ , such that  $\sup_x |R_\theta(x) - \tilde{R}_\theta^*(x)| \leq A\{N^{-3/2} + N^{-1/2}\theta^2\}$  under the conditions of lemma 4.2.4.

From  $\tilde{R}_\theta^*$  and  $\tilde{R}_\theta^*$  an expansion  $\tilde{\pi}_S(\theta)$  for the envelope power  $\pi_S(\theta)$  can be found easily. Define

$$(4.2.19) \quad n_1 = N^{1/2}\theta[E_0\psi_1^2(X_1)]^{1/2}, \quad n_2 = \frac{E_0\psi_1^4(X_1)}{[E_0\psi_1^2(X_1)]^2}, \quad n_3 = \frac{E_0\psi_2^2(X_1)}{[E_0\psi_1^2(X_1)]^2},$$

(cf. (3.4.24) and (3.4.26)) and

$$(4.2.20) \quad \tilde{\pi}_S(\theta) = 1 - \Phi(u_\alpha - n_1) + \Phi(u_\alpha - n_1) \left[ \frac{n_1(n_2 - 3)}{24N} (u_\alpha^2 - n_1 u_\alpha - 1) + \frac{n_1^3}{72N} (2n_2 - 3n_3) \right].$$

LEMMA 4.2.5. Let  $\{\delta_N\}$  be a sequence of positive real numbers with  $\lim_{N \rightarrow \infty} \delta_N = 0$  and assume that (4.2.5), (4.2.6) and (4.2.10) hold and suppose in addition that  $\epsilon' \leq \alpha \leq 1 - \epsilon'$  for some constant  $\epsilon' > 0$ . Then there exists  $A > 0$ , depending on  $N$ ,  $\theta$ ,  $\alpha$  and  $F$  only through  $\{\delta_N\}$ ,  $\epsilon$ ,  $C$ ,  $c$ ,  $n$  and  $\epsilon'$  and such that

$$(4.2.21) \quad |\pi_S(\theta) - \tilde{\pi}_S(\theta)| \leq A\{N^{-3/2} + N^{-1/2}\theta^2\}.$$

PROOF. From (4.2.9) it follows that the critical value  $\xi_\alpha$  satisfies

$$(4.2.22) \quad \xi_\alpha = u_\alpha - \frac{\eta_1(\eta_2 - 3)}{12N}(u_\alpha^2 - 1) + \frac{(\eta_2 - 3)}{24N}(u_\alpha^3 - 3u_\alpha) + O(N^{-3/2} + N^{-1/2}\theta^2).$$

Furthermore, we have for  $\pi_S(\theta)$

$$(4.2.23) \quad \pi_S(\theta) = 1 - R_\theta \left( \left[ \xi_\alpha + \frac{E_0 S - E_\theta S}{\sigma_0(S)} \right] \frac{\sigma_0(S)}{\sigma_1(S)} \right).$$

$E_0 S$  and  $\sigma_0(S)$  are given in (4.2.7),  $E_\theta S$  and  $\sigma_\theta(S)$  follow by changing  $\theta$  into  $(-\theta)$  in these expressions. Application of this and (4.2.22) to (4.2.23) leads to (4.2.21), with an additional remainder term of order  $N^{1/2}\theta^4\phi(N^{1/2}\theta)$ . This term can be omitted, as  $N^{1/2}\theta^4\phi(N^{1/2}\theta) = O(N^{-1/2}\theta^2)$ .  $\square$

REMARKS. 1. Conditions (4.2.6) and (4.2.10) determine a class of d.f.'s  $F$  for which expansion (4.2.20) holds uniformly. If we restrict attention to a fixed  $F$ , condition (4.2.6) can be weakened to

$$\limsup_{y \rightarrow 0} \int_{-\infty}^{\infty} |\psi_i^{5/i}(x+y)| f(x) dx < \infty, \quad i = 1, \dots, 5.$$

Furthermore, condition (4.2.10) can be omitted: if no uniformity is required, we only need that  $\psi_1(F^{-1}(y))$  is non-constant on  $(0,1)$ . This condition is always satisfied, since  $\psi_1(F^{-1}(y))$  is constant on  $(0,1)$  only for uniform d.f.'s. But these d.f.'s are already ruled out by the fact that their density is not everywhere positive and differentiable.

2. We have treated location alternatives with  $\theta = O(1)$ , since we shall need the results for such  $\theta$  in section 5.2. The typical case of interest, however, remains the case of contiguous location alternatives, where  $\theta = O(N^{-1/2})$ . Then  $\eta_1$ , and hence  $1/\{1 - \Phi(u_\alpha - \eta_1)\}$ , are  $O(1)$ .

As announced at the beginning of this section, we shall also give a power-expansion for the LMP test based on  $S^* = \sum_{j=1}^N S_j^*$ , where  $S_j^* = -\psi_1(X_j)$ . Under  $F(x)$  we have

$$E_0(S_1^*)^{2k} = E_0\psi_1^{2k}(X_1), \quad E_0(S_1^*)^{2k-1} = 0, \quad \text{for } k = 1, 2.$$

Under  $F(x-\theta)$  the necessary moments can be found by using

$$S_1^* = -\psi_1(X_1) = -\zeta_1(X_1) = -\zeta_1(Y_1) - \theta\zeta_2(Y_1) - \frac{\theta^2}{2}\zeta_3(Y_1) + \\ -\frac{\theta^3}{6}\zeta_4(Y_1) + [-\zeta_1(X_1) + \sum_{j=0}^3 \zeta_{j+1}(Y_1) \frac{\theta^j}{j!}],$$

where  $Y_1 = X_1 - \theta$  has d.f.  $F(x)$ . Proceeding in this way, we obtain expansions similar to (4.2.9), which are justified in a manner analogous to lemma 4.2.4. The final result is the following expansion  $\tilde{\pi}_{S^*}(\theta)$  for the power  $\pi_{S^*}(\theta)$ :

$$(4.2.24) \quad \tilde{\pi}_{S^*}(\theta) = 1 - \phi(u_\alpha - \eta_1) + \phi(u_\alpha - \eta_1) \left[ \frac{\eta_1(\eta_2 - 3)}{24N} (u_\alpha^2 - \eta_1 u_\alpha - 1) + \right. \\ \left. + \frac{\eta_1^3}{72N} (5\eta_2 - 12\eta_3 + 9) \right].$$

Under the assumptions of lemma 4.2.5 we have  $|\pi_{S^*}(\theta) - \tilde{\pi}_{S^*}(\theta)| \leq A\{N^{-3/2} + N^{-1/2}\theta^2\}$ . Comparison of (4.2.20) and (4.2.24) gives

$$\tilde{\pi}_S(\theta) - \tilde{\pi}_{S^*}(\theta) = \frac{\eta_1^3}{24N} (3\eta_3 - \eta_2 - 3)\phi(u_\alpha - \eta_1) = \\ = \frac{N^{1/2}\theta^3\sigma_0^2(\psi_1'(X_1))}{8[E_0\psi_1^2(X_1)]^{1/2}} \phi(u_\alpha - \eta_1) \geq 0,$$

where equality only occurs if  $\psi_1'(X_1)$  is constant a.s., i.e. if  $X_1$  is normally distributed. The fact that  $\tilde{\pi}_S(\theta) = \tilde{\pi}_{S^*}(\theta)$  here is obvious, since  $S$  and  $S^*$  are equivalent if  $X_1$  comes from  $\phi(x-\theta)$ .

#### 4.3. PERMUTATION TESTS

We shall start by showing how permutation tests can be derived in a natural

way (cf. Lehmann (1959)). Consider again the one sample problem:

$X_1, X_2, \dots, X_N$  are i.i.d. r.v.'s. Under the hypothesis they come from a d.f.  $F$ , which has a density  $f$  that is positive on  $R^1$  and symmetric around zero, under the alternative they come from a d.f.  $G$  which has a density  $g$  that is not symmetric around zero. Let  $X = (X_1, \dots, X_N)$ . In the previous section we considered LMP tests for this problem, which, of course, are not distribution-free. Here we shall restrict attention to distribution-free tests and search for the most powerful one in this restricted class.

If  $\phi$  is a distribution-free test, then  $\phi$  must have the same size  $\alpha$  for all  $F \in \mathcal{P}_X$ , the family of all d.f.'s with a continuous symmetric density, i.e.  $\phi$  has to be similar with respect to  $\mathcal{P}_X$ . A concept related to similarity is Neyman Structure (NS): a test  $\phi$  has NS with respect to a statistic  $T$  if  $T$  is sufficient for  $X$  with respect to  $\mathcal{P}_X$  and  $E(\phi(X)|T) = \alpha$  a.s. under  $\mathcal{P}_X$ . If  $\phi$  has NS, it is similar; the converse holds if  $T$  is also complete with respect to  $\mathcal{P}_X$ . Let  $Z = (Z_1, \dots, Z_N)$  denote again the vector of absolute order statistics of the  $X_j$ . As  $Z$  is sufficient and complete with respect to  $\mathcal{P}_X$ , the class of all similar tests for the one sample problem coincides with the class of all tests having  $E(\phi(X)|Z) = \alpha$  a.s. under  $H_0$ . This last condition can also be stated as

$$(4.3.1) \quad (2^N!)^{-1} \sum_{y \in S(x)} \phi(y) = \alpha \text{ a.s.},$$

where  $S(x)$  is the set of all  $y = (y_1, \dots, y_N)$ , giving rise to the same  $z$  as  $x$ .

Any test satisfying (4.3.1) is called a permutation test. In particular, every rank test satisfies (4.3.1) and therefore rank tests form a subclass of the family of permutation tests. This implies that the most powerful permutation test for a certain alternative is always at least as good as the most powerful rank test for that alternative. As they both possess the desirable property of being distribution-free, permutation tests are superior to rank tests. The only reason to prefer rank tests over permutation tests is of a practical nature: rank tests are much easier to apply.

In view of this relation between rank tests and permutation tests, it seems interesting to make deficiency comparisons between them. To this end we shall derive in this section a power expansion for the most powerful permutation test. First we shall derive the explicit form of its test-statistic if we consider a fixed alternative, under which the  $X_j$  come from a d.f.

G. Then we have for the conditional power

$$\begin{aligned} E(\phi(X)|Z) &= \sum_{y \in S(x)} \phi(y) \frac{P(Z=z, R=r, \text{sign } Y = \text{sign } y)}{P(Z=z)} = \\ &= \sum_{y \in S(x)} \phi(y) \frac{\prod_{j=1}^N g(y_j)}{\sum_{y \in S(x)} \prod_{j=1}^N g(y_j)}, \end{aligned}$$

where  $R = (R_1, \dots, R_N)$  is the vector of ranks for  $(|Y_1|, \dots, |Y_N|)$  and  $\text{sign } Y = (\text{sign } Y_1, \dots, \text{sign } Y_N)$ . Conditionally under (4.3.1),  $E(\phi(X)|Z)$  is maximal if

$$\phi(x) = \begin{cases} 1 & \text{for } \prod_{j=1}^N g(x_j) \geq c(Z), \\ 0 & \text{otherwise,} \end{cases}$$

where  $c(Z)$  depends on  $Z$  only. As  $\prod_{j=1}^N f(x_j)$  is constant over  $S(x)$ , the most powerful permutation test rejects for large values of  $\prod_{j=1}^N \{g(x_j)/f(x_j)\}$ , conditionally under  $Z$ .

We now restrict attention to contiguous location alternatives  $g(x) = f(x-\theta)$ , with  $0 \leq \theta \leq DN^{-1/2}$ , for some positive constant  $D$ . Then the most powerful permutation test can be based on  $(1/\theta) \sum_{j=1}^N \log\{f(X_j-\theta)/f(X_j)\}$ . As  $\sum_{j=1}^N \log\{f(-|X_j|-\theta)/f(-|X_j|)\} = \sum_{j=1}^N \log\{f(Z_j+\theta)/f(Z_j)\}$  is constant given  $Z = z$ , we can equivalently use

$$\frac{1}{2\theta} \sum_{j=1}^N \left[ \log \frac{f(X_j-\theta)}{f(X_j)} - \log \frac{f(-|X_j|-\theta)}{f(-|X_j|)} \right] = \frac{1}{2\theta} \sum_{X_j > 0} \log \frac{f(X_j-\theta)}{f(X_j+\theta)}.$$

A drawback of the above test is the fact that it is only optimal against one particular alternative  $\theta_1$ . For a composite hypothesis we may therefore prefer the LMP permutation test, which is based on

$$U = - \sum_{X_j > 0} \psi_1(X_j),$$

with  $\psi_1$  as defined in (4.2.1). The relation between  $U$  and the statistic of the most powerful permutation test is

$$\frac{1}{2\theta} \sum_{X_j > 0} \log \frac{f(X_j-\theta)}{f(X_j+\theta)} = U - \frac{\theta^2}{12} \sum_{X_j > 0} [\zeta_3(X_j - \nu_1 \theta) + \zeta_3(X_j + \nu_2 \theta)],$$

with  $0 \leq v_1, v_2 \leq 1$  and  $\zeta_3$  as defined in (4.2.1). This relation is the same as between  $S$  and  $S^*$  in the previous section. Note, however, that  $S-S^* = O(\theta)$  instead of  $O(\theta^2)$  as is the case here. This explains why  $\pi_{S^*}(\theta) - \pi_S(\theta) = O(N^{-1})$ , while the difference in the power of the two permutation tests is  $O(N^{-5/4})$ , as is shown in lemma 4.3.5. Hence, as concerns deficiencies, both tests perform equally well.

We shall proceed to give a power expansion for the test based on  $U$ . The unconditional distribution of  $U$  can be found easily, but since the critical value depends on  $Z$ , this does not lead to the power. We have to go through the following procedure: the distribution of  $U$ , conditionally on  $Z$ , has to be found, whereupon the conditional power can be evaluated as a function of  $Z$ . Then the unconditional power is found by taking the expectation with respect to  $Z$ .

In order to find the conditional distribution of  $U$ , it is useful to represent  $U$  as  $U = -\sum_{j=1}^N \psi_1(Z_j) V_j$ , where  $V_j = 1$  if the  $X_i$  corresponding to  $Z_j$  is positive, and  $V_j = 0$  otherwise. With this representation we can apply various results from Chapter 3. Just as in this chapter we have, conditionally on  $Z$ , the following situation: the  $-\psi_1(Z_j)$  are constant and  $V_1, \dots, V_N$  are independent with

$$(4.3.2) \quad P_\theta(V_j=1) = 1 - P_\theta(V_j=0) = \frac{f(Z_j-\theta)}{f(Z_j+\theta)+f(Z_j-\theta)}, \quad j = 1, \dots, N.$$

We introduce the following notation

$$A_j = -\psi_1(Z_j), \quad a_j = -\psi_1(z_j),$$

$$(4.3.3) \quad P_j = P_\theta(V_j=1|Z), \quad p_j = P_\theta(V_j=1|z),$$

$$Q_j = 2P_j - 1, \quad q_j = 2p_j - 1,$$

and  $A, a, P, p, Q, q$  are the corresponding  $N$ -dimensional vectors. Conditionally under  $Z = z$ ,  $2U$  has variance

$$(4.3.4) \quad \tau^2 = \sum_{j=1}^N (1 - q_j^2) a_j^2,$$

and the statistic  $2(U - \sum_{j=1}^N a_j p_j)/\tau$  has mean 0 and variance 1. Its third and fourth cumulants, multiplied by  $N^{1/2}$  and  $N$  respectively, are given by

$$(4.3.5) \quad \kappa_3 = -2N^{1/2} \left( \sum_{j=1}^N q_j (1-q_j^2) a_j^3 \right) / \tau^3, \quad \kappa_4 = -2N \left( \sum_{j=1}^N (1-q_j^2) (1-3q_j^2) a_j^4 \right) / \tau^4.$$

Define

$$(4.3.6) \quad R_\theta(x|z) = P_\theta \left( \frac{2(U - \sum_{j=1}^N a_j p_j)}{\tau} \leq x|z \right),$$

$$\gamma(\epsilon, \zeta, q) = \lambda \{x | \exists \text{ integer } j \text{ } |x - a_j| < \zeta, |q_j| \leq 1 - \epsilon\},$$

where  $0 < \epsilon < 1/2$ ,  $\zeta > 0$  and  $\lambda$  is Lebesgue measure.

The following lemma supplies an expansion for  $R_\theta(x|z)$ . From this point on, summation always runs from  $j = 1$  to  $j = N$ , unless stated otherwise.

LEMMA 4.3.1. *Let  $z$  and  $\theta$  be such that there exist positive numbers  $c$ ,  $C$ ,  $\delta$  and  $\epsilon$  for which the following conditions hold*

$$\frac{1}{N} \sum (1-q_j^2) a_j^2 \geq c, \quad \frac{1}{N} \sum a_j^4 \leq C, \quad \gamma(\epsilon, \zeta, q) \geq \delta N \zeta,$$

for some  $\zeta \geq N^{-3/2} \log N$ . Then there exists  $A > 0$ , depending on  $N$ ,  $z$  and  $\theta$  only through  $c$ ,  $C$ ,  $\delta$  and  $\epsilon$  and such that

$$(4.3.7) \quad R_\theta(x|z) = \Phi(x) + \phi(x) \left\{ \frac{\sum q_j a_j^3}{3(NB)^{3/2}} (x^2 - 1) + \frac{\sum a_j^4}{12(NB)^2} (x^3 - 3x) \right\} + R,$$

where  $B$  is an arbitrary positive constant and

$$(4.3.8) \quad |R| \leq A \{ N^{-5/4} + N^{-5/2} |\sum q_j a_j^3| |\tau^2 - NB| + N^{-3/2} |\sum q_j^3 a_j^3| + \\ + N^{-2} |\tau^2 - NB| + N^{-2} |\sum q_j^2 a_j^4| + N^{-2} |\sum q_j a_j^3| \}.$$

PROOF. Under the above conditions, theorem 3.2.1 can be applied to  $R_\theta(x|z)$ . This means that

$$(4.3.9) \quad R_\theta(x|z) = \tilde{R}_\theta(x|z) + O(N^{-5/4}),$$

where  $\tilde{R}_\theta$  is the Edgeworth expansion to  $O(N^{-1})$  for  $R_\theta$ , i.e.



$$(4.3.10) \quad \tilde{R}_\theta(x|z) = \phi(x) - \phi(x) \left\{ \frac{N^{-1/2} \kappa_3}{6} (x^2 - 1) + \frac{N^{-1} \kappa_4}{24} (x^3 - 3x) + \frac{N^{-1} \kappa_3^2}{72} (x^5 - 10x^3 + 15x) \right\}.$$

From (4.3.5) we obtain

$$\begin{aligned} \kappa_3 &= -2N^{1/2} \left( \frac{\sum q_j a_j^3 - \sum q_j^3 a_j^3}{(NB)^{3/2}} \right) \left( \frac{NB}{\tau^2} \right)^{3/2} = \\ &= -2 \frac{\sum q_j a_j^3}{NB^{3/2}} + O(N^{-2} |\sum q_j a_j^3| |\tau^2 - NB| + N^{-1} |\sum q_j^3 a_j^3|), \\ \kappa_4 &= -2N \left( \frac{\sum a_j^4 - 4 \sum q_j^2 a_j^4 + 3 \sum q_j^4 a_j^4}{(NB)^2} \right) \left( \frac{NB}{\tau^2} \right)^2 = - \frac{2 \sum a_j^4}{NB^2} + \\ &+ O(N^{-1} |\tau^2 - NB| + N^{-1} \sum q_j^2 a_j^4), \end{aligned}$$

$$\kappa_3^2 = \kappa_3 \kappa_3 = O(1) O(N^{-1} |\sum q_j a_j^3| + N^{-1} |\sum q_j^3 a_j^3|) = O(N^{-1} |\sum q_j a_j^3|).$$

Substitution of these expressions in (4.3.10) leads, in combination with (4.3.9), to the desired result.  $\square$

REMARK. The expansion in (4.3.7) is formally correct for any choice of  $B > 0$ . In future applications we shall choose  $B$  such that the expression in (4.3.8) is small. This is achieved by taking  $B = E_0 \psi_1^2(X_1)$ , which is the obvious thing to do in view of the definitions (4.3.3) and (4.3.4).

Let  $\pi(\theta|z)$  be the conditional power of the test based on  $U$  and let  $\alpha$  be the size of the test. In the next lemma an expansion for  $\pi(\theta|z)$  is given.

LEMMA 4.3.2. *Let  $z$  and  $\theta$  be such that the conditions of lemma 4.3.1 are satisfied and such that  $|\sum q_j a_j| \leq c' N^{1/2}$  for some constant  $c' > 0$ . If furthermore  $0 \leq \theta \leq DN^{-1/2}$ ,  $\epsilon' \leq \alpha \leq 1 - \epsilon'$  for positive constants  $D$  and  $\epsilon'$ , we have*

$$\pi(\theta|z) = \tilde{\pi}(\theta|z) + R,$$

where

$$\begin{aligned}
 (4.3.11) \quad \tilde{\pi}(\theta|z) = & 1 - \Phi(u_\alpha - \theta(NB)^{1/2}) - \phi(u_\alpha - \theta(NB)^{1/2}) \left[ \frac{\sum q_j a_j^3}{3(NB)^{3/2}} (u_\alpha + \right. \\
 & - \theta(NB)^{1/2})^2 - 1 + \frac{\sum a_j^4}{12(NB)^2} (-3u_\alpha^2 \theta(NB)^{1/2} + 3u_\alpha \theta^2 NB - \theta^3 (NB)^{3/2} + \\
 & + 3\theta(NB)^{1/2}) + \frac{1}{2} \frac{\sum q_j^2 a_j^2}{NB} (u_\alpha - \theta(NB)^{1/2}) + \frac{1}{2} \frac{(\sum a_j^2 - NB)}{NB} \theta(NB)^{1/2} + \\
 & - \frac{1}{8} \frac{(\sum a_j^2 - NB)^2}{(NB)^2} \theta(NB)^{1/2} \{3 + \theta(NB)^{1/2} (u_\alpha - \theta(NB)^{1/2})\} + \\
 & - \frac{\sum q_j a_j - \theta NB}{(NB)^{1/2}} + \frac{1}{2} \frac{(\sum q_j a_j - \theta NB)^2}{NB} (u_\alpha - \theta(NB)^{1/2}) + \\
 & \left. + \frac{1}{2} \frac{(\sum q_j a_j - \theta NB)(\sum a_j^2 - NB)}{(NB)^{3/2}} \{1 + \theta(NB)^{1/2} (u_\alpha - \theta(NB)^{1/2})\} \right],
 \end{aligned}$$

with B an arbitrary positive constant and

$$\begin{aligned}
 (4.3.12) \quad |R| \leq & A \{ N^{-5/4} + N^{-3} |\sum a_j^2 - NB|^{3+N-3/2} |\sum q_j a_j - \theta NB|^{3+N-3/2} (\sum q_j^2 a_j^2)^{3/2} + \\
 & + N^{-9/4} |\sum q_j a_j^3|^{3/2} + N^{-3/2} |\sum q_j^3 a_j^3| + N^{-2} |\sum q_j^2 a_j^4| \},
 \end{aligned}$$

where  $A > 0$  depends on  $N$ ,  $\alpha$ ,  $\theta$  and  $z$  only through  $c$ ,  $C$ ,  $\delta$ ,  $\epsilon$ ,  $c'$ ,  $D$  and  $\epsilon'$ .

PROOF. The conditions of lemma 4.3.1 are satisfied for some  $\theta \geq 0$ . Therefore they must be satisfied also for  $\theta = 0$ , because this is the most favourable case, as then all  $q_j$  are 0. From (4.3.7) and (4.3.8) it follows that

$$(4.3.13) \quad R_0(x|z) = \Phi(x) + \phi(x) \frac{\sum a_j^4}{12(NB)^2} (x^3 - 3x) + O(N^{-5/4} + N^{-2} |\sum a_j^2 - NB|).$$

Since  $\epsilon' \leq \alpha \leq 1 - \epsilon'$ ,  $u_\alpha = O(1)$  and therefore we derive from (4.3.13) that the conditional critical value  $\xi_\alpha$  of the test based on  $2(U - \sum p_j a_j)/\tau$ , satisfies

$$(4.3.14) \quad \xi_\alpha = u_\alpha - \frac{\sum a_j^4}{12(NB)^2} (u_\alpha^3 - 3u_\alpha) + O(N^{-5/4} + N^{-2} |\sum a_j^2 - NB|).$$

The connection between  $\pi(\theta|z)$  and  $\xi_\alpha$  is  $\pi(\theta|z) = P_\theta([\sum a_j] / (\sum a_j^2)^{1/2} > \xi_\alpha | z)$ . Together with (4.3.6) this gives

$$(4.3.15) \quad 1 - \pi(\theta|z) = R_\theta \left( \xi_\alpha \left( \frac{\sum a_j^2}{\tau^2} \right)^{1/2} - \frac{\sum q_j a_j}{\tau} | z \right).$$

Using (4.3.14) we obtain

$$\begin{aligned} \xi_\alpha \left( \frac{\sum a_j^2}{\tau^2} \right)^{1/2} - \frac{\sum q_j a_j}{\tau} &= u_\alpha^{-\theta(NB)}^{1/2} + \frac{1}{2} u_\alpha \frac{\sum q_j a_j^2}{NB} + \\ &- \frac{1}{12} (u_\alpha^3 - 3u_\alpha) \frac{\sum a_j^4}{(NB)^2} + \frac{1}{2} \theta(NB)^{1/2} \frac{\tau^2 - NB}{NB} + \\ &- \frac{3}{8} \theta(NB)^{1/2} \frac{(\tau^2 - NB)^2}{(NB)^2} - \frac{\sum q_j a_j^{-\theta NB}}{(NB)^{1/2}} + \\ &+ \frac{1}{2} \frac{(\sum q_j a_j^{-\theta NB})(\tau^2 - NB)}{(NB)^{3/2}} + R_1, \end{aligned}$$

where

$$\begin{aligned} R_1 &= O\{N^{-2}(\sum q_j a_j^2)^2 + N^{-2}|\sum q_j a_j^2| |\sum a_j^2 - NB| + N^{-2}\sum q_j a_j^2 + N^{-5/4} + \\ &+ N^{-2}|\sum a_j^2 - NB| + N^{-3}|\tau^2 - NB|^3 + N^{-5/2}|\sum q_j a_j^{-\theta NB}|(\tau^2 - NB)^2\}. \end{aligned}$$

By the following relations,  $R_1$  can be simplified considerably

$$N^{-5/2}|\sum q_j a_j^{-\theta NB}|(\tau^2 - NB)^2 \leq N^{-3/2}|\sum q_j a_j^{-\theta NB}|^3 + N^{-3}|\tau^2 - NB|^3,$$

$$N^{-2}|\sum a_j^2 - NB| \leq N^{-3/2} + N^{-3}|\sum a_j^2 - NB|^3,$$

$$N^{-2}\sum q_j a_j^2 |\sum a_j^2 - NB| \leq N^{-3/2}(\sum q_j a_j^2)^{3/2} + N^{-3}|\sum a_j^2 - NB|^3,$$

$$N^{-3}(\sum q_j a_j^2)^3 + N^{-2}(\sum q_j a_j^2)^2 + N^{-2}\sum q_j a_j^2 = O(N^{-3/2}(\sum q_j a_j^2)^{3/2}).$$

We also have

$$N^{-2}(\tau^2 - NB)^2 = N^{-2}(\sum a_j^2 - NB)^2 + O(N^{-2}\sum q_j a_j^2 |\sum a_j^2 - NB| + N^{-2}(\sum q_j a_j^2)^2),$$

$$\begin{aligned}
N^{-3/2}(\sum_{q_j a_j - \theta NB}(\tau^2 - NB)) &= N^{-3/2}(\sum_{q_j a_j - \theta NB})(\sum_{a_j^2 - NB}) + \\
+ O(N^{-3/2}|\sum_{q_j a_j - \theta NB}|\sum_{q_j a_j^2}) &= N^{-3/2}(\sum_{q_j a_j - \theta NB})(\sum_{a_j^2 - NB}) + \\
+ O(N^{-3/2}(\sum_{q_j a_j^2})^{3/2} + N^{-3/2}|\sum_{q_j a_j - \theta NB}|^3). &
\end{aligned}$$

Application of these results leads to

$$\begin{aligned}
(4.3.16) \quad \xi_\alpha \left(\frac{\sum_{a_j^2}}{\tau^2}\right)^{1/2} - \frac{\sum_{q_j a_j}}{\tau} &= u_\alpha - \theta(NB)^{1/2} + \frac{1}{2}(u_\alpha - \theta(NB)^{1/2}) \frac{\sum_{q_j a_j^2}}{NB} + \\
- \frac{1}{12}(u_\alpha^3 - 3u_\alpha) \frac{\sum_{a_j^4}}{(NB)^2} + \frac{1}{2}\theta(NB)^{1/2} \frac{\sum_{a_j^2 - NB}}{NB} - \frac{3}{8}\theta(NB)^{1/2} \frac{(\sum_{a_j^2 - NB})^2}{(NB)^2} + \\
- \frac{\sum_{q_j a_j - \theta NB}}{(NB)^{1/2}} + \frac{1}{2} \frac{(\sum_{q_j a_j - \theta NB})(\sum_{a_j^2 - NB})}{(NB)^{3/2}} + R_2, &
\end{aligned}$$

$$(4.3.17) \quad R_2 = O\{N^{-5/4} + N^{-3}|\sum_{a_j^2 - NB}|^3 + N^{-3/2}|\sum_{q_j a_j - \theta NB}|^3 + N^{-3/2}(\sum_{q_j a_j^2})^{3/2}\}.$$

According to (4.3.15),  $\pi(\theta|z)$  can be found by substituting expansion (4.3.16) in expansion (4.3.7) for  $R_\theta(x|z)$ . We shall first consider the  $\phi(x)$ -term, and after this the  $x^k\phi(x)$  terms, for  $k = 0, 1, 2, 3$ . The following expansion is used

$$\begin{aligned}
(4.3.18) \quad \phi\left(\xi_\alpha \left(\frac{\sum_{a_j^2}}{\tau^2}\right)^{1/2} - \frac{\sum_{q_j a_j}}{\tau}\right) &= \phi(u_\alpha - \theta(NB)^{1/2}) + \\
+ \phi(u_\alpha - \theta(NB)^{1/2})\left[\xi_\alpha \left(\frac{\sum_{a_j^2}}{\tau^2}\right)^{1/2} - \frac{\sum_{q_j a_j}}{\tau} - u_\alpha + \theta(NB)^{1/2} + \right. \\
- \left. \frac{1}{2}(u_\alpha - \theta(NB)^{1/2})\left\{\xi_\alpha \left(\frac{\sum_{a_j^2}}{\tau^2}\right)^{1/2} - \frac{\sum_{q_j a_j}}{\tau} - u_\alpha + \theta(NB)^{1/2}\right\}^2\right] + \\
+ O\left(\left|\xi_\alpha \left(\frac{\sum_{a_j^2}}{\tau^2}\right)^{1/2} - \frac{\sum_{q_j a_j}}{\tau} - u_\alpha + \theta(NB)^{1/2}\right|^3\right). &
\end{aligned}$$

After some calculations this leads to

$$\begin{aligned}
(4.3.19) \quad & \phi\left(\xi_\alpha \left(\frac{\sum a_j^2}{\tau^2}\right)^{1/2} - \frac{\sum q_j a_j}{\tau}\right) = \phi(u_\alpha - \theta(NB)^{1/2}) + \\
& + \phi(u_\alpha - \theta(NB)^{1/2}) \left[ \frac{1}{2}(u_\alpha - \theta(NB)^{1/2}) \frac{\sum q_j^2 a_j^2}{NB} - \frac{1}{12}(u_\alpha^3 - 3u_\alpha) \frac{\sum a_j^4}{(NB)^2} + \right. \\
& + \frac{1}{2}\theta(NB)^{1/2} \frac{\sum a_j^2 - NB}{NB} - \frac{1}{8}\theta(NB)^{1/2} \{3 + \theta(NB)^{1/2}(u_\alpha - \theta(NB)^{1/2})\} \cdot \\
& \frac{(\sum a_j^2 - NB)^2}{(NB)^2} - \frac{\sum q_j a_j - \theta NB}{(NB)^{1/2}} - \frac{1}{2}(u_\alpha - \theta(NB)^{1/2}) \frac{(\sum q_j a_j - \theta NB)^2}{NB} + \\
& \left. + \frac{1}{2} \{1 + \theta(NB)^{1/2}(u_\alpha - \theta(NB)^{1/2})\} \frac{(\sum q_j a_j - \theta NB)(\sum a_j^2 - NB)}{(NB)^{3/2}} \right] + O(R_2).
\end{aligned}$$

The remainder is still  $O(R_2)$  by virtue of the conditions  $|\sum q_j a_j| \leq c'N^{1/2}$ ,  $0 \leq \theta \leq DN^{-1/2}$ ,  $B$  constant and  $\sum a_j^4 \leq C$ . They ensure that every term occurring in (4.3.16) and (4.3.17), is  $O(1)$ .

A simpler version of (4.3.16) suffices for the  $x^k \phi(x)$  terms

$$\begin{aligned}
& \xi_\alpha \left(\frac{\sum a_j^2}{\tau^2}\right)^{1/2} - \frac{\sum q_j a_j}{\tau} = u_\alpha - \theta(NB)^{1/2} + R_3, \\
R_3 & = O(N^{-1} + N^{-1} |\sum q_j^2 a_j^2| + N^{-1} |\sum a_j^2 - NB| + N^{-1/2} |\sum q_j a_j - \theta NB|).
\end{aligned}$$

It follows that

$$\begin{aligned}
(4.3.20) \quad & \left(\xi_\alpha \left(\frac{\sum a_j^2}{\tau^2}\right)^{1/2} - \frac{\sum q_j a_j}{\tau}\right)^k \phi\left(\xi_\alpha \left(\frac{\sum a_j^2}{\tau^2}\right)^{1/2} - \frac{\sum q_j a_j}{\tau}\right) = \\
& = (u_\alpha - \theta(NB)^{1/2})^k \phi(u_\alpha - \theta(NB)^{1/2}) + O(R_3).
\end{aligned}$$

Inserting (4.3.19) and (4.3.20) in (4.3.7) leads to the required expression (4.3.11) for  $\pi(\theta|z)$ , with remainder

$$\begin{aligned}
& O\{R + (N^{-1} + N^{-3/2} |\sum q_j a_j^3|) R_3 + N^{-5/2} |\sum q_j a_j^3| |\tau^2 - NB| + N^{-3/2} |\sum q_j^3 a_j^3| + \\
& + N^{-2} |\sum q_j^2 a_j^4| + N^{-2} |\sum q_j a_j^3|\}.
\end{aligned}$$

Finally observe that

$$N^{-1}R_3 = O(R),$$

$$N^{-5/2} |\sum q_j a_j^3| |\tau^2 - NB| \leq N^{-9/4} |\sum q_j a_j^3|^{3/2} + N^{-3} |\tau^2 - NB|^3,$$

$$N^{-2} |\sum q_j a_j^3| |\sum q_j a_j^{-\theta NB}| \leq N^{-9/4} |\sum q_j a_j^3|^{3/2} + N^{-3/2} |\sum q_j a_j^{-\theta NB}|^3,$$

$$N^{-5/2} |\sum q_j a_j^3| |\sum q_j a_j^2| \leq N^{-3} (\sum q_j a_j^3)^2 + N^{-2} (\sum q_j a_j^2)^2,$$

$$N^{-3} (\sum q_j a_j^3)^2 + N^{-2} |\sum q_j a_j^3| = O(N^{-3/2} + N^{-9/4} |\sum q_j a_j^3|^{3/2}).$$

This shows that the remainder (4.3.12) has the required order.  $\square$

The next step is to show that under  $\theta$ , the set of  $Z$ -values for which the conditions of the previous lemma are not satisfied, has a sufficiently small probability. The following definition is analogous to (4.3.3)

$$\tilde{A}_j = -\psi_1(X_j), \quad \tilde{P}_j = \frac{f(X_j - \theta)}{f(X_j - \theta) + f(X_j + \theta)}, \quad \tilde{Q}_j = 2\tilde{P}_j - 1, \quad j = 1, \dots, N.$$

LEMMA 4.3.3. *Let  $f$  be symmetric around zero, positive on  $R^1$  and twice continuously differentiable. Assume that positive constants  $\epsilon'$ ,  $D$ ,  $\eta$ ,  $\tilde{c}$ ,  $\tilde{C}$  and  $\eta'$  exist, such that*

$$(4.3.21) \quad 0 \leq \theta \leq DN^{-1/2},$$

$$(4.3.22) \quad \epsilon' \leq \alpha \leq 1 - \epsilon',$$

$$(4.3.23) \quad \sup \left\{ \int_{-\infty}^{\infty} \psi_1^{10}(x+y)f(x)dx : |y| \leq \eta \right\} \leq \tilde{C},$$

$$(4.3.24) \quad \left| \frac{d}{dy} \psi_1(F^{-1}(y)) \right| \geq \tilde{c},$$

for  $y$  in a subinterval  $\tau$  of  $(0,1)$  with length at least  $\eta'$ . Then there exists  $A > 0$ , depending on  $N$ ,  $\theta$ ,  $F$  and  $\alpha$  only through  $\epsilon'$ ,  $D$ ,  $\tilde{c}$ ,  $\tilde{C}$  and  $\eta'$  and such that (4.3.11) and (4.3.12) hold for all  $z$ -values, except for those

in a set  $B$  with  $P_\theta(B) \leq AN^{-5/4}$ .

PROOF. We have to verify that the conditions of lemma 4.3.2 are satisfied, except on a set of probability of order  $N^{-5/4}$ , i.e. we have to show that there exist constants  $c, C, \delta, \varepsilon$  and  $c'$  such that

$$(4.3.25) \quad \frac{1}{N} \sum A_j^4 \leq c,$$

$$(4.3.26) \quad \frac{1}{N} \sum (1-Q_j^2) A_j^2 \geq c,$$

$$(4.3.27) \quad \gamma(\varepsilon, \zeta, Q) \geq \delta N \zeta, \text{ for some } \zeta \geq N^{-3/2} \log N,$$

$$(4.3.28) \quad |\sum Q_j A_j| \leq c' N^{1/2},$$

except on a set of probability  $O(N^{-5/4})$ .

First consider (4.3.25). As  $\psi_1$  is odd, we have  $\sum A_j^4 = \sum \tilde{A}_j^4$ . The  $\tilde{A}_j^4$  are i.i.d. r.v.'s and in view of (4.3.23) we have

$$\sup_{0 \leq \theta \leq DN^{-1/2}} E_\theta |\tilde{A}_j^{10}| \leq \tilde{C}.$$

From Chebyshev's inequality we obtain

$$(4.3.29) \quad P_\theta \left( \left| \frac{1}{N} \sum (\tilde{A}_j^4 - E_\theta \tilde{A}_j^4) \right| \geq d \right) \leq d^{-5/2} E_\theta \left| \frac{1}{N} \sum (\tilde{A}_j^4 - E_\theta \tilde{A}_j^4) \right|^{5/2}.$$

At this point we use an inequality which is given by Chung (1951) and due to Marcinkievitz, Zygmund and Chung: if  $Y_1, \dots, Y_N$  are independent r.v.'s, all having mean zero, we have, for all  $p \geq 1$ ,

$$E \left| \sum Y_j \right|^{2p} \leq \hat{C} N^{p-1} \sum E |Y_j|^{2p},$$

where the constant  $\hat{C}$  only depends on  $p$ . By taking  $Y_j = \tilde{A}_j^4 - E_\theta \tilde{A}_j^4$ ,  $j = 1, \dots, N$  and  $p = 5/4$ , it follows that

$$E_\theta \left| \frac{1}{N} \sum (\tilde{A}_j^4 - E_\theta \tilde{A}_j^4) \right|^{5/2} \leq \hat{C} N^{-9/4} \sum E_\theta |\tilde{A}_j^4 - E_\theta \tilde{A}_j^4|^{5/2} = \hat{C} N^{-5/4} E_\theta |\tilde{A}_1^4 - E_\theta \tilde{A}_1^4|^{5/2}.$$

Together with (4.3.23) and (4.3.29) this implies that  $\sum \tilde{A}_j^4 \leq N(\tilde{C} + d) = NC$ , except on a set  $B_1$  with  $P_\theta(B_1) = O([d^2 N]^{-5/4})$ .

The next condition to verify is (4.3.26). As was mentioned in theorem 3.3.1, (4.3.21) and (4.3.23) imply (3.2.16). For the present case this means that  $P_\theta(\epsilon \leq \tilde{P}_1 \leq 1-\epsilon) > 1-\delta'$  for some  $\delta'$  with  $0 < \delta' < \min(\delta/2, c^2 c^{-1})$ . In theorem 3.2.2 it was shown that under this condition (4.3.26) holds, except on a set of probability  $O(N^{-5/4})$ , provided that  $\sum A_j^2 \geq c^* N$ , for some  $c^* > c$ . By applying Chebyshev's inequality one shows that  $\sum A_j^2 = \sum \tilde{A}_j^2 \geq N(E_\theta \tilde{A}_1^2 - d)$ , except on a set  $B_2$ , with  $P_\theta(B_2) = O([d^2 N]^{-5/4})$ . Hence it only remains to prove that  $E_\theta \tilde{A}_1^2$  is positive. Since  $\theta \rightarrow 0$  as  $N \rightarrow \infty$ , (4.3.24) implies that

$$(4.3.30) \quad \left| \frac{d}{dy} \psi_1(\theta + F^{-1}(y)) \right| \geq \tilde{c},$$

on a subinterval of  $(0,1)$  with length at least  $\eta'/2$ . Therefore

$$E_\theta \tilde{A}_1^2 = \int_{-\infty}^{\infty} \psi_1^2(x) f(x-\theta) dx = \int_0^1 \psi_1^2(\theta + F^{-1}(y)) dy > 0.$$

The third condition we have to deal with is (4.3.27). Inspection of the sketch of the proof of theorem 3.2.1 shows that this condition only serves to prove that

$$(4.3.31) \quad \int_{b'N^{1/2} \leq |t| \leq bN^{3/2}} \frac{|\rho(t)|}{t} dt = O(N^{-3/2}),$$

where  $|\rho|$  is given by (cf. (3.2.8))

$$(4.3.32) \quad |\rho(t)| \leq \exp\{-\sum P_j(1-P_j)\left(1 - \frac{\cos A_j t}{\tau}\right)\} = \\ = \exp\{-2\sum \tilde{P}_j(1-\tilde{P}_j) \sin^2 \frac{\tilde{A}_j t}{2\tau}\}.$$

Instead of verifying (4.3.27) we shall prove (4.3.31) directly. From (4.3.30) and a similar argument as in lemma 4.2.4 it follows that

$$E_\theta \sin^2 \tilde{A}_1 t = \int_0^1 \sin^2 \{t \psi_1(\theta + F^{-1}(y))\} dy > 0,$$

for all  $t$  bounded away from zero. As  $\sin^2 \tilde{A}_1 t$  is bounded, all moments exist and an application of Chebyshev's inequality shows that  $N^{-1} \sum \sin^2 \tilde{A}_j t$  is also bounded away from zero for all such  $t$ , except on a set  $B_3$  with  $P_\theta(B_3) = O(N^{-r})$  for all  $r > 0$ . Note that  $B_3$  is in fact  $B_3(t)$ . To overcome this complication, we note that in view of the continuity of  $\sin \tilde{A}_j t$  in  $t$ , we can for



each  $t_0$  choose  $B_3(t)$  equal to  $B_3(t_0)$  for all  $t$  in some neighbourhood of  $t_0$ . If  $N^{-1} \sum \tilde{A}_j^2$  is bounded, which is the case except on a set  $B_4$  with  $P_\theta(B_4) = O(N^{-5/4})$ , the length of this neighbourhood can be bounded away from zero. Hence we can now conclude that  $N^{-1} \sum \sin^2 \tilde{A}_j t \geq c_1$  will hold on a set  $b_1 \leq |t| \leq b_2 N$ , with  $c_1, b_1, b_2$  positive constants, except on a set  $B_5$  which is a union of  $B_4$  and at most  $O(N)$  sets  $B_3(t)$ . In view of the above,  $P_\theta(B_5) = O(N^{-5/4})$ . Furthermore, we note that, with probability  $1 - O(N^{-5/4})$ ,  $\tau^2/N = \{(1 - Q_j^2)A_j^2\}/N$  is bounded from below as well as from above. It follows that for  $b'N^{1/2} \leq |t| \leq bN^{3/2}$  with probability  $1 - O(N^{-5/4})$

$$\frac{1}{N} \sum \sin^2 \frac{\tilde{A}_j t}{2\tau} \geq c_1.$$

Now there exists  $\tilde{\varepsilon} > 0$ , depending on  $c_1$ , such that at most a fraction  $c_1/2$  of the  $\tilde{P}_j$  does not lie in the interval  $(\tilde{\varepsilon}, 1 - \tilde{\varepsilon})$ , again with probability  $1 - O(N^{-5/4})$ . Hence, for all  $b'N^{1/2} \leq |t| \leq bN^{3/2}$ ,

$$\sum \tilde{P}_j (1 - \tilde{P}_j) \sin^2 \frac{\tilde{A}_j t}{2\tau} \geq \tilde{\varepsilon}(1 - \tilde{\varepsilon}) \left[ \sum \sin^2 \frac{\tilde{A}_j t}{2\tau} - \frac{c_1 N}{2} \right] \geq \frac{\tilde{\varepsilon}(1 - \tilde{\varepsilon})c_1 N}{2},$$

with probability  $1 - O(N^{-5/4})$ . Together with (4.3.32), this proves the validity of (4.3.31), except on a set  $B_6$  with  $P_\theta(B_6) = O(N^{-5/4})$ .

It remains to prove (4.3.28). As  $Q_j$  and  $A_j$  are both odd functions of  $Z_j$ , we have  $|\sum Q_j A_j| \leq \sum |Q_j A_j| \leq \sum |\tilde{Q}_j \tilde{A}_j|$ . Because

$$\begin{aligned} \tilde{Q}_1 \tilde{A}_1 &= -\psi_1(X_1) \frac{f(X_1 - \theta) - f(X_1 + \theta)}{f(X_1 - \theta) + f(X_1 + \theta)} = \\ &= 2\theta \psi_1(X_1) \int_0^1 \frac{f'(X_1 + v\theta)f(X_1 - v\theta) + f(X_1 + v\theta)f'(X_1 - v\theta)}{(f(X_1 + v\theta) + f(X_1 - v\theta))^2} dv, \end{aligned}$$

it follows that

$$\begin{aligned} |\tilde{Q}_1 \tilde{A}_1| &\leq 2\theta |\psi_1(X_1)| \int_0^1 \left\{ \frac{1}{2} \psi_1(X_1 + v\theta) + \frac{1}{2} \psi_1(X_1 - v\theta) \right\} dv \leq \\ &\leq \theta \{ |\psi_1^2(X_1)| + \int_0^1 \psi_1^2(X_1 + v\theta) dv + \int_0^1 \psi_1^2(X_1 - v\theta) dv \}. \end{aligned}$$

As  $0 \leq \theta \leq DN^{-1/2}$ , it remains to prove that, with probability  $1 - O(N^{-5/4})$

$$\sum_{j=1}^N \int_0^1 \psi_1^2(X_j + v\theta) dv \leq \tilde{C}N.$$

This is done in the same way as in which (4.3.25) is proved; therefore we

only mention that an application of Fubini's theorem shows that

$$\mathbb{E}_\theta \left| \int_0^1 \psi_1^2(X_{1+v\theta}) dv \right|^{5/2} \leq \sup_{|v| \leq 1} \mathbb{E}_\theta |\psi_1^5(X_j + v\theta)| \leq \tilde{C}. \quad \square$$

COROLLARY 4.3.1. *Under the conditions of lemma 4.3.3, the power  $\pi(\theta)$  of the permutation test based on  $U$ , satisfies*

$$(4.3.33) \quad |\pi(\theta) - \mathbb{E}_\theta \tilde{\pi}(\theta|Z)| \leq A \mathbb{E}_\theta R,$$

where  $\tilde{\pi}(\theta|z)$  and  $R$  are given in (4.3.11) and (4.3.12) and  $A$  is a positive constant, depending on  $N$ ,  $\theta$ ,  $F$  and  $\alpha$  only through  $\varepsilon'$ ,  $D$ ,  $\eta$ ,  $\tilde{c}$ ,  $\tilde{C}$  and  $\eta'$ .

PROOF. On  $B^c$   $\pi(\theta|z)$  satisfies  $\pi(\theta|z) = \tilde{\pi}(\theta|z) + R$ . Hence, for  $\pi(\theta) = \mathbb{E}_\theta \pi(\theta|Z)$  we obtain

$$|\pi(\theta) - \mathbb{E}_\theta (\tilde{\pi}(\theta|Z) + R)| \leq \mathbb{E}_\theta |\pi(\theta|Z) I_B(Z)| + \mathbb{E}_\theta |(\tilde{\pi}(\theta|Z) + R) I_B(Z)|,$$

where  $I_B$  is the indicator function of  $B$ . Now  $\pi(\theta|z)$ , being a probability, is bounded, and therefore  $\mathbb{E}_\theta |\pi(\theta|Z) I_B(Z)| = O(N^{-5/4})$ .

The term  $\mathbb{E}_\theta |(\tilde{\pi}(\theta|Z) + R) I_B(Z)|$  has to be treated with more care. First,  $\Phi(u_\alpha - \theta(NB))^{1/2}$  is bounded, and thus contributes  $O(N^{-5/4})$ . The next three terms of  $\tilde{\pi}(\theta|Z)$  can be split into a bounded part and a part that has the form of one of the remaining terms in  $\tilde{\pi}(\theta|Z)$ :

$$|N^{-3/2} \sum Q_j A_j^3| \leq N^{-3/2} \sum A_j^2 \sum |Q_j A_j| \leq N^{-2} (\sum A_j^2)^2 + N^{-1} (\sum |Q_j A_j|)^2 \leq$$

$$\leq 2(B^2 + \theta^2 NB^2) + 2N^{-2} (\sum A_j^2 - NB)^2 + 2N^{-1} (\sum |Q_j A_j| - \theta NB)^2,$$

$$N^{-2} \sum A_j^4 \leq N^{-2} (\sum A_j^2)^2 \leq 2B^2 + 2N^{-2} (\sum A_j^2 - NB)^2,$$

$$N^{-1} \sum Q_j^2 A_j^2 \leq N^{-1} (\sum |Q_j A_j|)^2 \leq 2\theta^2 NB^2 + 2N^{-1} (\sum |Q_j A_j| - \theta NB)^2.$$

The remaining terms of  $\tilde{\pi}(\theta|Z)$ , and the terms of  $R$ , can all be treated in the following way. Take for example the first term,  $N^{-1} (\sum A_j^2 - NB)$ . On the part of  $B$  where  $N^{-1} (\sum A_j^2 - NB) \leq 1$ , the contribution to the expectation is clearly  $O(N^{-5/4})$ . On the remaining part of  $B$  the contribution is

$O(N^{-3} E_{\theta} |\sum_j A_j^2 - NB|^3)$  because for any r.v.  $Y$  with d.f.  $H$ , and all  $p \geq 1$ ,

$$\int_{|y| \geq 1} |y| dH(y) \leq \int_{|y| \geq 1} |y|^p dH(y) \leq \int_{-\infty}^{\infty} |y|^p dH(y) = E|Y|^p.$$

Thus the total contribution of this term is  $O(N^{-5/4} + N^{-3} E_{\theta} |\sum_j A_j^2 - NB|^3)$ . But these terms already occur in  $E_{\theta} R$ . Inspection of the other terms shows that their contribution is always  $O(E_{\theta} R)$ .  $\square$

Our final task is the derivation of simple expressions for  $E_{\theta} \tilde{\pi}(\theta|Z)$  and  $E_{\theta} R$ . To this end,  $E_{\theta} \tilde{Q}_1^{r,s}$  has to be evaluated for various  $r$  and  $s$ , which is done by expanding  $\tilde{Q}_1^{r,s}$  around  $\theta = 0$  to the appropriate order. Note that  $E_{\theta} \tilde{Q}_1^{r,s}$  has a double dependence on  $\theta$ : explicitly in  $\tilde{Q}_1 = [f(X_1 - \theta) - f(X_1 + \theta)] / [f(X_1 - \theta) + f(X_1 + \theta)]$  and implicitly as  $X_1$  comes from  $F(x - \theta)$ . We introduce the following notation

$$h(x, \theta) = \frac{f(x+2\theta) - f(x)}{f(x+2\theta) + f(x)}, \quad q_{r,s}(x, \theta) = h^r(x, \theta) \psi_1^s(x + \theta).$$

Let  $q_{r,s}^{(1)}(x, \theta)$ ,  $h^{(1)}(x, \theta)$ ,  $\psi_1^{(1)}(x + \theta)$  denote the 1-th derivative of  $q_{r,s}(x, \theta)$ ,  $h(x, \theta)$  and  $\psi_1(x + \theta)$  respectively, with respect to  $\theta$ . In the next lemma we evaluate the necessary moments.

LEMMA 4.3.4. *Let  $f$  be five times differentiable and suppose that positive constants  $\tilde{C}$ ,  $D$  and  $\eta$  exist, such that  $0 \leq \theta \leq DN^{-1/2}$  and*

$$(4.3.34) \quad \sup \left\{ \int_{-\infty}^{\infty} |\psi_j(x+y)|^{5/j} f(x) dx : |y| \leq \eta \right\} \leq \tilde{C}, \quad j = 1, \dots, 5.$$

*Then there exists  $A > 0$ , depending on  $N$ ,  $\theta$  and  $F$  only through  $\tilde{C}$ ,  $D$  and  $\eta$  and such that*

$$(4.3.35) \quad \begin{aligned} E_{\theta} |\tilde{Q}_1^{r,s(5-r)}| &\leq AN^{-r/2}, \quad r = 0, 1, 2, \\ |E_{\theta} \tilde{Q}_1^{r,s(4-r)} - \theta^r E_{\theta} \tilde{A}_1^4| &\leq AN^{-(r+1)/2}, \quad r = 0, 1, 2, \\ |E_{\theta} \tilde{A}_1^2 - \{E_{\theta} \tilde{A}_1^2 + \frac{\theta^2}{3} E_{\theta} \psi_1^4(X_1)\}| &\leq AN^{-3/2}, \\ |E_{\theta} \tilde{Q}_1 \tilde{A}_1 - \{\theta E_{\theta} \tilde{A}_1^2 + \frac{\theta^3}{6} E_{\theta} (\frac{2}{3} \psi_1^4(X_1) - \psi_2^2(X_1))\}| &\leq AN^{-2}. \end{aligned}$$

PROOF. For  $l = 1, 2, 3, 4$  we have

$$\begin{aligned}
 \psi_1^{(1)}(x+\theta) &= \psi_2(x+\theta) - \psi_1^2(x+\theta), \\
 (4.3.36) \quad \psi_1^{(2)}(x+\theta) &= \psi_3(x+\theta) - 3\psi_1(x+\theta)\psi_2(x+\theta) + 2\psi_1^3(x+\theta), \\
 |\psi_1^{(l)}(x+\theta)| &\leq \bar{c} \sum_{j=1}^{l+1} |\psi_j^{1+j}(x+\theta)|,
 \end{aligned}$$

for some  $\bar{c} > 0$ . Differentiation of  $h$  yields that

$$\begin{aligned}
 h(x, 0) &= 0, \quad |h(x, \theta)| \leq 1, \\
 h^{(1)}(x, \theta) &= 4 \frac{f'(x+2\theta)}{f(x)} / \left( \frac{f(x+2\theta)}{f(x)} + 1 \right)^2, \quad h^{(1)}(x, 0) = \psi_1(x), \\
 h^{(2)}(x, \theta) &= 8 \frac{f''(x+2\theta)}{f(x)} / \left( \frac{f(x+2\theta)}{f(x)} + 1 \right)^2 + \\
 &\quad - 16 \left( \frac{f'(x+2\theta)}{f(x)} \right)^2 / \left( \frac{f(x+2\theta)}{f(x)} + 1 \right)^3, \\
 (4.3.37) \quad h^{(2)}(x, 0) &= 2(\psi_2(x) - \psi_1^2(x)), \\
 h^{(3)}(x, \theta) &= 16 \frac{f^{(3)}(x+2\theta)}{f(x)} / \left( \frac{f(x+2\theta)}{f(x)} + 1 \right)^2 + \\
 &\quad - 96 \frac{f'(x+2\theta)f''(x+2\theta)}{f^2(x)} / \left( \frac{f(x+2\theta)}{f(x)} + 1 \right)^3 + \\
 &\quad + 96 \left( \frac{f'(x+2\theta)}{f(x)} \right)^3 / \left( \frac{f(x+2\theta)}{f(x)} + 1 \right)^4, \\
 h^{(3)}(x, 0) &= 4\psi_3(x) - 12\psi_1(x)\psi_2(x) + 6\psi_1^3(x), \\
 |h^{(l)}(x, \theta)| &\leq \bar{c} \sum_{j=1}^l |\psi_j^{1+j}(x+2\theta)|, \quad l = 1, 2, 3, 4.
 \end{aligned}$$

Under  $F(x-\theta)$ ,  $X_1$  has the same distribution as  $X_1+\theta$  under  $F(x)$ . Hence

$$E_{\theta} \tilde{Q}_1^{\tilde{r}, \tilde{s}} = E_0 (-h)^{\tilde{r}}(X_1, \theta) (-\psi_1)^{\tilde{s}}(X_1+\theta) = E_0 q_{\tilde{r}, \tilde{s}}(X_1, \theta) (-1)^{\tilde{r}+\tilde{s}}.$$

We now prove the results in the first line of (4.3.35). Note that

$$E_{\theta} |\tilde{A}_1^{\tilde{s}}| = E_{\theta} |\psi_1^{\tilde{s}}(X_1)| = E_0 |\psi_1^{\tilde{s}}(X_1+\theta)| = O(1),$$

because of (4.3.34) and the fact that  $0 \leq \theta \leq DN^{-1/2}$ . Application of lemma 4.2.2 gives

$$E_{\theta} |\tilde{Q}_1 \tilde{A}_1^4| = E_0 |q_{1,4}(X_1, \theta)| \leq \theta \sup_{0 \leq v \leq 1} E_0 |q_{1,4}^{(1)}(X_1, v\theta)|,$$

since  $h(x, 0) = 0$ . Furthermore, for some  $C^* > 0$ ,

$$|q_{1,4}^{(1)}| \leq C^* \{ |h^{(1)} \psi_1^4| + |h \psi_1^3 \psi_1^{(1)}| \}.$$

Using (4.3.36), (4.3.37) and Hölder's inequality, one obtains

$$|q_{1,4}^{(1)}(x, v\theta)| \leq C^* \sum_{j=1}^2 \{ |\psi_j^{5/j}(x+v\theta)| + |\psi_j^{5/j}(x+2v\theta)| \}.$$

Hence, by (4.3.34),  $\sup_{0 \leq v \leq 1} E_0 |q_{1,4}^{(1)}(X_1, v\theta)| \leq C_1$ , for some  $C_1 > 0$ , and therefore  $E_{\theta} |\tilde{Q}_1 \tilde{A}_1^4| = O(N^{-1/2})$ . Similarly,

$$\begin{aligned} E_{\theta} |\tilde{Q}_1 \tilde{A}_1^3| &\leq \frac{\theta^2}{2} \sup_{0 \leq v \leq 1} E_0 |q_{2,3}^{(2)}(X_1, v\theta)| \leq \\ &\leq \frac{\theta^2}{2} C^* \sup_{0 \leq v \leq 1} E_0 \left[ \sum_{j=1}^3 \{ |\psi_j^{5/j}(X_1+v\theta)| + |\psi_j^{5/j}(X_1+2v\theta)| \} \right] = O(N^{-1}). \end{aligned}$$

The second line of (4.3.35) is proved by continuing in the same way. For  $r = 0, 1, 2$  we have

$$\begin{aligned} |E_{\theta} \tilde{Q}_1 \tilde{A}_1^{(4-r)} - \theta^r E_0 \tilde{A}_1^4| &= |E_0 q_{r,4-r}(X_1, \theta) - \theta^r E_0 \psi_1^4(X_1)| \leq \\ &\leq \frac{\theta^{r+1}}{(r+1)!} \sup_{0 \leq v \leq 1} |q_{r,4-r}^{(r+1)}(X_1, v\theta)| \leq \\ &\leq \frac{\theta^{(r+1)}}{(r+1)!} C^* \sup_{0 \leq v \leq 1} E_0 \left[ \sum_{j=1}^{r+2} \{ |\psi_j^{5/j}(X_1+v\theta)| + \right. \\ &\quad \left. + |\psi_j^{5/j}(X_1+2v\theta)| \} \right] = O(N^{-\frac{(r+1)}{2}}). \end{aligned}$$

$E_{\theta} \psi_1^2(X_1)$  is evaluated in the following way

$$\begin{aligned}
& |E_{\theta} \psi_1^2(X_1) - \{E_0 \psi_1^2(X_1) + 2\theta E_0[\psi_1 \psi_2(X_1) - \psi_1^3(X_1)]\} + \\
& + \theta^2 E_0[\psi_2^2(X_1) + \psi_1 \psi_3(X_1) - 5\psi_1^2 \psi_2(X_1) + 3\psi_1^4(X_1)]\} | \leq \\
& \leq \frac{\theta^3}{3!} \sup_{0 \leq v \leq 1} E_0 |q_{0,2}^{(3)}(X_1, v\theta)| \leq \frac{\theta^3}{3!} C^* \sup_{0 \leq v \leq 1} E_0 \left[ \sum_{j=1}^4 \{|\psi_j^{5/j}(X_1 + v\theta)| + \right. \\
& \left. + |\psi_j^{5/j}(X_1 + v\theta)|\} \right] = O(N^{-3/2}).
\end{aligned}$$

As  $\psi_1^2 \psi_2$  and  $\psi_1^3$  are odd, the corresponding moments are zero. Also, in section 4.2 we found by partial integration that  $E_0 \psi_1^2 \psi_2(X_1) = 2/3 E_0 \psi_1^4(X_1)$  and  $E_0 \psi_1 \psi_3(X_1) = 2/3 E_0 \psi_1^4(X_1) - E_0 \psi_2^2(X_1)$ . Application of these results yields the expression in (4.3.35) for  $E_{\theta} \tilde{A}_1^2 = E_{\theta} \psi_1^2(X_1)$ .

Finally  $E_{\theta} \tilde{Q}_1 \tilde{A}_1$  has to be found.

$$\begin{aligned}
& |E_{\theta} \tilde{Q}_1 \tilde{A}_1 - \sum_{j=0}^3 E_0 q_{1,1}^{(j)}(X_1, 0) \frac{\theta^j}{j!}| \leq \frac{\theta^4}{4!} \sup_{0 \leq v \leq 1} E_0 |q_{1,1}^{(4)}(X_1, v\theta)| \leq \\
& \leq \frac{\theta^4}{4!} C^* \sup_{0 \leq v \leq 1} E_0 \left[ \sum_{j=1}^5 \{|\psi_j^{5/j}(X_1 + v\theta)| + |\psi_j^{5/j}(X_1 + 2v\theta)|\} \right] = O(N^{-2}).
\end{aligned}$$

By applying (4.3.36) and (4.3.37),  $E_0 q_{1,1}^{(j)}(X_1, 0)$  can be obtained, which leads to the desired formula in (4.3.35).  $\square$

We are now in a position to give a simple expansion for  $\pi(\theta)$ .

**THEOREM 4.3.1.** *Suppose that positive constants  $D$ ,  $\eta$ ,  $\tilde{c}$ ,  $\tilde{C}$  and  $\varepsilon'$  exist such that (4.3.21), (4.3.22) and (4.3.24) hold and such that*

$$\sup \left\{ \int_{-\infty}^{\infty} |\psi_j^{m_j}(x+y)| f(x) dx : |y| \leq \eta \right\} \leq \tilde{C}, \quad j = 1, \dots, 5,$$

$$m_1 = 10, \quad m_j = 5/j, \quad j = 2, \dots, 5.$$

*Then there exists  $A > 0$  depending on  $N$ ,  $\theta$ ,  $F$  and  $\alpha$  only through  $D$ ,  $\eta$ ,  $\tilde{c}$ ,  $\tilde{C}$  and  $\varepsilon'$ , such that*

$$(4.3.38) \quad |\pi(\theta) - \tilde{\pi}(\theta)| \leq AN^{-5/4},$$

where

$$(4.3.39) \quad \tilde{\pi}(\theta) = 1 - \Phi(u_\alpha - \eta_1) + \frac{\eta_1 \phi(u_\alpha - \eta_1)}{N} \left[ -\frac{\eta_2}{12} u_\alpha^2 + \frac{(\eta_2 - 3)}{24} \eta_1 u_\alpha + \right. \\ \left. - \frac{(\eta_2 - 3)}{24} - \frac{(12\eta_3 - 5\eta_2 - 9)\eta_1^2}{72} \right],$$

and  $\eta_1$ ,  $\eta_2$  and  $\eta_3$  as defined in (4.2.19).

PROOF. The results of lemma 4.3.4 strongly suggest the choice of  $B = E_0 \psi_1^2(X_1)$  in  $E\tilde{\pi}(\theta|Z)$ . This is allowed, as  $E_0 \psi_1^2(X_1)$  is a positive constant as was proved in lemma 4.3.3. The conditions of the present theorem contain the conditions of lemmas 4.3.3 and 4.3.4. Hence we may use (4.3.35) in  $E(\tilde{\pi}(\theta|Z))$ . After some algebra we obtain  $\tilde{\pi}(\theta) = E_\theta \tilde{\pi}(\theta|Z) + O(N^{-3/2})$ . Hence, in order to prove (4.3.38), it suffices in view of (4.3.33), to prove that  $E_\theta R = O(N^{-5/4})$ . Consider the expression for  $R$  in (4.3.12). We begin with the last term

$$\begin{aligned} N^{-2} E_\theta |\sum Q_{jA_j}^{2,4}| &\leq N^{-1} E_\theta |\tilde{Q}_{1A_1}^{2,4}| \leq N^{-1} E_\theta |\tilde{Q}_{1A_1}^{4,4}| = O(N^{-3/2}), \\ N^{-3/2} E_\theta |\sum Q_{jA_j}^{3,3}| &\leq N^{-1/2} E_\theta |\tilde{Q}_{1A_1}^{3,3}| \leq N^{-1/2} E_\theta |\tilde{Q}_{1A_1}^{2,3}| = O(N^{-3/2}), \\ N^{-3/2} E_\theta |\sum Q_{jA_j}^{2,2}|^{3/2} &= N^{-3/2} E_\theta (\sum Q_{jA_j}^{4,4} + \sum_{i \neq j} \tilde{Q}_{iA_i}^{2,2} \tilde{Q}_{jA_j}^{2,2})^{3/4} \leq \\ &\leq N^{-3/2} E_\theta |\sum \tilde{Q}_{jA_j}^{3,3}| + N^{-3/2} (E_\theta \sum_{i \neq j} \tilde{Q}_{iA_i}^{2,2} \tilde{Q}_{jA_j}^{2,2})^{3/4} \leq \\ &\leq N^{-1/2} E_\theta |\tilde{Q}_{1A_1}^{2,3}| + (E_\theta \tilde{Q}_{1A_1}^{2,2})^{3/2} = O(N^{-3/2}), \\ N^{-9/4} E_\theta |\sum Q_{jA_j}^{3,3}|^{3/2} &\leq N^{-3/2} + N^{-5/2} E_\theta (\sum \tilde{Q}_{jA_j}^{3,3})^2 \leq \\ &\leq N^{-3/2} + N^{-3/2} E_\theta \tilde{Q}_{1A_1}^{2,6} + N^{-1/2} (E_\theta \tilde{Q}_{1A_1}^{3,3})^2 = O(N^{-3/2}). \end{aligned}$$

Another application of the inequality due to Marcinkievitz, Zygmund and Chung disposes of the remaining two terms. If  $Y_1, \dots, Y_N$  are i.i.d. r.v.'s

with mean zero, this inequality implies that

$$(4.3.40) \quad E|\sum Y_j^3| \leq \hat{C}N^{1/2} \sum E|Y_j|^3 = \hat{C}N^{3/2} E|Y_1|^3.$$

Hence

$$\begin{aligned} N^{-3} E_\theta |\sum A_j^2 - NB|^3 &= N^{-3} E_\theta |\sum (\tilde{A}_j^2 - E_\theta \tilde{A}_j^2)|^3 = O(N^{-3} E_\theta |\sum (\tilde{A}_j^2 - E_\theta \tilde{A}_j^2)|^3 + \\ &+ |E_\theta \tilde{A}_1^2 - E_\theta \tilde{A}_1^2|^3) = O(N^{-3/2} E_\theta |\tilde{A}_1^2 - E_\theta \tilde{A}_1^2|^3 + \\ &+ N^{-3/2} \{E_\theta (\tilde{A}_1^2 - E_\theta \tilde{A}_1^2)^2\}^{3/2} + N^{-3}) = O(N^{-3/2}), \end{aligned}$$

and in the same way  $N^{-3/2} E_\theta |\sum Q_j A_j - \theta NB|^3 = O(N^{-3/2})$ . Finally, the announced uniformity in (4.3.38) is an immediate consequence of the uniformity in the conditions.  $\square$

REMARK. The remainder is  $O(N^{-5/4})$  instead of  $O(N^{-3/2})$  only because of the fact that in lemma 4.3.1 the Edgeworth expansion of theorem 3.2.1 is used, which approximates to  $O(N^{-5/4})$ . If in theorem 3.2.1 condition  $\sum a_j^4 \leq CN$  is replaced by  $\sum |a_j|^5 \leq CN$ , the remainder becomes  $O(N^{-3/2})$ , as was remarked in chapter 3. Hence by changing of  $m_1 = 10$  to  $m_1 = 25/2$  one obtains a remainder of  $O(N^{-3/2})$ .

We conclude this section by considering again the permutation test based on  $U^* = (1/2\theta) \sum_{X_j > 0} \log \{f(X_j - \theta)/f(X_j + \theta)\}$ , instead of on  $U$ . Let  $\pi^*(\theta)$  be the power of the  $U^*$ -test and define

$$\begin{aligned} \tilde{A}_j^* &= \frac{1}{2\theta} \log \frac{f(X_j - \theta)}{f(X_j + \theta)}, \quad \tilde{a}_j^* = \frac{1}{2\theta} \log \frac{f(x_j - \theta)}{f(x_j + \theta)}, \\ A_j^* &= \frac{1}{2\theta} \log \frac{f(Z_j - \theta)}{f(Z_j + \theta)}, \quad a_j^* = \frac{1}{2\theta} \log \frac{f(z_j - \theta)}{f(z_j + \theta)}. \end{aligned}$$

In the following lemma we prove that  $\pi^*(\theta)$  agrees with  $\pi(\theta)$  up to  $O(N^{-5/4})$ .

LEMMA 4.3.5. *Under the conditions of theorem 4.3.1 we have*

$$|\pi^*(\theta) - \pi(\theta)| \leq AN^{-5/4},$$



where  $\tilde{\pi}(\theta)$  is given by (4.3.39).

PROOF. As  $U^* = \sum_j A_j^* V_j$  is of the same form as  $U = \sum_j A_j V_j$ , the proof consists of showing that all lemmas in this section continue to hold in the case of  $U^*$ , provided that some minor changes are made. Thus lemma 4.3.1 and 4.3.2 clearly hold for the  $U^*$ -test, if  $a_j$  is replaced by  $a_j^*$  everywhere. Lemma 4.3.3 remains valid without any changes at all, because  $\tilde{A}_j^* = -\frac{1}{2} \psi_1(X_j - v_1 \theta) - \frac{1}{2} \psi_1(X_j + v_2 \theta)$ ,  $0 \leq v_1, v_2 \leq 1$  and therefore (4.3.21) and (4.3.23) ensure that  $\sup_{0 \leq \theta \leq DN^{-\frac{1}{2}}} E_\theta |\tilde{A}_1^*|^{10} \leq C$  and  $E_\theta \tilde{A}_1^{*2} > 0$ . Corollary 4.3.1 also continues to hold in the same form. Hence it finally remains to adapt lemma 4.3.4. Expansion around  $\theta = 0$  shows that

$$\tilde{A}_1^* = -\psi_1(X_1) - \frac{\theta^2}{6} \{\psi_3(X_1) - 3\psi_1\psi_2(X_1) + 2\psi_1^3(X_1)\} + \dots, \quad (4.3.41)$$

$$(\tilde{A}_1^*)^2 = \psi_1^2(X_1) + \frac{\theta^2}{3} \{\psi_1\psi_3(X_1) - 3\psi_1^2\psi_2(X_1) + 2\psi_1^4(X_1)\} + \dots$$

In a similar way as in lemma 4.3.4 one shows that the first two statements in (4.3.35) remain valid if  $\tilde{A}_1$  is replaced by  $\tilde{A}_1^*$ . Moreover,

$$E_\theta (\tilde{A}_1^*)^2 = E_\theta \tilde{A}_1^2 + \frac{\theta^2}{3} E_0 \psi_1 \psi_3(X_1) + O(N^{-3/2}), \quad (4.3.42)$$

$$E_\theta (\tilde{Q}_1 \tilde{A}_1^*) = E_\theta (\tilde{Q}_1 \tilde{A}_1) + \frac{\theta^3}{6} E_0 \psi_1 \psi_3(X_1) + O(N^{-2}).$$

The expressions in (4.3.35) are used to obtain  $E_\theta \tilde{\pi}(\theta|Z)$  from (4.3.11). In (4.3.11),  $E_\theta (A_1^*)^2$  and  $E_\theta \tilde{Q}_1 \tilde{A}_1^*$  only occur in

$$E_\theta \left[ \frac{1}{2} \theta (NB)^{1/2} \frac{(\sum_j A_j^{*2} - NB)}{NB} - \frac{(\sum_j \tilde{Q}_j \tilde{A}_j^* - \theta NB)}{(NB)^{1/2}} \right].$$

But in view of (4.3.42), this expectation differs at most  $O(N^{-3/2})$  from the expectation of the same expression in  $\tilde{A}_j$  instead of  $\tilde{A}_j^*$ .  $\square$

#### 4.4. SCALE INVARIANT TESTS

One of the advantages of the use of rank tests is the fact that such tests are distribution-free. This was the motivation in the previous section to

compare their behaviour to that of most powerful distribution-free tests.

Another nice property of rank tests is their scale invariance. In analogy to the previous section it therefore seems worthwhile to compare LMP rank tests with most powerful scale invariant tests. Let again  $X_1, X_2, \dots, X_N$  be i.i.d. r.v.'s from a d.f.  $F(x-\theta)$ , where  $f = F'$  is symmetric around zero and positive on  $R^1$ . According to Hájek and Šidák (1967) the most powerful scale invariant test for  $\theta = 0$  against a simple alternative  $\theta > 0$  rejects the hypothesis for large values of

$$(4.4.1) \quad \left\{ \int_0^\infty \prod_{j=1}^N f(\lambda X_j - \theta) \lambda^{N-1} d\lambda \right\} / \left\{ \int_0^\infty \prod_{j=1}^N f(\lambda X_j) \lambda^{N-1} d\lambda \right\}.$$

In Hájek and Šidák (1967) it is also shown that in the normal case this test is equivalent with Student's one sample t-test. Unfortunately, for general  $f$  we are unable to find an expansion for the d.f. of the statistic in (4.4.1).

We conclude this section with a remark on the relation between the most powerful permutation test for the normal case and the t-test, being the most powerful scale invariant test for this case. Note that if  $f$  is the normal density, the permutation tests based on  $U$  and  $U^*$  are both equivalent to the permutation test based on  $\sum X_j$ . Under the conditions of theorem 4.3.1 its power satisfies, according to (4.3.38) and (4.3.39)

$$(4.4.2) \quad \pi(\theta) = 1 - \Phi(u_\alpha - N^{1/2}\theta) - \frac{\theta u_\alpha^2}{4N^{1/2}} \phi(u_\alpha - N^{1/2}\theta) + O(N^{-5/4}).$$

But this expansion also holds for the power of the t-test, as is shown by Hodges and Lehmann (1970). Hence, in particular, the normal permutation-test has deficiency zero with respect to the t-test if normal location alternatives are considered. This rather striking phenomenon can be made more transparent by looking directly at the two test statistics involved. For this approach the reader is referred to Albers, Bickel and van Zwet (1974).

#### 4.5. RANDOMIZED RANK SCORE TESTS

In the preceding sections of this chapter we considered tests that are only slightly better than the LMP rank test in the sense that the asymptotic relative efficiency (ARE) of the LMP rank test with respect to these tests

always equals one. Here we consider the opposite case: randomized rank score (RRS) tests are worse than LMP rank tests, but also have ARE 1 with respect to these tests.

RRS tests have been introduced by Bell and Doksum (1965) for the two sample problem. By proceeding analogously, we define a RRS test for the one sample problem. As before,  $X_1, \dots, X_N$  are i.i.d. r.v.'s from  $F(x-\theta)$ , where the known d.f.  $F$  has a density  $f$  that is symmetric around zero and positive on  $R^1$ . We also have an auxiliary independent sample  $X_1^*, \dots, X_N^*$  from  $F$ . Denote  $(X_1, \dots, X_N)$  as  $X$  and  $(X_1^*, \dots, X_N^*)$  as  $X^*$ . Let  $Z(Z^*)$  be the vector of order statistics for the absolute values  $|X_1|, \dots, |X_N| (|X_1^*|, \dots, |X_N^*|)$ . Now we test the hypothesis of symmetry against the restricted alternative  $F(x-\theta)$ ,  $\theta > 0$ , by rejecting the hypothesis for large values of

$$L = \sum A_j \bar{v}_j,$$

where  $A_j = -\psi_1(Z_j^*)$ ,  $\bar{v}_j = 1$  if the  $X_i$  corresponding to  $Z_j$  is positive and  $\bar{v}_j = -1$  otherwise.  $\sum$  always means  $\sum_{j=1}^N$ .

The statistic of the LMP rank test may be expressed as  $T = \sum (EA_j) \bar{v}_j$ . Note that a subscript in  $EA_j$  is superfluous since the  $X_j^*$  always come from  $F$ . Hence  $L$  can be interpreted as the randomized counterpart of  $T$ . An advantage of  $L$  over  $T$  lies in the fact that its computation requires no tables of  $EA_j$ . Moreover, in some cases,  $L$  has under the hypothesis a continuous, known and tabulated distribution, e.g. normal or  $\chi^2$ , whereas special tables are needed for the distribution of  $T$  under the hypothesis. One may suspect that the price for these advantages will be a loss of power of the RRS test. However, Bell and Doksum have shown that its ARE with respect to the LMP rank test equals one. Therefore it seems worthwhile to obtain an expansion for the power of the RRS test in order to obtain a comparison to  $o(N^{-1})$  instead of  $o(1)$ .

In deriving such an expansion, we exploit the resemblance between the RRS test, the LMP rank test and the LMP permutation test. For the LMP permutation test the scores are  $-\psi_1(Z_j)$ . These are also random and of the same form as the  $A_j$  used here. The typical difference, however, is that the permutation test scores are based on the sample itself, whereas the RRS test scores are found from a second, independent sample which comes from  $F$  under  $H_0$  as well as under  $H_1$ . The independence thus obtained between the  $A_j$  and

the  $\bar{V}_j$  enables us to apply the results of chapter 3 to a much larger extent than was possible for permutation tests.

Denote  $-\psi_1(z_j^*)$  as  $a_j$ ,  $-\psi_1(X_j^*)$  as  $\tilde{A}_j$  and  $-\psi_1(x_j^*)$  as  $\tilde{a}_j$ . Let  $V_j = (1 + \bar{V}_j)/2$ ,  $j = 1, \dots, N$ . Hence  $L = 2\sum A_j V_j - \sum A_j$ . Conditionally on  $Z^* = z^*$ ,  $L$  is equivalent to  $\sum a_j V_j$ . Moreover, in view of the independence of  $A_j$  and  $V_j$  we have  $P_\theta(V_j=1|z^*) = P_\theta(V_j=1)$ . Hence in the conditional situation we have exactly the same case as in chapter 3. Define

$$R_\theta(x|z^*) = P_\theta\left(\frac{L - \theta N E \tilde{A}_1^2}{(N E \tilde{A}_1^2)^{1/2}} \leq x|z^*\right),$$

$$\gamma(\zeta) = \lambda\{x|\exists_j |x - a_j| < \zeta\},$$

where  $\lambda$  denotes Lebesgue measure. The following lemma supplies an expansion for  $R_\theta(x|z^*)$ .

LEMMA 4.5.1. *Suppose that  $z^*$  is such that there exist positive constants  $c$ ,  $\bar{c}$  and  $\delta$  for which*

$$(4.5.1) \quad \sum a_j^4 \leq cN, \quad \sum a_j^2 \geq cN, \quad \gamma(\zeta) \geq \delta N \zeta,$$

for some  $\zeta \geq N^{-3/2} \log N$ . Moreover, assume that there are positive constants  $D$ ,  $\bar{C}$ ,  $\varepsilon$ ,  $\tilde{c}$  and  $\eta$  such that

$$(4.5.2) \quad 0 \leq \theta \leq DN^{-1/2},$$

$$(4.5.3) \quad \sup \left\{ \int_{-\infty}^{\infty} |\psi_j(x+y)|^{m_j} f(x) dx : |y| \leq \varepsilon \right\} \leq \bar{C}, \quad j = 1, \dots, 4,$$

$$m_1 = 6, \quad m_2 = 3, \quad m_3 = 4/3, \quad m_4 = 1,$$

$$(4.5.4) \quad \left| \frac{d}{dy} \psi_1(F^{-1}(y)) \right| \geq \tilde{c},$$

on a subinterval  $\tau$  of  $(0,1)$  with length at least  $\eta$ . Then there exists  $A > 0$ , depending on  $N$ ,  $z^*$ ,  $\theta$  and  $F$  only through  $c$ ,  $\bar{C}$ ,  $\delta$ ,  $D$ ,  $\bar{C}$ ,  $\varepsilon$ ,  $\tilde{c}$  and  $\eta$ , such that

$$\sup_x |R_\theta(x|z^*) - K_\theta(y_x)| \leq A \{N^{-5/4} + N^{-3/4} \theta^3 [\{E|A_j - EA_j|^3\}^{4/9}]^{9/4}\},$$

where  $K_\theta(x)$  is given by (3.3.4) and

$$y_x = [x - \theta \frac{[\sum_j (a_j EA_j - EA_j^2)]}{(NEA_1^2)^{1/2}}] \frac{[\sum_j \tilde{a}_j^2]^{-1/2}}{NEA_1^2}.$$

PROOF. In the first place we note that  $EA_1^2$  remains bounded away from zero in view of (4.5.4). Hence no problems arise from the fact that  $EA_1^2$  occurs in the denominator of  $(L - \theta NEA_1^2)/(NEA_1^2)^{1/2}$ .

Next we observe that  $P_\theta(x|z^*) = P_\theta(L/(\sum a_j^2)^{1/2} \leq \{x + \theta(NEA_1^2)^{1/2}\} \{\sum \tilde{a}_j^2 / NEA_1^2\}^{-1/2})$  and  $L = 2\sum a_j v_j - \sum a_j$ . In view of the conditions of the present lemma, we can apply theorem 3.3.1, where the argument of  $K_\theta$  is

$$[x + \theta(NEA_1^2)^{1/2}] \frac{[\sum_j \tilde{a}_j^2]^{-1/2}}{NEA_1^2} - \theta \frac{\sum_j a_j E_0(-\psi_1(z_j))}{(\sum a_j^2)^{1/2}}.$$

As  $E_0(-\psi_1(z_j)) = EA_j$ , the desired result follows.  $\square$

The obvious way to proceed is to show that the conditions of this lemma are satisfied for all  $z^*$ , except for those in a set B with sufficiently small probability. Then the unconditional distribution of L follows by taking expectations with respect to  $Z^*$ . However, as  $K_\theta(y_x)$  is not bounded on B, we first have to replace  $K_\theta(y_x)$  by an approximation that can be controlled on B. Define

$$(4.5.5) \quad \tilde{K}_\theta(x) = \phi(x) + \phi(x) \left[ \frac{\sum_j a_j^4}{12(NEA_1^2)^2} (x^3 - 3x) + \frac{\theta \sum_j a_j^3 EA_j}{3(NEA_1^2)^{3/2}} (x^2 - 1) + \right. \\ \left. + \frac{\theta^2}{2NEA_1^2} \{ \sum_j a_j^2 EA_j^2 - \sigma^2(\sum_j a_j A_j) \} x + \frac{\theta^3}{6(NEA_1^2)^{1/2}} \sum_j a_j E_0 \{ 3\psi_1^3(z_j) + \right. \\ \left. - 6\psi_1\psi_2(z_j) + \psi_3(z_j) \} - \frac{1}{2} x \frac{\sum_j (\tilde{a}_j^2 - EA_1^2)}{NEA_1^2} - \theta \frac{\sum_j (a_j EA_j - EA_j^2)}{(NEA_1^2)^{1/2}} + \right. \\ \left. - \frac{1}{8} \frac{(\sum_j (\tilde{a}_j^2 - EA_1^2))^2}{(NEA_1^2)^2} (x^3 - 3x) - \frac{\theta}{2} \frac{\sum_j (\tilde{a}_j^2 - EA_1^2) \sum_j (a_j EA_j - EA_j^2)}{(NEA_1^2)^{3/2}} (x^2 - 1) + \right. \\ \left. - \frac{\theta^2}{2} \frac{(\sum_j (a_j EA_j - EA_j^2))^2}{NEA_1^2} x \right],$$

$$(4.5.6) \quad R = N^{-5/4} + N^{-3} |\sum_j (\tilde{a}_j^2 - EA_1^2)|^3 + \theta^3 N^{-3/2} |\sum_j (a_j EA_j - EA_j^2)|^3.$$

LEMMA 4.5.2. Under the conditions of lemma 4.5.1 we have

$$\sup_x |K_\theta(y_x) - \tilde{K}_\theta(x)| \leq AR.$$

PROOF. As  $\sum(\tilde{a}_j^2 - EA_j^2)/(NEA_1^2)$  remains bounded away from -1, we get the following expansion for  $y_x$

$$(4.5.7) \quad y_x = \left\{ x - \theta \frac{\sum(a_j EA_j - EA_j^2)}{(NEA_1^2)^{1/2}} \right\} \left\{ 1 + \frac{\sum(\tilde{a}_j^2 - EA_1^2)}{NEA_1^2} \right\}^{-1/2} = \\ = x - \frac{1}{2} x \frac{\sum(\tilde{a}_j^2 - EA_1^2)}{NEA_1^2} - \theta \frac{\sum(a_j EA_j - EA_j^2)}{(NEA_1^2)^{1/2}} + \frac{3}{8} x \frac{(\sum(\tilde{a}_j^2 - EA_1^2))^2}{(NEA_1^2)^2} + \\ + \frac{\theta}{2} \frac{\sum(a_j EA_j - EA_j^2) \sum(\tilde{a}_j^2 - EA_1^2)}{(NEA_1^2)^{3/2}} + O(R[1+|x|]).$$

Under the conditions of lemma 4.5.1,  $1/y_x = O(1/x)$  as  $|x| \rightarrow \infty$ , and therefore

$$(4.5.8) \quad \phi(y_x) = \phi(x) + \phi(x) \left\{ (y_x - x) - \frac{1}{2} x (y_x - x)^2 \right\} + O(|y_x - x|^3 \phi(\frac{1}{2} x)).$$

As for all  $p > 0$ ,  $x^p \phi(x) = O(1)$ , (4.5.7) and (4.5.8) together show that

$$(4.5.9) \quad \phi(y_x) = \phi(x) + \phi(x) \left\{ -\frac{1}{2} x \frac{\sum(\tilde{a}_j^2 - EA_1^2)}{NEA_1^2} - \theta \frac{\sum(a_j EA_j - EA_j^2)}{(NEA_1^2)^{1/2}} + \right. \\ + \frac{1}{8} (3x - x^3) \frac{(\sum(\tilde{a}_j^2 - EA_1^2))^2}{(NEA_1^2)^2} + \frac{\theta}{2} (1 - x^2) \frac{\sum(\tilde{a}_j^2 - EA_1^2) \sum(a_j EA_j - EA_j^2)}{(NEA_1^2)^{3/2}} + \\ \left. - \frac{\theta^2}{2} x \frac{(\sum(a_j EA_j - EA_j^2))^2}{NEA_1^2} \right\} + O(R).$$

From (4.5.7) it is clear that

$$(4.5.10) \quad y_x = x + O(N^{-1} x |\sum(\tilde{a}_j^2 - EA_1^2)|) + \theta N^{-1/2} |\sum(a_j EA_j - EA_j^2)|.$$

By means of Hölder's inequality we show that the coefficients of the second order terms in  $K_\theta(x)$  are  $O(N^{-1})$ . Together with (4.5.10) this leads to

$$(4.5.11) \quad K_{\theta}(y_x) - \Phi(y_x) = K_{\theta}(x) - \Phi(x) + O(N^{-2} |\sum (\tilde{a}_j^2 - EA_j^2)| + \\ + \theta N^{-3/2} |\sum (a_j EA_j - EA_j^2)|) = K_{\theta}(x) - \Phi(x) + O(R).$$

Finally,

$$(4.5.12) \quad \theta^{(4-k)} (\sum a_j^2)^{-k/2} = (\theta)^{(4-k)} (NEA_1^2)^{-k/2} + O(N^{-2} |\sum (\tilde{a}_j^2 - EA_j^2)|) = \\ = \theta^{(4-k)} (NEA_1^2)^{-k/2} + O(R), \quad k = 1, 2, 3, 4.$$

Combination of (4.5.9), (4.5.11) and (4.5.12) leads to  $\tilde{K}_{\theta}(x)$ .  $\square$

Now we can give an expansion for the unconditional distribution of L. Define

$$R_{\theta}(x) = P_{\theta} \left( \frac{L - \theta NEA_1^2}{(NEA_1^2)^{1/2}} \leq x \right).$$

LEMMA 4.5.3. *Let there be positive constants D,  $\epsilon$ ,  $\tilde{C}$ ,  $\tilde{c}$  and  $\eta$  such that (4.5.2), (4.5.3) and (4.5.4) hold. Suppose in addition that  $\int_{-\infty}^{\infty} \psi_1^{10}(x) f(x) dx \leq \tilde{C}$ . Then there exists A > 0 depending on N,  $\theta$  and F only through D,  $\epsilon$ ,  $\tilde{C}$ ,  $\tilde{c}$  and  $\eta$ , such that*

$$(4.5.13) \quad \sup_x |R_{\theta}(x) - \tilde{E}K_{\theta}(x)| \leq A \{ N^{-3/4} \theta^3 [\sum \{E|A_j - EA_j|\}^3]^{4/9} + ER \},$$

where  $\tilde{K}_{\theta}(x)$  and R are given by (4.5.5) and (4.5.6) respectively.

PROOF. The scores  $A_j$  we use, have the same distribution as the scores  $-\psi_1(Z_j)$  for the LMP permutation test under the hypothesis and therefore we can apply lemma 4.3.3. This shows that under the conditions of the present lemma the results of lemma's 4.5.1 and 4.5.2 hold, except on a set B with  $P_{\theta}(B) = O(N^{-5/4})$ .

In view of corollary 4.3.1 it remains to prove that  $E|K_{\theta}(x)I_B| = O(ER)$ . To this end we note that

$$|\sum a_j^3 EA_j| \leq \sum a_j^4 + \sum (EA_j)^4 \leq \sum a_j^4 + \sum EA_j^4 = O(\sum a_j^4 + N).$$

Likewise  $|\sum a_j^2 EA_j^2|$  and  $|\sum a_j E_0 \{3\psi_1^3(Z_j) - 6\psi_1\psi_2(Z_j) + \psi_3(Z_j)\}|$  are of this order and  $\sigma^2(\sum a_j A_j) = O((\sum a_j^2)^2 + N^2)$ . Furthermore,  $\sum a_j^4 \leq (\sum a_j^2)^2 \leq 2N^2 \tilde{EA}_1^2 + 2\{\sum (\tilde{a}_j^2 - EA_1^2)\}^2$ . Hence the contribution of the second, third and fourth term of  $\tilde{K}_\theta$  to  $E|\tilde{K}_\theta I_B|$  is  $O(ER)$ . The same result can be proved for the remaining terms of  $\tilde{K}_\theta$  in a manner analogous to the proof for the corresponding terms in corollary 4.3.1.  $\square$

COROLLARY 4.5.1. *Under the conditions of lemma 4.5.3 we have*

$$(4.5.14) \quad \sup_x |R_\theta(x) - \tilde{R}_\theta(x)| \leq A \left\{ N^{-5/4} + N^{-3/4} \theta^3 [\sum \{E|A_j - EA_j\}^3]^{4/9} \right\}^{9/4} + \\ + N^{-1} \theta^2 E[\sum (A_j - EA_j)^2]^2 + N^{-1} \theta \{1 + \theta \sigma(\sum (EA_j) A_j)\} \{E[\sum (A_j - EA_j)^2]^2\}^{1/2},$$

where

$$(4.5.15) \quad \tilde{R}_\theta(x) = \phi(x) + \phi(x) \left\{ -\frac{N^{-1}}{24} \left( \frac{\tilde{EA}_1^4}{(EA_1^2)^2} - 3 \right) (x^3 - 3x) + \right. \\ + \frac{N^{-1/2} \theta}{12} \left\{ \frac{4N^{-1} \sum EA_j^3 EA_j - 3\tilde{EA}_1^4}{(EA_1^2)^{3/2}} + 3(EA_1^2)^{1/2} \right\} (x^2 - 1) + \\ + \frac{\theta^2}{2EA_1^2} \{ \sum (EA_j^2)^2 - 2\sigma^2(\sum (EA_j) A_j) \} x + \\ + \frac{\theta^3}{6(NEA_1^2)^{1/2}} \sum (EA_j) E_0 \{ 3\psi_1^3(Z_j) - 6\psi_1\psi_2(Z_j) + \psi_3(Z_j) \} + \\ \left. + \frac{\theta}{(NEA_1^2)^{1/2}} \sum \sigma^2(A_j) \right\}.$$

PROOF. In view of lemma 4.5.3 the essence of the proof is the evaluation of  $E\tilde{K}_\theta(x)$  and  $ER$  from (4.5.5) and (4.5.6). The first term in (4.5.5) that deserves attention is  $\sigma^2(\sum a_j A_j)$ . Its expectation with respect to  $Z^*$  is



$$\begin{aligned}
(4.5.16) \quad \mathbb{E}\sigma_0^2(\sum \psi_1(Z_j^*)\psi_1(Z_j)|Z^*) &= \mathbb{E}\mathbb{E}_0[\sum \psi_1(Z_j^*)\{\psi_1(Z_j)-\mathbb{E}_0\psi_1(Z_j)\}]^2 = \\
&= \mathbb{E}\mathbb{E}_0[\sum \mathbb{E}\psi_1(Z_j^*)\{\psi_1(Z_j)-\mathbb{E}_0\psi_1(Z_j)\} + \\
&+ \sum \{\psi_1(Z_j^*)-\mathbb{E}\psi_1(Z_j^*)\}\{\psi_1(Z_j)-\mathbb{E}_0\psi_1(Z_j)\}]^2 = \\
&= \sigma^2(\sum (\mathbb{E}A_j)A_j) + 2\mathbb{E}\mathbb{E}_0[\sum \mathbb{E}\psi_1(Z_j^*)\{\psi_1(Z_j)-\mathbb{E}_0\psi_1(Z_j)\}](\sum \{\psi_1(Z_j^*)-\mathbb{E}\psi_1(Z_j^*)\} \cdot \\
&\cdot \{\psi_1(Z_j)-\mathbb{E}_0\psi_1(Z_j)\})] + \mathbb{E}\mathbb{E}_0[\sum \{\psi_1(Z_j^*)-\mathbb{E}\psi_1(Z_j^*)\}\{\psi_1(Z_j)-\mathbb{E}_0\psi_1(Z_j)\}]^2.
\end{aligned}$$

By Schwarz' inequality the last term in (4.5.16) is at most  $\mathbb{E}[\sum (A_j - \mathbb{E}A_j)^2]^2$ ; by applying the same inequality to the second term we arrive at

$$\begin{aligned}
\mathbb{E}\sigma_0^2(\sum \psi_1(Z_j^*)\psi_1(Z_j)|Z^*) &= \sigma^2(\sum (\mathbb{E}A_j)A_j) + \\
&+ O(\sigma(\sum (\mathbb{E}A_j)A_j)[\mathbb{E}\{\sum (A_j - \mathbb{E}A_j)^2\}]^{1/2} + \mathbb{E}\{\sum (A_j - \mathbb{E}A_j)^2\}^2).
\end{aligned}$$

Now we treat the remaining terms in (4.5.5). We begin by noting that

$$\mathbb{E}[\sum (\tilde{A}_j^2 - \mathbb{E}\tilde{A}_j^2)] = 0, \quad \mathbb{E}[\sum (\tilde{A}_j^2 - \mathbb{E}\tilde{A}_j^2)]^2 = N(\mathbb{E}\tilde{A}_1^4 - (\mathbb{E}\tilde{A}_1^2)^2),$$

$$\mathbb{E}[\sum (A_j \mathbb{E}A_j - \mathbb{E}A_j^2)] = -\sum \sigma^2(A_j), \quad \mathbb{E}[\sum (\tilde{A}_j^2 - \mathbb{E}\tilde{A}_j^2)]^3 = O(N^{3/2}).$$

Next we observe that  $\sum (A_j \mathbb{E}A_j - \mathbb{E}A_j^2) = \frac{1}{2}\sum (\tilde{A}_j^2 - \mathbb{E}\tilde{A}_j^2) - \frac{1}{2}\sum (A_j - \mathbb{E}A_j)^2 - \frac{1}{2}\sum \sigma^2(A_j)$ .

This leads to

$$\begin{aligned}
\mathbb{E}\{\sum (A_j \mathbb{E}A_j - \mathbb{E}A_j^2)\sum (\tilde{A}_j^2 - \mathbb{E}\tilde{A}_j^2)\} &= \frac{1}{2}\mathbb{E}[\sum (\tilde{A}_j^2 - \mathbb{E}\tilde{A}_j^2)]^2 + \\
&- \frac{1}{2}\mathbb{E}\{\sum (A_j - \mathbb{E}A_j)^2\sum (\tilde{A}_j^2 - \mathbb{E}\tilde{A}_j^2)\} = \\
&= \frac{N}{2}\{\mathbb{E}\tilde{A}_1^4 - (\mathbb{E}\tilde{A}_1^2)^2\} + O(N^{1/2}\{\mathbb{E}[\sum (A_j - \mathbb{E}A_j)^2]^2\}^{1/2}).
\end{aligned}$$

Furthermore,

$$\begin{aligned}
 E[\sum(A_j EA_j - EA_j^2)]^2 &= E[\sum(EA_j)(A_j - EA_j) - \sum\sigma^2(A_j)]^2 = \\
 &= \sigma^2(\sum(EA_j)A_j) + (\sum\sigma^2(A_j))^2, \\
 E|\sum(A_j EA_j - EA_j^2)|^3 &= O(E|\sum(EA_j)(A_j - EA_j)|^3 + (\sum\sigma^2(A_j))^3) = \\
 &= O(N\{\sum\sigma^2(A_j)\}^2 + [\sum|EA_j|\{E|A_j - EA_j|^3\}^{1/3}]^3) = \\
 &= O(N^{3/4}[\sum\{E|A_j - EA_j|^3\}^{4/9}]^{9/4} + N\{\sum\sigma^2(A_j)\}^2).
 \end{aligned}$$

Substitution of these results in (4.5.13) gives (4.5.14) and (4.5.15) with an additional term of order  $N^{-1}\theta^2(\sum\sigma^2(A_j))^2$ . As

$$(\sum\sigma^2(A_j))^2 \leq E[\sum(A_j - EA_j)^2]^2,$$

this term may be omitted.  $\square$

A further simplification of (4.5.14) and (4.5.15) is achieved by applying theorem 3.3.2 and the results of section 3.4. In doing so, we restrict attention to a fixed d.f. F. One of the consequences of this restriction is that condition (4.5.3) can be given a weaker formulation, whereas condition (4.5.4) can be omitted altogether (cf. the first remark following lemma 4.2.5). Let  $\pi(\theta)$  be the power of the test based on L and define

$$\begin{aligned}
 (4.5.17) \quad L_\theta(x) &= \phi(x) + \frac{\phi(x)}{N} \left[ -\frac{(\eta_2 - 3)}{24}(x^3 - 3x) + \frac{\eta_1(\eta_2 + 3)}{12}(x^2 - 1) + \right. \\
 &\quad \left. + \frac{\eta_1^2(\eta_2 + 1)}{4}x + \frac{\eta_1^3(3\eta_3 + \eta_2)}{18} + \eta_1 \frac{\int_{-1}^{1-1/N} (\psi_1'(t))^2 t(1-t) dt}{E_0 \psi_1^2(X_1)} \right],
 \end{aligned}$$

$$(4.5.18) \quad \tilde{\pi}(\theta) = 1 - \Phi(u_\alpha - \eta_1) + \frac{\eta_1 \phi(u_\alpha - \eta_1)}{N} \left[ - \frac{(5\eta_2 - 3)}{24} (u_\alpha^2 - 1) + \right. \\ \left. + \frac{\eta_1(\eta_2 - 3)}{24} u_\alpha - \frac{(12\eta_3 - 5\eta_2 - 9)\eta_1^2}{72} - \frac{\int_{1/N}^{1-1/N} (\Psi_1'(t))^2 t(1-t) dt}{E_0 \psi_1^2(X_1)} \right],$$

where  $\Psi_1$ ,  $\eta_1$ ,  $\eta_2$ ,  $\eta_3$  are defined in (3.3.14), (3.4.24), (3.4.26). Let  $F$  be the class of d.f.'s  $F$ , defined by (3.4.5) and (3.4.6). We now arrive at our final result.

**THEOREM 4.5.1.** *Suppose that  $F$  is such that  $F \in \mathcal{F}$  and  $\int_{-\infty}^{\infty} \psi_1^{10}(x) f(x) dx < \infty$ . Let there be positive constants  $C$  and  $\varepsilon$  such that  $0 \leq \theta \leq CN^{-1/2}$  and  $\varepsilon \leq \alpha \leq 1 - \varepsilon$ . Then, for every fixed  $F$ ,  $C$  and  $\varepsilon$  there are positive numbers  $A$ ,  $\delta_1$ ,  $\delta_2, \dots$  such that  $\lim_{N \rightarrow \infty} \delta_N = 0$  and for every  $N$*

$$(4.5.19) \quad \sup_x |P_\theta \left( \frac{L_{-\theta NE} \psi_1^2(X_1)}{[NE \psi_1^2(X_1)]^{1/2}} \leq x \right) - L_\theta(x)| \leq \delta_N N^{-1} + \\ + AN^{-3/2} \int_{1/N}^{1-1/N} (\Psi_1'(t))^2 (t(1-t))^{1/2} dt,$$

$$(4.5.20) \quad |\pi(\theta) - \tilde{\pi}(\theta)| \leq \delta_N N^{-1} + AN^{-3/2} \int_{1/N}^{1-1/N} (\Psi_1'(t))^2 (t(1-t))^{1/2} dt.$$

**PROOF.** We check that under the conditions of this theorem the results of theorem 3.3.2 and lemma 4.5.3 hold. As was mentioned after theorem 3.4.2, the fact that  $F \in \mathcal{F}$  implies that  $\Psi_1'(t) = o([t(1-t)]^{-7/6})$  near 0 and 1. Hence theorem 3.3.2 holds and

$$N^{-3/4} \theta^3 [\sum \{E|A_j - EA_j|\}^3]^{4/9} 9^{1/4} = O(N^{-5/4}).$$

The other remainder terms in (4.5.14) are dealt with in an analogous fashion. One may show that the condition  $|\Psi_1'(t)| = o(t(1-t))^{-5/4}$  near 0 and 1, already suffices to ensure that  $\theta \sigma(\sum (EA_j)A_j) = o(1)$ ,  $E[\sum (A_j - EA_j)^2]^2 = o(N)$ , and hence that  $|R_\theta(x) - \tilde{R}_\theta(x)| = o(N^{-1})$ .

The fact that the conditions of the present theorem also imply the results of lemma 4.5.3 and corollary 4.5.1 is verified in the same way as in which it is shown that theorem 3.4.1 implies theorem 3.3.1. Application of the results in (3.4.16)-(3.4.23) to (4.5.15) yields (4.5.19). From this, (4.5.20) follows in the usual way.  $\square$

REMARK. Note that under the hypothesis the theorem asserts that

$$P_0\left(\frac{L}{[NE_0\psi_1^2(X_1)]^{1/2}} \leq x\right) = \Phi(x) - \phi(x) \frac{(n_2-3)}{24N}(x^3-3x) + o(N^{-1}).$$

This agrees with the fact that under  $H_0$  the statistics  $L$  and  $-\sum_{j=1}^N \psi_1(X_j)$  have the same distribution.

#### 4.6. DEFICIENCIES

In this section we obtain the deficiencies among the various tests considered. First we summarize the results of Chapter 3, section 4.2, 4.3 and 4.5.  $X_1, \dots, X_N$  are i.i.d. r.v.'s from a d.f.  $F(x-\theta)$ , where  $f = F'$  is symmetric around zero and positive on  $R^1$ . Denote the power

of the MP parametric test, based on $1/\theta \sum \log\{f(X_j-\theta)/f(X_j)\}$	by $\pi_1(\theta)$ ,
of the LMP parametric test, based on $-\sum \psi_1(X_j)$	by $\pi_2(\theta)$ ,
of the MP permutation test, based on $(1/2\theta) \sum_{X_j > 0} \log\{f(X_j-\theta)/f(X_j+\theta)\}$	by $\pi_3(\theta)$ ,
of the LMP permutation test, based on $-\sum_{X_j > 0} \psi_1(X_j)$	by $\pi_4(\theta)$ ,
of the LMP rank test, based on $-\sum E_0 \psi_1(Z_j) \bar{V}_j$	by $\pi_5(\theta)$ ,
of the LMP RRS test, based on $-\sum \psi_1(Z_j^*) \bar{V}_j$	by $\pi_6(\theta)$ .

Note that  $\pi_1(\theta)$  is the envelope power. The following two types of conditions will be imposed:

$$f \text{ is } k \text{ times differentiable and } \psi_j = \frac{f^{(j)}}{f} \text{ satisfies}$$

(4.6.1)

$$\limsup_{y \rightarrow 0} \int_{-\infty}^{\infty} |\psi_j(x+y)|^{m_j} f(x) dx < \infty, \quad j = 1, \dots, k,$$

where  $k$  is a positive integer and  $m_j > 0$ ,  $j = 1, \dots, k$ , and

$$(4.6.2) \quad \limsup_{t \rightarrow 0, 1} t(1-t) \left| \frac{\psi_1''(t)}{\psi_1'(t)} \right| < \frac{3}{2}, \quad \text{where } \Psi_1(t) = \psi_1(F^{-1}(\frac{1+t}{2})).$$

Now we introduce four classes of d.f.'s  $F$ , determined by such conditions

$$F_1 = \{F \mid (4.6.1) \text{ holds for } k = 5, m_j = \frac{5}{j}, j = 1, \dots, 5\},$$

$$F_2 = F_1 \cap \{F \mid \limsup_{y \rightarrow 0} \int_{-\infty}^{\infty} \psi_1^{10}(x+y) f(x) dx < \infty\},$$

$$F_3 = \{F \mid (4.6.1) \text{ holds for } k = 4, m_1 = 6, m_2 = 3, m_3 = \frac{4}{3}, \\ m_4 = 1; (4.6.2) \text{ holds}\},$$

$$F_4 = F_3 \cap \{F \mid \int_{-\infty}^{\infty} \psi_1^{10}(x)f(x)dx < \infty\}.$$

Finally, let  $K = (k_{ij})$  be the  $6 \times 4$  matrix

$$\begin{array}{cccc} (\eta_2-3)/24 & -(\eta_2-3)/24 & -(\eta_2-3)/24 & (2\eta_2-3\eta_3)/72 \\ (\eta_2-3)/24 & -(\eta_2-3)/24 & -(\eta_2-3)/24 & (5\eta_2-12\eta_3+9)/72 \\ -\eta_2/12 & (\eta_2-3)/24 & -(\eta_2-3)/24 & (5\eta_2-12\eta_3+9)/72 \\ -\eta_2/12 & (\eta_2-3)/24 & -(\eta_2-3)/24 & (5\eta_2-12\eta_3+9)/72 \\ -\eta_2/12 & (\eta_2-3)/24 & \eta_2/12 - \sum \sigma^2(\psi_1(U_{j:N})) / \{2E_0\psi_1^2(X_1)\} & (5\eta_2-12\eta_3+9)/72 \\ -(5\eta_2-3)/24 & (\eta_2-3)/24 & (5\eta_2-3)/24 - \sum \sigma^2(\psi_1(U_{j:N})) / \{E_0\psi_1^2(X_1)\} & (5\eta_2-12\eta_3+9)/72 \end{array}$$

After these preliminaries, the following theorem can be formulated.

**THEOREM 4.6.1.** *Suppose there are positive constants  $c, C$  and  $\epsilon$  such that  $c \leq N^{1/2}\theta \leq C$  and  $\epsilon \leq \alpha \leq 1-\epsilon$ . Moreover, assume that  $F \in F_1$  if  $i = 1, 2$ ,  $F \in F_2$  if  $i = 3, 4$ ,  $F \in F_3$  if  $i = 5$  and  $F \in F_4$  if  $i = 6$ . Then, for every fixed  $F, c, C$  and  $\epsilon$ , there exists positive numbers  $A, \delta_1, \delta_2, \dots$  such that  $\lim_{N \rightarrow \infty} \delta_N = 0$  and for every  $N$  and  $i = 1, 2, \dots, 6$*

$$(4.6.3) \quad \pi_i(\theta) = 1 - \Phi(u_\alpha - \eta_1) + \frac{\eta_1 \phi(u_\alpha - \eta_1)}{N} \{k_{i1} u_\alpha^2 + k_{i2} u_\alpha \eta_1 + k_{i3} + k_{i4} \eta_1^2\} + R_i,$$

where

$$\eta_1 = \theta (NE_0\psi_1^2(X_1))^{1/2}, \quad \eta_2 = \{E_0\psi_1^4(X_1)\} / \{E_0\psi_1^2(X_1)\}^2,$$

$$\eta_3 = \{E_0\psi_2^2(X_1)\} / \{E_0\psi_1^2(X_1)\}^2$$

and

$$|R_1| \leq AN^{-3/2}, \quad |R_2| \leq AN^{-3/2}, \quad |R_3| \leq AN^{-5/4},$$

$$|R_4| \leq AN^{-5/4}, \quad |R_5| \leq \delta_N N^{-1}, \quad |R_6| \leq \delta_N N^{-1}.$$

PROOF. The result is immediate in view of theorem 3.4.1, 3.4.2, lemma 4.2.5, theorem 4.3.1, lemma 4.3.5 and theorem 4.5.1. Note that here the result is given for a fixed  $F$ , not only in the case of rank tests and RRS tests, but also in the case of parametric and permutation tests (cf. the remark following lemma 4.2.5).  $\square$

REMARK. If  $F \in \mathcal{F}_3$ , the sum of variances occurring in  $k_{53}$  and  $k_{63}$ , can be written as

$$\sum \sigma^2\{\psi_1(U_{j:N})\} = \int_{1/N}^{1-1/N} (\psi_1'(t))^2 t(1-t) dt + R,$$

$$|R| \leq \delta_N + AN^{-1/2} \int_{1/N}^{1-1/N} (\psi_1'(t))^2 (t(1-t))^{1/2} dt.$$

Denote the deficiency of the test with power  $\pi_i(\theta)$  with respect to the test with power  $\pi_j(\theta)$  as  $d_N(i,j)$ ; if it exists, the corresponding asymptotic deficiency is denoted as  $d(i,j)$  ( $i, j = 1, \dots, 6$ ). In the following theorem we give  $d_N(i+1,i)$  and, if possible,  $d(i+1,i)$ ,  $i = 1, \dots, 5$ . As deficiencies are transitive, this suffices to find  $d_N(i,j)$  for all  $i$  and  $j$ .

THEOREM 4.6.2. *Under the conditions of theorem 4.6.1 we have*

$$d(2,1) = \eta_1^2(3\eta_3 - \eta_2 - 3)/12, \quad |d_N(2,1) - d(2,1)| \leq AN^{-1/2},$$

$$d(3,2) = (\eta_2 - 1)u_\alpha^2/4 - (\eta_2 - 3)\eta_1 u_\alpha/6, \quad |d_N(3,2) - d(3,2)| \leq AN^{-1/4},$$

$$d(4,3) = 0, \quad |d_N(4,3)| \leq AN^{-1/4},$$

$$d_N(5,4) = \{\sum \sigma^2(\psi_1(U_{j:N}))\} / \{E_0 \psi_1^2(X_1)\} - (\eta_2 - 1)/4 + \delta_N,$$

$$d_N(6,5) = (\eta_2 - 1)u_\alpha^2/4 + \{\sum \sigma^2(\psi_1(U_{j:N}))\} / \{E_0 \psi_1^2(X_1)\} - (\eta_2 - 1)/4 + \delta_N.$$

PROOF. As  $\phi\{u_\alpha - \theta(N + d_N)\}^{1/2} (E_0 \psi_1^2(X_1))^{1/2} = \phi(u_\alpha - \eta_1) - \eta_1 \phi(u_\alpha - \eta_1) d_N / (2N) + O(N^{-2})$ , it follows that  $d_N(i+1,i) = \{2N(\pi_i(\theta) - \pi_{i+1}(\theta))\} / \{\eta_1 \phi(u_\alpha - \eta_1)\} + O(N^{-1})$ . Application of (4.6.3) and the definition of  $K = (k_{ij})$  leads to the desired result.  $\square$

REMARK.  $d(2,1)$  is independent of  $\alpha$ ,  $d_N(5,4)$  only depends on  $\alpha$  and  $\theta$  through

$\delta_N$ ,  $d_N(6,5)$  only depends on  $\theta$  through  $\delta_N$ .

APPLICATION 1. We first consider the normal case  $F = \Phi$ . In section 3.4 it has already been verified that  $\Phi \in F_3$ . The other  $F_i$  also contain  $\Phi$ , as may be checked in the same way. Hence we may use all results of theorems 4.6.1 and 4.6.2. As  $\eta_1 = N^{1/2}\theta$ ,  $\eta_2 = 3$ ,  $\eta_3 = 2$ , it follows that

$$d(2,1) = 0, \quad d(3,2) = \frac{1}{2} u_\alpha^2, \quad d(4,3) = 0,$$

$$d_N(5,4) = \sum \sigma^2(\Phi^{-1}(\frac{1+U}{2} \frac{j:N}{N})) - \frac{1}{2} + \delta_N,$$

$$d_N(6,5) = \frac{1}{2}(u_\alpha^2 - 1) + \sum \sigma^2(\Phi^{-1}(\frac{1+U}{2} \frac{j:N}{N})) + \delta_N.$$

According to (3.4.34),

$$\sum \sigma^2(\Phi^{-1}(\frac{U}{2} \frac{j:N+1}{N})) = \int_0^{\Phi^{-1}(1-1/2N)} \frac{(2\Phi(x)-1)(1-\Phi(x))}{\phi(x)} dx + o(1) =$$

$$= 1/2 \log \log N + 0.288608... + o(1).$$

Hence  $d(5,4)$  and  $d(6,5)$  do not exist, but on the other hand,  $d_N(5,4)$  and  $d_N(6,5)$  are of order  $\log \log N$ .

In section 4.4 it is mentioned that the asymptotic deficiency  $d(3,t)$  of the normal permutation test with respect to the  $t$ -test equals zero. This result enables us to compare the  $t$ -test with the other tests we are considering. We have for example

$$(4.6.4) \quad d(t,1) = \frac{1}{2} u_\alpha^2,$$

$$(4.6.5) \quad d_N(5,t) = \sum \sigma^2(\Phi^{-1}(\frac{1+U}{2} \frac{j:N}{N})) - \frac{1}{2} + \delta_N.$$

The first result was already obtained by Hodges and Lehmann (1970).

A final remark on the normal case is that  $d_N(6,1) = 2d_N(5,1) + \delta_N$ : the deficiency of the RRS test with respect to the envelope power is twice the deficiency of the normal scores test with respect to the envelope power, apart from a term that tends to zero as  $N \rightarrow \infty$ .

APPLICATION 2. As a second example we take the logistic d.f.  $F(x) = 1/(1+e^{-x})$ .

As  $\psi_1 = 1 - 2F$ , one easily verifies that  $F \in \cap_{i=1}^4 F_i$ . We have  $n_1 = (N/3)^{1/2\theta}$ ,  $n_2 = n_3 = 9/5$ ,  $\sum \sigma^2(\psi_1(U_{j:N})) = 1/6 + o(1)$  and therefore  $d(i+1, i)$  exists for  $i = 1, \dots, 5$ :

$$d(2,1) = \frac{N\theta^2}{60}, \quad d(3,2) = \frac{u_\alpha^2}{5} + \frac{u_\alpha N^{1/2\theta}\sqrt{3}}{15}, \quad d(4,3) = 0,$$

$$d(5,4) = \frac{3}{10}, \quad d(6,5) = \frac{u_\alpha^2}{5} + \frac{3}{10}.$$

$d(2,1)$  and  $d(3,2)$  can be made more transparent by using the relation between  $\theta$  and the power of the test. If we want to achieve a certain power  $1 - \beta$  at level  $\alpha$ , it follows from  $\beta = \Phi(u_\alpha - \sqrt{3}N^{1/2\theta}/3) + O(N^{-1})$  that  $N^{1/2\theta} = \sqrt{3}(u_\alpha + u_\beta) + O(N^{-1})$  and therefore

$$d(2,1) = \frac{(u_\alpha + u_\beta)^2}{20}, \quad d_{3,2} = \frac{u_\alpha(2u_\alpha + u_\beta)}{5}.$$



## CHAPTER 5

## DEFICIENCIES OF SOME RELATED ESTIMATORS

## 5.1. INTRODUCTION

In this chapter we take advantage of the correspondence between some of the test statistics we considered in chapters 3 and 4, and some well-known estimators. This correspondence immediately gives expansions to  $o(N^{-1})$  for the distribution of these estimators. The expansions can be used to make deficiency comparisons between the estimators.

By applying certain generalizations of the Cramér-Rao bound - the so called Bhattacharyya bounds - we obtain a lower bound to  $o(N^{-1})$  for the variance of an unbiased estimator. We also derive the deficiency of the estimators considered with respect to this lower bound.

In section 5.2 maximum likelihood estimators are dealt with; in section 5.3 we consider Hodges-Lehmann estimators.

## 5.2. MAXIMUM LIKELIHOOD ESTIMATORS

Let  $X_1, \dots, X_N$  be i.i.d. r.v.'s from  $F(x-\theta)$ , where  $f = F'$  is symmetric around zero and positive on  $R^1$ . In section 4.2 we considered the test for  $\theta = 0$  against  $\theta > 0$  based on  $S^* = -\sum_{j=1}^N \psi_1(X_j)$  and derived the expansion  $\tilde{\pi}_{S^*}(\theta)$  for its power  $\pi_{S^*}(\theta)$  in (4.2.24). From this a similar expansion for the distribution of the maximum likelihood estimator (MLE)  $\hat{\theta}$  of  $\theta$  can be derived. Since  $\hat{\theta}$  is translation invariant, we may restrict attention to the case  $\theta = 0$ . Probabilities are then denoted as  $P_0$ , otherwise as  $P_\theta$ . Define

$$(5.2.1) \quad H_N(x) = P_0([\sum_{i=1}^N \psi_1^2(X_i)]^{1/2} \hat{\theta} \leq x),$$

$$(5.2.2) \quad \tilde{H}_N(x) = \phi(x) + \frac{x\phi(x)}{N} \left\{ -\frac{(\eta_2-3)}{24} + \frac{x^2}{72} (5\eta_2-12\eta_3+9) \right\},$$

where

$$\eta_2 = E_0 \psi_1^4(X_1) / \{E_0 \psi_1^2(X_1)\}^2 \quad \text{and} \quad \eta_3 = E_0 \psi_2^2(X_1) / \{E_0 \psi_1^2(X_1)\}^2.$$

LEMMA 5.2.1. Suppose that  $f$  is five times differentiable and

$$(5.2.3) \quad \limsup_{y \rightarrow 0} \int_{-\infty}^{\infty} |\psi_j(x+y)|^{5/j} f(x) dx < \infty, \quad j = 1, \dots, 5.$$

Moreover, assume that  $\psi_1$  is non-increasing. Then for every fixed d.f.  $F$  there exists  $A > 0$  such that for all  $x$

$$(5.2.4) \quad |H_N(x) - \tilde{H}_N(x)| \leq A\{N^{-3/2} + N^{-3/2} x^2\}.$$

PROOF. By definition,  $\hat{\theta}$  is the value of  $\theta$  for which  $\sum_{j=1}^N \psi_1(X_j - \theta) = 0$ . As  $\psi_1$  is non-increasing, the events  $\{\sum_{j=1}^N \psi_1(X_j - N^{-1/2}x) > 0\}$  and  $\{\hat{\theta} < N^{-1/2}x\}$  are essentially the same. Furthermore, under  $\theta = 0$ ,  $X_1 - N^{-1/2}x$  has the same distribution as  $X_1$  under  $\theta = -N^{-1/2}x$ . Together this shows that

$$(5.2.5) \quad P_{\left(-\frac{x}{\sqrt{N}}\right)} \left( \sum_{j=1}^N \psi_1(X_j) > 0 \right) \leq P_0(\hat{\theta} < \frac{x}{\sqrt{N}}) \leq P_{\left(-\frac{x}{\sqrt{N}}\right)} \left( \sum_{j=1}^N \psi_1(X_j) \geq 0 \right).$$

Let  $\{\delta_N\}$  be a sequence of positive real numbers defined by  $\delta_N = b \cdot N^{-p}$  for some  $b > 0$  and  $0 < p < \frac{1}{2}$ . We first restrict attention to the case where  $|x| < \delta_N N^{1/2}$ . Then, for  $\theta = -N^{-1/2}x$  and a fixed  $F$ , the conditions of lemmas 4.2.4 and 4.2.5 are satisfied (cf. the first remark after lemma 4.2.5). Hence we can use the expansion in (4.2.9) for the probabilities on the left and right side of (5.2.5). The first consequence of this is that these probabilities differ at most  $O(N^{-3/2} + N^{-3/2} x^2)$  and therefore

$$(5.2.6) \quad P_0(\hat{\theta} < \frac{x}{\sqrt{N}}) = P_{\left(-\frac{x}{\sqrt{N}}\right)}(S^* \leq 0) + O(N^{-3/2} + N^{-3/2} x^2).$$

Under  $\theta = 0$ , the symmetry of  $S^*$  gives  $P_0(S^* \leq 0) = \frac{1}{2} + O(N^{-3/2} + N^{-3/2} x^2)$ . Hence, if  $\pi_{S^*}(\theta, \alpha)$  denotes the power of the  $S^*$  test against the alternative  $\theta$  at level  $\alpha$ , (5.2.6) becomes

$$P_0(\hat{\theta} < \frac{x}{\sqrt{N}}) = 1 - \pi_{S^*}(-\frac{x}{\sqrt{N}}, \frac{1}{2}) + O(N^{-3/2} + N^{-3/2} x^2).$$

By replacing  $\pi_{S^*}$  by its expansion  $\tilde{\pi}_{S^*}$ , as given in (4.2.24), for  $\alpha = 1/2$ , we arrive at

$$P_0(\hat{\theta} < \frac{x}{\sqrt{N}}) = \tilde{H}_N(x(E_0 \psi_1^2(X_1))^{1/2}) + O(N^{-3/2} + N^{-3/2} x^2),$$

which proves (5.2.4) for  $|x| < \delta_N N^{1/2}$ .

From (5.2.2) it follows that  $|\tilde{H}_N(-x)| = |1 - \tilde{H}_N(x)| \leq A\{N^{-3/2} + N^{-3/2} x^2\}$  for  $|x| \geq \delta_N N^{1/2}$ . As  $H_N(x)$  is non-decreasing, this shows that (5.2.4) also holds for  $|x| \geq \delta_N N^{1/2}$ .  $\square$

REMARK. Linnik and Mitrofanova (1965) have given Edgeworth expansions to  $O(N^{-k})$  for the distribution of  $\hat{\theta}$  under rather restrictive conditions. Recently, Čibišev (1972) has announced results where such expansions are obtained under minimal assumptions.

We now have an expansion to  $o(N^{-1})$  for the distribution of  $\{NE_0\psi_1^2(X_1)\}^{1/2\hat{\theta}}$ . In order to answer the question, whether this also gives an expansion to  $o(N^{-1})$  for the variance of  $\{NE_0\psi_1^2(X_1)\}^{1/2\hat{\theta}}$ , we first consider the following frequently occurring situation (cf. Hodges and Lehmann (1956), Chernoff (1956)). A normal sequence of estimators  $T_N$  has an asymptotic distribution with variance  $\tau^2$ . Call  $\tau^2$  the variance of the asymptotic distribution. On the other hand, the variances of  $T_N$  tend to a limit  $\sigma^2$ , as  $N$  tends to infinity. Call  $\sigma^2$  the asymptotic variance. Now it is not necessarily true that  $\sigma^2 = \tau^2$ . We can only assert that  $\sigma^2 \geq \tau^2$  and strict inequality may occur. This arises, loosely speaking, when very large errors occur with very small probabilities. If one wants to take this possibility into account, one should use  $\sigma^2$  as a criterion of performance, otherwise one can use  $\tau^2$ .

In the present situation we define the following analogue of  $\tau^2$  for the normed sequence  $\{NE_0\psi_1^2(X_1)\}^{1/2\hat{\theta}}$

$$(5.2.7) \quad \tau_N^2 = \int_{-\infty}^{\infty} x^2 d\tilde{H}_N(x) - \left( \int_{-\infty}^{\infty} x d\tilde{H}_N(x) \right)^2.$$

Application of (5.2.2) shows that

$$(5.2.8) \quad \tau_N^2 = 1 + \frac{(n_3 - \frac{1}{3}n_2 - 1)}{N}.$$

On the other hand, we of course have the variance of  $\{NE_0\psi_1^2(X_1)\}^{1/2\hat{\theta}}$  which we shall denote by  $\sigma_N^2$ . Again one might use either  $\sigma_N^2$  or  $\tau_N^2$  as a criterion of performance. It is therefore desirable to have conditions under which  $\sigma_N^2 = \tau_N^2 + o(N^{-1})$ . Such conditions are supplied by the following lemma.

LEMMA 5.2.2. *Let there be constants  $k > 10$  and  $\tilde{c} > 0$  such that  $E_0|\psi_1(X_1)|^k < \infty$  and  $\psi_1^2 - \psi_2 \geq \tilde{c}$  on  $R^1$ . If condition (5.2.3) is also satisfied, we have, for every fixed d.f.  $F$ , a constant  $A > 0$  depending on  $N$  only through  $k$  and  $\tilde{c}$ , such that*

$$|\sigma_N^2 - \tau_N^2| \leq AN^{-\{1 + \frac{(k-10)}{2(k+2)}\}}.$$

PROOF. As  $-\psi'_1 = \psi_1^2 - \psi_2 \geq \tilde{c} > 0$ , we have for  $x > 0$

$$\begin{aligned} (5.2.9) \quad 1 - H_N(x\{E_0\psi_1^2(X_1)\}^{1/2}) &= P_0(\hat{\theta} > \frac{x}{\sqrt{N}}) = P_0(\sum_{j=1}^N \psi_1(X_j - \frac{x}{\sqrt{N}}) < 0) = \\ &= P_0(\sum_{j=1}^N \psi_1(X_j) < \frac{x}{\sqrt{N}} \sum_{j=1}^N \psi'_1(X_j - \rho_j \frac{x}{\sqrt{N}})) \leq P_0(\sum_{j=1}^N \psi_1(X_j) < -x\sqrt{N}\tilde{c}), \end{aligned}$$

where  $0 \leq \rho_j \leq 1$ ,  $j = 1, \dots, N$ . Application of Chebyshev's inequality leads to

$$(5.2.10) \quad P_0(\sum_{j=1}^N \psi_1(X_j) \leq -x\sqrt{N}\tilde{c}) \leq E_0 |N^{-1/2} \sum_{j=1}^N \psi_1(X_j)|^k (x\tilde{c})^{-k}.$$

As  $\psi_1(X_1), \dots, \psi_1(X_N)$  are i.i.d. r.v.'s with mean zero and  $E_0 |\psi_1(X_1)|^k < \infty$ , application of the inequality, due to Marcinkievitz, Zygmund and Chung (cf. 4.3.40), leads to

$$(5.2.11) \quad E_0 |N^{-1/2} \sum_{j=1}^N \psi_1(X_j)|^k < \infty.$$

Combining (5.2.9) - (5.2.11) we obtain that, for  $x > 0$ ,

$$(5.2.12) \quad 1 - H_N(x) = O(x^{-k}).$$

In view of the symmetry of  $f$ , we have for all  $x$

$$\begin{aligned} P_0(\hat{\theta} \leq \frac{x}{\sqrt{N}}) &= P_{(-\frac{x}{\sqrt{N}})}(\sum_{j=1}^N \psi_1(X_j) \geq 0) = \\ &= P_{(\frac{x}{\sqrt{N}})}(\sum_{j=1}^N \psi_1(X_j) \leq 0) = P_0(\hat{\theta} \geq -\frac{x}{\sqrt{N}}), \end{aligned}$$

and hence, for all  $x$ ,

$$(5.2.13) \quad H_N(-x) = 1 - H_N(x).$$

From (5.2.12) and (5.2.13) it follows that  $\int_{-\infty}^{\infty} x^2 dH_N(x) < \infty$ . But then (5.2.13) also implies that  $\int_{-\infty}^{\infty} x dH_N(x) = 0$  (i.e.,  $\hat{\theta}$  is unbiased), and hence  $\sigma_N^2 = \int_{-\infty}^{\infty} x^2 dH_N(x)$ . By another application of the symmetry of  $f$  and by

partial integration, this leads to

$$(5.2.14) \quad \sigma_N^2 = 2 \int_0^\infty x^2 dH_N(x) = 4 \int_0^\infty x(1-H_N(x))dx.$$

Let  $\{\rho_N\}$  be a sequence of positive real numbers with  $\rho_N \rightarrow \infty$  and  $\rho_N = o(N^{1/2})$ . In view of lemma 5.2.1 we have  $H_N(x) - \tilde{H}_N(x) = O(N^{-3/2} + N^{-3/2} x^2)$  for  $|x| < \rho_N$ . This leads to

$$\sigma_N^2 = 4 \int_0^{\rho_N} x(1-\tilde{H}_N(x))dx + 4 \int_{\rho_N}^\infty x(1-H_N(x))dx + O(N^{-3/2} \rho_N^4).$$

From (5.2.2) it follows that, for all  $x$ ,  $\tilde{H}_N(-x) = 1 - \tilde{H}_N(x)$  and hence  $\tau_N^2 = 4 \int_0^\infty x(1-\tilde{H}_N(x))dx$ . Furthermore, from (5.2.2) it is also clear that  $\int_{\rho_N}^\infty x(1-\tilde{H}_N(x))dx = O(\rho_N^3 \exp\{-\frac{1}{2}\rho_N^2\})$  and therefore

$$(5.2.15) \quad \sigma_N^2 - \tau_N^2 = 4 \int_{\rho_N}^\infty x(1-H_N(x))dx + O(\rho_N^3 e^{-\frac{1}{2}\rho_N^2} + N^{-3/2} \rho_N^4).$$

Application of (5.2.12) to the integral in (5.2.15) shows that

$$\sigma_N^2 - \tau_N^2 = O(\rho_N^{-k+2} + \rho_N^3 e^{-\frac{1}{2}\rho_N^2} + N^{-3/2} \rho_N^4).$$

For given  $k$ , it is most favourable to choose  $\rho_N = N^{\frac{3}{2}(k+2)^{-1}}$ . Then

$$\sigma_N^2 - \tau_N^2 = O(N^{-\frac{3}{2} \frac{k-2}{k+2}}),$$

which is the desired result.  $\square$

REMARK. Linnik and Mitrofanova (1965) prove that  $\sigma_N^2 = 1 + o(N^{-1/2})$ , under stronger conditions.

In the above lemma we have given conditions under which  $\sigma_N^2$  and  $\tau_N^2$  agree to  $o(N^{-1})$ . In the sequel we shall no longer consider both kinds of variances but always take  $\tau_N^2$  as our criterion of performance.

The expression for  $\tau_N^2$  in (5.2.8) can be used for deficiency comparisons between the MLE and other estimators. An application of this kind occurs in the next section, where the MLE is compared to Hodges-Lehmann estimators. In the present section another possibility is treated: here we use (5.2.8) for a comparison of  $\tau_N^2$  with certain lower bounds for the variance of an un-

biased estimator.

The first lower bound we consider is the well-known Cramér-Rao (CR) bound. The assumptions which are necessary to apply this bound are satisfied if the conditions of lemma 5.2.1 hold. We have for any unbiased estimator  $U_N(X_1, \dots, X_N)$  of  $\theta$  that

$$(5.2.16) \quad \sigma^2(\{NE_0\psi_1^2(X_1)\}^{1/2}U_N) \geq 1.$$

The variance of the asymptotic distribution of  $\{NE_0\psi_1^2(X_1)\}^{1/2}U_N$  may be smaller than this lower bound, but only on a subset of  $\theta \in R^1$  of Lebesgue measure zero. (cf. Bahadur (1964) and Rao (1965)). Comparison of (5.2.8) and (5.2.16) shows that the variance of the asymptotic distribution of  $\{NE_0\psi_1^2(X_1)\}^{1/2}\hat{\theta}$  is asymptotically equal to the CR bound and hence the MLE is optimal to this extent.

Now Bhattacharyya (1946) has developed a series of refinements of the CR bound, which can be applied if stronger conditions are satisfied than are necessary for the CR bound. The  $k$ -th Bhattacharyya (B) bound for  $\sigma^2(\{NE_0\psi_1^2(X_1)\}^{1/2}U_N)$  has the form  $1 + \sum_{j=1}^{k-1} a_j N^{-j}$ ; the first B bound is the CR bound and the  $(k+1)$ -th B bound is obtained from the  $k$ -th B bound by adding a term of order  $N^{-k}$ . Hence, for comparison to  $o(N^{-1})$ , the second B bound is needed. From Davis (1951) we obtain that the assumptions, needed for this bound, are satisfied if the conditions of lemma 5.2.1 hold. Furthermore, it follows that, in view of the symmetry of  $f$ ,  $a_1 \equiv 0$ , i.e. the second B bound coincides with the CR bound for the present case.

As  $\tau_N^2 = 1 + (\eta_3 - \frac{1}{3}\eta_2 - 1)/N$ , it follows that in general the MLE  $\hat{\theta}$  is not optimal to  $o(N^{-1})$ . The difference between  $\tau_N^2$  and the second B bound equals

$$(5.2.17) \quad \frac{\eta_3 - \frac{1}{3}\eta_2 - 1}{N} = \frac{\sigma_0^2(\psi_1'(X_1))}{N\{E_0\psi_1^2(X_1)\}^2} \geq 0,$$

where equality occurs only if  $\psi_1'$  is constant a.s., and this is the case if  $X_1$  has a normal distribution.

Finally we restate the result in terms of deficiencies. We have

$$\begin{aligned} \sigma^2(\hat{\theta}) &= \frac{\sigma_N^2}{NE_0\psi_1^2(X_1)} \geq \frac{1}{NE_0\psi_1^2(X_1)} + o(N^{-2}), \\ \frac{\tau_N^2}{NE_0\psi_1^2(X_1)} &= \frac{1}{NE_0\psi_1^2(X_1)} + \frac{\sigma_0^2(\psi_1'(X_1))}{N^2\{E_0\psi_1^2(X_1)\}^3}. \end{aligned}$$

This implies that the asymptotic deficiency of the MLE with respect to the second B bound is finite and equals

$$(5.2.18) \quad d = \frac{\sigma_0^2(\psi_1'(X_1))}{\{E_0\psi_1^2(X_1)\}^2}.$$

### 5.3. HODGES-LEHMANN ESTIMATORS

Let  $X_1, \dots, X_N$  be i.i.d. r.v.'s from  $F(x-\theta)$ , where  $f = F'$  is symmetric around zero and positive on  $R^1$ . Let  $Z_1, \dots, Z_N$  be the absolute order statistics of the  $X_1, \dots, X_N$  and define  $V_j = 1$  if the  $X_i$  corresponding to  $Z_j$  is positive, and  $V_j = 0$  otherwise, for  $j = 1, \dots, N$ . In chapter 3 we considered the test for the hypothesis of symmetry against  $F(x-\theta)$ ,  $\theta > 0$ , based on  $T = \sum_{j=1}^N a_j V_j$ , with  $a_j = EZ_j^*$ , where  $Z_1^*, \dots, Z_N^*$  are order statistics from a d.f.  $F^*$ .

From this rank test, Hodges and Lehmann (1963) have derived an estimator  $\tilde{\theta}$  of  $\theta$ . Define  $\mu = \frac{1}{2} \sum_{j=1}^N EZ_j^*$  and  $X-\theta = (X_1-\theta, \dots, X_N-\theta)$ . Under the hypothesis, the distribution of  $T$  is symmetric around  $\mu$ . Let  $\tilde{\theta}_1 = \sup \{\theta: T(X-\theta) > \mu\}$  and  $\tilde{\theta}_2 = \inf \{\theta: T(X-\theta) < \mu\}$ , then  $\tilde{\theta} = (\tilde{\theta}_1 + \tilde{\theta}_2)/2$  is the Hodges-Lehmann (HL) estimator. Hodges and Lehmann prove that  $\tilde{\theta}$  is translation invariant and distributed symmetrically around  $\theta$ . Hence it suffices to find the distribution of  $\tilde{\theta}$  for  $\theta = 0$ . The close connection between  $\tilde{\theta}$  and  $T$  makes it easy to find an expansion for the d.f. of  $\tilde{\theta}$  from the expansion for the d.f. of  $T$ , which we obtained in chapter 3. We restrict attention to the case where  $T$  is the locally most powerful rank test, i.e. where  $a_j = -E_0\psi_1(Z_j)$ . Define

$$(5.3.1) \quad \begin{aligned} K_N(x) &= P_0(\{NE_0\psi_1^2(X_1)\}^{1/2\tilde{\theta}} \leq x), \\ \tilde{K}_N(x) &= \Phi(x) + \frac{x\phi(x)}{N} \left\{ \frac{\eta_2}{12} - \frac{\sum_{j=1}^N \sigma_0^2(\psi_1(U_{j:N}))}{2E_0\psi_1^2(X_1)} + \frac{x^2}{72} (5\eta_2 - 12\eta_3 + 9) \right\}, \end{aligned}$$

where  $\eta_2$  and  $\eta_3$  are defined in (5.2.2),  $\psi_1(t) = \psi_1(F^{-1}(\frac{1+t}{2}))$  and  $U_{1:N} < \dots < U_{N:N}$  are order statistics of a sample of size  $N$  from the uniform distribution on  $(0,1)$ . Let  $F$  be the class of d.f.'s introduced in definition 3.4.1. The following lemma gives conditions under which  $|K_N - \tilde{K}_N|$  is  $o(N^{-1})$ .

LEMMA 5.3.1. *Suppose that  $F \in \mathcal{F}$  and that  $\psi_1$  is non-increasing. Then, for every fixed  $F$  and every positive constant  $C$ , there exist positive numbers*

$\delta_1, \delta_2, \dots$  such that  $\lim_{N \rightarrow \infty} \delta_N = 0$  and for all  $|x| \leq C$

$$|K_N(x) - \tilde{K}_N(x)| \leq \delta_N N^{-1}.$$

PROOF. From the construction of  $\tilde{\theta}$  and the monotonicity of  $\psi_1$  it follows that

$$(5.3.2) \quad P_0(T(X - \frac{X}{\sqrt{N}}) < \mu) \leq P_0(\tilde{\theta} \leq \frac{X}{\sqrt{N}}) \leq P_0(T(X - \frac{X}{\sqrt{N}}) \leq \mu),$$

cf. Hodges and Lehmann (1963). Note that expression (5.3.2) is completely analogous to (5.2.5). The remaining part of the proof is therefore analogous to the proof of lemma 5.2.1. We only mention that under the conditions of the present lemma, theorem 3.4.2 can be applied.  $\square$

In analogy to the previous section, we use as a criterion of performance for  $\tilde{\theta}$

$$(5.3.3) \quad \begin{aligned} \tilde{\tau}_N^2 &= \int_{-\infty}^{\infty} x^2 d\tilde{K}_N(x) - \left( \int_{-\infty}^{\infty} x d\tilde{K}_N(x) \right)^2 = \\ &= 1 + \frac{1}{N} \left\{ \eta_3 - \frac{7}{12} \eta_2 - \frac{3}{4} + \frac{\sum_{j=1}^N \sigma_0^2(\psi_1(U_{j:N}))}{E_0 \psi_1^2(X_1)} \right\}. \end{aligned}$$

Comparison of (5.2.8) and (5.3.3) leads to

$$(5.3.4) \quad \tilde{\tau}_N^2 - \tau_N^2 = \frac{\sum_{j=1}^N \sigma_0^2(\psi_1(U_{j:N}))}{NE_0 \psi_1^2(X_1)} - \frac{(\eta_2 - 1)}{4N}.$$

It follows that the deficiency of the HL estimator  $\tilde{\theta}$  with respect to the MLE  $\hat{\theta}$  equals

$$(5.3.5) \quad d_N = \frac{\sum_{j=1}^N \sigma_0^2(\psi_1(U_{j:N}))}{E_0 \psi_1^2(X_1)} - \frac{(\eta_2 - 1)}{4} + o(1).$$

Note that  $d_N$  equals the deficiency of the LMP rank test with respect to the LMP test for the size  $\alpha = 1/2$  (cf. Albers, Bickel and van Zwet (1974)). Finally, the deficiency of the HL estimator with respect to the second B bound is the sum of the deficiencies in (5.2.18) and (5.3.5).



## CHAPTER 6

## FINITE SAMPLE COMPUTATIONS

## 6.1. INTRODUCTION

In section 3.4 we derived expansions for the power of one sample linear rank tests based on exact or approximate scores against contiguous location alternatives. In section 6.2 to 6.4 we investigate the performance of these expansions as approximations to the finite sample power. In particular we compare these expansions to the usual normal approximations.

In section 4.6 deficiencies of the above rank tests with respect to various other types of tests for the one sample problem have been obtained. Section 6.5 is devoted to the comparison of these expressions with deficiencies for finite sample sizes that are obtained numerically. We focus attention on the normal case and consider the test based on the sample mean, the t-test and the one sample normal scores test.

## 6.2. THE NORMAL SCORES AND THE WILCOXON TEST AGAINST NORMAL AND LOGISTIC ALTERNATIVES

Here we shall consider the one sample normal scores (NS) test and the one sample Wilcoxon (W) test, both against normal (N) location alternatives  $G(x) = \Phi(x-\theta)$  and logistic (L) location alternatives  $G(x) = 1/[1+\exp(-x+\theta)]$ . We assume that  $\theta$  is non-negative and  $\theta = O(N^{-1/2})$ . From section 3.4 we have

$$(6.2.1) \quad \pi_{NS,N}(\theta) = 1 - \Phi\left(u_{\alpha} - \eta_1 + \frac{\eta_1}{4N}\{u_{\alpha}^2 + \right. \\ \left. - 1 + 2 \sum_{j=1}^N \sigma^2(\Phi^{-1}(\frac{1+U_{j:N}}{2}))\}\right) + o(N^{-1}),$$

$$(6.2.2) \quad \pi_{W,L}(\theta) = 1 - \Phi\left(u_{\alpha} - \eta_2 + \frac{\eta_2}{20N}\{3u_{\alpha}^2 + u_{\alpha}\eta_2 + 2 + \eta_2^2\}\right) + o(N^{-1}),$$

$$(6.2.3) \quad \pi_{NS,L}(\theta) = 1 - \Phi\left(u_{\alpha} - \eta_3 + \frac{\eta_3}{12N}\{u_{\alpha}^2 + u_{\alpha}\eta_3(-5+2\pi) + (23-12\sqrt{2}) + \right. \\ \left. + \eta_3^2(12 \arctan \frac{23\sqrt{2}+1}{20} - 4\pi) - 6 \sum_{j=1}^N \sigma^2(\Phi^{-1}(\frac{1+U_{j:N}}{2}))\}\right) + o(N^{-1}),$$

$$\begin{aligned}
(6.2.4) \quad \pi_{W,N}(\theta) &= 1 - \Phi\left(u_\alpha - \eta_4 + \frac{\eta_4}{N} \left\{ u_\alpha^2 \left( \frac{6 \arctan \frac{1}{4}\sqrt{2}}{\pi} - \frac{9}{20} \right) + \right. \right. \\
&+ u_\alpha \eta_4 \left( \frac{49}{20} - \frac{2}{3}\sqrt{3} - \frac{12 \arctan \frac{1}{4}\sqrt{2}}{\pi} \right) + \left( \frac{44}{20} - \sqrt{2} - \frac{6 \arctan \frac{1}{4}\sqrt{2}}{\pi} \right) + \\
&\left. \left. + \eta_4^2 \left( \frac{6 \arctan \frac{1}{4}\sqrt{2}}{\pi} + \frac{2}{3}\sqrt{3} - \frac{43}{20} + \frac{\pi}{9} \right) \right\} \right) + o(N^{-1}),
\end{aligned}$$

where  $U_{1:N} < \dots < U_{N:N}$  are order statistics from the uniform distribution on  $(0,1)$  and

$$(6.2.5) \quad \eta_1 = N^{1/2}\theta, \quad \eta_2 = \left(\frac{N}{3}\right)^{1/2}\theta, \quad \eta_3 = \left(\frac{N}{\pi}\right)^{1/2}\theta, \quad \eta_4 = \left(\frac{3N}{\pi}\right)^{1/2}\theta,$$

$$\begin{aligned}
(6.2.6) \quad \sum_{j=1}^N \sigma^2\left(\Phi^{-1}\left(\frac{1+U_{j:N}}{2}\right)\right) &= \int_0^{\Phi^{-1}\left(1-\frac{1}{2N}\right)} \frac{(2\Phi(x)-1)(1-\Phi(x))}{\phi(x)} dx + o(1) = \\
&= \frac{1}{2} \log \log N + 0.288608\dots + o(1).
\end{aligned}$$

Upon evaluation of the coefficients, (6.2.3) and (6.2.4) become

$$\begin{aligned}
(6.2.7) \quad \pi_{NS,L}(\theta) &= 1 - \Phi\left(u_\alpha - \eta_3 + \frac{\eta_3}{N} \left\{ 0.08333\{u_\alpha^2 + 0.10693 u_\alpha \eta_3 + \right. \right. \\
&+ 0.50245 + 0.05565 \eta_3^2 - \frac{1}{2} \sum_{j=1}^N \sigma^2\left(\Phi^{-1}\left(\frac{1+U_{j:N}}{2}\right)\right) \left. \left. \right\} \right) + o(N^{-1}).
\end{aligned}$$

$$\begin{aligned}
(6.2.8) \quad \pi_{W,N}(\theta) &= 1 - \Phi\left(u_\alpha - \eta_4 + \frac{\eta_4}{N} \left\{ 0.19904 u_\alpha^2 - 0.00278 u_\alpha \eta_4 + \right. \right. \\
&\left. \left. + 0.13675 + 0.00281 \eta_4^2 \right\} \right) + o(N^{-1}).
\end{aligned}$$

We shall now investigate how well the exact power is approximated by (6.2.1) - (6.2.4) for small samples. We shall also compare this approximation to the usual normal approximation, which approximates to  $o(1)$  instead of  $o(N^{-1})$ . The necessary results about exact powers of the tests involved can be found in papers by Klotz (1963) and Thompson, Govindarajulu and Eisenstat (1967).

Klotz gives the small sample power for the normal scores test and the Wilcoxon test against normal alternatives, for sample sizes  $N = 5(1)10$  at significance levels  $\alpha = k/2^N$  for various integers  $k$  and for shifts

$\theta = 1/4(1/4)3/2(1/2)3$ . The non-standard levels  $k/2^N$  are necessary to avoid randomization. In short, his method consists of selecting the  $k = \alpha \cdot 2^N$  orderings  $V = (V_1, \dots, V_N)$  that give rise to the largest values of  $T = \sum_{j=1}^N E J(U_{j:N}) V_j$  and adding the probabilities associated with these orderings. The evaluation of such probabilities involves the evaluation of an  $N$ -dimensional integral. By using a recursive scheme, this problem can be reduced to the computation of  $N$  one-dimensional integrals. In this way Klotz can go as far as  $N = 10$ , obtaining exact results in four decimal places.

For larger sample sizes the amount of computation that is necessary for this method, becomes prohibitive and one has to turn to Monte Carlo methods, as is done in the paper by Thompson et al. (1967).

They give the power for the normal scores and the Wilcoxon test against both normal and logistic alternatives for  $N = 10, 20$ ,  $\alpha = 0.01, 0.025$  and  $0.05$  and  $\theta = 1/4(1/4)1,3/2$ . In their paper these results are collected in table 4.1, where it is indicated that the test sizes considered are  $\alpha = 0.01, 0.025$  and  $0.10$ . However, the last value should be  $0.05$ . This is not only evident from the numerical results obtained, but also from a remark elsewhere in the paper.

The method used by Thompson et al. is the following: first the required critical values are found by using Edgeworth approximations up to an appropriate order under the hypothesis. Then, for each combination of  $N$ ,  $\theta$  and  $\alpha$  under consideration, 1000 trials are conducted. Each trial consists of drawing a random sample of size  $N$  from the standard normal or logistic distribution, shifting it over  $\theta$ , forming the Wilcoxon and normal scores test statistics and counting the number of samples for which the tests reject the hypothesis. This procedure results in unbiased power estimates with standard deviation at most 0.016.

In order to compute (6.2.1) to (6.2.4), we can use (6.2.6) for the sum of variances occurring in (6.2.1) and (6.2.3). Another possibility is to use values that are obtained numerically. Klotz (1963) gives a table of  $E\Phi^{-1}\{(1+U_{j:N})/2\}$  for  $N = 5(1)10$ , whereas Thompson et al. (1967) give  $\sum_{j=1}^N [E\Phi^{-1}\{(1+U_{j:N})/2\}]^2$  for  $N = 10(1)20, 30, 50$  and  $100$ . As

$$\sum_{j=1}^N \sigma^2 \Phi^{-1}\{(1+U_{j:N})/2\} = N - \sum_{j=1}^N [E\Phi^{-1}\{(1+U_{j:N})/2\}]^2,$$

these results enable us to find the sum of variances for various sample sizes. For the sample sizes we shall consider, we list in table 6.2.1 the numerical values and the values supplied by the second approximation in (6.2.6).

Table 6.2.1

Approximations of  $\sum_{j=1}^N \sigma^2 \phi^{-1} \left( \frac{1+U_{j:N}}{2} \right)$

N	numerical approximation	$\frac{1}{2} \log \log N + 0.288608$
5	0.693	0.527
6	0.724	0.580
7	0.752	0.621
8	0.777	0.655
9	0.794	0.682
10	0.810	0.706
20	0.911	0.837
50	1.022	0.971
100	1.080	1.052

Tables 6.2.2 - 6.2.7 (p.128-p.133) give the results of the comparison of the approximations (6.2.1) - (6.2.4) to the normal approximation and to the results of Klotz and Thompson et al. We have used the numerically obtained values for  $\sum_{j=1}^N \sigma^2 \phi^{-1} \{ (1+U_{j:N})/2 \}$  in (6.2.1) and (6.2.3).

Inspection of these tables shows that (6.2.1) - (6.2.4) supply excellent approximations for all N,  $\alpha$  and  $\theta$  under consideration. They always constitute a substantial improvement over the usual normal approximation, which yields values that are systematically too large. This bias is corrected by the  $O(N^{-1})$  term, as may also be seen from (6.2.1) to (6.2.4).

Note that by absorbing the  $N^{-1}$ -term in the argument of  $\phi$ , we have arrived in (6.2.1) - (6.2.4) at slightly different, but asymptotically equivalent versions of (3.4.30) - (3.4.33). As  $\{1-\phi(x)-\epsilon\phi(x)\} - \{1-\phi(x+\epsilon)\} \approx -\epsilon^2 x\phi(x)/2$  for small  $\epsilon$ , it follows that the approximations considered in the present chapter will tend to be slightly smaller (larger) than those from chapter 3 if the power which is approximated is larger (smaller) than 1/2. Since both versions tend to overestimate the exact power, this seems to indicate that the formulas (6.2.1) - (6.2.4) are slightly preferable.

### 6.3. THE WILCOXON TEST AGAINST CAUCHY ALTERNATIVES

In the previous section we have considered cases where the normal approximation already gives reasonable results. Adding terms of order  $N^{-1}$  merely constitutes an improvement, however substantial it may be, over an already

rather satisfactory approximation. In view of this, it seems more interesting to consider situations where the normal approximation performs very badly. If in such a case the approximation to  $o(N^{-1})$  does give reasonable results, we have found an approximation for a situation where none was available yet.

A case in which the normal approximation leads to very bad results occurs for example if Wilcoxon's test is used against location alternatives from a Cauchy distribution. For this case, an expansion to  $o(N^{-1})$  can be justified. The standard Cauchy distribution has density  $f(x) = 1/\{\pi(1+x^2)\}$ . Hence, in the notation of chapter 3,  $\psi_1(x) = -2x/(1+x^2)$ ,  $\psi_2(x) = 2(3x^2-1)/(1+x^2)^2$  and  $\psi_3(x) = -24x(x^2-1)/(1+x^2)^3$ . Furthermore,  $F(x) = (\arctan x)/\pi + 1/2$  and therefore  $F^{-1}\{(1+t)/2\} = \text{tg}(\pi t/2)$ ,  $-\Psi_1(t) = \sin \pi t$ . Finally, for Wilcoxon's test, we have  $J(t) = t$ . From these facts it can easily be verified that the conditions of theorem 3.4.1 are satisfied. After elementary integrations we find that the power  $\pi_{W,C}(\theta)$  satisfies  $\pi_{W,C}(\theta) = \tilde{\pi}_{W,C}(\theta) + o(N^{-1})$ , where

$$(6.3.1) \quad \tilde{\pi}_{W,C}(\theta) = 1 - \Phi\left(u_\alpha - \eta + \frac{\eta}{N} \left\{ u_\alpha^2 \left( \frac{11}{20} - \frac{6}{\pi} \right) + \right. \right. \\ \left. \left. + u_\alpha \eta \left( \frac{12}{\pi} - \frac{21}{20} \right) + \left( \frac{6}{\pi} - \frac{4}{5} \right) + \eta^2 \left( \frac{\pi^2}{9} + \frac{7}{20} - \frac{6}{\pi} \right) \right\} \right),$$

with  $\eta = (3N)^{1/2} \theta / \pi$ .

The exact power results for this case are obtained from a paper by Arnold (1965). Using the same approach as Klotz (1963), Arnold gives the power for Wilcoxon's test against alternatives from  $t$ -distributions with  $\frac{1}{2}$ , 1, 2 and 4 degrees of freedom. Note that the Cauchy distribution is the  $t_1$ -distribution and that the normal distribution is the  $t_\infty$ -distribution. The sample sizes considered are  $N = 5(1)10$ , the levels are  $\alpha = k/2^N$  for various  $k$  and the shifts are  $\mu = 1/4, 1/2, 1, 2$  and 3. To obtain a better comparison with the case of normal alternatives, Arnold has scaled all densities if he considers in such a way that  $\int_{1.645}^{\infty} f(x) dx = 0.05$ . Here  $1.645 = \Phi^{-1}(0.95)$ . As  $\frac{1}{\pi} \int_{6.314}^{\infty} (1+x^2)^{-1} dx = 0.05$ , this means for the Cauchy case that the standard density  $1/\{\pi(1+x^2)\}$  is replaced by  $f(x/\sigma)/\sigma$ , with  $1/\sigma = 6.314/1.645 = 3.838$ . Since the power for a shift  $\mu$  and a density  $f(x/\sigma)/\sigma$  under the hypothesis is the same as the power for a shift  $\mu/\sigma$  and a density  $f(x)$

under the hypothesis, the results from Arnold's paper and those obtained from (6.3.1) become comparable by inserting  $\theta = \mu/\sigma = 3.838\mu$  in (6.3.1), instead of  $\theta = \mu$ .

In analogy to the previous section we compare the exact results with the normal approximation and with (6.3.1). The normal approximation is very bad. It tends to 1 too fast for increasing  $\mu$ . However, the expansion in (6.3.1) leads to even worse results: as  $\mu$  increases, this approximation tends to 0 very fast. A typical result is

Table 6.3.1

$$N = 8, \alpha = 0.05469 = 14/256$$

$\mu$	power			
	exact	normal appr.	(6.3.1)	(6.3.1)'
1/4	.35	.46	.32	.32
1/2	.57	.92	.06	.47
1	.74	1.00	.00	1.00
2	.86	1.00	.00	1.00
3	.90	1.00	.00	1.00

Apparently we have not succeeded in finding a useful approximation by considering higher order terms. The expansion (6.3.1) obviously has a very local character. It will only give reasonable results for very small values of  $\theta$ . Here "small" means that these  $\theta$ -values give rise to values of  $\pi(\theta)$  considerably below 0.5, and this region is of little practical interest. The local character of (6.3.1) is borne out by computation of the coefficients in this expansion. We find

$$(6.3.2) \quad \tilde{\pi}_{W,C}(\theta) = 1 - \Phi\left(u_{\alpha} - 0.55N^{1/2}\theta + N^{-1/2}\theta\{-0.03u_{\alpha}^2 + 0.05u_{\alpha}N^{1/2}\theta - 0.10 + 0.14N\theta^2\}\right) \pm 0.00\dots$$

This shows that  $\tilde{\pi}_{W,C}(\theta)$  reaches its maximum for  $\theta \approx \left(\frac{0.55}{0.42}\right)^{1/2} \approx 1.1$ . As any approximation of the power  $\pi_{W,C}(\theta)$  should be increasing, it follows that  $\tilde{\pi}_{W,C}(\theta)$  is certainly unreliable for  $\theta \geq 1.1$ . Since  $\theta = 3.838\mu$ , we have that (6.3.1) is certainly bad for  $\mu \geq 0.3$ , which agrees with table 6.3.1.

A similar inspection of (6.2.7) and (6.2.8) shows that here the coefficient of  $N^{1/2}\theta^3\phi(u_\alpha - \eta)$  is much smaller than is the case in (6.3.2). Hence, these approximations first reach their maximum in  $\theta$  if  $\theta$  is large, i.e. if  $\pi(\theta)$  is already very close to 1. This explains their excellent performance.

Just as in section 6.2, we note that in (6.3.1) we have used the  $\{1-\Phi(x+\epsilon)\}$ -version of the expansion. Again, we can also consider the variant  $1-\Phi(x)-\epsilon\phi(x)$ , which has been denoted by (6.3.1)' in table 6.3.1. Clearly, in the present case this last version is better than the first. At least, it is less bad; in particular it is slightly better than the normal approximation instead of being worse.

#### 6.4. THE SIGN TEST AGAINST NORMAL, LOGISTIC AND DOUBLE EXPONENTIAL ALTERNATIVES

In section 3.5 an expansion was derived for the power  $\pi(\theta)$  of the sign test. Let  $\tau = N^{1/2}(2F(\theta)-1)$  and  $\gamma_\alpha = (N+1+N^{1/2}u_\alpha)/2 - [(N+1+N^{1/2}u_\alpha)/2]$ , where  $[y]$  denotes the integer part of  $y$ . Then, for all  $\theta$  such that  $\tau$  is bounded,

$$(6.4.1) \quad \pi(\theta) = 1 - \Phi\left(u_\alpha - \tau + \frac{\tau}{12N}\{u_\alpha^2 + u_\alpha\tau - 3\tau^2 + 24\gamma_\alpha(1-\gamma_\alpha) - 3\}\right) + O(N^{-3/2}).$$

Here we shall investigate the performance of (6.4.1) as an approximation to the exact power, in the case of normal, logistic and double exponential alternatives. The last type of alternatives is considered since the sign test is the locally most powerful rank test against such alternatives. For these three alternatives we have  $\tau = (2N/\pi)^{1/2} \int_0^\theta \exp(-x^2/2) dx$ ,  $\tau = N^{1/2}(1-e^{-\theta})/(1+e^{-\theta})$  and  $\tau = N^{1/2}(1-e^{-\theta})$ , respectively.

The paper by Thompson et al. (1967), discussed in the previous section, also contains Monte Carlo estimates of the sign test power against these three types of alternatives. The values of  $N$ ,  $\alpha$  and  $\theta$  that are considered, are the same as in the previous section. We compare approximation (6.4.1) with these estimates, and also with the normal approximation. The results are collected in tables 6.4.1-6.4.3 (p.134-p.136). It appears that (6.4.1) is better than the normal approximation, but the improvement is less striking than in section 6.2. For a number of combinations of  $N$ ,  $\alpha$  and  $\theta$  for which both approximations perform rather well (e.g. where the absolute error is less than 0.03), the normal approximation is even slightly better. However, in cases where larger errors occur (e.g. larger than 0.06), approximation (6.4.1) is always substantially better than the normal approximation. The explanation of the fact that including terms of order  $N^{-1}$  is less effective than in section 6.2, probably lies in the pronounced lattice character of the sign test statistic.

## 6.5. DEFICIENCIES BETWEEN TESTS FOR THE NORMAL CASE

In chapter 4 an approximation to  $o(1)$  for the deficiency of the locally most powerful rank test against alternatives  $F(x-\theta)$  was found with respect to various other tests that are optimal in some sense for the one sample problem. Here we shall go into the question to what extent such an asymptotic expansion is useful as a prediction of the deficiency for finite sample sizes.

We shall restrict attention to normal alternatives. This case is very interesting, as some of the competitors of the locally most powerful rank test, i.e. the normal scores test, are well-known. In the first place, the parametric tests based on  $\sum_{j=1}^N \log\{f(X_j-\theta)/f(X_j)\}$  and  $\sum_{j=1}^N -\psi_1(X_j)$  respectively, both reduce to the test based on the sample mean  $\bar{X} = N^{-1} \sum_{j=1}^N X_j$ . Furthermore, in the normal case the teststatistic of the locally most powerful scale invariant test is explicitly known: it is the t-test statistic. Let  $d_N(NS, \bar{X})(d_N(NS, t))$  denote the deficiency of the normal scores test with respect to the  $\bar{X}$ -test (t-test) based on  $N$  observations. Now we have from (4.6.4) and (4.6.5) that

$$(6.5.1) \quad d_N(NS, \bar{X}) = \tilde{d}_N(NS, \bar{X}) + o(1), \quad d_N(NS, t) = \tilde{d}_N(NS, t) + o(1),$$

where

$$(6.5.2) \quad \tilde{d}_N(NS, \bar{X}) = \tilde{d}_N(NS, t) + \frac{1}{2}u_\alpha^2, \quad \tilde{d}_N(NS, t) = \sum_{j=1}^N \sigma^2 \phi^{-1}\left(\frac{1+U_{j:N}}{2}\right) - \frac{1}{2}.$$

The remaining part of this section is devoted to the comparison of  $d_N(NS, \bar{X})$  with  $\tilde{d}_N(NS, \bar{X})$  and of  $d_N(NS, t)$  with  $\tilde{d}_N(NS, t)$ . For the approximations  $\tilde{d}_N$  we can use the values of  $\sum_{j=1}^N \sigma^2 \phi^{-1}\{(1+U_{j:N})/2\}$ , given in the first column of table 6.2.1. The exact values are obtained as follows: if the power of the normal scores test for a certain sample size  $N$  is available, we determine the sample size  $k_N$  for which the  $\bar{X}$ -test (or the t-test) reaches the same power, and this gives  $d_{k_N} = N - k_N$ . Here the role of  $N$  and  $k_N$  has been interchanged because of the fact that for the normal scores test only a limited number of exact power values is available, whereas the exact power of the other two tests can be obtained rather easily for any sample size. For convenience we compare  $d_{k_N}$  with  $\tilde{d}_N$  rather than with  $\tilde{d}_{k_N}$ . As  $d_{k_N} - \tilde{d}_{k_N} = o(1)$  and  $\tilde{d}_N - \tilde{d}_{k_N} = O(N^{-1} \log \log N)$ , this modification seems rather harmless, which



impression is confirmed by the numerical results.

In the above  $k_N$  is treated as a continuous variable, which is interpreted as follows: for non-integer  $k_N$ , we select sample size  $[k_N]$  or  $[k_N]+1$  with probability  $1-k_N+[k_N]$  and  $k_N-[k_N]$  respectively. Here  $[y]$  means the integer part of  $y$ . This yields an expected sample size  $k_N$  and an expected power  $(1-k_N+[k_N])\pi_{[k_N]} + (k_N-[k_N])\pi_{[k_N]+1}$ .

The first series of comparisons is based on the exact values of the power of the normal scores test, obtained by Klotz (1963) (cf. section 6.2). The power of the  $\bar{X}$ -test is  $1-\Phi(u_\alpha - N^{1/2}\theta)$  and hence  $d_{k_N}(NS, \bar{X})$  can be evaluated in a straightforward manner. Klotz also tabulated the exact efficiency  $e_{k_N}(NS, t) = k_N/N$  of the normal scores test with respect to the t-test. Now  $d_{k_N}(NS, t)$  immediately follows from the relation  $d_{k_N}(NS, t) = N\{1 - e_{k_N}(NS, t)\}$ . The results are collected in table 6.5.1. (p.137). Note that  $\tilde{d}_N(NS, \bar{X})$  only depends on  $N$  and  $\alpha$  and that  $\tilde{d}_N(NS, t)$  only depends on  $N$ . Here and in the sequel, we use for  $\sum_{j=1}^N \sigma^2 \Phi^{-1}\{(1+U_{j:N})/2\}$  the numerically obtained values from table 6.2.1. The agreement of the exact and asymptotic results appears to be satisfactory already at these small sample sizes.

The results of Thompson et al. (1967) from section 6.2 for the power of the normal scores test against normal alternatives, can also be used for deficiency computations. Here we deal with Monte Carlo estimates instead of with exact values. This leads to values of  $d_{k_N}(NS, \bar{X})$  and  $d_{k_N}(NS, t)$  which are also subject to error. As in general  $d_{k_N} = N - k_N$  is much smaller than  $N$ , the relative error in  $d_{k_N}$  will be much larger than the relative error in the power estimates. To give an impression of the reliability of the obtained values of  $d_N$ , we also evaluate the values of  $d_N$  for the power estimates plus or minus their standard deviation. We only use the power values for  $N = 20$ : for  $N = 10$  we already have the exact results of Klotz, which are much more informative.

The necessary power values of the  $\bar{X}$ -test are again immediately given by  $1-\Phi(u_\alpha - N^{1/2}\theta)$ . For the power values of the t-test we proceed in the following way: the critical values involved are found from Owen (1962). Furthermore, Resnikoff and Lieberman (1957) have tabulated the non-central t-distribution function for various degrees of freedom  $f$  and various non-centrality parameters  $\delta$ . These  $\delta$ 's are of the form  $(f+1)^{1/2}u_\alpha$ , whereas we need  $(f+1)^{1/2}k/4$ , for  $k = 1, \dots, 6$ . Hence the necessary power values cannot be found directly from these tables. Using the description of the method of

computation that is contained in the introduction to the tables, a program was written to obtain the power values for the values of  $\delta$  considered here. According to Resnikoff and Lieberman the accuracy is four decimal places, which amply suffices for our purposes. The deficiency results are collected in table 6.5.2. (p.138). They are not very conclusive, as the estimates of the exact deficiency appear to be very crude, but again it seems that the asymptotic results to reasonable predictions.

For sample sizes larger than 20, no results about the power of the normal scores test are available in literature. Yet it seems desirable to have some idea about the agreement between the  $d_{k_N}$  and  $\tilde{d}_N$  for such sample sizes. Therefore, we use the simulation method described by Thompson et al. (1967) to obtain estimates for the power of the normal scores test for sample sizes larger than 20.

In section 6.2 we already mentioned that this method involves conducting  $\tilde{N}$  simulations, each consisting of drawing a random sample of size  $N$  from the standard normal distribution, shifting it over  $\theta$ , computing  $T = \sum_{j=1}^N E\Phi^{-1}\{(1+U_{j:N})/2\}V_j$  and counting the number of samples for which  $T$  exceeds the critical value  $c_\alpha$ . Here we supply some more details. In the first place, we restrict attention to the case where  $\tilde{N} = 1600$  and  $N = 50$ . For this value of  $N$ , the scores  $E\Phi^{-1}\{(1+U_{j:N})/2\}$  can be found in tables by Govindarajulu and Eisenstat (1964). As these values are exact to five decimal places, their contribution to the error in the power estimates can be neglected.

For  $N$  as large as 50 it is impracticable to evaluate the exact critical values  $c_\alpha$  and we have to use Edgeworth expansions, as advised and tabulated by Thompson et al. Denote these approximate values as  $c'_\alpha$ . The portion of the error in the power estimate, due to the use of  $c'_\alpha$ , may be estimated as follows: according to Thompson et al. the use of  $c'_\alpha$  instead of  $c_\alpha$  causes an error of at most 2% in the test size  $\alpha$ , i.e.

$$(6.5.3) \quad |P_0(T > c'_\alpha) - P_0(T > c_\alpha)| \leq 0.02\alpha.$$

The distribution of  $(T - E_0 T)/\sigma_0(T)$  is asymptotically standard normal and therefore  $(c_\alpha - E_0 T)/\sigma_0(T) \approx u_\alpha$ . Hence

$$(6.5.4) \quad P_0(T > c'_\alpha) - P_0(T > c_\alpha) \approx \Phi\left(\frac{c_\alpha - E_0 T}{\sigma_0(T)}\right) - \Phi\left(\frac{c'_\alpha - E_0 T}{\sigma_0(T)}\right) \approx \frac{c_\alpha - c'_\alpha}{\sigma_0(T)} \phi(u_\alpha).$$

Denote the exact power  $P_\theta(T > c_\alpha)$  as  $\pi$ , the approximate power  $P_\theta(T > c'_\alpha)$  as  $\pi'$  and let  $u_\pi = \Phi^{-1}(1-\pi)$ . In analogy to (6.5.4) we have

$$(6.5.5) \quad \pi' - \pi \approx \frac{c_\alpha - c'_\alpha}{\sigma_\theta(T)} \phi(u_\pi).$$

As  $\sigma_\theta^2(T) \approx \sigma_0^2(T) = N/4$ , (6.5.3)-(6.5.5) lead to the following upper bound for the error that is caused by the use of inexact critical values

$$(6.5.6) \quad |\pi' - \pi| \leq 0.02\alpha \frac{\phi(u_\pi)}{\phi(u_\alpha)}.$$

The main source of error remains of course the fact that we use simulation methods to find  $\pi$ . For  $i = 1, \dots, \tilde{N}$ , define the r.v.'s  $\hat{\pi}_i$  by

$$\hat{\pi}_i = \begin{cases} 1, & T > c'_\alpha \text{ for the } i^{\text{th}} \text{ sample,} \\ 0, & \text{otherwise.} \end{cases}$$

Moreover, define  $\hat{\pi} = \frac{1}{\tilde{N}} \sum_{i=1}^{\tilde{N}} \hat{\pi}_i$ .  $\hat{\pi}$  clearly is an unbiased estimate of  $\pi'$ . If all samples are drawn independently, we have as an unbiased estimate for its variance

$$(6.5.7) \quad \hat{\sigma}^2(\hat{\pi}) = \frac{\hat{\pi}(1-\hat{\pi})}{\tilde{N}}.$$

The variance of  $\hat{\pi}$  can be reduced by using the antithetic method that is also applied by Thompson et al.. Here we independently draw  $\tilde{N}/2$  samples  $X_1, \dots, X_{\tilde{N}/2}$  from  $\Phi(x-\theta)$  and form the other  $\tilde{N}/2$  samples by taking  $2\theta - X_1, \dots, 2\theta - X_{\tilde{N}/2}$  for each of the first  $\tilde{N}/2$  samples. Note that if  $X_1$  has d.f.  $\Phi(x-\theta)$ , this is also the case for  $2\theta - X_1$ . Now the  $\hat{\pi}_i$  form  $\tilde{N}/2$  pairs of dependent r.v.'s. Each pair has the same covariance, which we denote as  $\text{Covar}$ , and the pairs are mutually independent. The variance of  $\hat{\pi}$  is  $\{\pi'(1-\pi') + \text{Covar}\}/\tilde{N}$ . From the construction of the pairs it is clear that  $\text{Covar} < 0$  and therefore this method decreases the variance of  $\hat{\pi}$ . Computation of  $\text{Covar}$  is too complicated, but the estimated values of  $\text{Covar}$  that are obtained from our numerical results, indicate that the reduction is rather small. In view of this, the conjecture of Thompson et al. that  $\hat{\sigma}^2(\hat{\pi})$  becomes about ten times smaller seems far too optimistic. The main advantage of the antithetic method lies here in the reduction of the time, needed to form the  $\tilde{N}$  samples.

The expressions in (6.5.2) for  $\tilde{d}_N(NS, \bar{X})$  and  $\tilde{d}_N(NS, t)$  suggest that the normal scores test is only slightly worse than the  $\bar{X}$ -test and the t-test. Hence we may expect the critical regions of the three tests to be much the same. As the power of the  $\bar{X}$ -test and the t-test can be obtained with high accuracy for all  $N$ ,  $\alpha$  and  $\theta$ , the above resemblance enables us to achieve a further reduction of  $\sigma^2(\hat{\pi})$ . For each of the  $\tilde{N}$  samples that is drawn, we not only note whether  $T > c'_\alpha$ , but also whether the  $\bar{X}$ -test and the t-test exceed their respective critical values. To be precise, we define for  $i, j, k = 0, 1$  the r.v.'s  $\hat{\pi}_{ijk}$  as  $\hat{\pi}_{ijk} = \tilde{N}^{-1} \times$  the number of samples for which the  $\bar{X}$ -test does (does not) reject the hypothesis if  $i = 0(1)$ , for which the t-test does (does not) reject the hypothesis if  $j = 0(1)$  and for which the normal scores test does (does not) reject the hypothesis if  $k = 0(1)$ . Moreover, let  $\hat{\pi}_{ij.} = \hat{\pi}_{ij0} + \hat{\pi}_{ij1}$ , and define  $\hat{\pi}_{i.k}$  and  $\hat{\pi}_{.jk}$  analogously. Then we have

$$\hat{\pi} = \hat{\pi}_{000} + \hat{\pi}_{100} + \hat{\pi}_{010} + \hat{\pi}_{110} = \hat{\pi}_t + \hat{\pi}_{.10} - \hat{\pi}_{.01} =$$

(6.5.8)

$$= \hat{\pi}_{\bar{X}} + \hat{\pi}_{1.0} - \hat{\pi}_{0.1},$$

where  $\hat{\pi}_t$  and  $\hat{\pi}_{\bar{X}}$  are unbiased estimates of the power  $\pi_t$  of the t-test and the power  $\pi_{\bar{X}}$  of the  $\bar{X}$ -test, respectively. Since  $\pi_t$  and  $\pi_{\bar{X}}$  can be obtained exactly, we can improve on  $\hat{\pi}$  by considering the following two estimates

$$\hat{\pi}_1 = \pi_t + \hat{\pi}_{.10} - \hat{\pi}_{.01} = \hat{\pi} + (\pi_t - \hat{\pi}_t),$$

(6.5.9)

$$\hat{\pi}_2 = \pi_{\bar{X}} + \hat{\pi}_{1.0} - \hat{\pi}_{0.1} = \hat{\pi} + (\pi_{\bar{X}} - \hat{\pi}_{\bar{X}}),$$

Note that  $\hat{\pi}_1$  and  $\hat{\pi}_2$  are also unbiased estimates of  $\pi$ . From the close resemblance of the critical regions of the three tests it follows that  $\hat{\pi}_{.10}$ ,  $\hat{\pi}_{.01}$ ,  $\hat{\pi}_{1.0}$  and  $\hat{\pi}_{0.1}$  estimate very small probabilities. Hence, their variances are also small. In view of this and of (6.5.9), we may expect that  $\sigma^2(\hat{\pi}_1)$  and  $\sigma^2(\hat{\pi}_2)$  are considerably smaller than  $\sigma^2(\hat{\pi})$  and hence this approach yields another reduction of  $\sigma^2(\hat{\pi})$ . Unbiased estimates for the variances of  $\hat{\pi}_1$  and  $\hat{\pi}_2$  are

$$\hat{\sigma}^2(\hat{\pi}_1) = \tilde{N}^{-1} \{ \hat{\pi}_{.10}(1-\hat{\pi}_{.10}) + \hat{\pi}_{.01}(1-\hat{\pi}_{.01}) + 2\hat{\pi}_{.10}\hat{\pi}_{.01} \},$$

(6.5.10)

$$\hat{\sigma}^2(\hat{\pi}_2) = \tilde{N}^{-1} \{ \hat{\pi}_{1.0}(1-\hat{\pi}_{1.0}) + \hat{\pi}_{0.1}(1-\hat{\pi}_{0.1}) + 2\hat{\pi}_{1.0}\hat{\pi}_{0.1} \}.$$

As our ultimate estimate we use  $\hat{\pi}_3 = (\hat{\pi}_1 + \hat{\pi}_2)/2$ . An unbiased estimate for  $\sigma^2(\hat{\pi}_3)$  is

$$\hat{\sigma}^2(\hat{\pi}_3) = \{ \hat{\sigma}^2(\hat{\pi}_1) + \hat{\sigma}^2(\hat{\pi}_2) \} / 4 + \{ \hat{\pi}_{110} - \hat{\pi}_{.10}\hat{\pi}_{1.0} + \hat{\pi}_{001} +$$

(6.5.11)

$$- \hat{\pi}_{.01}\hat{\pi}_{0.1} + \hat{\pi}_{.10}\hat{\pi}_{0.1} + \hat{\pi}_{1.0}\hat{\pi}_{.01} \} / (2\tilde{N}).$$

Together with (6.5.5) this leads to the following estimated standard deviation of the obtained power estimates.

$$(6.5.12) \quad \left\{ \left[ 0.02\alpha \frac{\phi(u_\pi)}{\phi(u_\alpha)} \right]^2 + \hat{\sigma}^2(\hat{\pi}_3) \right\}^{1/2}.$$

The numerical results are given in tables 6.5.3 and 6.5.4 (p.139). We also give the values of  $d_N$  that are obtained if we use the power estimates plus or minus their standard deviation. Again the agreement between finite and asymptotic results is satisfactory.

The general conclusions of this section are that the asymptotic results seem to provide a reasonable approximation of the exact values and that the normal scores test requires only very few additional observations to attain the same power against normal alternatives as the  $\bar{X}$ -test or the t-test.

Table 6.2.2

Power of the normal scores test against normal alternatives. The upper, middle and lower numbers give the exact values obtained by Klotz, the approximations (6.2.1), and the normal approximations, respectively.

N	$\theta$ $\alpha$	1/4	1/2	3/4	1	5/4	3/2
5	.06250	.145	.278	.450	.629	.781	.888
		.146	.284	.465	.654	.810	.913
		.165	.339	.557	.759	.896	.966
6	.04688	.125	.263	.450	.646	.807	.911
		.126	.268	.465	.670	.834	.933
		.144	.326	.564	.780	.917	.977
7	.05469	.155	.332	.556	.760	.897	.966
		.156	.336	.566	.775	.910	.973
		.174	.390	.649	.852	.956	.991
8	.05469	.167	.369	.614	.818	.936	.983
		.168	.373	.623	.829	.944	.987
		.186	.426	.698	.890	.973	.996
8	.07422	.212	.438	.684	.866	.958	.991
		.213	.441	.692	.875	.964	.993
		.230	.488	.750	.917	.982	.997
9	.02734	.104	.271	.515	.750	.904	.973
		.104	.274	.524	.765	.917	.979
		.121	.337	.629	.860	.966	.995
9	.03711	.132	.326	.581	.803	.932	.983
		.133	.329	.590	.816	.942	.988
		.150	.388	.679	.888	.975	.997
9	.04883	.164	.380	.642	.847	.953	.990
		.165	.384	.650	.857	.960	.992
		.182	.438	.724	.910	.982	.998
10	.00098	.006	.025	.077	.178	.327	.501
		.006	.027	.092	.229	.439	.669
		.011	.064	.234	.526	.804	.950
10	.00977	.048	.158	.362	.612	.820	.937
		.048	.160	.373	.636	.845	.954
		.061	.225	.514	.796	.947	.992
10	.02441	.102	.283	.544	.785	.928	.983
		.102	.285	.553	.798	.938	.987
		.119	.348	.656	.883	.976	.997
10	.05273	.186	.431	.706	.894	.974	.996
		.186	.434	.712	.901	.978	.997
		.204	.485	.774	.939	.990	.999
10	.09668	.288	.572	.820	.950	.991	.999
		.289	.575	.825	.953	.992	.999
		.305	.610	.858	.969	.996	1.000

Table 6.2.3

Power of the Wilcoxon test against normal alternatives. The upper, middle and lower numbers give the exact values obtained by Klotz, the approximations (6.2.4), and the normal approximations, respectively.

N	$\theta$ $\alpha$	1/4	1/2	3/4	1	5/4	3/2
5	.06250	.145	.278	.450	.629	.781	.888
		.146	.283	.462	.650	.805	.909
		.162	.329	.542	.742	.884	.959
6	.04688	.125	.263	.450	.646	.807	.911
		.126	.268	.464	.669	.832	.931
		.141	.316	.547	.763	.906	.972
7	.05469	.155	.332	.556	.760	.897	.966
		.155	.334	.563	.771	.907	.972
		.170	.379	.632	.837	.948	.989
8	.05469	.167	.369	.614	.818	.936	.983
		.167	.370	.619	.825	.941	.986
		.181	.413	.681	.878	.968	.995
8	.07422	.212	.436	.683	.865	.958	.991
		.211	.437	.686	.870	.961	.992
		.225	.475	.735	.906	.978	.997
9	.02734	.104	.271	.515	.750	.904	.973
		.104	.275	.525	.765	.916	.979
		.117	.324	.609	.844	.959	.993
9	.03711	.132	.326	.581	.804	.933	.984
		.132	.328	.589	.814	.940	.987
		.146	.375	.660	.874	.970	.995
9	.04883	.164	.379	.640	.846	.953	.990
		.164	.381	.646	.853	.957	.992
		.178	.424	.706	.899	.978	.997
10	.00098	.006	.025	.077	.178	.327	.501
		.006	.031	.105	.262	.489	.719
		.010	.060	.218	.497	.778	.938
10	.00977	.048	.158	.362	.612	.820	.937
		.049	.164	.382	.646	.853	.957
		.059	.215	.493	.775	.937	.989
10	.02441	.102	.282	.544	.785	.928	.983
		.102	.286	.554	.798	.937	.987
		.115	.335	.636	.869	.971	.996
10	.05273	.184	.427	.701	.891	.973	.996
		.185	.430	.707	.897	.976	.996
		.199	.470	.758	.929	.988	.999
10	.09668	.286	.567	.815	.948	.990	.999
		.286	.568	.817	.949	.991	.999
		.299	.596	.845	.963	.995	1.000

Table 6.2.4

Power of the normal scores test against normal alternatives. The upper, middle and lower numbers give the Monte Carlo powers obtained by Thompson et al., the approximations (6.2.1), and the normal approximations, respectively.

N	$\theta$	1/4	1/2	3/4	1	3/2
	$\alpha$					
10	.01	.041	.160	.365	.622	.954
		.049	.163	.377	.640	.956
		.062	.228	.518	.798	.992
10	.025	.086	.285	.554	.794	.990
		.104	.289	.559	.802	.988
		.121	.352	.660	.885	.997
10	.05	.178	.425	.697	.896	.998
		.179	.423	.702	.895	.997
		.196	.475	.766	.935	.999
20	.01	.112	.406	.781	.955	-
		.098	.396	.778	.964	1.000
		.113	.464	.848	.984	1.000
20	.025	.199	.551	.874	.988	-
		.182	.558	.885	.988	1.000
		.200	.609	.918	.994	1.000
20	.05	.296	.683	.940	.995	-
		.282	.689	.941	.996	1.000
		.299	.723	.956	.998	1.000



Table 6.2.5

Power of the Wilcoxon test against normal alternatives. The upper, middle and lower numbers give the Monte Carlo powers obtained by Thompson et al., the approximations (6.2.4), and the normal approximations, respectively.

N	$\alpha$	$\theta$				
		1/4	1/2	3/4	1	3/2
10	.01	.041	.158	.366	.621	.952
		.050	.166	.386	.650	.958
		.060	.217	.497	.778	.990
10	.025	.084	.288	.556	.793	.990
		.104	.290	.559	.802	.987
		.118	.339	.640	.871	.996
10	.05	.171	.412	.690	.893	.998
		.178	.419	.697	.891	.996
		.192	.460	.749	.926	.999
20	.01	.106	.396	.768	.953	-
		.097	.392	.774	.962	1.000
		.109	.444	.829	.980	1.000
20	.025	.185	.520	.868	.982	-
		.180	.550	.879	.986	1.000
		.193	.589	.906	.992	1.000
20	.05	.276	.650	.924	.993	-
		.278	.679	.936	.995	1.000
		.290	.705	.949	.997	1.000

Table 6.2.6

Power of the normal scores test against logistic alternatives. The upper, middle and lower numbers give the Monte Carlo powers obtained by Thompson et al., the approximations (6.2.3), and the normal approximations, respectively.

N	$\theta$	1/4	1/2	3/4	1	3/2
	$\alpha$					
10	.01	.029	.063	.133	.239	.486
		.028	.066	.131	.226	.467
		.030	.076	.162	.294	.637
10	.025	.056	.144	.244	.373	.632
		.062	.130	.233	.364	.636
		.065	.143	.267	.430	.763
10	.05	.122	.220	.353	.507	.755
		.112	.212	.346	.498	.762
		.115	.226	.380	.555	.849
20	.01	.046	.144	.301	.510	.849
		.043	.131	.293	.504	.850
		.045	.143	.332	.578	.928
20	.025	.103	.238	.458	.653	.929
		.089	.229	.437	.657	.927
		.092	.242	.473	.713	.966
20	.05	.166	.358	.560	.763	.963
		.152	.337	.568	.771	.965
		.155	.351	.598	.810	.984

Table 6.2.7

Power of the Wilcoxon test against logistic alternatives. The upper, middle and lower numbers give the Monte Carlo powers obtained by Thompson et al., the approximations (6.2.2), and the normal approximations, respectively.

N	$\theta$	1/4	1/2	3/4	1	3/2
	$\alpha$					
10	.01	.028	.061	.131	.240	.486
		.028	.065	.132	.231	.489
		.031	.079	.169	.308	.660
10	.025	.055	.139	.243	.373	.637
		.062	.131	.237	.374	.662
		.066	.148	.277	.447	.782
10	.05	.121	.230	.356	.516	.770
		.112	.215	.355	.512	.785
		.117	.232	.391	.572	.863
20	.01	.044	.133	.301	.520	.860
		.043	.134	.302	.522	.872
		.046	.150	.348	.601	.939
20	.025	.093	.232	.447	.648	.926
		.090	.234	.450	.677	.941
		.094	.252	.491	.733	.972
20	.05	.152	.338	.557	.762	.964
		.154	.345	.583	.789	.973
		.159	.362	.615	.826	.987

Table 6.4.1

Power of the sign test against normal alternatives. The upper, middle and lower values give the Monte Carlo powers obtained by Thompson et al., the approximations (6.4.1), and the normal approximations, respectively.

N	$\theta$	1/4	1/2	3/4	1	3/2
	$\alpha$					
10	.01	.042	.128	.284	.487	.832
		.041	.123	.270	.455	.741
		.044	.132	.275	.433	.660
10	.025	.071	.207	.386	.591	.921
		.085	.213	.403	.601	.846
		.091	.227	.409	.579	.782
10	.05	.148	.328	.555	.761	.967
		.152	.334	.558	.749	.928
		.154	.332	.534	.696	.863
20	.01	.070	.263	.558	.823	-
		.071	.264	.567	.815	.978
		.074	.270	.547	.766	.939
20	.025	.137	.403	.719	.915	-
		.138	.408	.721	.908	.993
		.141	.402	.686	.863	.972
20	.05	.211	.524	.826	.961	-
		.220	.535	.818	.951	.997
		.223	.527	.788	.920	.987

Table 6.4.2

Power of the sign test against logistic alternatives. The upper, middle and lower values give the Monte Carlo powers obtained by Thompson et al., the approximations (6.4.1), and the normal approximations, respectively.

N	$\theta$	1/4	1/2	3/4	1	3/2
	$\alpha$					
10	.01	.026	.058	.112	.192	.409
		.025	.056	.108	.184	.385
		.027	.060	.116	.194	.375
10	.025	.049	.110	.177	.288	.533
		.056	.110	.191	.296	.530
		.059	.118	.204	.309	.519
10	.05	.095	.191	.298	.449	.691
		.104	.190	.305	.437	.684
		.105	.192	.304	.427	.642
20	.01	.031	.100	.219	.400	.739
		.037	.104	.228	.399	.737
		.038	.109	.235	.398	.696
20	.025	.079	.189	.351	.552	.875
		.079	.191	.364	.561	.856
		.080	.194	.360	.542	.811
20	.05	.129	.290	.480	.678	.936
		.137	.289	.488	.683	.918
		.138	.291	.483	.663	.884

Table 6.4.3

Power of the sign test against double exponential alternatives. The upper, middle and lower numbers give the Monte Carlo powers obtained by Thompson et al., the approximations (6.4.1), and the normal approximations, respectively.

N	$\theta$	1/4	1/2	3/4	1	3/2
	$\alpha$					
10	.01	.050	.135	.261	.404	.662
		.048	.130	.249	.381	.604
		.052	.140	.255	.372	.552
10	.025	.097	.226	.381	.510	.773
		.097	.223	.377	.525	.737
		.104	.237	.385	.516	.690
10	.05	.176	.369	.535	.691	.890
		.170	.347	.530	.680	.856
		.172	.344	.509	.638	.792
20	.01	.072	.256	.501	.712	.951
		.086	.280	.528	.732	.926
		.091	.285	.513	.692	.874
20	.025	.155	.424	.670	.858	.979
		.163	.428	.686	.852	.971
		.166	.421	.655	.807	.935
20	.05	.243	.547	.779	.914	.992
		.254	.555	.791	.915	.987
		.256	.546	.763	.881	.966

Table 6.5.1

Deficiencies under normal alternatives of the normal scores test with respect to the  $\bar{X}$ -test and the t-test. The upper and lower numbers give  $d_{k_N}(NS, \bar{X})$  and  $d_{k_N}(NS, t)$ , respectively.

N	$\theta$ $\alpha$	1/4	1/2	3/4	1	5/4	3/2	$\tilde{d}_N(NS, \bar{X})$	$\tilde{d}_N(NS, t)$
5	0.06250	1.364 0.070	1.413 0.080	1.463 0.090	1.501 0.095	1.536 0.100	1.566 0.105	1.370	0.193
6	0.04688	1.582 0.102	1.650 0.120	1.711 0.138	1.771 0.150	1.835 0.162	1.909 0.168	1.629	0.224
7	0.05469	1.480 0.119	1.541 0.133	1.595 0.154	1.642 0.168	1.690 0.175	1.744 0.182	1.534	0.252
8	0.05469	1.520 0.160	1.577 0.184	1.630 0.200	1.677 0.216	1.726 0.224	1.782 0.232	1.559	0.277
8	0.07422	1.310 0.176	1.358 0.200	1.401 0.216	1.440 0.232	1.479 0.240	1.521 0.256	1.321	0.277
9	0.02734	2.036 0.162	2.104 0.180	2.175 0.207	2.240 0.225	2.300 0.243	2.352 0.252	2.141	0.294
9	0.03711	1.818 0.180	1.886 0.198	1.954 0.225	2.031 0.234	2.098 0.261	2.160 0.279	1.888	0.294
9	0.04883	1.623 0.180	1.682 0.198	1.739 0.234	1.796 0.243	1.865 0.261	1.945 0.261	1.666	0.294
10	0.00098	4.629 0.710	4.835 0.800	5.061 0.920	5.293 1.040	5.510 1.180	5.719 1.310	5.104	0.310
10	0.00977	2.804 0.190	2.912 0.210	3.018 0.240	3.110 0.260	3.218 0.280	3.309 0.290	3.037	0.310
10	0.02441	2.125 0.170	2.211 0.200	2.287 0.230	2.357 0.250	2.423 0.280	2.494 0.290	2.252	0.310
10	0.05273	1.574 0.210	1.638 0.230	1.696 0.250	1.753 0.270	1.820 0.290	1.875 0.310	1.621	0.310
10	0.09668	1.171 0.250	1.216 0.280	1.254 0.300	1.294 0.320	1.327 0.340	1.333 -	1.156	0.310

Table 6.5.2

Deficiencies under normal alternatives of the normal scores test with respect to the  $\bar{X}$ -test and the t-test. The upper numbers give  $d_{k_N}(NS, \bar{X})$  if one uses the power estimate plus its standard deviation, the power estimate itself and the power estimate minus its standard deviation, respectively. The lower numbers similarly give  $d_{k_N}(NS, t)$ .

N	$\theta$	1/4	1/2	3/4	$\tilde{d}_N(NS, \bar{X})$	$\tilde{d}_N(NS, t)$
	$\alpha$					
20	0.01	-2.8, 0.2, 3.2	1.9, 2.5, 3.2	2.2, 2.9, 3.4	3.118	0.411
		-5.3, -2.4, 0.6	-0.9, -0.2, 0.5	-0.5, 0.1, 0.7		
20	0.025	-2.0, 0.0, 2.1	1.8, 2.5, 3.2	1.9, 2.8, 3.8	2.333	0.411
		-3.9, -1.9, 0.2	-0.1, 0.6, 1.2	-0.1, 0.8, 1.6		
20	0.05	-1.4, 0.3, 1.9	1.2, 2.0, 2.7	0.0, 1.7, 3.2	1.764	0.411
		-2.7, -1.0, 0.6	-0.2, 0.6, 1.3	-1.5, 0.4, 1.7		

Table 6.5.3

Power of the normal scores test against normal alternatives. The first number gives the power estimate, the second gives its standard deviation.

N	$\theta$	0.40		$\frac{3}{10} \sqrt{2}$	
	$\alpha$				
50	0.05	0.871	0.004	0.907	0.004
50	0.025	0.781	0.005	0.833	0.005
50	0.01	0.660	0.006	0.714	0.006



Table 6.5.4

Deficiencies under normal alternatives of the normal scores test with respect to the  $\bar{X}$ -test and the t-test. The organization of this table is the same as for table 6.5.2.

N	$\theta$ $\alpha$	0.40	$\frac{3}{10} \sqrt{2}$	$\tilde{d}_N(NS, \bar{X})$	$\tilde{d}_N(NS, t)$
50	0.01	2.6, 3.1, 3.7 -0.2, 0.4, 0.9	3.0, 3.6, 4.1 0.3, 0.8, 1.2	3.23	0.52
50	0.025	2.5, 3.2, 3.7 0.7, 1.3, 1.9	1.7, 2.4, 3.0 -0.1, 0.5, 1.1	2.45	0.52
50	0.05	1.1, 1.8, 2.5 -0.2, 0.5, 1.2	0.3, 1.0, 1.8 -0.9, -0.2, 0.5	1.87	0.52

- [1] ALBERS, W., BICKEL, P.J. and VAN ZWET, W.R. (1974). *Asymptotic expansions for the power of distributionfree tests in the one-sample problem*. Mathematical Centre Report SW 30/74, Amsterdam. Appeared in *Ann. Statist.* 4, 108-156 (1976).
- [2] ARNOLD, H.J. (1965). Small sample power of the one sample Wilcoxon test for non-normal shift alternatives. *Ann. Math. Statist.* 36, 1767-1778.
- [3] BAHADUR, R.R. (1964). On Fisher's bound for asymptotic variances. *Ann. Math. Statist.* 35, 1545-1552.
- [4] BELL, C.B. and DOKSUM, K.A. (1965). Some new distribution-free statistics. *Ann. Math. Statist.* 36, 203-214.
- [5] BHATTACHARYYA, A. (1946). On some analogues of the amount of information and their use in statistical information. *Sankhyā* A8, 1-14.
- [6] BICKEL, P.J. (1974). Edgeworth expansions in nonparametric statistics. *Ann. Statist.* 2, 1-21.
- [7] BICKEL, P.J. and VAN ZWET, W.R. (1974). Asymptotic expansions for the power of distributionfree tests in the two-sample problem. Appeared in *Ann. Statist.* 6, 937-1004 (1978).
- [8] CHERNOFF, H. (1956). Large sample theory: parametric case. *Ann. Math. Statist.* 27, 1-22.
- [9] CHUNG, K.L. (1951). The strong law of large numbers. *Proceedings of the second Berkeley Symposium on mathematical statistics and probability*, 341-352.
- [10] ČIBIŠOV, D.M. (1972). Asymptotic expansions for maximum likelihood estimates. *Teor. Verojatnost. i Primenen* 17, 387-388.
- [11] CRAMÉR, H. (1946). *Mathematical methods of statistics*. Princeton University Press, Princeton.
- [12] DAVIS, R.C. (1951). Note on uniformly best unbiased estimates. *Ann. Math. Statist.* 22, 440-446.
- [13] DIEUDONNÉ, J. (1960). *Foundations of modern analysis*. Academic Press, New York.
- [14] FELLER, W. (1966). *An introduction to probability theory and its applications 2*. Wiley, New York.

- [15] FELLINGHAM, S.A. and STOKER, D.J. (1964). An approximation for the exact distribution of Wilcoxon's test for symmetry. *J. Amer. Statist. Ass.* 59, 899-905.
- [16] FISHER, R.A. (1925). Theory of statistical estimation. *Proc. Cambridge Philos. Soc.* 22, 700-725.
- [17] GOVINDARAJULU, Z. and EISENSTAT, S. (1964). Best estimates of location and scale parameters of a chi (1 d.f.) distribution using ordered observations, Case Institute of Technology, Cleveland, Ohio, Report 206.
- [18] HÁJEK, J. and ŠIDÁK, Z. (1967). *Theory of rank tests*. Academic Press, New York.
- [19] HODGES, J.L., JR. and FIX, E. (1955). Significance probabilities of the Wilcoxon test. *Ann. Math. Statist.* 26, 301-312.
- [20] HODGES, J.L., JR. and LEHMANN, E.L. (1956). Two approximations to the Robbins-Monro process. *Proceedings of the third Berkeley Symposium on mathematical statistics and probability, vol. 1*, 95-105.
- [21] HODGES, J.L., JR. and LEHMANN, E.L. (1963). Estimates of location based on rank tests. *Ann. Math. Statist.* 34, 598-611.
- [22] HODGES, J.L., JR. and LEHMANN, E.L. (1970). Deficiency. *Ann. Math. Statist.* 41, 783-801.
- [23] KLOTZ, J. (1963). Small sample power and efficiency for the one sample Wilcoxon and normal scores tests. *Ann. Math. Statist.* 34, 624-632.
- [24] LEHMANN, E.L. (1959). *Testing statistical hypotheses*. Wiley, New York.
- [25] LINNIK, Yu.V. and MITROFANOVA, N.M. (1965). Some asymptotic expansions for the distribution of the maximum likelihood estimate. *Sankhyā* A27, 73-82.
- [26] MOLENAAR, W. (1970). *Approximations to the Poisson, binomial and hypergeometric distribution functions*. Mathematical Centre, Amsterdam.
- [27] OWEN, D.B. (1962). *Handbook of statistical tables*. Pergamon Press, London.

- [28] RAO, C.R. (1965). *Linear statistical inference and its applications*. Wiley, New York.
- [29] RESNIKOFF, G.J. and LIEBERMAN, G.J. (1957). *Tables of the non-central  $t$ -distribution*. Stanford University Press, Stanford.
- [30] ROGERS, W.F. (1971). Exact null distribution and asymptotic expansions for rank test statistics. Technical Report, Stanford University.
- [31] SOBEL, M. and WEISS, G.H. (1970). Play-the-Winner sampling for selecting the better of two binomial populations. *Biometrika* 57, 357-366.
- [32] SUNDRUM, R. (1954). A further approximation to the distribution of Wilcoxon's statistic. *J.R. Statist. Soc.* B16, 266-268.
- [33] TITCHMARSH, E.C. (1939). *The theory of functions*. Oxford University Press, London.
- [34] THOMPSON, R., GOVINDARAJULU, Z. and DOKSUM, K.A. (1967). Distribution and power of the absolute normal scores test. *J. Amer. Statist. Ass.* 62, 966-975.
- [35] WITTING, H. (1960). A generalized Pitman efficiency. *Ann. Math. Statist.* 31, 405-414.

## INDEX

approximate scores	47
asymptotic deficiency	1
asymptotic expansion	2
asymptotic relative efficiency (ARE)	1
 Bhattacharyya bounds	 112
 Cauchy distribution	 119
contiguous alternatives	44
contiguous location alternatives	40
Cramér-Rao bound	112
cumulant	31
 deficiency	 1
double-exponential distribution	52
 Edgeworth expansion	 31
envelope power	59
exact scores	47
exponential distribution	11
 geometric distribution	 11
 Hodges-Lehmann estimators	 113
 lattice distribution	 8
linear rank test	32
locally most powerful (LMP)	59
locally most powerful (parametric) test	59
locally most powerful permutation test	72
locally most powerful rank test	47
location alternatives	5
logistic alternatives	53

Marcinkievitz-Zygmund-Chung inequality	81
maximum likelihood estimator (MLE)	107
Monte Carlo methods	117
most powerful (parametric) test	59
normal scores test	54
one sample linear rank test	32
one sample normal scores test	54
one sample problem	4
one sample van der Waerden test	55
one sample Wilcoxon test	53
permutation test	71
Poisson distribution	10
randomized rank score (RRS) test	93
scores	32
sign test	55
2x2 table	11
uniformly most powerful unbiased (UMPU) test	9
van der Waerden test	55
Wilcoxon's (signed rank) test	53

## TITLES IN THE SERIES MATHEMATICAL CENTRE TRACTS

(An asterisk before the MCT number indicates that the tract is under preparation).

A leaflet containing an order form and abstracts of all publications mentioned below is available at the Mathematisch Centrum, Kruislaan 413, 1098 SJ Amsterdam, The Netherlands. Orders should be sent to the same address.

- 
- MCT 1 T. VAN DER WALT, *Fixed and almost fixed points*, 1963.  
ISBN 90 6196 002 9.
- MCT 2 A.R. BLOEMENA, *Sampling from a graph*, 1964. ISBN 90 6196 003 7.
- MCT 3 G. DE LEVE, *Generalized Markovian decision processes, part I: Model and method*, 1964. ISBN 90 6196 004 5.
- MCT 4 G. DE LEVE, *Generalized Markovian decision processes, part II: Probabilistic background*, 1964. ISBN 90 6196 005 3.
- MCT 5 G. DE LEVE, H.C. TIJMS & P.J. WEEDA, *Generalized Markovian decision processes, Applications*, 1970. ISBN 90 6196 051 7.
- MCT 6 M.A. MAURICE, *Compact ordered spaces*, 1964. ISBN 90 6196 006 1.
- MCT 7 W.R. VAN ZWET, *Convex transformations of random variables*, 1964.  
ISBN 90 6196 007 X.
- MCT 8 J.A. ZONNEVELD, *Automatic numerical integration*, 1964.  
ISBN 90 6196 008 8.
- MCT 9 P.C. BAAYEN, *Universal morphisms*, 1964. ISBN 90 6196 009 6.
- MCT 10 E.M. DE JAGER, *Applications of distributions in mathematical physics*, 1964. ISBN 90 6196 010 X.
- MCT 11 A.B. PAALMAN-DE MIRANDA, *Topological semigroups*, 1964.  
ISBN 90 6196 011 8.
- MCT 12 J.A.Th.M. VAN BERCKEL, H. BRANDT CORSTIUS, R.J. MOKKEN & A. VAN WIJNGAARDEN, *Formal properties of newspaper Dutch*, 1965.  
ISBN 90 6196 013 4.
- MCT 13 H.A. LAUWERIER, *Asymptotic expansions*, 1966, out of print; replaced by MCT 54.
- MCT 14 H.A. LAUWERIER, *Calculus of variations in mathematical physics*, 1966. ISBN 90 6196 020 7.
- MCT 15 R. DOORNBOS, *Slippage tests*, 1966. ISBN 90 6196 021 5.
- MCT 16 J.W. DE BAKKER, *Formal definition of programming languages with an application to the definition of ALGOL 60*, 1967.  
ISBN 90 6196 022 3.

- MCT 17 R.P. VAN DE RIET, *Formula manipulation in ALGOL 60, part 1*, 1968. ISBN 90 6196 025 8.
- MCT 18 R.P. VAN DE RIET, *Formula manipulation in ALGOL 60, part 2*, 1968. ISBN 90 6196 038 X.
- MCT 19 J. VAN DER SLOT, *Some properties related to compactness*, 1968. ISBN 90 6196 026 6.
- MCT 20 P.J. VAN DER HOUWEN, *Finite difference methods for solving partial differential equations*, 1968. ISBN 90 6196 027 4.
- MCT 21 E. WATTEL, *The compactness operator in set theory and topology*, 1968. ISBN 90 6196 028 2.
- MCT 22 T.J. DEKKER, *ALGOL 60 procedures in numerical algebra, part 1*, 1968. ISBN 90 6196 029 0.
- MCT 23 T.J. DEKKER & W. HOFFMANN, *ALGOL 60 procedures in numerical algebra, part 2*, 1968. ISBN 90 6196 030 4.
- MCT 24 J.W. DE BAKKER, *Recursive procedures*, 1971. ISBN 90 6196 060 6.
- MCT 25 E.R. PAËRL, *Representations of the Lorentz group and projective geometry*, 1969. ISBN 90 6196 039 8.
- MCT 26 EUROPEAN MEETING 1968, *Selected statistical papers, part I*, 1968. ISBN 90 6196 031 2.
- MCT 27 EUROPEAN MEETING 1968, *Selected statistical papers, part II*, 1969. ISBN 90 6196 040 1.
- MCT 28 J. OOSTERHOFF, *Combination of one-sided statistical tests*, 1969. ISBN 90 6196 041 X.
- MCT 29 J. VERHOEFF, *Error detecting decimal codes*, 1969. ISBN 90 6196 042 8.
- MCT 30 H. BRANDT CORSTIUS, *Exercises in computational linguistics*, 1970. ISBN 90 6196 052 5.
- MCT 31 W. MOLENAAR, *Approximations to the Poisson, binomial and hypergeometric distribution functions*, 1970. ISBN 90 6196 053 3.
- MCT 32 L. DE HAAN, *On regular variation and its application to the weak convergence of sample extremes*, 1970. ISBN 90 6196 054 1.
- MCT 33 F.W. STEUTEL, *Preservation of infinite divisibility under mixing and related topics*, 1970. ISBN 90 6196 061 4.
- MCT 34 I. JUHÁSZ, A. VERBEEK & N.S. KROONENBERG, *Cardinal functions in topology*, 1971. ISBN 90 6196 062 2.
- MCT 35 M.H. VAN EMDEN, *An analysis of complexity*, 1971. ISBN 90 6196 063 0.
- MCT 36 J. GRASMAN, *On the birth of boundary layers*, 1971. ISBN 90 6196 064 9.
- MCT 37 J.W. DE BAKKER, G.A. BLAAUW, A.J.W. DULJVESTIJN, E.W. DIJKSTRA, P.J. VAN DER HOUWEN, G.A.M. KAMSTEEG-KEMPER, F.E.J. KRUSEMAN ARETZ, W.L. VAN DER POEL, J.P. SCHAAP-KRUSEMAN, M.V. WILKES & G. ZOUTENDIJK, *MC-25 Informatica Symposium 1971*. ISBN 90 6196 065 7.



- MCT 38 W.A. VERLOREN VAN THEMAAT, *Automatic analysis of Dutch compound words*, 1971. ISBN 90 6196 073 8.
- MCT 39 H. BAVINCK, *Jacobi series and approximation*, 1972. ISBN 90 6196 074 6.
- MCT 40 H.C. TIJMS, *Analysis of (s,S) inventory models*, 1972. ISBN 90 6196 075 4.
- MCT 41 A. VERBEEK, *Superextensions of topological spaces*, 1972. ISBN 90 6196 076 2.
- MCT 42 W. VERVAAT, *Success epochs in Bernoulli trials (with applications in number theory)*, 1972. ISBN 90 6196 077 0.
- MCT 43 F.H. RUYMGAART, *Asymptotic theory of rank tests for independence*, 1973. ISBN 90 6196 081 9.
- MCT 44 H. BART, *Meromorphic operator valued functions*, 1973. ISBN 90 6196 082 7.
- MCT 45 A.A. BALKEMA, *Monotone transformations and limit laws* 1973. ISBN 90 6196 083 5.
- MCT 46 R.P. VAN DE RIET, *ABC ALGOL, A portable language for formula manipulation systems, part 1: The language*, 1973. ISBN 90 6196 084 3.
- MCT 47 R.P. VAN DE RIET, *ABC ALGOL, A portable language for formula manipulation systems, part 2: The compiler*, 1973. ISBN 90 6196 085 1.
- MCT 48 F.E.J. KRUSEMAN ARETZ, P.J.W. TEN HAGEN & H.L. OUDSHOORN, *An ALGOL 60 compiler in ALGOL 60, Text of the MC-compiler for the EL-X8*, 1973. ISBN 90 6196 086 X.
- MCT 49 H. KOK, *Connected orderable spaces*, 1974. ISBN 90 6196 088 6.
- MCT 50 A. VAN WIJNGAARDEN, B.J. MAILLOUX, J.E.L. PECK, C.H.A. KOSTER, M. SINTZOFF, C.H. LINDSEY, L.G.L.T. MEERTENS & R.G. FISHER (eds), *Revised report on the algorithmic language ALGOL 68*, 1976. ISBN 90 6196 089 4.
- MCT 51 A. HORDIJK, *Dynamic programming and Markov potential theory*, 1974. ISBN 90 6196 095 9.
- MCT 52 P.C. BAAYEN (ed.), *Topological structures*, 1974. ISBN 90 6196 096 7.
- MCT 53 M.J. FABER, *Metrizability in generalized ordered spaces*, 1974. ISBN 90 6196 097 5.
- MCT 54 H.A. LAUWERIER, *Asymptotic analysis, part 1*, 1974. ISBN 90 6196 098 3.
- MCT 55 M. HALL JR. & J.H. VAN LINT (eds), *Combinatorics, part 1: Theory of designs, finite geometry and coding theory*, 1974. ISBN 90 6196 099 1.
- MCT 56 M. HALL JR. & J.H. VAN LINT (eds), *Combinatorics, part 2: Graph theory, foundations, partitions and combinatorial geometry*, 1974. ISBN 90 6196 100 9.
- MCT 57 M. HALL JR. & J.H. VAN LINT (eds), *Combinatorics, part 3: Combinatorial group theory*, 1974. ISBN 90 6196 101 7.

- MCT 58 W. ALBERS, *Asymptotic expansions and the deficiency concept in statistics*, 1975. ISBN 90 6196 102 5.
- MCT 59 J.L. MIJNHEER, *Sample path properties of stable processes*, 1975. ISBN 90 6196 107 6.
- MCT 60 F. GÖBEL, *Queueing models involving buffers*, 1975. ISBN 90 6196 108 4.
- \*MCT 61 P. VAN EMDE BOAS, *Abstract resource-bound classes, part 1*, ISBN 90 6196 109 2.
- \*MCT 62 P. VAN EMDE BOAS, *Abstract resource-bound classes, part 2*, ISBN 90 6196 110 6.
- MCT 63 J.W. DE BAKKER (ed.), *Foundations of computer science*, 1975. ISBN 90 6196 111 4.
- MCT 64 W.J. DE SCHIPPER, *Symmetric closed categories*, 1975. ISBN 90 6196 112 2.
- MCT 65 J. DE VRIES, *Topological transformation groups 1 A categorical approach*, 1975. ISBN 90 6196 113 0.
- MCT 66 H.G.J. PIJLS, *Locally convex algebras in spectral theory and eigenfunction expansions*, 1976. ISBN 90 6196 114 9.
- \*MCT 67 H.A. LAUWERIER, *Asymptotic analysis, part 2*, ISBN 90 6196 119 X.
- MCT 68 P.P.N. DE GROEN, *Singularly perturbed differential operators of second order*, 1976. ISBN 90 6196 120 3.
- MCT 69 J.K. LENSTRA, *Sequencing by enumerative methods*, 1977. ISBN 90 6196 125 4.
- MCT 70 W.P. DE ROEVER JR., *Recursive program schemes: Semantics and proof theory*, 1976. ISBN 90 6196 127 0.
- MCT 71 J.A.E.E. VAN NUNEN, *Contracting Markov decision processes*, 1976. ISBN 90 6196 129 7.
- MCT 72 J.K.M. JANSEN, *Simple periodic and nonperiodic Lamé functions and their applications in the theory of conical waveguides*, 1977. ISBN 90 6196 130 0.
- MCT 73 D.M.R. LEIVANT, *Absoluteness of intuitionistic logic*, 1979. ISBN 90 6196 122 X.
- MCT 74 H.J.J. TE RIELE, *A theoretical and computational study of generalized aliquot sequences*, 1976. ISBN 90 6196 131 9.
- MCT 75 A.E. BROUWER, *Treelike spaces and related connected topological spaces*, 1977. ISBN 90 6196 132 7.
- MCT 76 M. REM, *Associations and the closure statement*, 1976. ISBN 90 6196 135 1.
- MCT 77 W.C.M. KALLENBERG, *Asymptotic optimality of likelihood ratio tests in exponential families*, 1977. ISBN 90 6196 134 3.
- MCT 78 E. DE JONGE & A.C.M. VAN ROOIJ, *Introduction to Riesz spaces*, 1977. ISBN 90 6196 133 5.

- MCT 79 M.C.A. VAN ZUIJLEN, *Empirical distributions and rank statistics*, 1977. ISBN 90 6196 145 9.
- MCT 80 P.W. HEMKER, *A numerical study of stiff two-point boundary problems*, 1977. ISBN 90 6196 146 7.
- MCT 81 K.R. APT & J.W. DE BAKKER (eds), *Foundations of computer science II*, part 1, 1976. ISBN 90 6196 140 8.
- MCT 82 K.R. APT & J.W. DE BAKKER (eds), *Foundations of computer science II*, part 2, 1976. ISBN 90 6196 141 6.
- MCT 83 L.S. BENTHEM JUTTING, *Checking Landau's "Grundlagen" in the AUTOMATH system*, 1979. ISBN 90 6196 147 5.
- MCT 84 H.L.L. BUSARD, *The translation of the elements of Euclid from the Arabic into Latin by Hermann of Carinthia (?) books vii-xii*, 1977. ISBN 90 6196 148 3.
- MCT 85 J. VAN MILL, *Supercompactness and Wallman spaces*, 1977. ISBN 90 6196 151 3.
- MCT 86 S.G. VAN DER MEULEN & M. VELDHORST, *Torrix I, A programming system for operations on vectors and matrices over arbitrary fields and of variable size*. 1978. ISBN 90 6196 152 1.
- \*MCT 87 S.G. VAN DER MEULEN & M. VELDHORST, *Torrix II*, ISBN 90 6196 153 X.
- MCT 88 A. SCHRIJVER, *Matroids and linking systems*, 1977. ISBN 90 6196 154 8.
- MCT 89 J.W. DE ROEVER, *Complex Fourier transformation and analytic functionals with unbounded carriers*, 1978. ISBN 90 6196 155 6.
- MCT 90 L.P.J. GROENEWEGEN, *Characterization of optimal strategies in dynamic games*, 1981. ISBN 90 6196 156 4.
- MCT 91 J.M. GEYSEL, *Transcendence in fields of positive characteristic*, 1979. ISBN 90 6196 157 2.
- MCT 92 P.J. WEEDA, *Finite generalized Markov programming*, 1979. ISBN 90 6196 158 0.
- MCT 93 H.C. TIJMS & J. WESSELS (eds), *Markov decision theory*, 1977. ISBN 90 6196 160 2.
- MCT 94 A. BIJLSMA, *Simultaneous approximations in transcendental number theory*, 1978. ISBN 90 6196 162 9.
- MCT 95 K.M. VAN HEE, *Bayesian control of Markov chains*, 1978. ISBN 90 6196 163 7.
- MCT 96 P.M.B. VITÁNYI, *Lindenmayer systems: Structure, languages, and growth functions*, 1980. ISBN 90 6196 164 5.
- \*MCT 97 A. FEDERGRUEN, *Markovian control problems; functional equations and algorithms*, . ISBN 90 6196 165 3.
- MCT 98 R. GEEL, *Singular perturbations of hyperbolic type*, 1978. ISBN 90 6196 166 1.

- MCT 99 J.K. LENSTRA, A.H.G. RINNOOY KAN & P. VAN EMDE BOAS, *Interfaces between computer science and operations research*, 1978. ISBN 90 6196 170 X.
- MCT 100 P.C. BAAYEN, D. VAN DULST & J. OOSTERHOFF (eds), *Proceedings bicentennial congress of the Wiskundig Genootschap, part 1*, 1979. ISBN 90 6196 168 8.
- MCT 101 P.C. BAAYEN, D. VAN DULST & J. OOSTERHOFF (eds), *Proceedings bicentennial congress of the Wiskundig Genootschap, part 2*, 1979. ISBN 90 6196 169 6.
- MCT 102 D. VAN DULST, *Reflexive and superreflexive Banach spaces*, 1978. ISBN 90 6196 171 8.
- MCT 103 K. VAN HARN, *Classifying infinitely divisible distributions by functional equations*, 1978. ISBN 90 6196 172 6.
- MCT 104 J.M. VAN WOUWE, *Go-spaces and generalizations of metrizability*, 1979. ISBN 90 6196 173 4.
- \*MCT 105 R. HELMERS, *Edgeworth expansions for linear combinations of order statistics*, . ISBN 90 6196 174 2.
- MCT 106 A. SCHRIJVER (ed.), *Packing and covering in combinatorics*, 1979. ISBN 90 6196 180 7.
- MCT 107 C. DEN HEIJER, *The numerical solution of nonlinear operator equations by imbedding methods*, 1979. ISBN 90 6196 175 0.
- MCT 108 J.W. DE BAKKER & J. VAN LEEUWEN (eds), *Foundations of computer science III, part 1*, 1979. ISBN 90 6196 176 9.
- MCT 109 J.W. DE BAKKER & J. VAN LEEUWEN (eds), *Foundations of computer science III, part 2*, 1979. ISBN 90 6196 177 7.
- MCT 110 J.C. VAN VLIET, *ALGOL 68 transput, part I: Historical review and discussion of the implementation model*, 1979. ISBN 90 6196 178 5.
- MCT 111 J.C. VAN VLIET, *ALGOL 68 transput, part II: An implementation model*, 1979. ISBN 90 6196 179 3.
- MCT 112 H.C.P. BERBEE, *Random walks with stationary increments and renewal theory*, 1979. ISBN 90 6196 182 3.
- MCT 113 T.A.B. SNIJDERS, *Asymptotic optimality theory for testing problems with restricted alternatives*, 1979. ISBN 90 6196 183 1.
- MCT 114 A.J.E.M. JANSSEN, *Application of the Wigner distribution to harmonic analysis of generalized stochastic processes*, 1979. ISBN 90 6196 184 X.
- MCT 115 P.C. BAAYEN & J. VAN MILL (eds), *Topological Structures II, part 1*, 1979. ISBN 90 6196 185 5.
- MCT 116 P.C. BAAYEN & J. VAN MILL (eds), *Topological Structures II, part 2*, 1979. ISBN 90 6196 186 6.
- \*MCT 117 P.J.M. KALLENBERG, *Branching processes with continuous state space*, 1979. ISBN 90 6196 188 2.

- MCT 118 P. GROENEROOM, *Large deviations and asymptotic efficiencies*, 1980. ISBN 90 6196 190 4.
- MCT 119 F. J. PETERS, *Sparse matrices and substructures, with a novel implementation of finite element algorithms*, 1980. ISBN 90 6196 192 0.
- MCT 120 W.P.M. DE RUYTER, *On the asymptotic analysis of large-scale ocean circulation*, 1980. ISBN 90 6196 192 9.
- MCT 121 W.H. HAEMERS, *Eigenvalue techniques in design and graph theory*, 1980. ISBN 90 6196 194 7.
- MCT 122 J.C.P. BUS, *Numerical solution of systems of nonlinear equations*, 1980. ISBN 90 6196 195 5.
- MCT 123 I. YUHÁSZ, *Cardinal functions in topology - ten years later*, 1980. ISBN 90 6196 196 3.
- MCT 124 R.D. GILL, *Censoring and stochastic integrals*, 1980. ISBN 90 6196 197 1.
- MCT 125 R. EISING, *2-D systems, an algebraic approach*, 1980. ISBN 90 6196 198 X.
- MCT 126 G. VAN DER HOEK, *Reduction methods in nonlinear programming*, 1980. ISBN 90 6196 199 8.
- MCT 127 J.W. KLOP, *Combinatory reduction systems*, 1980. ISBN 90 6196 200 5.
- MCT 128 A.J.J. TALMAN, *Variable dimension fixed point algorithms and triangulations*, 1980. ISBN 90 6196 201 3.
- MCT 129 G. VAN DER LAAN, *Simplicial fixed point algorithms*, 1980. ISBN 90 6196 202 1.
- MCT 130 P.J.W. TEN HAGEN et al., *ILP Intermediate language for pictures*, 1980. ISBN 90 6196 204 8.
- MCT 131 R.J.R. BACK, *Correctness preserving program refinements: Proof theory and applications*, 1980. ISBN 90 6196 207 2.
- MCT 132 H.M. MULDER, *The interval function of a graph*, 1980. ISBN 90 6196 208 0.
- MCT 133 C.A.J. KLAASSEN, *Statistical performance of location estimators*, 1981. ISBN 90 6196 209 9.
- MCT 134 J.C. VAN VLIET & H. WUPPER (eds), *Proceedings international conference on ALGOL 68*, 1981. ISBN 90 6196 210 2.
- MCT 135 J.A.G. GROENENDIJK, T.M.V. JANSSEN & M.J.B. STOKHOF (eds), *Formal methods in the study of language, part I*, 1981. ISBN 90 6196 211 0.
- MCT 136 J.A.G. GROENENDIJK, T.M.V. JANSSEN & M.J.B. STOKHOF (eds), *Formal methods in the study of language, part II*, 1981. ISBN 90 6196 213 7.
- MCT 137 J. TELGEN, *Redundancy and linear programs*, 1981. ISBN 90 6196 215 3.
- MCT 138 H.A. LAUWERIER, *Mathematical models of epidemics*, 1981. ISBN 90 6196 216 1.
- MCT 139 J. VAN DER WAL, *Stochastic dynamic programming, successive approximations and nearly optimal strategies for Markov decision processes and Markov games*, 1980. ISBN 90 6196 218 8.

- MCT 140 J.H. VAN GELDROP, *A mathematical theory of pure exchange economies without the no-critical-point hypothesis*, 1981.  
ISBN 90 6196 219 6.
- MCT 141 G.E. WELTERS, *Abel-Jacobi isogenies for certain types of Fano three-folds*, 1981.  
ISBN 90 6196 227 7.
- MCT 142 H.R. BENNETT & D.J. LUTZER (eds), *Topology and order structures*, part 1, 1981.  
ISBN 90 6196 228 5.
- MCT 143 H. SCHUMACHER, *Dynamic feedback in finite- and infinite-dimensional linear systems*, 1981.  
ISBN 90 6196 229 3.
- MCT 144 P. EIJGENRAAM, *The solution of initial value problems using interval arithmetic. Formulation and analysis of an algorithm*, 1981.  
ISBN 90 6196 230 7.
- MCT 145 A.J. BRENTJES, *Multi-dimensional continued fraction algorithms*, 1981. ISBN 90 6196 231 5.
- MCT 146 C. VAN DER MEE, *Semigroup and factorization methods in transport theory*, 1982. ISBN 90 6196 233 1.

An asterisk before the number means "to appear".