











ALGOL 60 PROCEDURES IN NUMERICAL ALGEBRA

PART 1

T. J. DEKKER



MATHEMATICAL CENTRE TRACTS
22

ALGOL 60 PROCEDURES
IN NUMERICAL ALGEBRA

PART 1

BY

T.J. DEKKER

2nd Edition

MATHEMATISCH CENTRUM AMSTERDAM

PREFACE

We here present a set of ALGOL 60 procedures for solving systems of linear equations, for inverting matrices and for solving linear least-squares problems. The procedures use single-length scalar-product procedures and no iterations are applied for improving the solutions. In the future, we plan to present a corresponding set using double-length scalar-product procedures and applying iterations for improving the solutions, and a set for calculating eigenvalues and eigenvectors of matrices.

The procedures have been tested on an Electrologica X8 computer by means of the "MC ALGOL 60 system for the X8" of the Mathematical Centre, Amsterdam, written by F. E. J. Kruseman Aretz. Only the procedures mca 2000 to 2005 of section 200 are available as EL X8-machine-code procedures. The texts of the procedures have been edited by an ALGOL editing program written by H. L. Oudshoorn, H. N. Glorie and G. C. J. M. Nogarède[12].

In the second edition some minor errors have been corrected. Three of these corrections concern errors in the texts of rnkelm (mca 2110), detbnd (mca 2120) and detsolbnd (mca 2122) which were related to the handling of the row norms in the pivot selection.

ACKNOWLEDGEMENTS

The author is indebted to Prof. Dr. Ir. A. van Wijngaarden and Prof. Dr. F. E. J. Kruseman Aretz for invaluable suggestions and criticisms, to B. J. Mailloux M. Sc. for carefully reading the manuscript and suggesting several improvements, to P. Beertema, F. J. van de Bosch, W. Hoffmann, R. van der Horst, D. P. de Jong, L. Meertens, R. P. Nederpelt Lazarom, J. van der Velden and R. Wiggers for their help in writing the ALGOL texts and in testing the procedures, to Mrs. E. Binnenmarsch for typing the manuscript, and to D. Zwarst and J. Suiker for the printing and the binding of this booklet.

CONTENTS

| | |
|--|---------|
| PREFACE | 1 |
| ACKNOWLEDGEMENTS | " |
| CONTENTS | 2 - 3 |
| NOTATIONS | 4 |
| DEFINITIONS | " |
| INTRODUCTION | 5 - 6 |
| CHAPTER 20 VECTOR OPERATIONS | 7 - 17 |
| Section 200 Scalar products | 8 - 11 |
| mca 2000 vecvec | 8 - 9 |
| mca 2001 matvec | " |
| mca 2002 tamvec | " |
| mca 2003 matmat | " |
| mca 2004 tammatt | " |
| mca 2005 mattam | " |
| mca 2006 seqvec | 10 - 11 |
| mca 2008 scaprd1 | " |
| Section 201 Elimination | 12 - 13 |
| mca 2010 elmvec | " |
| mca 2011 elmveccol | " |
| mca 2012 elmcolvec | " |
| mca 2013 elmcol | " |
| mca 2014 elmrow | " |
| mca 2019 maxelmrow | " |
| Section 202 Interchanging | 14 - 15 |
| mca 2020 ichvec | " |
| mca 2021 ichcol | " |
| mca 2022 ichrow | " |
| mca 2023 ichrowcol | " |
| mca 2024 ichseqvec | " |
| mca 2025 ichseq | " |
| Section 203 Rotation | 16 - 17 |
| mca 2031 rotcol | " |
| mca 2032 rotrow | " |
| CHAPTER 21 LINEAR SYSTEMS AND MATRIX INVERSION | 18 - 39 |
| Section 210 Triangular decomposition with partial pivoting | 19 - 23 |
| mca 2100 det | 20 - 21 |
| mca 2101 sol | " |
| mca 2102 detsol | " |
| mca 2103 inv | 22 - 23 |
| mca 2104 detinv | " |

| | |
|---|---------|
| Section 211 Elimination with complete pivoting | 24 - 31 |
| mca 2110 rnkelm | 26 - 27 |
| mca 2111 solelm | 28 - 29 |
| mca 2112 rnkssolelm | " |
| mca 2113 solhom | " |
| mca 2114 invelm | 30 - 31 |
| Section 212 Band matrices | 33 - 39 |
| mca 2120 detbnd | 34 - 35 |
| mca 2121 solbnd | 36 - 37 |
| mca 2122 detsolbnd | 38 - 39 |
| CHAPTER 22 POSITIVE DEFINITE SYMMETRIC LINEAR SYSTEMS AND MATRIX INVERSION | 40 - 69 |
| Section 220 Cholesky decomposition without pivoting | 41 - 47 |
| mca 2200 detsym2 | 42 - 43 |
| mca 2201 solsym2 | " |
| mca 2202 detsolsym2 | " |
| mca 2203 invsym2 | " |
| mca 2204 detinvsym2 | " |
| mca 2205 detsym1 | 44 - 45 |
| mca 2206 solsym1 | " |
| mca 2207 detsolsym1 | " |
| mca 2208 invsym1 | 46 - 47 |
| mca 2209 detinvsym1 | " |
| Section 221 Cholesky decomposition with pivoting | 48 - 59 |
| mca 2210 rnksym20 | 50 - 51 |
| mca 2211 solsym20 | " |
| mca 2212 rnksolsym20 | 52 - 53 |
| mca 2213 invsym20 | " |
| mca 2214 rnkinvsym20 | " |
| mca 221a solsymhom20 | 54 - 55 |
| mca 2215 rnksym10 | 56 - 57 |
| mca 2216 solsym10 | 58 - 59 |
| mca 2217 rnksolsym10 | " |
| mca 2218 invsym10 | " |
| mca 2219 rnkinvsym10 | " |
| Section 222 Cholesky decomposition for band matrices | 61 - 63 |
| mca 2220 detsymbnd | 62 - 63 |
| mca 2221 solsymbnd | " |
| mca 2222 detsolsymbnd | " |
| Section 224 Least-squares problems | 65 - 69 |
| mca 2240 lsqdec | 66 - 67 |
| mca 2241 lsqsol | 68 - 69 |
| mca 2242 lsqdglinv | " |
| mca 2243 lsqdecsol | " |
| REFERENCES | 70 |
| APPENDIX. TIMES FOR THE MC ALGOL 60 SYSTEM FOR THE X8 | 71 - 73 |

NOTATIONS

References are given between the square brackets "[" and "]".
 ":" denotes the integer division symbol "÷" [1,3.3.4.2].
 "goto" denotes the same symbol as "go to", [1,4.3].
 "0" denotes a null vector; the number of elements will be clear from the context.
 "min" ("max") denotes the function whose value is the minimum (maximum) value of its two operands.
 The prime symbol "'" denotes transposition of a matrix.
 "M" denotes the matrix considered and "n" denotes its order, unless stated otherwise.

DEFINITIONS

The "dimension" of an array is the number of its subscripts (see also [1,5.2.3.2.]). Thus we speak about "one-dimensional" and "two-dimensional" arrays. The first subscript of a two-dimensional array is called the "row index" and the second the "column index".

The i -th "row" ("column") of a two-dimensional array is the set of its elements for which the row (column) index equals i .

The "upper triangle" of a matrix or of a two-dimensional array is the set of its elements for which the first subscript does not exceed the second.

A "unit triangular" matrix is a triangular matrix whose diagonal elements are equal to 1.

We use the following vector norms [2, p.80] [3, p.55]: the "one-norm", i.e. the sum of the absolute values of the elements of the vector; the "Euclidean norm", i.e. the square root of the sum of their squares; and the "infinity-norm", i.e. their maximum absolute value.

We use the following matrix norms: the "infinity-norm", i.e. the maximum one-norm of its rows; the "maximum-norm", i.e. the maximum absolute value of its elements. The maximum-norm of a positive semidefinite symmetric matrix obviously equals the maximum of its diagonal elements.

The "machine precision" is the largest number, p , for which $1 + p = 1$ on the computer (about 10^{-12} for the X8).

A "relative tolerance" is a tolerance relative to some vector or matrix norm. Relative tolerances must be chosen smaller than one, and should be chosen not smaller than the machine precision.

INTRODUCTION

Chapter 20 contains a set of procedures for vector operations. Most of these are used in the subsequent chapters; some procedures of section 201 and 203 will be used in procedures for calculating eigenvalues and eigenvectors (to be published). A vector is given either in a one-dimensional array or in a row or column of a two-dimensional array. In the former case, we often use the same name for the vector as for the array if the whole array is used for the vector.

Chapter 21 contains a set of procedures for solving linear systems and for inverting matrices. The matrix is given in a two-dimensional array, the columns and rows of which correspond with the columns and rows of the matrix. A band matrix, however, is given in a one-dimensional array. For details, see sections 212 and (for the positive-definite symmetric case) 222.

Chapter 22 deals with the special case of positive-definite symmetric matrices and, moreover, contains a section for solving linear least-squares problems. Of symmetric matrices and upper-triangular matrices only the upper triangle will be given or delivered, either in the upper triangle of a two-dimensional array (in which case the remaining part of the array is not used) or in a one-dimensional array [8]. In the latter case, the (i, j) -th element of the matrix is, for $i < j$, the array element whose subscript equals $(j - 1) \times j : 2 + i$. Thus, the memory space occupied by the matrix is cut nearly in half. A drawback is that the elements in a row of the upper triangle are not linear functions of the running subscript, so that special procedures for the vector operations are needed.

In each chapter, we give a survey of its contents and some numerical considerations and comparisons. The chapters are subdivided into sections in each of which we give a more detailed survey of its contents and explain the numerical methods used. Each section contains one or more procedures. For each procedure, we give a description in which the required data, the delivered results and the (directly or indirectly) used nonlocal procedures are mentioned. The data of the procedures are given by actual parameters whose corresponding formal parameters are either specified real or integer and called by value, or specified array or integer array and called by name. The results of the procedures are delivered either as the value assigned to the procedure identifier (of type real or integer), or in arrays corresponding to formal parameters specified array or integer array. Some arrays are used as well for data as for results. For each formal parameter specified array or integer array, we give the "minimal declaration", i.e. a declaration with the appropriate number of bound pairs, where each pair indicates the range which is actually used by the procedure. Of course, the declaration of the corresponding actual parameter may contain smaller lower bounds and

greater upper bounds. (The descriptions of the procedures of sections 202 and 203 contain two minimal declarations of the same parameter with the meaning that the declaration of the corresponding actual parameter has to include both of them.) Sometimes not all elements of the array indicated by the minimal declaration are used. In the descriptions, we always mention which part is used for the data and which part for the results, so that it is always clear which elements are not used at all and which elements are left unchanged. Only in some cases shall we explicitly state that some elements are left intact, if it is important for the applications.

CHAPTER 20

VECTOR OPERATIONS

This chapter contains procedures for calculating scalar products, for adding a multiple of a vector to another vector, for interchanging two vectors and for rotating two vectors.

Each of the vectors is given by means of a pair of subscript bounds and either a two-dimensional array identifier with a row or column number or a one-dimensional array identifier. The procedures which use only one-dimensional arrays, have some extra facilities. Some of these procedures handle rows of upper-triangular or symmetric matrices given in a one-dimensional array, which are represented in a special way, viz. with linearly increasing spacing of the successive elements. Compared with the technique of defining the vectors by means of subscripted variables explicitly depending on a bound variable, as is common for scalar product procedures [1,4.7.2, and 5.4.2.] [5] [6] [10], our technique is less flexible, but more efficient (at least in the MC ALGOL 60 system for the X8); moreover, our procedures may well be written as machine-code procedures in which the elements of the vectors are selected more efficiently on account of their equidistant (or linearly increasing) spacing in the memory. As to the lesser flexibility, instead of one procedure we need a set of procedures for the most important applications. As to the machine-code procedures, if explicit bound variables were used, the more efficient element selection would be possible only if the subscripts were linear (or, in the case of linearly increasing spacing, quadratic) functions of the bound variable, but this requirement is easily violated and difficult to check.

Our procedures avoid this difficulty.

```

comment mca 2000;
real procedure vecvec(l, u, shift, a, b); value l, u, shift;
integer l, u, shift; array a, b;
begin integer k; real s;
    s:= 0;
    for k:= 1 step 1 until u do s:= a[k] × b[shift + k] + s;
    vecvec:= s
end vecvec;

```

```

comment mca 2001; Matvec
real procedure matvec(l, u, i, a, b); value l, u, i; integer l, u, i;
array a, b;
begin integer k; real s;
    s:= 0;
    for k:= 1 step 1 until u do s:= a[i,k] × b[k] + s; matvec:= s
end matvec;

```

```

comment mca 2002;
real procedure tamvec(l, u, i, a, b); value l, u, i; integer l, u, i;
array a, b;
begin integer k; real s;
    s:= 0;
    for k:= 1 step 1 until u do s:= a[k,i] × b[k] + s; tamvec:= s
end tamvec;

```

```

comment mca 2003; Matmat
real procedure matmat(l, u, i, j, a, b); value l, u, i, j;
integer l, u, i, j; array a, b;
begin integer k; real s;
    s:= 0;
    for k:= 1 step 1 until u do s:= a[i,k] × b[k,j] + s; matmat:= s
end matmat;

```

```

comment mca 2004;
real procedure tammat(l, u, i, j, a, b); value l, u, i, j;
integer l, u, i, j; array a, b;
begin integer k; real s;
    s:= 0;
    for k:= 1 step 1 until u do s:= a[k,i] × b[k,j] + s; tammat:= s
end tammat;

```

```

comment mca 2005;
real procedure mattam(l, u, i, j, a, b); value l, u, i, j;
integer l, u, i, j; array a, b;
begin integer k; real s;
    s:= 0;
    for k:= 1 step 1 until u do s:= a[i,k] × b[j,k] + s; mattam:= s
end mattam;

```

Section 200 Scalar products

The procedures of this section calculate the scalar product of two vectors, each of which is given either as (a part of) a one-dimensional array or as row or column of a two-dimensional array. If the lower bound of the running subscript is greater than the upper bound, then 0 is delivered as scalar product.

The lower and upper bound of the running subscript are given by two parameters;

vecvec and seqvec feature the additional possibility of shifting the range of the running subscript of the second vector; in scaprd1, moreover, the spacings of the vectors are arbitrary constants; in seqvec the spacing of the successive elements of the first vector increases linearly. (The latter procedure is used for symmetric or upper-triangular matrices given in one-dimensional arrays.)

In the MC ALGOL 60 system for the X8 the procedures mca 2000 to 2005 are available as machine-code procedures (which are about 7 times faster than the corresponding equivalent ALGOL procedures, see Appendix).

Description mca 2000

vecvec := scalar product of the vectors given in array a[1:u] and array b[shift + 1 : shift + u].

Description mca 2001

matvec := scalar product of the row vector given in array a[i:i, 1:u] and the vector given in array b[1:u].

Description mca 2002

tanvec := scalar product of the column vector given in array a[1:u, 1:1] and the vector given in array b[1:u].

Description mca 2003

matmat := scalar product of the row vector given in array a[i:i, 1:u] and the column vector given in array b[1:u, j:j].

Description mca 2004

tamat := scalar product of the column vectors given in array a[1:u, 1:1] and array b[1:u, j:j].

Description mca 2005

mattam := scalar product of the row vectors given in array a[i:i, 1:u] and array b[j:j, 1:u].

```
comment mca 2006;  
real procedure seqvec(l, u, il, shift, a, b); value l, u, il, shift;  
integer l, u, il, shift; array a, b;  
begin real s;  
  s:= 0;  
  for l:= 1 step 1 until u do  
    begin s:= a[il] × b[l + shift] + s; il:= il + 1 end;  
  seqvec:= s  
end seqvec;
```

```
comment mca 2008;  
real procedure scaprd1(la, sa, lb, sb, n, a, b);  
value la, sa, lb, sb, n; integer la, sa, lb, sb, n; array a, b;  
begin integer k;  
  real s;  
  s:= 0;  
  for k:= 1 step 1 until n do  
    begin s:= a[la] × b[lb] + s; la:= la + sa; lb:= lb + sb end;  
  scaprd1:= s  
end scaprd1;
```

Description mca 2006

seqvec := scalar product of the vectors given in
array a[il : il + (u + 1 - 1) × (u - 1) : 2] and
array b[shift + 1 : shift + u], where the elements of the first vector
are a[il + (j + 1 - 1) × (j - 1) : 2] for j = 1, ..., u.

Description mca 2008

scaprd := scalar product of the vectors given in
array a[min (la, la + (n - 1) × sa) : max (la, la + (n - 1) × sa)]
and array b[min (lb, lb + (n - 1) × sb) : max (lb, lb + (n - 1) × sb)],
where the elements of the vectors are
a[la + (j - 1) × sa] and b[lb + (j - 1) × sb] for j = 1, ..., n.

12

```
comment mca 2010;  
procedure elmvec(l, u, shift, a, b, x); value l, u, shift, x;  
integer l, u, shift; real x; array a, b;  
for l:= 1 step 1 until u do a[l]:= a[l] + b[l + shift] × x;
```

```
comment mca 2011;  
procedure elmveccol(l, u, i, a, b, x); value l, u, i, x;  
integer l, u, i; real x; array a, b;  
for l:= 1 step 1 until u do a[l]:= a[l] + b[l,i] × x;
```

```
comment mca 2012;  
procedure elmcolvec(l, u, i, a, b, x); value l, u, i, x;  
integer l, u, i; real x; array a, b;  
for l:= 1 step 1 until u do a[l,i]:= a[l,i] + b[l] × x;
```

```
comment mca 2013;  
procedure elmcol(l, u, i, j, a, b, x); value l, u, i, j, x;  
integer l, u, i, j; real x; array a, b;  
for l:= 1 step 1 until u do a[l,i]:= a[l,i] + b[l,j] × x;
```

```
comment mca 2014;  
procedure elmrow(l, u, i, j, a, b, x); value l, u, i, j, x;  
integer l, u, i, j; real x; array a, b;  
for l:= 1 step 1 until u do a[i,l]:= a[i,l] + b[j,l] × x;
```

```
comment mca 2019;  
integer procedure maxelmrow(l, u, i, j, a, b, x); value l, u, i, j, x;  
integer l, u, i, j; real x; array a, b;  
begin integer k;  
  real r, s;  
  s:= 0;  
  for k:= 1 step 1 until u do  
    begin r:= a[i,k]:= a[i,k] + b[j,k] × x; if abs(r) > s then  
      begin s:= abs(r); l:= k end  
    end;  
  maxelmrow:= l  
end maxelmrow;
```

Section 201 Elimination

The procedures of this section perform a Gaussian elimination on a vector. More precisely, a multiple of one vector is added to another vector. Each vector is given either as a one-dimensional array or as row or column of a two-dimensional array. The lower and upper bound of the running subscript are given by two parameters; `elmvec` features the additional possibility of shifting the range of the running subscript of one vector.

Description mca 2010

`elmvec` adds x times the vector given in array `b[shift + 1 : shift + u]` to the vector given in array `a[1:u]`.

Description mca 2011

`elmveccol` adds x times the column vector given in array `b[1:u, i:i]` to the vector given in array `a[1:u]`.

Description mca 2012

`elmcolvec` adds x times the vector given in array `b[1:u]` to the column vector given in array `a[1:u, i:i]`.

Description mca 2013

`elmcol` adds x times the column vector given in array `b[1:u, j:j]` to the column vector given in array `a[1:u, i:i]`.

Description mca 2014

`elmrow` adds x times the row vector given in array `b[j:j, 1:u]` to the row vector given in array `a[i:i, 1:u]`.

Description mca 2019

`maxelmrow` adds x times the row vector given in array `b[j:j, 1:u]` to the row vector given in array `a[i:i, 1:u]`.
Moreover, `maxelmrow:=` the value of the second subscript of an element of the new row vector in array `a` which is of maximum absolute value. If, however, $l > u$, then `maxelmrow:= 1`.

14

```
comment mca 2020;  
procedure ichvec(l, u, shift, a); value l, u, shift;  
integer l, u, shift; array a;  
begin real r;  
    for l:= 1 step 1 until u do  
        begin r:= a[l]; a[l]:= a[l + shift]; a[l + shift]:= r end  
end ichvec;
```

```
comment mca 2021;  
procedure ichcol(l, u, i, j, a); value l, u, i, j; integer l, u, i, j;  
array a;  
begin real r;  
    for l:= 1 step 1 until u do  
        begin r:= a[l,i]; a[l,i]:= a[l,j]; a[l,j]:= r end  
end ichcol;
```

```
comment mca 2022;  
procedure ichrow(l, u, i, j, a); value l, u, i, j; integer l, u, i, j;  
array a;  
begin real r;  
    for l:= 1 step 1 until u do  
        begin r:= a[i,l]; a[i,l]:= a[j,l]; a[j,l]:= r end  
end ichrow;
```

```
comment mca 2023;  
procedure ichrowcol(l, u, i, j, a); value l, u, i, j;  
integer l, u, i, j; array a;  
begin real r;  
    for l:= 1 step 1 until u do  
        begin r:= a[i,l]; a[i,l]:= a[l,j]; a[l,j]:= r end  
end ichrowcol;
```

```
comment mca 2024;  
procedure ichseqvec(l, u, il, shift, a); value l, u, il, shift;  
integer l, u, il, shift; array a;  
begin real r;  
    for l:= 1 step 1 until u do  
        begin r:= a[il]; a[il]:= a[l + shift]; a[l + shift]:= r;  
            il:= il + 1  
        end  
end ichseqvec;
```

```
comment mca 2025;  
procedure ichseq(l, u, il, shift, a); value l, u, il, shift;  
integer l, u, il, shift; array a;  
begin real r;  
    for l:= 1 step 1 until u do  
        begin r:= a[il]; a[il]:= a[il + shift]; a[il + shift]:= r;  
            il:= il + 1  
        end  
end ichseq;
```


Section 202 Interchanging

The procedures of this section interchange the elements of two vectors. Each vector is given either as (a part of) a one-dimensional array or as row or column of a two-dimensional array. The lower and upper bound of the running subscript are given by two parameters; `ichvec` and `ichseqvec` feature the additional possibility of shifting the range of the running subscript of the second vector; in `ichseqvec` and `ichseq` the spacing of the successive elements of one or both of the vectors increases linearly. (The latter procedures are used for symmetric matrices given in one-dimensional arrays.)

Description mca 2020

`ichvec` interchanges the elements of the vector given in array `a[1:u]` and array `a[shift + 1 : shift + u]`.

Description mca 2021

`ichcol` interchanges the elements of the column vectors given in array `a[1:u, i:i]` and array `a[1:u, j:j]`.

Description mca 2022

`ichrow` interchanges the elements of the row vectors given in array `a[i:i, 1:u]` and array `a[j:j, 1:u]`.

Description mca 2023

`ichrowcol` interchanges the elements of the row vector given in array `a[i:i, 1:u]` and the column vector given in array `a[1:u, j:j]`.

Description mca 2024

`ichseqvec` interchanges the elements of the vectors given in array `a[il : il + (u + 1 - 1) × (u - 1) : 2]` and array `a[shift + 1 : shift + u]`, where the elements of the first vector are `a[il + (j + 1 - 1) × (j - 1) : 2]` for $j = 1, \dots, u$.

Description mca 2025

`ichseq` interchanges the elements of the vectors given in array `a[il : il + (u + 1 - 1) × (u - 1) : 2]` and array `a[shift + il : shift + il + (u + 1 - 1) × (u - 1) : 2]`, where the elements of the vectors are `a[il + (j + 1 - 1) × (j - 1) : 2]` and `a[shift + il + (j + 1 - 1) × (j - 1) : 2]` for $j = 1, \dots, u$.

16

```
comment mca 2031;  
procedure rotcol(l, u, i, j, a, c, s); value l, u, i, j, c, s;  
integer l, u, i, j; real c, s; array a;  
begin real x, y;  
  for l:= 1 step 1 until u do  
    begin x:= a[l,i]; y:= a[l,j]; a[l,i]:= x × c + y × s;  
      a[l,j]:= y × c - x × s  
    end  
end rotcol;
```

```
comment mca 2032;  
procedure rotrow(l, u, i, j, a, c, s); value l, u, i, j, c, s;  
integer l, u, i, j; real c, s; array a;  
begin real x, y;  
  for l:= 1 step 1 until u do  
    begin x:= a[i,l]; y:= a[j,l]; a[i,l]:= x × c + y × s;  
      a[j,l]:= y × c - x × s  
    end  
end rotrow;
```

Section 203 Rotation

The procedures of this section perform a rotation on two vectors, x and y (say); i.e. the two vectors are replaced by $cx + sy$ and $cy - sx$, where c and s are two given real values. Each vector is given as row or column of a two-dimensional array. The lower and upper bound of the running subscript are given by two parameters. (These procedures are to be used in procedures for calculating eigenvalues and eigenvectors (to be published).)

Description mca 2031

rotcol replaces the column vector x given in array $a[1:u, 1:1]$ and the column vector y given in array $a[1:u, j:j]$ by the vectors $cx + sy$ and $cy - sx$.

Description mca 2032

rotrow replaces the row vector x given in array $a[i:i, 1:u]$ and the row vector y given in array $a[j:j, 1:u]$ by the vectors $cx + sy$ and $cy - sx$.

CHAPTER 21

LINEAR SYSTEMS AND MATRIX INVERSION

This chapter contains procedures for solving systems of linear equations and for matrix inversion. Moreover, section 211 contains a procedure for calculating the rank of a matrix and a procedure for solving a homogeneous linear system.

In section 210 triangular decomposition with partial pivoting is used and in section 211 Gaussian elimination with complete pivoting. The procedures of section 212 solve linear systems whose matrices are in band form, by means of Gaussian elimination with partial pivoting.

In exceptional cases, partial pivoting may yield useless results, even for well-conditioned matrices; this may occur only for large matrices whose order (or in the case of band matrices, band width) is not much smaller than the number of binary digits in the number representation [2, p.97] [3, p.212]. Complete pivoting, however, always yields good results for well-conditioned matrices; a "condition number" (i. e. a norm of the matrix times a norm of its inverse) is a measure of the relative accuracy of the solution [2, p.91]. Moreover, complete pivoting is indispensable for calculating the rank of a matrix and for solving homogeneous systems.

For large order n the computation time for solving linear systems and for matrix inversion is proportional to n cubed. Complete pivoting requires some extra time for the pivot selection, which, for large n , is a nearly constant (small) fraction of the total computation time. In the MC ALGOL 60 system for the X8, the partial pivoting procedures of section 210 are much faster than the complete pivoting procedures of section 211, because the procedures mca 2000 to 2005 are available in machine-code.

The procedures of section 212, for solving linear systems whose matrices are in band form, save a considerable amount of computation time and memory space, if the band width is much smaller than n . For large matrices, the computation time is proportional to $n \times$ band width \times the number of diagonals on or to the left of the main diagonal.

Section 210 Triangular decomposition with partial pivoting

This section contains procedures for solving linear systems and for inverting matrices:

detsol solves a system of linear equations and calculates the determinant of the system;
 detinv calculates the inverse and the determinant of a matrix;
 det calculates the determinant of a matrix;
 sol solves a system of linear equations and inv inverts a matrix, provided the matrix is given in the triangularly decomposed form produced by det. One call of det followed by several calls of sol may be used to solve several linear systems having the same matrix but different right hand sides.

The method used in det is triangular decomposition with stabilizing row interchanges, also called "partial pivoting" [2, p.115] [3, p.201] [4] [5] [6].

The method yields a lower-triangular matrix L and a unit upper-triangular matrix U such that the product LU equals the given matrix M with permuted rows.

The process is performed in n steps. The k-th step, $k = 1, \dots, n$, produces the k-th column of L; subsequently, the "pivot" is selected in this column; the pivotal row and the k-th row of M (and thus also of L) are interchanged; finally, the k-th row of U is produced. That element of the k-th column of L is chosen as pivot, whose absolute value divided by the Euclidean norm of the corresponding row of M is maximal. Thus, matrix M is "equilibrated" in this pivoting strategy such that the rows effectively obtain unit Euclidean norm. No test for singularity of M is performed.

The determinant equals the product of the diagonal elements of L or minus this value, if the number of proper interchanges is odd.

After the triangular decomposition, sol obtains the solution x of the linear system $Mx = b$ by first permuting the elements of b in the same way as the rows of M, then calculating y such that Ly equals b with permuted elements (forward substitution), and finally calculating x such that $Ux = y$ (back substitution).

The method used in inv is as follows. The inverse, X, of the product LU is calculated from the conditions that XL be a unit upper-triangular matrix and UX a lower-triangular matrix [4, p.34-38]. Subsequently, in correspondence with the interchanges applied on the rows of M, the same interchanges are carried out in reverse order on the columns of X, in order to obtain the inverse of M.

```

comment mca 2100;
real procedure det(a, n, p); value n; integer n; array a;
integer array p;
begin integer i, k, k1, pk;
  real d, r, s;
  array v[1:n];
  for i:= 1 step 1 until n do v[i]:= 1 / sqrt(mattam(1, n, i, i,
a, a)); d:= 1;
  for k:= 1 step 1 until n do
begin r:= - 1; k1:= k - 1;
  for i:= k step 1 until n do
begin a[i,k]:= a[i,k] - matmat(1, k1, i, k, a, a);
  s:= abs(a[i,k]) × v[i]; if s > r then
begin r:= s; pk:= i end
end lower;
p[k]:= pk; v[pk]:= v[k]; s:= a[pk,k]; d:= s × d;
if pk ≠ k then
begin d:= - d; ichrow(1, n, k, pk, a) end;
for i:= k + 1 step 1 until n do a[k,i]:= (a[k,i] - matmat(1,
k1, k, i, a, a)) / s
end lu;
det:= d
end det;

```

```

comment mca 2101;
procedure sol(a, n, p, b); value n; integer n; array a, b;
integer array p;
begin integer k, pk;
  real r;
  for k:= 1 step 1 until n do
begin r:= b[k]; pk:= p[k];
  b[k]:= (b[pk] - matvec(1, k - 1, k, a, b)) / a[k,k];
  if pk ≠ k then b[pk]:= r
end;
for k:= n step - 1 until 1 do b[k]:= b[k] - matvec(k + 1, n, k,
a, b)
end sol;

```

```

comment mca 2102;
real procedure detsol(a, n, b); value n; integer n; array a, b;
begin integer array p[1:n];
  detsol:= det(a, n, p); sol(a, n, p, b)
end detsol;

```

Description mca 2100

det:= determinant of the n-th order matrix M given in array a[1:n, 1:n], and the triangular decomposition of M is performed. The resulting lower-triangular matrix and unit upper-triangular matrix with its unit diagonal omitted are overwritten on a. The pivotal indices are delivered in integer array p[1:n]. det uses mattam, matmat and ichrow (chapter 20).

Description mca 2101

sol should be called after det and solves the linear system $Mx = b$, where M is the n-th order matrix whose triangularly decomposed form and pivotal indices, as produced by det, are given in array a[1:n, 1:n] and integer array p[1:n], and where b is the vector given as array b[1:n]. The solution vector x is overwritten on b. sol leaves a and p intact, so that, after one call of det, several calls of sol may follow for solving several systems having the same matrix but different right hand sides. sol uses matvec (mca 2001).

Description mca 2102

detsol:= determinant of the n-th order matrix M given in array a[1:n, 1:n], and the triangular decomposition of M is performed. Moreover the linear system $Mx = b$ is solved, where vector b is given as array b[1:n]. The solution vector x is overwritten on b, and the triangularly decomposed form of M is overwritten on a. detsol uses det, sol and, indirectly, also mattam, matmat, matvec and ichrow (chapter 20).

```

comment mca 2103;
procedure inv(a, n, p); value n; integer n; array a; integer array p;
begin integer j, k, k1;
  real r;
  array v[1:n];
  for k:= n step - 1 until 1 do
    begin k1:= k + 1;
      for j:= n step - 1 until k1 do
        begin a[j,k1]:= v[j]; v[j]:= - matmat(k1, n, k, j, a, a) end;
        r:= a[k,k];
        for j:= n step - 1 until k1 do
          begin a[k,j]:= v[j]; v[j]:= - matmat(k1, n, j, k, a, a) / r
          end;
        v[k]:= (1 - matmat(k1, n, k, k, a, a)) / r
      end;
    for j:= n step - 1 until 1 do a[j,1]:= v[j];
    for k:= n - 1 step - 1 until 1 do
      begin k1:= p[k]; if k1 ≠ k then ichcol(1, n, k, k1, a) end
    end inv;

```

```

comment mca 2104;
real procedure detinv(a, n); value n; integer n; array a;
begin integer array p[1:n];
  detinv:= det(a, n, p); inv(a, n, p)
end detinv;

```


Description mca 2103

inv should be called after det and calculates the inverse of the matrix whose triangularly decomposed form and pivotal indices, as produced by det, are given in array a[1:n, 1:n] and integer array p[1:n]. The calculated inverse is overwritten on a. inv uses matmat and ichcol (chapter 20).

Description mca 2104

detinv:= determinant of the n-th order matrix given in array a[1:n, 1:n]. Moreover, the inverse of this matrix is calculated and overwritten on a.

detinv uses det, inv and, indirectly, also matam, matmat, ichrow and ichcol (chapter 20).

Section 211 Elimination with complete pivoting

This section contains procedures for calculating the rank of a matrix, for solving linear systems and for inverting matrices:

rnksolelm solves a system of linear equations and calculates the determinant of the system;

invelm calculates the inverse and the determinant of a matrix;

rnkelm calculates the rank of a rectangular matrix;

solelm solves a system of linear equations and solhom calculates a solution of a homogeneous system, provided the matrix is given in the Gaussian-eliminated form produced by rnkelm.

One call of rnkelm followed by several calls of solelm may be used to solve several linear systems having the same matrix but different right-hand sides. By means of successive calls of solhom one can obtain a complete linearly independent set of solutions of a homogeneous system.

The method used, in rnkelm, is Gaussian elimination with complete pivoting [2, p.97] [3, p.212] which yields a lower-triangular matrix L and a unit upper-triangular matrix U such that the product LU equals the given matrix M with permuted rows and columns. Let M have n rows and m columns. The elimination is performed in at most $\min(n, m)$ steps. In the k -th step, $k = 1, \dots, \min(n, m)$, a "pivot" is selected from the remaining submatrix having $n - k + 1$ rows and $m - k + 1$ columns; then the pivotal row is interchanged with the k -th row and the pivotal column with the k -th column; subsequently, the k -th "unknown" is eliminated in the last $n - k$ rows. The pivot is selected in such a way that its absolute value divided by the one-norm of the corresponding row of M is maximal. Thus, matrix M is "equilibrated" in this pivoting strategy such that the rows effectively obtain unit one-norm. If all elements of the remaining submatrix are smaller in absolute value than a given relative tolerance times the one-norm of the corresponding row of M , then the elimination is discontinued and the previous step number is delivered as the rank of M .

After the Gaussian elimination, solelm obtains the solution x of the linear system $Mx = b$ by calling sol (mca 2101) and then interchanging the elements of the solution vector produced by sol in "reverse correspondence" with the interchanges of the columns of M ; i.e. the same interchanges are carried out in reverse order.

In solhom, a solution x of the homogeneous system $Mx = 0$, where M is an $n \times m$ matrix of rank r , is obtained as follows. Let \bar{V} be the r -th order upper-triangular matrix consisting of the first r columns of the matrix U produced by `rnkelm` and W the $r \times (m - r)$ matrix consisting of the remaining part of the first r rows of U (the other rows of U are negligible). First, the system $Vy =$ minus the k -th column of W , where k is a given positive integer $< m - r$, is solved (back substitution). Then the vector y and the k -th unit $(m - r)$ -vector are combined to form a single m -vector; its elements are then interchanged in reverse correspondence with the interchanges of the columns of M , in order to obtain a solution vector x . The solution vectors thus obtained for $k = 1, \dots, m - r$ form a complete set of linearly independent solution vectors of the homogeneous system.

The method used in `invelm` is Gauss-Jordan elimination with complete pivoting. This method introduces the zeros not only below but also above the main diagonal. At each stage, the element of greatest absolute value of the remaining submatrix is chosen as pivot. (Here, the matrix is not equilibrated, because this would not leave the inverse invariant.) After completing the elimination, the rows and columns are interchanged in reverse correspondence with the interchanges of the columns and rows of M .

```

comment mca 2110;
integer procedure rnkelm(a, n, m, aux, ri, ci); value n, m;
integer n, m; array aux; integer array ri, ci;
begin integer i, j, p, q, r, r1, jcrit;
  real crit, rnorm, max, aid, det, eps, minpiv, pivot;
  array norm[1:n];
  crit:= 0;
  for p:= 1 step 1 until n do
  begin rnorm:= max:= abs(a[p,1]); jcrit:= 1;
    for q:= 2 step 1 until m do
    begin aid:= abs(a[p,q]); rnorm:= rnorm + aid;
      if aid > max then
      begin max:= aid; jcrit:= q end
    end;
    norm[p]:= rnorm:= 1 / rnorm; if max × rnorm > crit then
    begin crit:= max × rnorm; i:= p; j:= jcrit end
  end;
  eps:= aux[0]; det:= 1; minpiv:= crit;
  for r:= 1 step 1 until n do
  begin if crit < eps then goto rank; r1:= r + 1;
    if crit < minpiv then minpiv:= crit; if i ≠ r then
    begin det:= - det; norm[i]:= norm[r]; ichrow(1, m, r, i, a)
    end;
    if j ≠ r then
    begin det:= - det; ichcol(1, n, r, j, a) end;
    ri[r]:= i; ci[r]:= j; pivot:= a[r,r]; det:= det × pivot;
    crit:= 0; if r1 < m then
    begin for q:= r1 step 1 until m do a[r,q]:= a[r,q] / pivot;
      for p:= r1 step 1 until n do
      begin jcrit:= maxelmrow(r1, m, p, r, a, a, - a[p,r]);
        aid:= abs(a[p,jcrit]) × norm[p]; if aid > crit then
        begin crit:= aid; i:= p; j:= jcrit end
      end
    end
  end
end elimination;
r:= n + 1;
rank: rnkelm:= r - 1; aux[1]:= 1 / minpiv; aux[2]:= crit;
aux[3]:= det
end rnkelm;

```

Description mca 2110

rnkelm:= rank, r , of the $n \times m$ matrix M given in array $a[1:n, 1:m]$.
In array $aux[0:3]$ one must give a relative tolerance, $aux[0]$.
The Gaussian-eliminated form of M is overwritten on a , and the pivotal row and column indices are delivered in integer array $ri, ci[1:r]$.
Moreover,
 $aux[1]$:= reciprocal of the minimum absolute value of "pivot / one-norm of the corresponding row of M ";
 $aux[2]$:= maximum absolute value of "element of the remaining $(n - r) \times (m - r)$ submatrix / one-norm of the corresponding row of M " if $r < \min(n, m)$, and otherwise 0;
 $aux[3]$:= determinant of the principal submatrix of order r .
rnkelm uses maxelmrow, ichrow and ichcol (chapter 20).

```

comment mca 2111;
procedure solelm(a, n, ri, ci, b); value n; integer n; array a, b;
integer array ri, ci;
begin integer r, cir;
    real w;
    sol(a, n, ri, b);
    for r:= n step - 1 until 1 do
    begin cir:= ci[r]; if cir  $\neq$  r then
        begin w:= b[r]; b[r]:= b[cir]; b[cir]:= w end
    end
end solelm;

```

```

comment mca 2112;
integer procedure rnksolelm(a, n, aux, b); value n; integer n;
array a, aux, b;
begin integer rank;
    integer array ri, ci[1:n];
    rank:= rnksolelm:= rnkelm(a, n, n, aux, ri, ci);
    if rank = n then solelm(a, n, ri, ci, b)
end rnksolelm;

```

```

comment mca 2113;
procedure solhom(a, rank, m, k, ci, x); value rank, m, k;
integer rank, m, k; array a, x; integer array ci;
begin integer r, rk;
    real w;
    rk:= rank + k;
    for r:= rank step - 1 until 1 do x[r]:= - (matvec(r + 1, rank,
    r, a, x) + a[r,rk]);
    for r:= rank + 1 step 1 until m do x[r]:= 0; x[rk]:= 1;
    for r:= rank step - 1 until 1 do
    begin k:= ci[r]; if k  $\neq$  r then
        begin w:= x[r]; x[r]:= x[k]; x[k]:= w end
    end
end solhom;

```

Description mca 2111

solelm should be called after rnkelm (but only if the rank delivered equals n), and solves the linear system $Mx = b$, where M is the n -th order matrix whose Gaussian-eliminated form and pivotal row and column indices, as produced by rnkelm, are given in array $a[1:n, 1:n]$ and integer array $ri, ci[1:n]$, and where b is the vector given as array $b[1:n]$.

The solution vector x is overwritten on b , and sol leaves the other data invariant, so that, after one call of rnkelm, several calls of solelm may follow for solving several systems having the same matrix but different right-hand sides.

solelm uses sol (mca 2101) and, indirectly, also matvec (mca 2001).

Description mca 2112

rnksolelm:= rank, r, of the n -th order matrix M given in array $a[1:n, 1:n]$.

In array $aux[0:3]$ one must give a tolerance, $aux[0]$.

If $r = n$, the linear system $Mx = b$ is solved, where b is the vector given as array $b[1:n]$, and the solution vector x is overwritten on b . The Gaussian-eliminated form of M is overwritten on a .

Moreover,

$aux[1] :=$ reciprocal of the minimum absolute value of "pivot / one-norm of the corresponding row of M ";

$aux[2] := 0$;

$aux[3] :=$ determinant of M .

If, however, $r < n$, then no solution is calculated and the effect of rnksolelm is the same as that of rnkelm.

rnksolelm uses rnkelm, solelm and, indirectly, also sol (mca 2101), matvec, maxelmrow, ichrow and ichcol (chapter 20).

Description mca 2113

solhom should be called after rnkelm and calculates the k -th solution vector of the homogeneous system $Mx = 0$, where M is the matrix whose Gaussian-eliminated form (or rather its first rank rows) and pivotal column indices, as produced by rnkelm, are given in array $a[1:rank, 1:m]$ and integer array $ci[1:rank]$, rank being the rank of M delivered by rnkelm. The given integer k must satisfy $1 \leq k \leq m - rank$.

The solution vector is delivered as array $x[1:m]$.

Calling solhom consecutively with $k = 1, \dots, m - rank$, one obtains a complete set of linearly independent solution vectors of the homogeneous system.

solhom uses matvec (mca 2001).

```

comment mca 2114;
real procedure invelm(a, n, aux); value n; integer n; array a, aux;
begin integer p, q, r, i, j;
  real t, w, det, pivot, co, tol, max;
  integer array ri, ci[1:n];
  i:= j:= 1; pivot:= abs(a[1,1]);
  for p:= 1 step 1 until n do
  for q:= 1 step 1 until n do if abs(a[p,q]) > pivot then
  begin i:= p; j:= q; pivot:= abs(a[p,q]) end;
  max:= pivot; det:= 1; co:= 0; tol:= aux[0] × max;
  for r:= 1 step 1 until n do
  begin if pivot < tol then
  begin det:= 0; aux[1]:= - r + 1; goto exit end;
  if i ≠ r then
  begin det:= - det; ichrow(1, n, r, i, a) end;
  if j ≠ r then
  begin det:= - det; ichcol(1, n, r, j, a) end;
  ri[r]:= i; ci[r]:= j; w:= a[r,r]; det:= det × w;
  a[r,r]:= 1 / w;
  for q:= n step - 1 until r + 1, r - 1 step - 1 until 1 do
  a[r,q]:= a[r,q] / w; pivot:= 0;
  for p:= 1 step 1 until r - 1 do
  begin t:= - a[p,r]; a[p,r]:= 0; elmrow(1, n, p, r, a, a, t)
  end;
  for p:= r + 1 step 1 until n do
  begin t:= - a[p,r]; a[p,r]:= 0; elmrow(1, r, p, r, a, a, t);
  q:= maxelmrow(r + 1, n, p, r, a, a, t);
  if abs(a[p,q]) > pivot then
  begin i:= p; j:= q; pivot:= abs(a[p,q]) end
  end
  end elimination;
  for p:= n step - 1 until 1 do
  begin for q:= 1 step 1 until n do if abs(a[p,q]) > co then co:=
  abs(a[p,q]); r:= ci[p]; if r ≠ p then ichrow(1, n, p, r, a)
  end;
  for q:= n step - 1 until 1 do
  begin r:= ri[q]; if r ≠ q then ichcol(1, n, q, r, a) end;
  aux[1]:= max × co;
  exit: invelm:= det
  end invelm;

```


Description mca 2114

invelm:= determinant of the n-th order matrix M given in array a[1:n, 1:n]. In array aux[0:1] one must give a relative tolerance, aux[0].

The inverse of M is calculated and overwritten on a, and aux[1]:= the product of the maximum-norm of M and that of its calculated inverse, this being a condition number of M.

If, however, M is singular (more precisely, if, at some stage, the absolute value of the pivot is smaller than aux[0] × the maximum-norm of M), then the calculation is discontinued, invelm:= 0, and aux[1]:= minus the rank of M.

invelm uses elmrow, maxelmrow, ichrow and ichcol (chapter 20).

Section 212 Band matrices

The procedures of this section solve a system of linear equations whose matrix is in band form and / or calculate the determinant of a band matrix:

detsolbnd solves a system of linear equations and calculates the determinant of the system;

detbnd calculates the determinant;

solbnd solves a system of linear equations whose matrix is given in the Gaussian-eliminated form produced by detbnd.

One call of detbnd followed by several calls of solbnd may be used to solve several linear systems having the same matrix, but different right-hand sides.

The method used is Gaussian elimination with stabilizing row interchanges (partial pivoting) [2, p.94] [3, p.204] [7]. Complete pivoting is superfluous if the band width is small (certainly if it is much smaller than the number of binary digits in the number representation).

If the given matrix M has lw nonzero codiagonals to the left and rw to the right of the main diagonal, then the Gaussian elimination yields a unit lower-triangular band matrix L of Gaussian multipliers having lw nonzero codiagonals, and an upper-triangular band matrix U of the resulting equivalent system having $lw + rw$ nonzero codiagonals. The Gaussian elimination is performed in n steps. In the k -th step, $k = 1, \dots, n$, a "pivot" is selected in the k -th column of the remaining submatrix of order $n - k + 1$ (which column contains at most $lw + 1$ nonzero elements); then the pivotal row is interchanged with the k -th row; subsequently, the k -th "unknown" is eliminated in the last $n - k$ rows (at most the first lw rows of which are involved). The pivot is selected in such a way that its absolute value divided by the Euclidean norm of the corresponding row of M , is maximal. Thus, matrix M is "equilibrated" in this pivoting strategy such that the rows effectively obtain unit Euclidean norm.

If M is singular (i.e. if, in some step, the absolute value of the pivot is smaller than a given relative tolerance times the Euclidean norm of the corresponding row of M), then the elimination is discontinued and 0 is delivered as the determinant value.

The solution x of the linear system $Mx = b$ is obtained by carrying out the corresponding eliminations on b , thereby yielding a vector y (say), and then solving the system $Ux = y$ (back substitution).

```

comment mca 2120;
real procedure detbnd(a, n, lw, rw, aux, m, p); value n, lw, rw;
integer n, lw, rw; integer array p; array a, m, aux;
begin integer i, j, k, kk, kk1, pk, mk, ik, lw1, f, q, w, w1, w2, iw,
nrw;
real r, s, norm, eps, min, det;
array v[1:n];
f:=lw; det:=1; w1:=lw+rw; w:=w1+1; w2:=w-2; iw:=0;
nrw:=n-rw; lw1:=lw+1; q:=lw-1;
for i:=2 step 1 until lw do
begin q:=q-1; iw:=iw+w1;
for j:=iw-q step 1 until iw do a[j]:=0
end;
norm:=0; iw:=-w2; q:=nrw+w-1; j:=rw-1;
for i:=1 step 1 until n do
begin iw:=lw+w; if i<lw1 then iw:=iw-1;
r:=v[i]:=sqrt(vecvec(iw, iw+(if i<lw then j+i else
if i>nrw then q-i else w1), 0, a, a));
if r>norm then norm:=r
end;
eps:=aux[0]; min:=1; kk:=-w1; mk:=-lw;
if f>nrw then w2:=w2+nrw-f;
for k:=1 step 1 until n do
begin if f<n then f:=f+1; ik:=kk:=kk+w; mk:=mk+lw;
s:=abs(a[kk])/v[k]; pk:=k; kk1:=kk+1;
for i:=k+1 step 1 until f do
begin ik:=ik+w1; m[mk+i-k]:=r:=a[ik]; a[ik]:=0;
r:=abs(r)/v[i]; if r>s then
begin s:=r; pk:=i end
end;
if s<min then min:=s; if s<eps then
begin detbnd:=0; aux[1]:=1-k; aux[2]:=s; goto end end;
if f>nrw then w2:=w2-1; p[k]:=pk; if pk≠k then
begin v[pk]:=v[k]; pk:=pk-k;
ichvec(kk1, kk1+w2, pk×w1, a); det:=-det;
r:=m[mk+pk]; m[mk+pk]:=a[kk]; a[kk]:=r
end
else r:=a[kk]; det:=r×det; iw:=kk1; lw1:=f-k+mk;
for i:=mk+1 step 1 until lw1 do
begin m[i]:=s:=m[i]/r; iw:=iw+w1;
elmvec(iw, iw+w2, kk1-iw, a, a, -s)
end
end;
aux[1]:=1/min; detbnd:=det; aux[2]:=min;
end;
end detbnd;

```

Description mca 2120

detbnd:= determinant of the n -th order band matrix M having lw codiagonals to the left and rw to the right of the main diagonal, and which is given in array $a[1 : (lw + rw) \times (n - 1) + n]$ in such a way that the (i, j) -th element of M is $a[(lw + rw) \times (i - 1) + j]$ for $i = 1, \dots, n$ and $j = \max(1, i - lw), \dots, \min(n, i + rw)$. The values of the remaining elements of a are irrelevant. In array $aux[0:2]$, one must give a relative tolerance, $aux[0]$.

The upper-triangular band matrix U resulting from the Gaussian elimination is delivered in a such that the (i, j) -th element of U is $a[(lw + rw) \times (i - 1) + j]$ for $i = 1, \dots, n$ and $j = i, \dots, \min(n, i + lw + rw)$; the matrix L , of Gaussian multipliers, is delivered in array $m[1 : lw \times (n - 2) + 1]$ such that the i -th multiplier of the j -th step is $m[lw \times (j - 1) + i - j]$ for $j = 1, \dots, n - 1$ and $i = j + 1, \dots, \min(n, j + lw)$; the pivotal indices are delivered in integer array $p[1:n]$.

Moreover,

$aux[2]$:= minimum absolute value of "pivot / Euclidean norm of the corresponding row of M ";

$aux[1]$:= $1 / aux[2]$.

If, however, M is singular, then the Gaussian elimination is discontinued, $detbnd := 0$, $aux[1]$:= minus the previous step number, and $aux[2]$:= absolute value of the last (rejected) pivot / Euclidean norm of the corresponding row of M .

detbnd uses vecvec, elmvec and ichvec (chapter 20).

```

comment mca 2121;
procedure solbnd(a, n, lw, rw, m, p, b); value n, lw, rw;
integer n, lw, rw; integer array p; array a, b, m;
begin integer f, i, k, kk, w, w1, w2, shift;
  real s;
  f:= lw; shift:= - lw; w1:= lw - 1;
  for k:= 1 step 1 until n do
  begin if f < n then f:= f + 1; shift:= shift + w1; i:= p[k];
    s:= b[i]; if i ≠ k then
    begin b[i]:= b[k]; b[k]:= s end;
    elmvec(k + 1, f, shift, b, m, - s)
  end;
  w1:= lw + rw; w:= w1 + 1; kk:= (n + 1) × w - w1; w2:= - 1;
  shift:= n × w1;
  for k:= n step - 1 until 1 do
  begin kk:= kk - w; shift:= shift - w1;
    if w2 < w1 then w2:= w2 + 1;
    b[k]:= (b[k] - vecvec(k + 1, k + w2, shift, b, a)) / a[kk]
  end
end solbnd;

```

Description mca 2121

solbnd should be called after detbnd (but only if the determinant is not zero), and solves the linear system $Mx = b$, where M is the n -th order band matrix having lw codiagonals to the left and rw to the right of the main diagonal, and whose Gaussian-eliminated form and pivotal row indices, as produced by detbnd, are given in array $a[1 : (lw + rw) \times (n - 1) + n]$, $m[1 : lw \times (n - 2) + 1]$ and integer array $p[1:n]$, and where b is the vector given as array $b[1:n]$. The solution vector x is overwritten on b . solbnd leaves a , m and p invariant, so that, after one call of detbnd, several calls of solbnd may follow for solving several systems having the same band matrix but different right-hand sides. solbnd uses vecvec and elmvec (chapter 20).

```

comment mca 2122;
real procedure detsolbnd(a, n, lw, rw, aux, b); value n, lw, rw;
integer n, lw, rw; array a, b, aux;
begin integer i, j, k, kk, kk1, pk, ik, lw1, f, q, w, w1, w2, iw,
nrw, shift;
real r, s, norm, eps, min, det; array m[0:lw], v[1:n];
f:= lw; det:= 1; w1:= lw + rw; w:= w1 + 1; w2:= w - 2; iw:= 0;
nrw:= n - rw; lw1:= lw + 1; q:= lw - 1;
for i:= 2 step 1 until lw do
begin q:= q - 1; iw:= iw + w1;
for j:= iw - q step 1 until iw do a[j]:= 0
end;
norm:= 0; iw:= - w2; q:= nrw + w - 1; j:= rw - 1;
for i:= 1 step 1 until n do
begin iw:= iw + w; if i < lw1 then iw:= iw - 1;
r:= v[i]:= sqrt(vecvec(iw, iw + (if i < lw then j + i else
if i > nrw then q - i else w1), 0, a, a));
if r > norm then norm:= r
end;
eps:= aux[0]; min:= 1; kk:= - w1;
if f > nrw then w2:= w2 + nrw - f;
for k:= 1 step 1 until n do
begin if f < n then f:= f + 1; ik:= kk:= kk + w;
s:= abs(a[kk]) / v[k]; pk:= k; kk1:= kk + 1;
for i:= k + 1 step 1 until f do
begin ik:= ik + w1; m[i - k]:= r:= a[ik]; a[ik]:= 0;
r:= abs(r) / v[i]; if r > s then
begin s:= r; pk:= i end
end;
if s < min then min:= s; if s < eps then
begin detsolbnd:= 0; aux[1]:= 1 - k; aux[2]:= s; goto end
end;
if f > nrw then w2:= w2 - 1; if pk ≠ k then
begin v[pk]:= v[k]; r:= b[k]; b[k]:= b[pk]; b[pk]:= r;
pk:= pk - k; ichvec(kk1, kk1 + w2, pk × w1, a);
det:= - det; r:= m[pk]; m[pk]:= a[kk]; a[kk]:= r
end
else r:= a[kk]; det:= r × det; iw:= kk1; lw1:= f - k;
for i:= 1 step 1 until lw1 do
begin m[i]:= s:= m[i] / r; iw:= iw + w1;
elmvec(iw, iw + w2, kk1 - iw, a, a, - s);
b[k + i]:= b[k + i] - b[k] × s
end
end;
aux[1]:= 1 / min; detsolbnd:= det; aux[2]:= s;
kk:= (n + 1) × w - w1; w2:= - 1; shift:= n × w1;
for k:= n step - 1 until 1 do
begin kk:= kk - w; shift:= shift - w1;
if w2 < w1 then w2:= w2 + 1;
b[k]:= (b[k] - vecvec(k + 1, k + w2, shift, b, a)) / a[kk]
end;
end;
end detsolbnd;

```

Description mca 2122

detsolbnd:= determinant of the n -th order band matrix M having lw codiagonals to the left and rw to the right of the main diagonal, and which is given in array $a[1 : (lw + rw) \times (n - 1) + n]$ in such a way that the (i, j) -th element of M is $a[(lw + rw) \times (i - 1) + j]$ for $i = 1, \dots, n$ and $j = \max(1, i - lw), \dots, \min(n, i + rw)$. The values of the remaining elements of a are irrelevant. In array $aux[0:2]$, one must give a relative tolerance, $aux[0]$.

The solution vector x of the linear system $Mx = b$, where b is the vector given as array $b[1:n]$, is calculated and overwritten on b . The upper-triangular band matrix U resulting from the Gaussian elimination is overwritten on a (in the same way as in detbnd).

Moreover,

$aux[2]$:= minimum absolute value of "pivot / Euclidean norm of the corresponding row of M ";

$aux[1]$:= $1 / aux[2]$.

If, however, M is singular, then the Gaussian elimination is discontinued, $detbnd := 0$, $aux[1]$:= minus the previous step number, $aux[2]$:= absolute value of the last (rejected) pivot / Euclidean norm of the corresponding row of M , and no solution vector is calculated. detsolbnd uses vecvec, elmvec and ichvec (chapter 20).

CHAPTER 22

POSITIVE DEFINITE SYMMETRIC LINEAR SYSTEMS AND MATRIX INVERSION.

This chapter contains procedures for solving systems of linear equations and for matrix inversion, provided the matrix is positive definite symmetric. Moreover section 221 contains procedures for calculating the rank and solving a homogeneous system whose matrix is positive definite symmetric, and section 224 contains procedures for solving linear least squares problems.

In sections 220 and 222 (the latter section deals with band matrices) the ordinary Cholesky method is used and in section 221 Cholesky with pivoting along the main diagonal. The latter method is indispensable for determining the rank of singular positive semidefinite symmetric matrices and for solving homogeneous systems.

Section 224 uses Householder transformations with pivoting [10].

For large order n , the computation time for solving linear systems and for matrix inversion is proportional to n cubed, and about one half of the time required for the general case (see chapter 21).

Pivoting along the main diagonal requires extra computation time which is proportional to n squared, and, thus, small with respect to the total time for (very) large n .

The procedures exist in two versions; one version uses the upper triangle of a two-dimensional array for the matrix and the other a one-dimensional array, so that, in the latter case, the memory space occupied by the matrix is cut nearly in half [8]. In the one-dimensional array, the elements of the upper triangle of the matrix are equidistant in the columns but not in the rows, so that special procedures for handling these rows are needed.

In the MC ALGOL 60 system for the X8, a large positive definite symmetric matrix given in a two-dimensional array is inverted faster than one given in a one-dimensional array, because the procedures mca 2000 to 2005 are available in machine-code, but mca 2006 is not. A similar statement holds for solving many (at least $n/2$, say) linear systems having the same positive definite symmetric matrix, but different right-hand sides.

The procedures of section 222, for solving linear systems with positive definite symmetric band matrices, save a considerable amount of computation time and memory space, if the band width is much smaller than n .

For large matrices, the computation time is proportional to $n \times$ the square of the band width.

Section 220 Cholesky decomposition without pivoting

This section contains procedures for solving linear systems and for inverting matrices, provided the matrices are positive definite symmetric:

detsolsym2 and detsolsym1 solve a system of linear equations and calculate the determinant of the system;

detinvsym2 and detinvsym1 calculate the determinant and inverse of a matrix;

detsym2 and detsym1 calculate the determinant of a matrix.

The other procedures of this section are to be used in combination with detsym2 or detsym1 for solving a linear system (or several systems having the same matrix but different right-hand sides) or for inverting a matrix.

The method used is Cholesky's square-root method without pivoting [2, p.117] [3, p.229] [4] [5] [8]. If the given symmetric matrix M is positive definite, then the method yields an upper-triangular matrix U , the "Cholesky matrix" of M , such that $U'U$ equals M ; moreover, the determinant of M is delivered, calculated as the product of the squares of the diagonal elements of U (and, thus, always positive). The process is completed in n stages, each stage producing a row of U . However, the process is discontinued if at some stage, k , the k -th diagonal element of M minus the sum of the squared elements of the k -th column of U (the sqrt of this quantity being the k -th diagonal element of U) is not positive, meaning that M , perhaps modified by rounding errors, is not positive definite. In that case, instead of the determinant, minus the last stage number k is delivered.

The solution of the linear system $U'Ux = b$ is obtained by solving $U'y = b$ (forward substitution) and $Ux = y$ (back substitution).

The inverse, X , of $U'U$ is obtained from the condition that UX be a lower-triangular matrix whose main diagonal elements are the reciprocals of the diagonal elements of U [4, p. 34-38].

The procedures mca 2200 - 2204 use the upper triangle of a two-dimensional array $a[1:n, 1:n]$ in which the upper triangle of M or U must be given and the upper triangle of U or X is delivered. Thus, $a[i, j]$ is the (i, j) -th element of the matrix only for $i \leq j$. The elements $a[i, j]$ for $i > j$ are neither used nor changed.

The procedures mca 2205 - 2209 use a one-dimensional array $a[1 : (n + 1) \times n : 2]$ in which the upper triangle of M or U must be given and the upper triangle of U or X is delivered in such a way that the (i, j) -th element of the matrix is $a[(j - 1) \times j : 2 + i]$ for $1 \leq i \leq j \leq n$.

```

comment mca 2200;
real procedure detsym2(a, n); value n; integer n; array a;
begin integer k, j; real r, d;
  d:= 1;
  for k:= 1 step 1 until n do
    begin r:= a[k,k] - tammat(1, k - 1, k, k, a, a); if r < 0 then
      begin detsym2:= - k; goto end end;
      d:= r × d; a[k,k]:= r:= sqrt(r);
      for j:= k + 1 step 1 until n do a[k,j]:= (a[k,j] - tammat(1,
        k - 1, j, k, a, a)) / r
    end;
    detsym2:= d;
  end;
end detsym2;

```

```

comment mca 2201;
procedure solsym2(a, n, b); value n; integer n; array a, b;
begin integer i;
  for i:= 1 step 1 until n do b[i]:= (b[i] - tamvec(1, i - 1, i,
    a, b)) / a[i,i];
  for i:= n step - 1 until 1 do b[i]:= (b[i] - matvec(i + 1, n, i,
    a, b)) / a[i,i]
end solsym2;

```

```

comment mca 2202;
real procedure detsolsym2(a, n, b); value n; integer n; array a, b;
begin real det;
  detsolsym2:= det:= detsym2(a, n);
  if det > 0 then solsym2(a, n, b)
end detsolsym2;

```

```

comment mca 2203;
procedure invsym2(a, n); value n; integer n; array a;
begin real r; integer i, j, i1; array u[1:n];
  for i:= n step - 1 until 1 do
    begin r:= 1 / a[i,i]; i1:= i + 1;
      for j:= i1 step 1 until n do u[j]:= a[i,j];
      for j:= n step - 1 until i1 do a[i,j]:= - (tamvec(i1, j, j,
        a, u) + matvec(j + 1, n, j, a, u)) × r;
      a[i,i]:= (r - matvec(i1, n, i, a, u)) × r
    end
  end invsym2;

```

```

comment mca 2204;
real procedure detinvsym2(a, n); value n; integer n; array a;
begin real det;
  detinvsym2:= det:= detsym2(a, n); if det ≥ 0 then invsym2(a, n)
end detinvsym2;

```

Description mca 2200

detsym2:= determinant of the n -th order positive definite symmetric matrix M whose upper triangle is given in array $a[1:n, 1:n]$. Moreover, the Cholesky matrix of M is calculated and overwritten on the upper triangle of a .

If, however, M is not positive definite, the Cholesky decomposition is discontinued and detsym2:= minus the last stage number.

detsym2 uses tamm (mca 2004).

Description mca 2201

solsym2 solves the n -th order linear system $U^*Ux = b$, where U is the upper-triangular matrix, given in the upper triangle of array $a[1:n, 1:n]$, and b is the vector given as array $b[1:n]$.

The solution vector x is overwritten on b .

If U is the Cholesky matrix of a positive definite symmetric matrix M , as produced by detsym2, then the calculated solution vector x is the solution of the linear system $Mx = b$.

solsym2 leaves the elements of a invariant, so that after one call of detsym2 several calls of solsym2 may follow for solving several linear systems having the same matrix but different right-hand sides. solsym2 uses matvec and tamvec (section 200).

Description mca 2202

detsolsym2:= determinant of the n -th order positive definite symmetric matrix M whose upper-triangle is given in array $a[1:n, 1:n]$.

Moreover, detsolsym2 solves the linear system $Mx = b$, where the vector b is given as array $b[1:n]$. The solution vector x is overwritten on b and the Cholesky matrix of M is overwritten on the upper triangle of a . If, however, M is not positive definite, then the Cholesky decomposition is discontinued, no solution is calculated, and detsolsym2:= minus the last stage number.

detsolsym2 uses detsym2, solsym2 and, indirectly, also matvec, tamvec and tamm (section 200).

Description mca 2203

invsym2 calculates the inverse, X , of the matrix U^*U , where U is the upper-triangular matrix given in the upper triangle of

array $a[1:n, 1:n]$. The upper triangle of X is overwritten on a .

invsym2 uses matvec and tamvec (section 200).

Description mca 2204

detinvsym2:= determinant of the n -th order positive definite symmetric matrix M whose upper triangle is given in array $a[1:n, 1:n]$.

Moreover, the upper triangle of the inverse of M is calculated and overwritten on a .

If, however, M is not positive definite, the Cholesky decomposition is discontinued and detinvsym2:= minus the last stage number.

detinvsym2 uses detsym2, invsym2 and, indirectly, also matvec, tamvec and tamm (section 200).

44

```
comment mca 2205;  
real procedure detsym1(a, n); value n; integer n; array a;  
begin integer i, j, k, kk, kj, low, up;  
  real d, r;  
  d:= 1; kk:= 0;  
  for k:= 1 step 1 until n do  
    begin kk:= kk + k; low:= kk - k + 1; up:= kk - 1;  
      r:= a[kk] - vecvec(low, up, 0, a, a); if r <= 0 then  
        begin detsym1:= - k; goto end end;  
      d:= d × r; a[kk]:= r:= sqrt(r); kj:= kk + k;  
      for j:= k + 1 step 1 until n do  
        begin a[kj]:= (a[kj] - vecvec(low, up, kj - kk, a, a)) / r;  
          kj:= kj + j  
        end  
      end  
    end;  
  detsym1:= d;  
end;  
end detsym1;
```

```
comment mca 2206;  
procedure solsym1(a, n, b); value n; integer n; array a, b;  
begin integer i, ii;  
  ii:= 0;  
  for i:= 1 step 1 until n do  
    begin ii:= ii + i;  
      b[i]:= (b[i] - vecvec(1, i - 1, ii - i, b, a)) / a[ii]  
    end;  
  for i:= n step - 1 until 1 do  
    begin b[i]:= (b[i] - seqvec(i + 1, n, ii + i, 0, a, b)) / a[ii];  
      ii:= ii - i  
    end  
  end solsym1;
```

```
comment mca 2207;  
real procedure detsolsym1(a, n, b); value n; integer n; array a, b;  
begin real det;  
  detsolsym1:= det:= detsym1(a, n);  
  if det > 0 then solsym1(a, n, b)  
end detsolsym1;
```

Description mca 2205

detsym1 := determinant of the n -th order positive definite symmetric matrix M whose upper triangle is given in array $a[1 : (n + 1) \times n : 2]$.

Moreover, the Cholesky matrix of M is calculated and overwritten on a . If, however, M is not positive definite, the Cholesky decomposition is discontinued and detsym1 := minus the last stage number.

detsym1 uses vecvec (mca 2000).

Description mca 2206

solsym1 solves the n -th order linear system $U'Ux = b$, where U is an upper-triangular matrix, given in array $a[1 : (n + 1) \times n : 2]$ and b is given as array $b[1:n]$.

The solution vector x is overwritten on b .

If U is the Cholesky matrix of a positive definite symmetric matrix M , as produced by detsym1, then the calculated solution vector x is the solution of the linear system $Mx = b$.

solsym1 leaves the elements of a invariant, so that after one call of detsym1 several calls of solsym1 may follow for solving several linear systems having the same matrix but different right-hand sides.

solsym1 uses vecvec and seqvec (section 200).

Description mca 2207

detsolsym1 := determinant of the n -th order positive definite symmetric matrix M whose upper triangle is given in array $a[1 : (n + 1) \times n : 2]$.

Moreover, detsolsym1 solves the linear system $Mx = b$, where the vector b is given as array $b[1:n]$.

The solution vector x is overwritten on b and the Cholesky matrix of M is overwritten on a .

If, however, M is not positive definite, then the Cholesky decomposition is discontinued, no solution is calculated, and detsolsym1 := minus the last stage number.

detsolsym1 uses detsym1, solsym1 and, indirectly, also vecvec and seqvec (section 200).

```

comment mca 2208;
procedure invsym1(a, n); value n; integer n; array a;
begin integer i, ii, i1, j, ij, jj;
    real r;
    array u[1:n];
    ii := (n + 1) × n : 2;
    for i := n step - 1 until 1 do
    begin r := 1 / a[ii]; i1 := i + 1; ij := ii + i;
        for j := i1 step 1 until n do
        begin u[j] := a[ij]; ij := ij + j end;
        for j := n step - 1 until i1 do
        begin jj := ij - i; ij := ij - j;
            a[ij] := - (vecvec(i1, j, jj - j, u, a) + seqvec(j + 1,
                n, jj + j, 0, a, u)) × r
        end;
        a[ii] := (r - seqvec(i1, n, ii + 1, 0, a, u)) × r; ii := ii - i
    end
end invsym1;

```

```

comment mca 2209;
real procedure detinvsym1(a, n); value n; integer n; array a;
begin real det;
    detinvsym1 := det := detsym1(a, n); if det ≥ 0 then invsym1(a, n)
end detinvsym1;

```

Description mca 2208

`invsym1` calculates the inverse, X , of the matrix $U'U$, where U is an upper-triangular matrix, given in array $a[1 : (n + 1) \times n : 2]$. The upper triangle of X is overwritten on a . `invsym1` uses `vecvec` and `seqvec` (section 200).

Description mca 2209

`detinvsym1` := determinant of the n -th order positive definite symmetric matrix M whose upper triangle is given in array $a[1 : (n + 1) \times n : 2]$.

Moreover, the upper triangle of the inverse of M is calculated and overwritten on a .

If, however, M is not positive definite, then the Cholesky decomposition is discontinued and `detinvsym1` := minus the last stage number.

`detinvsym1` uses `detsym1`, `invsym1` and, indirectly, also `vecvec` and `seqvec` (section 200).

Section 221 Cholesky decomposition with pivoting

This section contains procedures for calculating the rank of matrices, for solving linear systems and for inverting matrices, provided the matrices are positive definite symmetric:

rnksym20 and rnksym10 calculate the rank of a matrix;

rnksolsym20 and rnksolsym10 moreover solve a linear system, and

rnkinvsym20 and rnkinvsym10 invert a matrix;

solsymhom solves a system of homogeneous linear equations.

The other procedures of this section are to be used in combination with rnksym20 or rnksym10 for solving a linear system (or several linear systems having the same matrix but different right-hand sides) or for inverting a matrix.

The method used is Cholesky's square-root method (see section 220) with pivoting along the main diagonal. If the given symmetric matrix M is positive semidefinite, then the method yields an upper-triangular matrix U , the "pivot-Cholesky matrix" of M , such that the product $U'U$ equals M with permuted rows and columns. If the rank, r , of M is smaller than the order n , then the last $n - r$ rows of U (nearly) vanish.

The process is performed in at most n stages. At the k -th stage, the k -th row and column are interchanged with the $p[k]$ -th row and column (thus preserving the symmetry), the k -th "pivotal index" $p[k]$ being chosen in such a way that the diagonal elements of U turn out to be monotonically nonincreasing, and then the k -th row of U is produced. The process is terminated if at some stage, k , the k -th pivot (i. e. the maximum diagonal element of the remaining submatrix of order $n - k + 1$, the sqrt of this quantity being the k -th diagonal element of U) is negative or smaller than some tolerance, viz. a given relative tolerance times the maximum diagonal element of M (the maximum diagonal element being equal to the maximum-norm of M , if M is positive semidefinite). If no such k exists, then n is delivered as the rank of M . Otherwise, the maximum absolute value of the elements of the remaining submatrix of order $n - k + 1$ is calculated. If this is smaller than twice the tolerance, then $k - 1$ is delivered as the rank of the positive semidefinite matrix M ; otherwise, M is apparently not positive semidefinite, and, instead of the rank, the value $-k$ is delivered.

The solution of the linear system $Mx = b$ is obtained by first interchanging the elements of b in the same way as the rows (and columns) of M , subsequently performing forward and back substitution (see section 220) and finally carrying out the same interchanges in reverse order ("reverse correspondence") on the elements of the solution vector.

The inverse of $U'U$ is obtained by calling invsym2 or invsym1 (section 220). Then, the inverse of M is obtained by interchanging the rows and columns in reverse correspondence with the interchanges of the rows and columns of M .

A homogeneous linear system $Mx = 0$, where M is an n -th order positive semidefinite symmetric matrix of rank r , is obtained as follows. Let V be the r -th order upper-triangular matrix consisting of the first r columns of the pivot-Cholesky matrix U of M and W the $r \times (n - r)$ matrix consisting of the remaining part of the first r rows of U (the other rows of U are negligible). First, the system $VY = -W$ is solved (back substitution). Then Y and the $(n - r)$ -th order identity matrix are combined to form a single $r \times n$ matrix; its rows are then interchanged in reverse correspondence with the interchanges of the rows and columns of M . The columns of the resulting matrix form a complete linearly independent set of solution vectors of the homogeneous system.

The procedures mca 2210 - 2214 and 221a use the upper triangle of a two-dimensional array $a[1:n, 1:n]$ in which the upper triangle of M or U must be given and the upper triangle of U or X is delivered. Thus $a[i, j]$ is the (i, j) -th element of the matrix only for $i \leq j$. The elements $a[i, j]$ for $i > j$ are neither used nor changed. (Only mca 221a delivers a rectangular matrix involving some elements in the lower triangle of a as well.)

The procedures mca 2215 - 2219 use a one-dimensional array $a[1 : (n + 1) \times n : 2]$ in which the upper triangle of M or U must be given and the upper triangle of U or X is delivered, in such a way that the (i, j) -th element of the matrix is $a[(j - 1) \times j : 2 + i]$ for $1 \leq i \leq j \leq n$.

```

comment mca 2210;
integer procedure rnksym20(a, n, p, aux); value n; integer n;
integer array p; array a, aux;
begin integer k, i, j, pk; real w, max, m, t, r, d, norm, epsnorm;
  d:= 1; norm:= 0;
  for i:= 1 step 1 until n do if a[i,i] > norm then norm:= a[i,i];
  epsnorm:= aux[0] X norm; aux[1]:= norm; m:= 0;
  for k:= 1 step 1 until n do
  begin max:= epsnorm;
    for j:= k step 1 until n do if a[j,j] > max then
    begin max:= a[j,j]; pk:= j end;
    if max < epsnorm then
    begin for i:= k step 1 until n do
      begin t:= abs(a[i,i]); if t > m then m:= t;
        for j:= i + 1 step 1 until n do
        begin t:= a[i,j]:= a[i,j] - tammat(1, k - 1, i, j,
          a, a); t:= abs(t); if t > m then m:= t
        end
      end;
    end;
    goto end
  end;
  pk:= pk; d:= d X max; if pk ≠ k then
  begin ichcol(1, k - 1, k, pk, a);
    ichrowcol(k + 1, pk - 1, k, pk, a);
    ichrow(pk + 1, n, k, pk, a); a[pk,pk]:= a[k,k]
  end;
  a[k,k]:= r:= sqrt(max);
  for j:= k + 1 step 1 until n do
  begin w:= a[k,j]:= (a[k,j] - tammat(1, k - 1, k, j, a, a)) /
    r; a[j,j]:= a[j,j] - w X w
  end
  end;
  k:= n + 1;
end: aux[2]:= m; aux[3]:= d;
rnksym20:= if m ≤ 2 X epsnorm then k - 1 else - k
end rnksym20;

```

```

comment mca 2211;
procedure solsym20(a, n, p, b); value n; integer n; integer array p;
array a, b;
begin integer i, pi; real r;
  for i:= 1 step 1 until n do
  begin r:= b[i]; pi:= p[i];
    b[i]:= (b[pi] - tamvec(1, i - 1, i, a, b)) / a[i,i];
    if pi ≠ i then b[pi]:= r
  end;
  for i:= n step - 1 until 1 do b[i]:= (b[i] - matvec(i + 1, n, i,
  a, b)) / a[i,i];
  for i:= n step - 1 until 1 do
  begin pi:= p[i]; if pi ≠ i then
  begin r:= b[i]; b[i]:= b[pi]; b[pi]:= r end
  end
end solsym20;

```

Description mca 2210

rnksym20:= rank, r , of the n -th order positive semi-definite symmetric matrix M whose upper triangle is given in array $a[1:n, 1:n]$.

In array $aux[0:3]$, one must give a relative tolerance, $aux[0]$.

The pivot-Cholesky matrix of M is overwritten on the upper triangle of a , and the pivotal indices are delivered in integer array $p[1:r]$.

Moreover,

$aux[1]$:= the maximum diagonal element of M ;

$aux[2]$:= the maximum absolute value of the elements of the remaining submatrix of order $n - r$ if $r < n$, and otherwise 0;

$aux[3]$:= determinant of the principal submatrix of order r .

However, if M is not positive semidefinite, then $rnksym20$:= minus the last stage number.

$rnksym20$ uses $tammat$, $ichcol$, $ichrow$ and $ichrowcol$ (chapter 20).

Description mca 2211

solsym20 should be called after $rnksym20$ (but only if the rank equals n), and solves the n -th order linear system $Mx = b$, where M is the positive definite symmetric matrix whose pivot-Cholesky matrix and pivotal indices, as produced by $rnksym20$, are given in the upper triangle of array $a[1:n, 1:n]$ and in integer array $p[1:n]$, and where b is the vector given as array $b[1:n]$.

The solution vector x is overwritten on b .

$solsym20$ leaves a and p invariant, so that after one call of $rnksym20$ several calls of $solsym20$ may follow for solving several linear systems having the same matrix but different right-hand sides.

$solsym20$ uses $matvec$ and $tamvec$ (section 200).

```

comment mca 2212;
integer procedure rnksolsym20(a, n, b, aux); value n; integer n;
array a, b, aux;
begin integer rank;
  integer array p[1:n];
  rnksolsym20:= rank:= rnksym20(a, n, p, aux);
  if rank = n then solsymb20(a, n, p, b)
end rnksolsym20;

```

```

comment mca 2213;
procedure invsym20(a, n, p); value n; integer n; integer array p;
array a;
begin integer i, j, pi;
  real r;
  invsym2(a, n);
  for i:= n step - 1 until 1 do
    begin pi:= p[i]; if pi ≠ i then
      begin ichcol(1, i - 1, i, pi, a);
        ichrowcol(i + 1, pi - 1, i, pi, a);
        ichrow(pi + 1, n, i, pi, a); r:= a[i,i];
        a[i,i]:= a[pi,pi]; a[pi,pi]:= r
      end
    end
  end invsym20;

```

```

comment mca 2214;
integer procedure rnkinvsym20(a, n, aux); value n; integer n;
array a, aux;
begin integer rank;
  integer array p[1:n];
  rnkinvsym20:= rank:= rnksym20(a, n, p, aux);
  if rank = n then invsym20(a, n, p)
end rnkinvsym 20;

```

Description mca 2212

`rnksolsym20` := rank, r , of the n -th order positive semidefinite symmetric matrix M whose upper triangle is given in array `a[1:n, 1:n]`. In array `aux[0:3]` one must give a relative tolerance, `aux[0]`.

If $r = n$, the linear system $Mx = b$ is solved, where b is the vector given as array `b[1:n]`, and the solution vector x is overwritten on b . The pivot-Cholesky matrix of M is overwritten on the upper triangle of `a`. Moreover,

`aux[1]` := the maximum diagonal element of M ;

`aux[2]` := 0;

`aux[3]` := determinant of M .

However, if $1 < r < n$ (M positive semidefinite) or $r < 0$ (M not positive semidefinite), then no solution vector is calculated and the results of `rnksolsym20` are the same as those of `rnksym20`.

`rnksolsym20` uses `rnksym20`, `solsym20` and, indirectly, also `matvec`, `tamvec`, `tammat`, `ichcol`, `ichrow` and `ichrowcol` (chapter 20).

Description mca 2213

`invsym20` should be called after `rnksym20` (but only if the rank equals n), and calculates the inverse, X , of the n -th order positive definite symmetric matrix M whose pivot-Cholesky matrix and pivotal indices, as produced by `rnksym20`, are given in the upper triangle of array `a[1:n, 1:n]` and in integer array `p[1:n]`.

The upper triangle of X is overwritten on `a`.

`invsym20` uses `invsym2` (mca 2203), `ichcol`, `ichrow`, `ichrowcol` (section 202) and, indirectly, also `matvec` and `tamvec` (section 200).

Description mca 2214

`rnkinvsym20` := rank, r , of the n -th order positive semi-definite symmetric matrix M whose upper triangle is given in array `a[1:n, 1:n]`. In array `aux[0:3]` one must give a relative tolerance, `aux[0]`.

If $r = n$, the upper triangle of the inverse of M is calculated and overwritten on `a`.

Moreover,

`aux[1]` := the maximum diagonal element of M ;

`aux[2]` := 0;

`aux[3]` := determinant of M .

However, if $1 < r < n$ (M positive semidefinite) or $r < 0$ (M not positive semidefinite), then no inverse is calculated and the results of `rnkinvsym20` are the same as those of `rnksym20`.

`rnkinvsym20` uses `rnksym20`, `invsym20` and, indirectly, also `invsym2` (mca 2203), `matvec`, `tamvec`, `tammat`, `ichcol`, `ichrow` and `ichrowcol` (chapter 20).

54

```
comment mca 221a;  
integer procedure solsymhom20(a, n, aux); value n; integer n;  
array a, aux;  
begin integer i, pj, j, rank;  
  real r;  
  integer array p[1:n];  
  solsymhom20:= rank:= rnksym20(a, n, p, aux); if rank  $\geq$  0 then  
  begin for i:= rank + 1 step 1 until n do  
    for j:= rank + 1 step 1 until n do a[i,j]:= if i = j then 1  
    else 0;  
    for i:= rank step - 1 until 1 do  
    begin r:= - a[i,i];  
      for j:= rank + 1 step 1 until n do a[i,j]:= (a[i,j] +  
        matmat(i + 1, rank, i, j, a, a)) / r  
    end;  
    for j:= rank step - 1 until 1 do  
    begin pj:= p[j]; if pj  $\neq$  j then ichrow(rank + 1, n, j, pj, a)  
    end  
  end  
end solsymhom20;
```

Description mca 221a

`solsymhom20` solves the homogeneous linear system whose n -th order positive semidefinite symmetric matrix M is given in the upper triangle of array `a[1:n, 1:n]`.

In array `aux[0:3]` one must give a relative tolerance, `aux[0]`.

`solsymhom20:=rank, r`, of M , calculated by means of `rnksym20`.

Subsequently, a complete set of $n - r$ linearly independent solution vectors of the homogeneous linear system is calculated and delivered in the last $n - r$ columns of `a`. The first r columns of the pivot-Cholesky matrix of M are delivered in the first r columns of `a`.

Moreover, the same results are delivered in `aux` as by `rnksym20`.

However, if `rnksym20` delivers a negative (integral) value, indicating that M is not positive semidefinite, no solution of the homogeneous system is calculated and only the results of `rnksym20` are delivered.

`solsymhom20` uses `rnksym20`, `matmat`, `ichrow` and, indirectly, also `tammat`, `ichcol` and `ichrowcol` (chapter 20).


```

comment mca 2215;
integer procedure rnksym10(a, n, p, aux); value n; integer n;
integer array p; array a, aux;
begin integer k, pk, kk, kj, pp, i, j, jj, t, low, up;
real norm, epsnorm, m, max, d, r, w;
d:= 1; norm:= 0; kk:= 0;
for k:= 1 step 1 until n do
begin kk:= kk + k; if a[kk] > norm then norm:= a[kk] end;
epsnorm:= aux[0] × norm; aux[1]:= norm; m:= 0; kk:= 0;
for k:= 1 step 1 until n do
begin max:= epsnorm; t:= kk;
for j:= k step 1 until n do
begin t:= t + j; if a[t] > max then
begin max:= a[t]; pk:= j; pp:= t end
end;
if max < epsnorm then
begin for i:= k step 1 until n do
begin kk:= kk + i; low:= kk - i + 1; up:= low + k - 2;
r:= abs(a[kk]); if r > m then m:= r; kj:= kk + i;
for j:= i + 1 step 1 until n do
begin r:= a[kj]:= a[kj] - vecvec(low, up, kj - kk,
a, a); r:= abs(r); if r > m then m:= r;
kj:= kj + j
end
end;
goto end
end;
kk:= kk + k; low:= kk - k + 1; up:= kk - 1; p[k]:= pk;
d:= d × max; if pk ≠ k then
begin ichvec(low, up, pp - pk - kk + k, a);
ichseqvec(k + 1, pk - 1, kk + k, pp - pk, a);
ichseq(pk + 1, n, pp + k, pk - k, a); a[pp]:= a[kk]
end;
a[kk]:= r:= sqrt(max); kj:= kk + k; jj:= kk;
for j:= k + 1 step 1 until n do
begin w:= a[kj]:= (a[kj] - vecvec(low, up, kj - kk, a, a)) /
r; jj:= jj + j; a[jj]:= a[jj] - w × w; kj:= kj + j
end
end;
k:= n + 1;
end: aux[2]:= m; aux[3]:= d;
rnksym10:= if m < 2 × epsnorm then k - 1 else - k
end rnksym10;

```

Description mca 2215

rnksym10:= rank, r , of the n -th order positive semidefinite symmetric matrix M whose upper triangle is given in array $a[1 : (n + 1) \times n : 2]$. In array $aux[0:3]$ one must give a relative tolerance, $aux[0]$.

The pivot-Cholesky matrix of M is overwritten on a and the pivotal indices are delivered in integer array $p[1:r]$.

Moreover,

$aux[1]$:= the maximum diagonal element of M ;

$aux[2]$:= the maximum absolute value of the elements of the remaining submatrix of order $n - r$ if $r < n$, and otherwise 0;

$aux[3]$:= determinant of the principal submatrix of order r .

However, if M is not positive semidefinite, then rnksym10:= minus the last stage number.

rnksym10 uses vecvec, ichvec, ichseqvec and ichseq (chapter 20).

```

comment mca 2216;
procedure solsym10(a, n, p, b); value n; integer n; array a, b;
integer array p;
begin integer i, ii, pi; real s;
  ii:= 0;
  for i:= 1 step 1 until n do
  begin s:= b[i]; pi:= p[i]; ii:= ii + i;
    b[i]:= (b[pi] - vecvec(1, i - 1, ii - i, b, a)) / a[ii];
    if pi  $\neq$  i then b[pi]:= s
  end;
  for i:= n step - 1 until 1 do
  begin b[i]:= (b[i] - seqvec(i + 1, n, ii + i, 0, a, b)) / a[ii];
    ii:= ii - i
  end;
  for i:= n step - 1 until 1 do
  begin pi:= p[i]; if pi  $\neq$  i then
    begin s:= b[i]; b[i]:= b[pi]; b[pi]:= s end
  end
end solsym10;

```

```

comment mca 2217;
integer procedure rnksolsym10(a, n, b, aux); value n; integer n;
array a, b, aux;
begin integer rank; integer array p[1:n];
  rnksolsym10:= rank:= rnksym10(a, n, p, aux);
  if rank = n then solsym10(a, n, p, b)
end rnksolsym10;

```

```

comment mca 2218;
procedure invsym10(a, n, p); value n; integer n; integer array p;
array a;
begin integer i, ii, pi, pp; real r;
  invsym1(a, n); ii:= (n + 1)  $\times$  n : 2;
  for i:= n step - 1 until 1 do
  begin pi:= p[i]; if pi  $\neq$  i then
    begin pp:= (pi + 1)  $\times$  pi : 2;
      ichvec(ii - i + 1, ii - 1, pp - pi - ii + i, a);
      ichseqvec(i + 1, pi - 1, ii + i, pp - pi, a);
      ichseq(pi + 1, n, pp + i, pi - i, a); r:= a[ii];
      a[ii]:= a[pp]; a[pp]:= r
    end;
    ii:= ii - i
  end
end invsym10;

```

```

comment mca 2219;
integer procedure rnkinvsym10(a, n, aux); value n; integer n;
array a, aux;
begin integer rank; integer array p[1:n];
  rnkinvsym10:= rank:= rnksym10(a, n, p, aux);
  if rank = n then invsym10(a, n, p)
end rnkinvsym 10;

```

Description mca 2216

`solsym10` should be called after `rnksym10` (but only if the rank equals n), and solves the n -th order linear system $Mx = b$, where M is the positive definite symmetric matrix whose pivot-Cholesky matrix and pivotal indices, as produced by `rnksym10`, are given in array `a[1 : (n + 1) × n : 2]` and in integer array `p[1:n]`, and where b is the vector given as array `b[1:n]`.

The solution vector x is overwritten on b .

`solsym10` leaves a and p invariant, so that after one call of `rnksym10` several calls of `solsym10` may follow for solving several linear systems having the same matrix but different right-hand sides. `solsym10` uses `vecvec` and `seqvec` (section 200).

Description mca 2217

`rnksolsym10:= rank, r`, of the n -th order positive semidefinite symmetric matrix M whose upper triangle is given in array `a[1 : (n + 1) × n : 2]`.

In array `aux[0:3]` one must give a relative tolerance, `aux[0]`.

If $r = n$, the linear system $Mx = b$ is solved, where b is the vector given as array `b[1:n]`, and the solution vector x is overwritten on b . The pivot-Cholesky matrix of M is overwritten on a .

Moreover, `aux[1]:=` the maximum diagonal element of M ; `aux[2]:= 0`; `aux[3]:=` determinant of M .

However, if $1 < r < n$ (M positive semidefinite) or $r < 0$ (M not positive semidefinite), then no solution vector is calculated and the results of `rnksolsym10` are the same as those of `rnksym10`.

`rnksolsym10` uses `rnksym10`, `solsym10` and, indirectly, also `vecvec`, `seqvec`, `ichvec`, `ichseqvec` and `ichseq` (chapter 20).

Description mca 2218

`invsym10` should be called after `rnksym10` (but only if the rank equals n), and then calculates the inverse, X , of the n -th order positive definite symmetric matrix M whose pivot-Cholesky matrix and pivotal indices, as produced by `rnksym10`, are given in array `a[1 : (n + 1) × n : 2]` and in integer array `p[1:n]`.

The upper triangle of X is overwritten on a .

`invsym10` uses `invsym1` (mca 2208), `ichvec`, `ichseqvec`, `ichseq` (section 202) and, indirectly, also `vecvec` and `seqvec` (section 200).

Description mca 2219

`rnkinvsym10:= rank, r`, of the n -th order positive semidefinite symmetric matrix M whose upper triangle is given in array `a[1 : (n + 1) × n : 2]`.

In array `aux[0:3]` one must give a relative tolerance, `aux[0]`.

If $r = n$, the upper triangle of the inverse of M is calculated and overwritten on a .

Moreover, `aux[1]:=` the maximum diagonal element of M ; `aux[2]:= 0`; `aux[3]:=` determinant of M .

However, if $1 < r < n$ (M positive semidefinite) or $r < 0$ (M not positive semidefinite), then no inverse is calculated and the results of `rnkinvsym10` are the same as those of `rnksym10`.

`rnkinvsym10` uses `rnksym10`, `invsym10` and, indirectly, also `invsym1` (mca 2208), `vecvec`, `seqvec`, `ichvec`, `ichseqvec` and `ichseq` (chapter 20).

Section 222 Cholesky decomposition for band matrices

The procedures of this section solve a system of linear equations and / or calculate the determinant of a matrix, provided the matrix is a positive definite symmetric band matrix:

detsolsymbnd solves a system of linear equations and calculates the determinant of the system;

detsymbnd calculates the determinant of a matrix;

solsymbnd solves a linear system whose matrix is given in the Cholesky-decomposed form produced by detsymbnd.

One call of detsymbnd followed by several calls of solsymbnd may be used to solve several linear systems having the same matrix but different right-hand sides.

The method used is Cholesky's square-root method without pivoting (see section 220 and [9]). If the given symmetric band matrix M is positive definite, then the method yields an upper-triangular band matrix U , the "Cholesky matrix" of M , such that $U'U$ equals M ; moreover, the determinant of M is delivered, calculated as the product of the squares of the diagonal elements of U (and, thus, always positive). The number of nonzero diagonals of U is the same as that of the upper triangle of M . The process is completed in n stages, each stage producing a row of U . However, the process is discontinued if at some stage, k , the k -th diagonal element of M minus the sum of the squared elements of the k -th column of U (the sqrt of this quantity being the k -th diagonal element of U) is not positive, meaning that M , perhaps modified by rounding errors, is not positive definite. In that case, instead of the determinant, minus the last stage number k is delivered.

The solution of the linear system $U'Ux = b$ is obtained by solving $U'y = b$ (forward substitution) and $Ux = y$ (back substitution).

The procedures of this section use a one-dimensional array $a[1 : (n - 1) \times w + n]$ for the upper triangle of M and U , where n is the order of the matrix and w the number of non-zero codiagonals above the main diagonal; the (i, j) -th element of matrix M or U is $a[(j - 1) \times w + i]$ for $j = 1, \dots, n$ and $i = \max(1, j - w), \dots, j$; the other elements of a are neither used nor changed.

```

comment mca 2220;
real procedure detsymbnd(a, n, w); value n, w; integer n, w; array a;
begin integer j, k, jmax, kk, kj, w1, start;
  real r, det;
  det:= 1; jmax:= w; w1:= w + 1; kk:= - w;
  for k:= 1 step 1 until n do
    begin if k + w > n then jmax:= jmax - 1; kk:= kk + w1;
      start:= kk - k + 1;
      r:= a[kk] - vecvec(if k < w1 then start else kk - w, kk -
        1, 0, a, a); if r < 0 then
        begin detsymbnd:= - k; goto end end;
      det:= r × det; a[kk]:= r:= sqrt(r); kj:= kk;
      for j:= 1 step 1 until jmax do
        begin kj:= kj + w;
          a[kj]:= (a[kj] - vecvec(if k + j < w1 then start else kk
            - w + j, kk - 1, kj - kk, a, a)) / r
        end
      end;
    detsymbnd:= det;
  end:
end detsymbnd;

```

```

comment mca 2221;
procedure solsymbnd(a, n, w, b); value n, w; integer n, w; array a, b;
begin integer i, k, imax, kk, w1;
  kk:= - w; w1:= w + 1;
  for k:= 1 step 1 until n do
    begin kk:= kk + w1;
      b[k]:= (b[k] - vecvec(if k < w1 then 1 else k - w, k - 1, kk
        - k, b, a)) / a[kk]
    end;
  imax:= - 1;
  for k:= n step - 1 until 1 do
    begin if imax < w then imax:= imax + 1;
      b[k]:= (b[k] - scaprd1(kk + w, w, k + 1, 1, imax, a, b)) /
        a[kk]; kk:= kk - w1
    end
  end
end solsymbnd;

```

```

comment mca 2222;
real procedure detsolsymbnd(a, n, w, b); value n, w; integer n, w;
array a, b;
begin real det;
  detsolsymbnd:= det:= detsymbnd(a, n, w);
  if det > 0 then solsymbnd(a, n, w, b)
end detsolsymbnd;

```

Description mca 2220

detsymbnd:= determinant of the n-th order positive definite symmetric band matrix M having w codiagonals on each side of the main diagonal, and whose upper triangle is given in array a[1 : (n - 1) × w + n]. Moreover, the Cholesky matrix of M is calculated and delivered in a. If, however, M is not positive definite, then the Cholesky decomposition is discontinued, and detsymbnd:= minus the last stage number. detsymbnd uses vecvec (mca 2000).

Description mca 2221

solsymbnd solves the n-th order linear system $U'Ux = b$, where b is the vector given as array b[1 : n], and U is the upper-triangular band matrix having w codiagonals, and which is given in array a[1 : (n - 1) × w + n].

The solution vector x is overwritten on b.

If U is the Cholesky matrix of a positive definite symmetric band matrix M, as produced by detsymbnd, then the calculated solution vector x is the solution of the linear system $Mx = b$.

solsymbnd leaves the elements of a invariant, so that after one call of detsymbnd several calls of solsymbnd may follow for solving several linear systems having the same matrix but different right-hand sides. solsymbnd uses vecvec and scaprd1 (section 200).

Description mca 2222

detsolsymbnd:= determinant of the n-th order positive definite symmetric band matrix M having w codiagonals on each side of the main diagonal, and whose upper triangle is given in array a[1 : (n - 1) × w + n].

Moreover, the solution vector x of the linear system $Mx = b$, where b is the vector given as array b[1:n], is calculated and overwritten on b, and the Cholesky matrix of M is delivered in a.

If, however, M is not positive definite, then the Cholesky decomposition is discontinued, no solution is calculated, and detsolsymbnd:= minus the last stage number.

detsolsymbnd uses detsymbnd, solsymbnd and, indirectly, also vecvec and scaprd1 (section 200).

Section 224 Least-squares problems

This section contains procedures for solving linear least-squares problems:

lsqdecsol calculates the solution x of a least-squares problem $Mx = b$ and, moreover, the main diagonal of the inverse of the product $M'M$; lsqdec performs the Householder triangularisation of M and calculates its rank;

lsqsol and lsqdglinv are to be used in combination with lsqdec for solving a linear least-squares problem (or several problems having the same matrix M but different right-hand sides) or for calculating the main diagonal of the inverse of $M'M$.

Apart from some changes and adaptations to our vector procedures, lsqdec and lsqsol have been derived from [10]. However, our procedures do not perform iterations for improving the solution; these iterations would be of limited value, as is pointed out in [11].

The method is Householder triangularisation with column interchanges. Let M have n rows and m columns; lsqdec produces an n -th order orthogonal matrix Q and an $n \times m$ upper-triangular matrix R such that R equals QM with permuted columns. Matrix Q is the product of at most m orthogonal symmetric n -th order "Householder matrices", which are of the form $I - sww'$, where I is the identity matrix, w a column vector and s a scalar. Matrix M is reduced to R in (at most) m stages. In the k -th stage, the desired zeroes are introduced in the k -th column of the matrix as follows: first the "pivotal" column, i. e. the column having maximum Euclidean norm, is selected from the remaining $(n - k + 1) \times (m - k + 1)$ submatrix, and the pivotal and the k -th columns are interchanged; then the k -th Householder matrix is calculated and postmultiplied by the remaining submatrix.

The k -th Householder matrix is chosen such that this postmultiplication introduces the desired zeroes in the k -th column, and the first $k - 1$ elements of w are zero.

If at some stage k the Euclidean norm of the pivotal column is smaller than some tolerance, viz. a given relative tolerance times the maximum of the Euclidean norms of the columns of M , then the process is discontinued, and $k - 1$ is delivered as the rank of M ; otherwise, the rank equals m .

In lsqsol, the least-squares solution x of the problem $Mx = b$ is obtained by first calculating $y = Qb$, then solving the triangular system consisting of the first m equations of $Rx = y$ (back substitution), and finally interchanging the elements of x in "reverse correspondence" with the interchanges of the columns of M , i. e. the same interchanges are carried out in reverse order. As by-product the last $n - m$ elements of y are delivered; the sum of the squares of these elements is approximately equal to the square of the Euclidean norm of the residue vector $Mx - b$.

In lsqdglinv, the main diagonal of $M'M$ is obtained by calculating the inverse of R , from this the main diagonal of the inverse of $R'R$, and then interchanging the calculated diagonal elements in reverse correspondence with the interchanges of the columns of M .


```

comment mca 2240;
integer procedure lsqdec(a, n, m, aux, aid, ci); value n, m;
integer n, m; array a, aux, aid; integer array ci;
begin integer j, k, kpiv;
  real beta, sigma, norm, w, eps, akk, aidk;
  array sum[1:m];
  norm:= 0; lsqdec:= m;
  for k:= 1 step 1 until m do
  begin w:= sum[k]:= tammat(1, n, k, k, a, a);
    if w > norm then norm:= w
  end;
  w:= aux[1]:= sqrt(norm); eps:= aux[0] × w;
  for k:= 1 step 1 until m do
  begin sigma:= sum[k]; kpiv:= k;
    for j:= k + 1 step 1 until m do if sum[j] > sigma then
    begin sigma:= sum[j]; kpiv:= j end;
    if kpiv ≠ k then
    begin sum[kpiv]:= sum[k]; ichcol(1, n, k, kpiv, a) end;
    ci[k]:= kpiv; akk:= a[k,k]; sigma:= tammat(k, n, k, k, a, a);
    w:= sqrt(sigma); aidk:= aid[k]:= if akk < 0 then w else -w;
    if w < eps then
    begin lsqdec:= k - 1; goto enddec end;
    beta:= 1 / (sigma - akk × aidk); a[k,k]:= akk - aidk;
    for j:= k + 1 step 1 until m do
    begin elmcol(k, n, j, k, a, a, -beta × tammat(k, n, k, j,
    a, a)); sum[j]:= sum[j] - a[k,j] ↑ 2
    end
  end for k;
enddec: aux[2]:= w
end lsqdec;

```

Description mca 2240

lsqdec:= rank, r , of the $n \times m$ matrix M given in array $a[1:n, 1:m]$.
In array $aux[0:2]$ one must give a relative tolerance, $aux[0]$.
The pivotal column indices are delivered in integer array $ci[1 : r]$,
the (r first) diagonal elements of the upper-triangular matrix R in
array $aid[1 : r]$, and the other elements of the upper triangle of R in
array a , together with the vectors w of the Householder matrices.
Moreover,
 $aux[1]$:= the maximum Euclidean norm of the columns of M ,
 $aux[2]$:= the absolute value of the r -th diagonal element of R .
lsqdec uses tamm, elmcol and ichcol (chapter 20).

```

comment mca 2241;
procedure lsqsol(a, n, m, aid, ci, b); value n, m; integer n, m;
array a, aid, b; integer array ci;
begin integer k, cik;
  real w;
  for k:= 1 step 1 until m do elmveccol(k, n, k, b, a, tamvec(k,
n, k, a, b) / (aid[k] × a[k,k]));
  for k:= m step - 1 until 1 do b[k]:= (b[k] - matvec(k + 1, m, k,
a, b)) / aid[k];
  for k:= m step - 1 until 1 do
  begin cik:= ci[k]; if cik ≠ k then
    begin w:= b[k]; b[k]:= b[cik]; b[cik]:= w end
  end
end lsqsol;

```

```

comment mca 2242;
procedure lsqdglinv(a, m, aid, ci, diag); value m; integer m;
array a, aid, diag; integer array ci;
begin integer j, k, cik;
  real w;
  for k:= 1 step 1 until m do
  begin diag[k]:= 1 / aid[k];
    for j:= k + 1 step 1 until m do diag[j]:= - tamvec(k, j - 1,
j, a, diag) / aid[j]; diag[k]:= vecvec(k, m, 0, diag, diag)
  end;
  for k:= m step - 1 until 1 do
  begin cik:= ci[k]; if cik ≠ k then
    begin w:= diag[k]; diag[k]:= diag[cik]; diag[cik]:= w end
  end
end lsqdglinv;

```

```

comment mca 2243;
integer procedure lsqdecsol(a, n, m, aux, diag, b); value n, m;
integer n, m; array a, aux, diag, b;
begin integer rank;
  array aid[1:m];
  integer array ci[1:m];
  rank:= lsqdecsol:= lsqdec(a, n, m, aux, aid, ci);
  if rank = m then
  begin lsqdglinv(a, m, aid, ci, diag); lsqsol(a, n, m, aid, ci, b)
  end
end lsqdecsol;

```

Description mca 2241

lsqsol should be called after lsqdec (but only if the rank equals m), and calculates the least-squares solution x of $Mx = b$, where b is the vector given as array $b[1:n]$, and M is the $n \times m$ matrix whose Householder-triangularised form R , with the vectors w of the Householder matrices and the pivotal indices, as produced by lsqdec, are given in array $a[1:n, 1:m]$, aid $[1:m]$ and integer array $ci[1:m]$. The solution vector x is overwritten on the first m elements of b , and the last $n - m$ elements of y are overwritten on the last $n - m$ elements of b .

lsqsol leaves the elements of a , aid and ci intact, so that, after one call of lsqdec, several calls of lsqsol may follow for solving several least-squares problems having the same matrix M but different right-hand sides b .

lsqsol uses matvec, tamvec and elmveccol (chapter 20).

Description mca 2242

lsqdglinv should be called after lsqdec (but only if the rank equals m), and calculates the main diagonal of the inverse of M^tM , where M is the matrix whose Householder-triangularised form R with the pivotal indices, as produced by lsqdec, are given in array $a[1:m, 1:m]$, aid $[1:m]$ and integer array $ci[1:m]$.

The calculated main diagonal is delivered in array $diag[1:m]$; the elements of a , aid and ci are left intact.

lsqdglinv uses vecvec and tamvec (section 200).

Description mca 2243

lsqdecsol:=rank, r, of the $n \times m$ matrix M given in array $a[1:n, 1:m]$. In array $aux[0:2]$ one must give a relative tolerance, aux $[0]$.

If $r = m$, then the least-squares solution x of $Mx = b$, where b is the vector given as array $b[1:n]$, is calculated and overwritten on the first m elements of b ; the last $n - m$ elements of y are overwritten on the last $n - m$ elements of b ; moreover, the main diagonal of the inverse of M^tM is delivered in array $diag[1:m]$.

However, if $r < m$, then no solution and main diagonal are calculated, and the elements of b and $diag$ are left unchanged.

In either case,

aux $[1]$:= the maximum Euclidean norm of the columns of M ,

aux $[2]$:= the absolute value of the r -th diagonal element of R , and the elements of a are altered.

lsqdecsol uses lsqdec, lsqsol, lsqdglinv and, indirectly also vecvec, matvec, tamvec, tammatt, elmveccol, elmcol and ichcol (chapter 20).

REFERENCES

1. P. Naur (ed.), Revised report on the algorithmic language ALGOL 60 (1962).
2. J. H. Wilkinson, Rounding errors in algebraic processes (London 1963).
3. J. H. Wilkinson, The algebraic eigenvalue problem (Oxford 1965).
4. L. Fox, Practical solution of linear equations and inversion of matrices, Nat. Bur. Stand. AMS 39 (edited by O. Taussky), p. 1 - 54 (1951).
5. T. J. Dekker, Evaluation of determinants, solution of systems of linear equations and matrix inversion (MR63, Mathematical Centre, Amsterdam, 1963).
6. H. C. Thacher, Crout with pivoting II, Alg. 43, Comm. ACM 4 (1961) p. 176.
7. J. Garwick, solution of a linear system with a band coefficient matrix, BIT 3 (1963) 207 - 208.
8. R. S. Martin, G. Peters and J. H. Wilkinson, Symmetric decomposition of a positive definite matrix, Num. Mat. 7 (1965) 362 - 383.
9. R. S. Martin and J. H. Wilkinson, Symmetric decomposition of positive definite band matrices, Num. Mat. 7 (1965) 355 - 361.
10. P. Businger and G. H. Golub, Linear least squares solution by Householder transformations, Num. Mat. 7 (1965) 269 - 276.
11. G. H. Golub and J. H. Wilkinson, Note on the iterative refinement of least squares solution, Num. Mat. 9 (1966) 139 - 148.
12. H. L. Oudshoorn, H. N. Glorie and G. C. J. M. Nogaerde, ALGOL editor (MR98, Mathematical Centre, Amsterdam, 1968).

APPENDIX

TIMES FOR THE MC ALGOL 60 SYSTEM FOR THE X8.

In this appendix we give practical formulas for the computation times in milliseconds of the procedures published above. The coefficients of these formulas have been obtained from tests on an Electrologica X8 computer using the MC ALGOL 60 system for the X8, in which system the procedures mca 2000 to 2005 are available as machine-code procedures. For comparison we moreover give the formulas for the computation times of the nonmachine-code ALGOL 60 procedures mca 2000 to 2005 and of the procedures of section 210 using them. The coefficients of the time formulas have a relative precision of at most one or two digits.

CHAPTER 20 VECTOR OPERATIONS

Here n is the number of elements used in each vector, thus, $n = u - 1 + 1$, except for scaprd1.

Section 200 Scalar products

| | machine-code | ALGOL 60 |
|------------------|-----------------------|----------------------|
| mca 2000 vecvec | $.085 \times n + 1.1$ | $.46 \times n + .9$ |
| mca 2001 matvec | $.085 \times n + 1.2$ | $.54 \times n + .9$ |
| mca 2002 tamvec | $.085 \times n + 1.2$ | $.54 \times n + .9$ |
| mca 2003 matmat | $.085 \times n + 1.4$ | $.63 \times n + 1.0$ |
| mca 2004 tammat | $.085 \times n + 1.4$ | $.63 \times n + 1.0$ |
| mca 2005 mattam | $.085 \times n + 1.4$ | $.63 \times n + 1.0$ |
| mca 2006 seqvec | | $.50 \times n + 1.0$ |
| mca 2008 scaprd1 | | $.53 \times n + 1.1$ |

Section 201 Elimination

| | | |
|--------------------|--|----------------------|
| mca 2010 elmvec | | $.61 \times n + 1.1$ |
| mca 2011 elmveccol | | $.69 \times n + 1.1$ |
| mca 2012 elmcolvec | | $.78 \times n + 1.1$ |
| mca 2013 elmcol | | $.87 \times n + 1.2$ |
| mca 2014 elmrow | | $.87 \times n + 1.2$ |
| mca 2019 maxelmrow | | $.94 \times n + 1.3$ |

Section 202 Interchanging

| | | |
|--------------------|--|----------------------|
| mca 2020 ichvec | | $.81 \times n + .7$ |
| mca 2021 ichcol | | $1.14 \times n + .8$ |
| mca 2022 ichrow | | $1.14 \times n + .8$ |
| mca 2023 ichrowcol | | $1.14 \times n + .8$ |
| mca 2024 ichseqvec | | $.84 \times n + .8$ |
| mca 2025 ichseq | | $.84 \times n + .8$ |

Section 203 Rotation

| | | |
|-----------------|--|-----------------------|
| mca 2031 rotcol | | $1.40 \times n + 1.3$ |
| mca 2032 rotrow | | $1.40 \times n + 1.3$ |

CHAPTER 21 LINEAR SYSTEMS AND MATRIX INVERSION

The formulas for this chapter hold for nonsingular matrices, unless stated otherwise.

Section 210 Triangular decomposition with partial pivoting

| | mca 2000 to 2005 in machine-code | mca 2000 to 2005 in ALGOL 60 |
|-----------------|---|--|
| mca 2100 det | $(.033 \times n + 3.4) \times n \uparrow 2$ | $(.22 \times n + 2.9) \times n \uparrow 2$ |
| mca 2101 sol | $(.094 \times n + 4.5) \times n$ | $(.55 \times n + 2.9) \times n$ |
| mca 2102 detsol | $(.033 \times n + 3.5) \times n \uparrow 2$ | $(.22 \times n + 3.5) \times n \uparrow 2$ |
| mca 2103 inv | $(.061 \times n + 3.4) \times n \uparrow 2$ | $(.43 \times n + 1.9) \times n \uparrow 2$ |
| mca 2104 detinv | $(.094 \times n + 6.8) \times n \uparrow 2$ | $(.65 \times n + 4.8) \times n \uparrow 2$ |

Section 211 Elimination with complete pivoting

The formula for rnkelm holds, if n and m are nearly equal and the rank equals $\min(n, m)$; the formula for solhom holds, provided the rank is not much smaller than m .

| | |
|--------------------|--|
| mca 2110 rnkelm | $(.51 \times n + 4.2) \times n \times (m - n/3)$ |
| mca 2111 solelm | $(.100 \times n + 4.8) \times n$ |
| mca 2112 rnksolelm | $(.34 \times n + 2.9) \times n \uparrow 2$ |
| mca 2113 solhom | $(.046 \times \text{rank} + 2.9) \times \text{rank}$ |
| mca 2114 invelm | $(.95 \times n + 5.7) \times n \uparrow 2$ |

Section 212 Band matrices

Here w is the band width, thus, $w = lw + rw + 1$.
The formulas for this section hold only if w is much smaller than n .

| | |
|--------------------|--|
| mca 2120 detbnd | $(.56 \times lw + 2.3) \times w \times n$ |
| mca 2121 solbnd | $(.7 \times lw + .1 \times rw + 4.2) \times n$ |
| mca 2122 detsolbnd | $(.55 \times lw + 2.9) \times w \times n$ |

CHAPTER 22 POSITIVE DEFINITE SYMMETRIC LINEAR SYSTEMS AND MATRIX
INVERSION

The formulas for this chapter hold for nonsingular matrices, unless stated otherwise.

Section 220 Cholesky decomposition without pivoting

| | | |
|----------|------------|---|
| mca 2200 | detsym2 | $(.016 \times n + 1.2) \times n \uparrow 2$ |
| mca 2201 | solsym2 | $(.092 \times n + 4.1) \times n$ |
| mca 2202 | detsolsym2 | $(.016 \times n + 1.3) \times n \uparrow 2$ |
| mca 2203 | invsym2 | $(.032 \times n + 1.7) \times n \uparrow 2$ |
| mca 2204 | detinvsym2 | $(.048 \times n + 2.9) \times n \uparrow 2$ |
| mca 2205 | detsym1 | $(.016 \times n + 1.0) \times n \uparrow 2$ |
| mca 2206 | solsym1 | $(.30 \times n + 3.3) \times n$ |
| mca 2207 | detsolsym1 | $(.016 \times n + 1.3) \times n \uparrow 2$ |
| mca 2208 | invsym1 | $(.104 \times n + 1.4) \times n \uparrow 2$ |
| mca 2209 | detinvsym1 | $(.120 \times n + 2.4) \times n \uparrow 2$ |

Section 221 Cholesky decomposition with pivoting

The formula for solsym20 holds, if the rank equals $n - 1$.

| | | |
|----------|-------------|---|
| mca 2210 | rnksym20 | $(.016 \times n + 2.6) \times n \uparrow 2$ |
| mca 2211 | solsym20 | $(.092 \times n + 5.6) \times n$ |
| mca 2212 | rnksolsym20 | $(.016 \times n + 2.7) \times n \uparrow 2$ |
| mca 2213 | invsym20 | $(.032 \times n + 2.8) \times n \uparrow 2$ |
| mca 2214 | rnkinvsym20 | $(.048 \times n + 5.4) \times n \uparrow 2$ |
| mca 221a | solsymhom20 | $(.016 \times n + 2.9) \times n \uparrow 2$ |
| mca 2215 | rnksym10 | $(.016 \times n + 2.0) \times n \uparrow 2$ |
| mca 2216 | solsym10 | $(.30 \times n + 4.8) \times n$ |
| mca 2217 | rnksolsym10 | $(.016 \times n + 2.3) \times n \uparrow 2$ |
| mca 2218 | invsym10 | $(.104 \times n + 2.2) \times n \uparrow 2$ |
| mca 2219 | rnkinvsym10 | $(.120 \times n + 4.2) \times n \uparrow 2$ |

Section 222 Cholesky decomposition for band matrices

The formulas for this section hold only if w is much smaller than n .

| | | |
|----------|--------------|---|
| mca 2220 | detsymbnd | $(.036 \times w + 2.4) \times w \times n$ |
| mca 2221 | solsymbnd | $(.67 \times w + 4.8) \times n$ |
| mca 2222 | detsolsymbnd | $(.036 \times w + 3.1) \times w \times n$ |

Section 224 Least-squares problems

The formulas for this section hold, if $n \geq m$ and the rank equals m .

| | | |
|----------|-----------|--|
| mca 2240 | lsqdec | $(.50 \times m + 2.2) \times m \times (n - m/3)$ |
| mca 2214 | lsqsol | $(.75 \times m + 1.0) \times n$ |
| mca 2242 | lsqdglinv | $(.016 \times m + 1.1) \times m \uparrow 2$ |
| mca 2243 | lsqdecsol | $(.50 \times m + 4.1) \times m \times (n - m/3)$ |