

Complexity of the positive semidefinite matrix completion problem with a rank constraint

M. E.-Nagy^{*1}, M. Laurent^{†1,2}, and A. Varvitsiotis^{‡1}

¹Centrum Wiskunde & Informatica (CWI), Amsterdam, The Netherlands.

²Tilburg University, Tilburg, The Netherlands.

September 19, 2012

Abstract

We consider the decision problem asking whether a partial rational symmetric matrix with an all-ones diagonal can be completed to a full positive semidefinite matrix of rank at most k . We show that this problem is \mathcal{NP} -hard for any fixed integer $k \geq 2$. Equivalently, for $k \geq 2$, it is \mathcal{NP} -hard to test membership in the rank constrained elliptope $\mathcal{E}_k(G)$, i.e., the set of all partial matrices with off-diagonal entries specified at the edges of G , that can be completed to a positive semidefinite matrix of rank at most k . Additionally, we show that deciding membership in the convex hull of $\mathcal{E}_k(G)$ is also \mathcal{NP} -hard for any fixed integer $k \geq 2$.

1 Introduction

Geometric representations of graphs are widely studied within a broad range of mathematical areas, ranging from combinatorial matrix theory, linear algebra, discrete geometry, and combinatorial optimization. They arise typically when labeling the nodes by vectors assumed to satisfy certain properties. For instance, one may require that the vectors labeling adjacent nodes are at distance 1, leading to unit distance graphs. Or one may require that

^{*}M.E.Nagy@cwi.nl

[†]M.Laurent@cwi.nl

[‡]A.Varvitsiotis@cwi.nl

the vectors labeling adjacent nodes are orthogonal, leading to orthogonal representations of graphs. One may furthermore ask, e.g., that nonadjacent nodes receive vector labels that are not orthogonal. Many other geometric properties of orthogonal labelings and other types of representations related, e.g., to Colin de Verdière type graph parameters, are of interest and have been investigated (see [9]). A basic question is to determine the smallest possible dimension of such vector representations. There is a vast literature, we refer in particular to the surveys [11, 12, 21] and further references therein for additional information.

In this note we revisit orthogonal representations of graphs, in the wider context of Gram representations of weighted graphs. We show some complexity results for the following notion of Gram dimension, which has been considered in [19, 20].

Definition 1.1 *Given a graph $G = (V = [n], E)$ and $x \in \mathbb{R}^E$, a Gram representation of x in \mathbb{R}^k consists of a set of unit vectors $v_1, \dots, v_n \in \mathbb{R}^k$ such that*

$$v_i^\top v_j = x_{ij} \quad \forall \{i, j\} \in E.$$

The Gram dimension of x , denoted as $\text{gd}(G, x)$, is the smallest integer k for which x has such a Gram representation in \mathbb{R}^k (assuming it has one in some space).

As we restrict our attention to Gram representations of $x \in \mathbb{R}^E$ by unit vectors, all coordinates of x should lie in the interval $[-1, 1]$, so that we can parametrize x as

$$x = \cos(\pi a), \quad \text{where } a \in [0, 1]^E.$$

In other words, the inequality $\text{gd}(G, x) \leq k$ means that (G, a) can be isometrically embedded into the spherical metric space $(\mathbf{S}^{k-1}, d_{\mathbf{S}})$, where \mathbf{S}^{k-1} is the unit sphere in the Euclidean space \mathbb{R}^k and $d_{\mathbf{S}}$ is the spherical distance:

$$d_{\mathbf{S}}(u, v) = \arccos(u^\top v) / \pi \quad \forall u, v \in \mathbf{S}^{k-1}.$$

Moreover, there are also tight connections with graph realizations in the Euclidean space (cf. [6, 7]); see Section 2.3 for a brief discussion and Section 3.2 for further results.

Determining the Gram dimension can also be reformulated in terms of finding low rank positive semidefinite matrix completions of partial matrices, as we now see. We use the following notation: \mathcal{S}^n denotes the set of

symmetric $n \times n$ matrices and \mathcal{S}_+^n is the cone of positive semidefinite (psd) matrices in \mathcal{S}^n . The subset

$$\mathcal{E}_n = \{X \in \mathcal{S}_+^n : X_{ii} = 1 \ \forall i \in [n]\},$$

consisting of all positive semidefinite matrices with an all-ones diagonal (aka the correlations matrices), is known as the *elliptope*. Given a graph $G = ([n], E)$, π_E denotes the projection from \mathcal{S}^n onto the subspace \mathbb{R}^E indexed by the edges of G . Then, the projection $\mathcal{E}(G) = \pi_E(\mathcal{E}_n)$ is known as the *elliptope* of the graph G . Given an integer $k \geq 1$, define the *rank constrained elliptope*

$$\mathcal{E}_{n,k} = \{X \in \mathcal{E}_n : \text{rank}(X) \leq k\},$$

and, for any graph G , its projection $\mathcal{E}_k(G) = \pi_E(\mathcal{E}_{n,k})$. Then the points x in the elliptope $\mathcal{E}(G)$ correspond precisely to those vectors $x \in \mathbb{R}^E$ that admit a Gram representation by unit vectors. Moreover, $x \in \mathcal{E}_k(G)$ precisely when it has a Gram representation by unit vectors in \mathbb{R}^k ; that is:

$$x \in \mathcal{E}_k(G) \iff \text{gd}(G, x) \leq k.$$

The elements of $\mathcal{E}(G)$ can be seen as the G -partial symmetric matrices, i.e., the partial matrices whose entries are specified at the off-diagonal positions corresponding to edges of G and whose diagonal entries are all equal to 1, that can be completed to a positive semidefinite matrix. Hence the problem of deciding membership in $\mathcal{E}(G)$ can be reformulated as the problem of testing whether a given G -partial matrix can be completed to a psd matrix. Moreover, for fixed $k \geq 1$, the membership problem in $\mathcal{E}_k(G)$ is the problem of deciding whether a given G -partial matrix has a psd completion of rank at most k . Using the notion of Gram dimension this can be equivalently formalized as:

Given a graph $G = (V, E)$ and $x \in \mathbb{Q}^E$, decide whether $\text{gd}(G, x) \leq k$.

A first main result of our paper is to prove \mathcal{NP} -hardness of this problem for any fixed $k \geq 2$ (cf. Theorems 3.2 and 3.3). Additionally, we consider the problem of testing membership in the convex hull of the rank constrained elliptope:

Given a graph $G = (V, E)$ and $x \in \mathbb{Q}^E$, decide whether $x \in \text{conv } \mathcal{E}_k(G)$.

The study of this problem is motivated by the relevance of the convex set $\text{conv } \mathcal{E}_k(G)$ to the maximum cut problem and to the rank constrained Grothendieck problem. Indeed, for $k = 1$, $\text{conv } \mathcal{E}_1(G)$ coincides with the cut polytope of G and it is well known that linear optimization over the cut

polytope is \mathcal{NP} -hard [13]. For any $k \geq 2$, the worst case ratio of optimizing a linear function over the ellipptope $\mathcal{E}(G)$ versus the rank constrained ellipptope $\mathcal{E}_k(G)$ (equivalently, versus the convex hull $\text{conv } \mathcal{E}_k(G)$) is known as the *rank k Grothendieck constant* of the graph G (see [8] for results and further references). It is believed that linear optimization over $\text{conv } \mathcal{E}_k(G)$ is also hard for any fixed k (cf., e.g., the quote of Lovász [21, p. 61]). We show that the strong membership problem in $\text{conv } \mathcal{E}_k(G)$ is \mathcal{NP} -hard, thus providing some evidence of hardness of optimization (cf. Theorem 4.2).

Contents of the paper. In Section 2 we present some background geometrical facts about cut and metric polytopes, about ellipptopes, and about Euclidean graph realizations. In Section 3 we show \mathcal{NP} -hardness of the membership problem in $\mathcal{E}_k(G)$ for any fixed $k \geq 2$; we use two different reductions depending whether $k = 2$ or $k \geq 3$. In Section 4 we show \mathcal{NP} -hardness of the membership problem in the convex hull of $\mathcal{E}_k(G)$ for any fixed $k \geq 2$. In Section 2.3 we discuss links to complexity results for Euclidean graph realizations, and in Section 5 we conclude with some open questions.

Some notation. Throughout $K_n = ([n], E_n)$ is the complete graph on n nodes; C_n denotes the circuit of length n , with node set $[n]$ and with edges the pairs $\{i, i+1\}$ for $i \in [n]$ (indices taken modulo n), and its set of edges is again denoted as C_n for simplicity. Given a graph $G = (V, E)$, its *suspension graph* ∇G is the new graph obtained from G by adding a new node, called the *apex* node and often denoted as 0, which is adjacent to all the nodes of G . A *minor* of G is any graph which can be obtained from G by iteratively deleting edges or nodes and contracting edges.

2 Preliminaries

We recall here some basic geometric facts about metric and cut polyhedra, about ellipptopes, and about Euclidean graph realizations.

2.1 Metric and cut polytopes

First we recall the definition of the *metric polytope* $\text{MET}(G)$ of a graph $G = (V, E)$. As a motivation recall the following basic 3D geometric result: Given a matrix $X = (x_{ij})$ in the ellipptope \mathcal{E}_3 , parametrized as before by $x_{ij} = \cos(\pi a_{ij})$ where $a_{ij} \in [0, 1]$, then $X \succeq 0$ if and only if the a_{ij} 's satisfy the following *triangle inequalities*:

$$a_{ij} \leq a_{ik} + a_{jk}, \quad a_{ij} + a_{ik} + a_{jk} \leq 2 \quad (1)$$

for distinct $i, j, k \in \{1, 2, 3\}$. (See e.g. [5]). The elliptope \mathcal{E}_3 (or rather, its bijective image $\mathcal{E}(K_3)$) is illustrated in Figure 1.

The metric polytope of the complete graph $K_n = ([n], E_n)$ is the polyhedron in \mathbb{R}^{E_n} defined by the above $4\binom{n}{3}$ triangle inequalities (1). More generally, the metric polytope of a graph $G = ([n], E)$ is the polyhedron $\text{MET}(G)$ in \mathbb{R}^E , which is defined by the following linear inequalities (in the variable $a \in \mathbb{R}^E$):

$$0 \leq a_e \leq 1 \quad \forall e \in E, \tag{2}$$

$$a(F) - a(C \setminus F) \leq |F| - 1 \tag{3}$$

for all circuits C of G and for all odd cardinality subsets $F \subseteq C$.

As is well known, the inequality (2) defines a facet of $\text{MET}(G)$ if and only if the edge e does not belong to a triangle of G , while (3) defines a facet of $\text{MET}(G)$ if and only if the circuit C has no chord (i.e., two non-consecutive nodes on C are not adjacent in G). In particular, for $G = K_n$, $\text{MET}(K_n)$ is defined by the triangle inequalities (1), obtained by considering only the inequalities (3) where C is a circuit of length 3. Moreover, $\text{MET}(G)$ coincides with the projection of $\text{MET}(K_n)$ onto the subspace \mathbb{R}^E indexed by the edge set of G . (See [10] for details.)

A main motivation for considering the metric polytope is that it gives a tractable linear relaxation of the cut polytope. Recall that the rank 1 matrices in the elliptope \mathcal{E}_n are of the form uu^\top for all $u \in \{\pm 1\}^n$, they are sometimes called the *cut matrices* since they correspond to the cuts of the complete graph K_n . Then the *cut polytope* $\text{CUT}(G)$ is defined as the projection onto \mathbb{R}^E of the convex hull of the cut matrices:

$$\text{CUT}(G) = \pi_E(\text{conv}(\mathcal{E}_{n,1})). \tag{4}$$

It is always true that $\text{CUT}(G) \subseteq \text{MET}(G)$, moreover equality holds if and only if G has no K_5 minor [4]. Linear optimization over the cut polytope models the maximum cut problem, well known to be \mathcal{NP} -hard [13], and testing membership in the cut polytope $\text{CUT}(K_n)$ or, equivalently, in the convex hull of the rank constrained elliptope $\mathcal{E}_{n,1}$, is an \mathcal{NP} -complete problem [2].

2.2 Elliptopes

From the above discussion about the elliptope \mathcal{E}_3 and the metric polytope, we can derive the following necessary condition for membership in the elliptope $\mathcal{E}(G)$ of a graph G , which turns out to be sufficient when G has no K_4 minor.

Proposition 2.1 [17] For any graph $G = (V, E)$,

$$\mathcal{E}(G) \subseteq \left\{ x \in [-1, 1]^E : \frac{1}{\pi} \arccos x \in \text{MET}(G) \right\}.$$

Moreover, equality holds if and only if G has no K_4 minor.

This result permits, in particular, to characterize membership in the elliptope $\mathcal{E}(C_n)$ of a circuit.

Corollary 2.1 [5] Consider a vector $x = \cos(\pi a) \in \mathbb{R}^{C_n}$ with $a \in [0, 1]^{C_n}$. Then, $x \in \mathcal{E}(C_n)$ if and only if a satisfies the linear inequalities

$$a(F) - a(C_n \setminus F) \leq |F| - 1 \quad \forall F \subseteq C_n \quad \text{with } |F| \text{ odd.} \quad (5)$$

We also recall the following result of [19] which characterizes membership in the rank constrained elliptope $\mathcal{E}_k(C_n)$ of a circuit in the case $k = 2$; see Lemma 3.2 for an extension to arbitrary graphs.

Lemma 2.1 [19] For $x \in [-1, 1]^{C_n}$, $x \in \mathcal{E}_2(C_n)$ if and only if there exists $\epsilon \in \{\pm 1\}^{C_n}$ such that $\epsilon^\top \arccos x \in 2\pi\mathbb{Z}$.

We conclude with some observations about the elliptope $\mathcal{E}(C_n)$ of a circuit. Figure 1 shows the elliptope $\mathcal{E}(C_3)$. Points x on the boundary of $\mathcal{E}(C_3)$ have $\text{gd}(C_3, x) = 2$ except $\text{gd}(C_3, x) = 1$ at the four corners (corresponding to the four cuts of K_3), while points in the interior of $\mathcal{E}(C_3)$ have $\text{gd}(C_3, x) = 3$.

Now let $n \geq 4$. Let $x = \cos(\pi a) \in \mathcal{E}(C_n)$, thus $a \in [0, 1]^{C_n}$ satisfies the inequalities (5). It is known that $\text{gd}(C_n, x) \leq 3$ (see [19], or derive it directly by triangulating C_n and applying Lemma 4.1 below). Moreover, x lies in the interior of $\mathcal{E}(C_n)$ if and only if x has a positive definite completion or, equivalently, a lies in the interior of the metric polytope $\text{MET}(C_n)$.

If x lies on the boundary of $\mathcal{E}(C_n)$ then, either (i) $a_e \in \{0, 1\}$ for some edge e of C_n , or (ii) a satisfies an inequality (5) at equality. In case (i), $\text{gd}(C_n, x)$ can be equal to 1 (x is a cut), 2, or 3. In case (ii), by Lemma 2.1, $\text{gd}(C_n, x) \leq 2$ since $a(F) - a(C_n \setminus F) = |F| - 1 \in 2\mathbb{Z}$ for some $F \subseteq C_n$. If x is in the interior of $\mathcal{E}(C_n)$ then $\text{gd}(C_n, x) \in \{2, 3\}$.

As an illustration, for $n = 4$, consider the vectors $x_1 = (0, 0, 0, 1)^\top$, $x_2 = (0, \sqrt{3}/2, \sqrt{3}/2, \sqrt{3}/2)^\top$, $x_3 = (0, 0, 0, 0)^\top$ and $x_4 = (0, 0, 0, 1/2)^\top \in \mathbb{R}^{C_4}$. Then both x_1 and x_2 lie on the boundary of $\mathcal{E}(C_4)$ with $\text{gd}(C_4, x_1) = 3$ and $\text{gd}(C_4, x_2) = 2$, and both x_3 and x_4 lie in the interior of $\mathcal{E}(C_4)$ with $\text{gd}(C_4, x_3) = 2$ and $\text{gd}(C_4, x_4) = 3$.




Figure 1: The elliptope $\mathcal{E}(C_3)$.

2.3 Euclidean graph realizations

In this section we recall some basic facts about Euclidean graph realizations.

Definition 2.1 *Given a graph $G = ([n], E)$ and $d \in \mathbb{R}_+^E$, a Euclidean (distance) representation of d in \mathbb{R}^k consists of a set of vectors $p_1, \dots, p_n \in \mathbb{R}^k$ such that*

$$\|p_i - p_j\|^2 = d_{ij} \quad \forall \{i, j\} \in E.$$

Then, $\text{ed}(G, d)$ denotes the smallest integer $k \geq 1$ for which d has a Euclidean representation in \mathbb{R}^k (assuming d has a Euclidean representation in some space).

Then the problem of interest is to decide whether a given vector $d \in \mathbb{Q}_+^E$ admits a Euclidean representation in \mathbb{R}^k . Formally, for fixed $k \geq 1$, we consider the following problem:

Given a graph $G = (V, E)$ and $d \in \mathbb{Q}_+^E$, decide whether $\text{ed}(G, d) \leq k$.

This problem has been extensively studied (e.g. in [6, 7]) and its complexity is well understood. In particular, using a reduction from the 3SAT problem, Saxe [26] shows the following complexity result.

Theorem 2.1 [26] *For any fixed $k \geq 1$, deciding whether $\text{ed}(G, d) \leq k$ is \mathcal{NP} -hard, already when restricted to weights $d \in \{1, 2\}^E$.*

We now recall a well known connection between Euclidean and Gram realizations. Given a graph $G = (V, E)$ and its suspension graph ∇G , consider the one-to-one map $\phi : \mathbb{R}^{V \cup E} \mapsto \mathbb{R}^{E(\nabla G)}$, which maps $x \in \mathbb{R}^{V \cup E}$ to $\varphi(x) = d \in \mathbb{R}^{E(\nabla G)}$ defined by

$$d_{0i} = x_{ii} \ (i \in [n]), \quad d_{ij} = x_{ii} + x_{jj} - 2x_{ij} \ (\{i, j\} \in E). \quad (6)$$

Then the vectors $u_1, \dots, u_n \in \mathbb{R}^k$ form a Gram representation of x if and only if the vectors $u_0 = 0, u_1, \dots, u_n$ form a Euclidean representation of $d = \varphi(x)$ in \mathbb{R}^k . This implies the following:

Lemma 2.2 *Let $G = (V, E)$ be a graph and $x \in \mathcal{E}(G)$. Then,*

$$\text{gd}(G, x) = \text{ed}(\nabla G, \varphi(x)).$$

As we will see in the next section, this connection will enable us to recover the above result of Saxe for the case $k \geq 3$ from results about the Gram dimension (cf. Corollary 3.1).

3 Testing membership in $\mathcal{E}_k(G)$

In this section we discuss the complexity of testing membership in the rank constrained ellipsope $\mathcal{E}_k(G)$. Specifically, for fixed $k \geq 1$ we consider the following problem:

Given a graph $G = (V, E)$ and $x \in \mathbb{Q}^E$, decide whether $\text{gd}(G, x) \leq k$.

In the language of matrix completions this corresponds to deciding whether a rational G -partial matrix has a psd completion of rank at most k .

For $k = 1$, $x \in \mathcal{E}_1(G)$ if and only if $x \in \{\pm 1\}^E$ corresponds to a cut of G , and it is an easy exercise that this can be decided in polynomial time. In this section we show that the problem is \mathcal{NP} -hard for any $k \geq 2$. It turns out that we have to use different reductions for the cases $k \geq 3$ and $k = 2$.

3.1 The case $k \geq 3$

First we consider the problem of testing membership in $\mathcal{E}_k(G)$ when $k \geq 3$. We show this is an \mathcal{NP} -hard problem, already when $G = \nabla^{k-3}H$ is the suspension of a planar graph H and $x = \mathbf{0}$, the all-zero vector.

The key idea is to relate the parameter $\text{gd}(G, \mathbf{0})$ to the chromatic number $\chi(G)$ (the minimum number of colors needed to color the nodes of G in such a way that adjacent nodes receive distinct colors). It is easy to check that

$$\text{gd}(G, \mathbf{0}) \leq \chi(G), \quad (7)$$

with equality if $\chi(G) \leq 2$ (i.e., if G is a bipartite graph). For $k \geq 3$ the inequality (7) can be strict. This is the case, e.g., for orthogonality graphs of Kochen-Specker sets (see [16]).

However, Peeters [22, Theorem 3.1] gives a polynomial time reduction of the problem of deciding 3-colorability of a graph to that of deciding $\text{gd}(G, \mathbf{0}) \leq 3$. Namely, given a graph G , he constructs (in polynomial time) a new graph G' having the property that

$$\chi(G) \leq 3 \iff \chi(G') \leq 3 \iff \text{gd}(G', \mathbf{0}) \leq 3. \quad (8)$$

The graph G' is obtained from G by adding for each pair of distinct nodes $i, j \in V$ the gadget graph H_{ij} shown in Figure 2. Moreover, using a more




Figure 2: The gadget graph H_{ij} .

involved construction, Peeters [23] constructs (in polynomial time) from any graph G a new *planar* graph G' satisfying (8). As the problem of deciding whether a given planar graph is 3-colorable is \mathcal{NP} -complete (see [27]) we have the following result.

Theorem 3.1 [23] *It is \mathcal{NP} -hard to decide whether $\text{gd}(G, \mathbf{0}) \leq 3$, already for the class of planar graphs.*

This hardness result can be extended to any fixed $k \geq 3$ using the suspension operation on graphs. The *suspension* graph $\nabla^p G$ is obtained from G by adding p new nodes that are pairwise adjacent and that are adjacent to all the nodes of G . It is easy observation that

$$\text{gd}(\nabla^p G, \mathbf{0}) = \text{gd}(G, \mathbf{0}) + p. \quad (9)$$

Theorem 3.1 combined with equation (9) implies:

Theorem 3.2 *Fix $k \geq 3$. It is \mathcal{NP} -hard to decide whether $\text{gd}(G, \mathbf{0}) \leq k$, already for graphs of the form $G = \nabla^{k-3} H$ where H is a planar graph.*

As an application we can recover the complexity result of Saxe from Theorem 2.1 for the case $k \geq 3$.

Corollary 3.1 *For fixed $k \geq 3$, it is an \mathcal{NP} -hard problem to decide whether $\text{ed}(G, d) \leq k$, already when $G = \nabla^{k-2}H$ with H planar and d is $\{1, 2\}$ -valued (more precisely, all edges adjacent to a given apex node have weight 1 and all other edges have weight 2).*

Proof. This follows directly from Lemma 2.2 combined with Theorem 3.2: By Lemma 2.2, $\text{gd}(\nabla^{k-3}H, \mathbf{0}) = \text{ed}(\nabla^{k-2}H, \varphi(\mathbf{0}))$ and observe that the image $d = \varphi(\mathbf{0})$ of the zero vector under the map φ from (6) satisfies: $d_{0i} = 1$ and $d_{ij} = 2$ for all nodes i, j of $\nabla^{k-3}H$. \square

3.2 The case $k = 2$

In this section we show \mathcal{NP} -hardness of testing membership in $\mathcal{E}_2(G)$. Our strategy to show this result is as follows: Given a graph $G = (V, E)$ with edge weights $d \in \mathbb{R}_+^E$, define the new edge weights $x = \cos(d) \in \mathbb{R}^E$. We show a close relationship between the two problems of testing whether $\text{ed}(G, d) \leq 1$, and whether $\text{gd}(G, x) \leq 2$ (or, equivalently, $x \in \mathcal{E}_2(G)$). More precisely, we show that each of these two properties can be characterized in terms of the existence of a ± 1 -signing of the edges of G satisfying a suitable ‘flow conservation’ type property; moreover, both are equivalent when the edge weights d are small enough.

As a motivation, let us consider first the case when $G = C_n$ is a circuit of length n . Say, weight d_i (resp., $x_i = \cos d_i$) is assigned to the edge $(i, i + 1)$ for $i \in [n]$ (setting $n + 1 = 1$). Then the following property holds:

$$\text{ed}(C_n, d) \leq 1 \iff \exists \epsilon \in \{\pm 1\}^n \text{ such that } \epsilon^\top d = 0. \quad (10)$$

This is the key fact used by Saxe [26] for showing \mathcal{NP} -hardness of the problem of testing $\text{ed}(C_n, d) \leq 1$ by reducing it from the Partition problem for $d = (d_1, \dots, d_n) \in \mathbb{Z}_+^n$. Lemma 2.1 shows the analogous result for the Gram dimension:

$$\text{gd}(C_n, \cos d) \leq 2 \iff \exists \epsilon \in \{\pm 1\}^n \text{ such that } \epsilon^\top d \in 2\pi\mathbb{Z}. \quad (11)$$

We now observe that these two characterizations extend for an arbitrary graph G . To formulate the result we need to fix an (arbitrary) orientation \vec{G} of G . Let $P = (u_0, u_1, \dots, u_{k-1}, u_k)$ be a walk in G , i.e., $\{u_i, u_{i+1}\} \in E$

for all $i \leq k-1$. For $\epsilon \in \{\pm 1\}^E$, we define the following weighted sum along the edges of P :

$$\phi_{d,\epsilon}(P) = \sum_{i=0}^{k-1} d_{u_i, u_{i+1}} \epsilon_{u_i u_{i+1}} \eta_i, \quad (12)$$

setting $\eta_i = 1$ if the edge $\{u_i, u_{i+1}\}$ is oriented in \tilde{G} from u_i to u_{i+1} and $\eta_i = -1$ otherwise.

Lemma 3.1 *Consider a graph $G = (V, E)$ with edge weights $d \in \mathbb{R}_+^E$ and fix an orientation \tilde{G} of G . The following assertions are equivalent.*

- (i) $\text{ed}(G, d) \leq 1$.
- (ii) *There exists an edge-signing $\epsilon \in \{\pm 1\}^E$ for which the function $\phi_{d,\epsilon}$ from (12) satisfies: $\phi_{d,\epsilon}(C) = 0$ for all circuits C of G .*

Proof. Assume that (i) holds. Let $f : V \rightarrow \mathbb{R}$ satisfying $|f(u) - f(v)| = d_{uv}$ for all $\{u, v\} \in E$. If the edge $\{u, v\}$ is oriented from u to v in \tilde{G} , let $\epsilon_{uv} \in \{\pm 1\}$ such that $f(v) - f(u) = d_{uv} \epsilon_{uv}$. This defines an edge-signing $\epsilon \in \{\pm 1\}^E$; we claim that (ii) holds for this edge-signing. For this, pick a circuit $C = (u_0, u_1, \dots, u_k = u_0)$ in G . By construction of the edge-signing, the term $\epsilon_{u_i u_{i+1}} d_{u_i u_{i+1}} \eta_i$ is equal to $f(u_{i+1}) - f(u_i)$ for all i . This implies that $\phi_{d,\epsilon}(C) = \sum_{i=0}^{k-1} f(u_{i+1}) - f(u_i) = 0$ and thus (ii) holds. Conversely, assume (ii) holds. We may assume that G is connected (else apply the following to each connected component). Fix an arbitrary node $u_0 \in V$. We define the function $f : V \rightarrow \mathbb{R}$ by setting $f(u_0) = 0$ and, for $u \in V \setminus \{u_0\}$, $f(u) = \phi_{d,\epsilon}(P)$ where P is a walk from u_0 to u . It is easy to verify that since (ii) holds this definition does not depend on the choice of P . We claim that f is a Euclidean embedding of (G, d) into \mathbb{R} . For this, pick an edge $\{u, v\} \in E$; say, it is oriented from u to v in \tilde{G} . Pick a walk P from u_0 to u , so that $Q = (P, v)$ is a walk from u_0 to v . Then, $f(u) = \phi_{d,\epsilon}(P)$, $f(v) = \phi_{d,\epsilon}(Q) = \phi_{d,\epsilon}(P) + d_{uv} \epsilon_{uv} = f(u) + d_{uv} \epsilon_{uv}$, which implies that $|f(v) - f(u)| = d_{uv}$. \square

Lemma 3.2 *Consider a graph $G = (V, E)$ with edge weights $d \in \mathbb{R}_+^E$ and fix an orientation \tilde{G} of G . The following assertions are equivalent.*

- (i) $\text{gd}(G, \cos d) \leq 2$.
- (ii) *There exists an edge-signing $\epsilon \in \{\pm 1\}^E$ for which the function $\phi_{d,\epsilon}$ from (12) satisfies: $\phi_{d,\epsilon}(C) \in 2\pi\mathbb{Z}$ for all circuits C of G .*

Proof. Assume (i) holds. That is, there exists a labeling of the nodes $u \in V$ by unit vectors $g(u) = (\cos f(u), \sin f(u))$ where $f(u) \in [0, 2\pi]$. For any edge $\{u, v\} \in E$, we have: $\cos d_{uv} = g(u)^\top g(v) = \cos(f(u) - f(v))$. If $\{u, v\}$ is oriented from u to v , define $\epsilon \in \{\pm 1\}$ such that $f(v) - f(u) - \epsilon_{uv}d_{uv} \in 2\pi\mathbb{Z}$. This defines an edge-signing $\epsilon \in \{\pm 1\}^E$ which satisfies (ii) (same argument as in the proof of Lemma 3.1).

Conversely, assume (ii) holds. Analogously to the proof of Lemma 3.1, fix a node $u_0 \in V$ and consider the unit vectors $g(u_0) = (1, 0)$ and $g(u) = (\cos(\phi_{d,\epsilon}(P_u)), \sin(\phi_{d,\epsilon}(P_u)))$, where P_u is a walk from $u_0 \in V$ to $u \in V \setminus \{u_0\}$; one can verify that these vectors form a Gram realization of $(G, \cos d)$. \square

Corollary 3.2 *Consider a graph $G = (V, E)$ with edge weights $d \in \mathbb{R}_+^E$ satisfying $\sum_{e \in E} d_e < 2\pi$. Then, $\text{ed}(G, d) \leq 1 \iff \text{gd}(G, \cos d) \leq 2$.*

Proof. Note that if C is a circuit of G , then $\phi_{d,\epsilon}(C) \in 2\pi\mathbb{Z}$ implies $\phi_{d,\epsilon}(C) = 0$, since $|\phi_{d,\epsilon}(C)| \leq \sum_{e \in E} d_e < 2\pi$. The result now follows directly by applying Lemmas 3.1 and 3.2. \square

We can now show \mathcal{NP} -hardness of testing membership in the rank constrained ellipsope $\mathcal{E}_2(G)$. For this we use the result of Theorem 2.1 for the case $k = 1$: Given edge weights $d \in \{1, 2\}^E$, it is \mathcal{NP} -hard to decide whether $\text{ed}(G, d) \leq 1$.

Theorem 3.3 *Given a graph $G = (V, E)$ and rational edge weights $x \in \mathbb{Q}^E$, it is \mathcal{NP} -hard to decide whether $x \in \mathcal{E}_2(G)$ or, equivalently, $\text{gd}(G, x) \leq 2$.*

Proof. Fix edge weights $d \in \{1, 2\}^E$. We reduce the problem of testing whether $\text{ed}(G, d) \leq 1$ to the problem of testing whether $\text{gd}(G, \cos(\alpha d)) \leq 2$, where α is chosen in such a way that $\cos \alpha \in \mathbb{Q}$ and $\alpha < 1/(\sum_{e \in E} d_e)$. For this, set $D = \sum_{e \in E} d_e$ and define the angle $\alpha > 0$ by

$$\cos \alpha = \frac{16D^2 - 1}{16D^2 + 1} \in \mathbb{Q}, \quad \sin \alpha = \frac{8D}{16D^2 + 1} \in \mathbb{Q}.$$

Then, $\sin \alpha < 1/(2D) \leq 0.5 < \sin 1$, which implies that $\alpha < 2 \sin \alpha \leq 1/D$ and thus $\alpha < 1/D = 1/(\sum_{e \in E} d_e)$.

As $d_e \in \{1, 2\}$, $\cos(\alpha d_e) \in \{\cos \alpha, \cos(2\alpha) = 2 \cos^2 \alpha - 1\}$ is rational valued for all edges $e \in E$. As $\sum_{e \in E} \alpha d_e < 1 < 2\pi$, Corollary 3.2 shows that $\text{gd}(G, \cos(\alpha d)) \leq 2$ is equivalent to $\text{ed}(G, \alpha d) \leq 1$ and thus to $\text{ed}(G, d) \leq 1$. This concludes the proof. \square

We conclude with a remark about the complexity of the Gram dimension of weighted circuits.

Remark 3.1 Consider the case when $G = C_n$ is a circuit and the edge weights $d \in \mathbb{Z}_+^{C_n}$ are integer valued. Relation (10) shows that $\text{ed}(C_n, d) \leq 1$ if and only if the sequence $d = (d_1, \dots, d_n)$ can be partitioned, thus showing \mathcal{NP} -hardness of the problem of testing $\text{ed}(C_n, d) \leq 1$.

As in the proof of Theorem 3.3 let us choose α such that $\cos \alpha, \sin \alpha \in \mathbb{Q}$ and $\alpha < 1/(\sum_{i=1}^n d_i)$; then $\cos(t\alpha) \in \mathbb{Q}$ for all $t \in \mathbb{Z}$. The analogous relation (11) holds, which shows that $\text{gd}(C_n, \cos(\alpha d)) \leq 2$ if and only if the sequence $d = (d_1, \dots, d_n)$ can be partitioned. However, it is not clear how to use this fact in order to show \mathcal{NP} -hardness of the problem of testing $\text{gd}(C_n, x) \leq 2$. Indeed, although all $\cos(\alpha d_i)$ are rational valued, the difficulty is that it is not clear how to compute $\cos(\alpha d_i)$ in time polynomial in the bit size of d_i (while it can be shown to be polynomial in d_i).

Finally we point out the following link to the construction of Aspnes et al. [1, §IV]. Consider the edge weights $x = \cos(\alpha d) \in \mathbb{R}^{C_n}$ for the circuit C_n and $y = \varphi(x)$ for its suspension ∇C_n , which is the wheel graph W_{n+1} . Thus $y_{0i} = 1$ and $y_{i,i+1} = 2 - 2\cos(\alpha d_i) = 4\sin^2(\alpha d_i/2)$ for all $i \in [n]$. Taking square roots we find the edge weights used in [1] to claim \mathcal{NP} -hardness of realizing weighted wheels (that have the property of admitting unique (up to congruence) realizations in the plane). As explained above in the proof of Theorem 3.3, if we suitably choose α we can make sure that all $\sin(\alpha d_i/2)$ be rational valued, while [1] uses real numbers. However, it is not clear how to control their bit sizes, and thus how to deduce \mathcal{NP} -hardness.

4 Testing membership in $\text{conv } \mathcal{E}_k(G)$

In the previous section we showed that testing membership in the rank constrained elliptope $\mathcal{E}_k(G)$ is an \mathcal{NP} -hard problem for any fixed $k \geq 2$. A related question is to determine the complexity of optimizing a linear objective function over $\mathcal{E}_k(G)$ or, equivalently, over its convex hull $\text{conv } \mathcal{E}_k(G)$. This question has been raised, in particular, by Lovász [21, p.61] and more recently in [8], and we will come back to it in Section 5. In turn, this is related to the problem of testing membership in the convex hull $\text{conv } \mathcal{E}_k(G)$ which we address in this section. Specifically, for any fixed $k \geq 1$ we consider the following problem:

Given a graph $G = (V, E)$ and $x \in \mathbb{Q}^E$, decide whether $x \in \text{conv } \mathcal{E}_k(G)$.

For $k = 1$, $\text{conv } \mathcal{E}_1(G)$ coincides with the cut polytope of G , for which the membership problem is \mathcal{NP} -complete [2]. In this section we will show that this problem is \mathcal{NP} -hard for any fixed $k \geq 2$. The key fact to prove

hardness is to consider the membership problem in $\text{conv } \mathcal{E}_k(G)$ for extreme points of the ellipsope $\mathcal{E}(G)$.

For a convex set K recall that a point $x \in K$ is an *extreme point* of K if $x = \lambda y + (1 - \lambda)z$ where $0 < \lambda < 1$ and $y, z \in K$ implies that $x = y = z$. The set of extreme points of K is denoted by $\text{ext } K$. Clearly, for $x \in \text{ext } \mathcal{E}(G)$,

$$x \in \text{conv } \mathcal{E}_k(G) \iff x \in \mathcal{E}_k(G). \quad (13)$$

Our strategy for showing hardness of membership in $\text{conv } \mathcal{E}_k(G)$ is as follows: Given a graph $G = (V, E)$ and a rational vector $x \in \mathcal{E}(G)$, we construct (in polynomial time) a new graph $\widehat{G} = (\widehat{V}, \widehat{E})$ (containing G as a subgraph) and a new rational vector $\widehat{x} \in \mathbb{Q}^{\widehat{E}}$ (extending x) satisfying the following properties:

$$\widehat{x} \in \text{ext } \mathcal{E}(\widehat{G}), \quad (14)$$

$$x \in \mathcal{E}_k(G) \iff \widehat{x} \in \mathcal{E}_k(\widehat{G}). \quad (15)$$

Combining these two conditions with (13), we deduce:

$$x \in \mathcal{E}_k(G) \iff \widehat{x} \in \mathcal{E}_k(\widehat{G}) \iff \widehat{x} \in \text{conv } \mathcal{E}_k(\widehat{G}). \quad (16)$$




Figure 3: The graph \widehat{C}_5 .

Given $G = (V, E)$, the construction of the new graph $\widehat{G} = (\widehat{V}, \widehat{E})$ is as follows: For each edge $\{i, j\}$ of G , we add a new node v_{ij} , adjacent to the two nodes i and j . Let C_{ij} denote the clique on $\{i, j, v_{ij}\}$ and set $\widehat{V} = V \cup \{v_{ij} : \{i, j\} \in E\}$. Then \widehat{G} has node set \widehat{V} and its edge set is the union of all the cliques C_{ij} for $\{i, j\} \in E$. As an illustration Figure 3 shows the graph \widehat{C}_5 .

Given $x \in \mathbb{Q}^E$, the construction of the new vector $\widehat{x} \in \mathbb{Q}^{\widehat{E}}$ is as follows: For each edge $\{i, j\} \in E$,

$$\widehat{x}_{ij} = x_{ij}, \quad (17)$$

$$\widehat{x}_{i,v_{ij}} = 4/5, \widehat{x}_{j,v_{ij}} = 3/5 \quad \text{if } x_{ij} = 0, \quad (18)$$

$$\widehat{x}_{i,v_{ij}} = x_{ij}, \widehat{x}_{j,v_{ij}} = 2x_{ij}^2 - 1 \quad \text{if } x_{ij} \neq 0. \quad (19)$$

We will use the following result characterizing the extreme points of the elliptope \mathcal{E}_3 .

Theorem 4.1 [14] *A matrix $X = (x_{ij}) \in \mathcal{E}_3$ is an extreme point of \mathcal{E}_3 if either $\text{rank}(X) = 1$, or $\text{rank}(X) = 2$ and $|x_{ij}| < 1$ for all $i \neq j \in \{1, 2, 3\}$.*

We also need the following well known (and easy to check) result permitting to construct points in the elliptope of clique sums of graphs.

Lemma 4.1 *Given two graphs $G_l = (V_l, E_l)$ ($l = 1, 2$), where $V_1 \cap V_2$ is a clique in both G_1, G_2 , the graph $G = (V_1 \cup V_2, E_1 \cup E_2)$ is called their clique sum. Given $x_l \in \mathbb{R}^{E_l}$ ($l = 1, 2$) such that $(x_1)_{ij} = (x_2)_{ij}$ for $i, j \in V_1 \cap V_2$, let $x = (x_{ij}) \in \mathbb{R}^E$ be their common extension, defined as $x_{ij} = (x_l)_{ij}$ if $i, j \in V_l$. Then, for any integer $k \geq 1$,*

$$x \in \mathcal{E}_k(G) \iff x_1 \in \mathcal{E}_k(G_1) \text{ and } x_2 \in \mathcal{E}_k(G_2).$$

We can now show that our construction for \widehat{x} satisfies the two properties (14) and (15).

Lemma 4.2 *Given a graph $G = (V, E)$ and $x \in \mathbb{Q}^E$, let $\widehat{G} = (\widehat{V}, \widehat{E})$ be defined as above and let $\widehat{x} \in \mathbb{Q}^{\widehat{E}}$ be defined by (17)-(19). For fixed $k \geq 2$ we have that $x \in \mathcal{E}_k(G)$ if and only if $\widehat{x} \in \mathcal{E}_k(\widehat{G})$ and $\widehat{x} \in \text{ext } \mathcal{E}(\widehat{G})$.*

Proof. Sufficiency follows trivially so it remains to prove necessity. Observe that, for each edge $\{i, j\} \in E$, the restriction $\widehat{x}_{C_{ij}}$ of \widehat{x} to the clique C_{ij} is an extreme point of $\mathcal{E}(C_{ij})$. Indeed, applying Theorem 4.1, we find that the following matrices

$$\begin{pmatrix} 1 & 0 & 3/5 \\ 0 & 1 & 4/5 \\ 3/5 & 4/5 & 1 \end{pmatrix}, \begin{pmatrix} 1 & x_{ij} & x_{ij} \\ x_{ij} & 1 & 2x_{ij}^2 - 1 \\ x_{ij} & 2x_{ij}^2 - 1 & 1 \end{pmatrix} \quad \text{where } x_{ij} \in [-1, 1] \setminus \{0\}$$

are extreme points of \mathcal{E}_3 so have rank at most 2. By construction, \widehat{G} is obtained as the clique sum of G with the cliques C_{ij} . Hence, by Lemma 4.1, we deduce that $\widehat{x} \in \mathcal{E}_k(\widehat{G})$.

Finally, we show that \widehat{x} is an extreme point of $\mathcal{E}(\widehat{G})$, which follows from the fact that each $\widehat{x}_{C_{ij}}$ is an extreme point of $\mathcal{E}(C_{ij})$, combined with the

fact that the cliques C_{ij} (for $\{i, j\} \in E$) cover the graph G . Indeed, assume $\hat{x} = \sum_{i=1}^m \lambda_i \hat{x}_i$ where $\lambda_i > 0$, $\sum_{i=1}^m \lambda_i = 1$ and $\hat{x}_i \in \mathcal{E}(\hat{G})$. Taking the projection onto the clique C_{ij} and using the fact that $\hat{x}_{C_{ij}} \in \text{ext } \mathcal{E}(C_{ij})$ we deduce that, for all k , $(\hat{x}_k)_{C_{ij}} = \hat{x}_{C_{ij}}$ for all $\{i, j\} \in E$ and thus $\hat{x} = \hat{x}_k$. \square

Combining Theorems 3.2 and 3.3 with Lemma 4.2 and relation (16) we deduce the following complexity result.

Theorem 4.2 *For any fixed $k \geq 2$, testing membership in $\text{conv } \mathcal{E}_k(G)$ is an \mathcal{NP} -hard problem.*

5 Concluding remarks

In this note we have shown \mathcal{NP} -hardness of the membership problem in the rank constrained elliptope $\mathcal{E}_k(G)$ and in its convex hull $\text{conv } \mathcal{E}_k(G)$, for any fixed $k \geq 2$. As mentioned earlier, it would be interesting to settle the complexity status of linear optimization over $\text{conv } \mathcal{E}_k(G)$. The case $k = 1$ is settled: Then $\text{conv } \mathcal{E}_1(G)$ is the cut polytope and both the membership problem and the linear optimization problem are \mathcal{NP} -complete. For $k \geq 2$, the convex set $\text{conv } \mathcal{E}_k(G)$ is in general non-polyhedral. Hence the right question to ask is about the complexity of the *weak* optimization problem. It follows from general results about the ellipsoid method (see, e.g., [15] for details) that the weak optimization problem and the weak membership problems for $\text{conv } \mathcal{E}_k(G)$ have the same complexity status. Although we could prove that the (strong) membership problem in $\text{conv } \mathcal{E}_k(G)$ is \mathcal{NP} -hard, we do not know whether this is also the case for the *weak* membership problem.

A second question of interest is whether the problems belong to \mathcal{NP} . Indeed it is not clear how to find *succinct* certificates for membership in $\mathcal{E}(G)$ or in $\mathcal{E}_k(G)$. For one thing, even if the given partial matrix x is rational valued and is completable to a psd matrix, it is not known whether it admits a *rational* completion. (A positive result has been shown in [18] in the case of chordal graphs, and for graphs with minimum fill-in 1). In a more general setting, it is not known whether the problem of testing feasibility of a semidefinite program belongs to \mathcal{NP} . On the positive side it is known that this problem belongs to \mathcal{NP} if and only if it belongs to $\text{co-}\mathcal{NP}$ [25] and that it can be solved in polynomial time when fixing the dimension or the number of constraints [24].

Acknowledgements. We thank A. Schrijver for useful discussions and a referee for drawing our attention to the paper by Aspnes et al. [1].

References

- [1] J. Aspnes, T. Eren, D.K. Goldenberg, A.S. Morse, W. Whiteley, Y.R. Yang, B.D.O. Anderson, P.N. Belhumeur. A theory of network localization. *IEEE Transactions on Mobile Computing*, **5(12)**:1663–1678, 2006.
- [2] D. Avis and M. Deza. The cut cone, L1 embeddability, complexity, and multicommodity flows. *Networks*, **21(6)**:595–617, 1991.
- [3] M. Abramowitz and I.A. Stegun (Eds.). Orthogonal Polynomials. *Ch. 22 in Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, 9th printing. New York: Dover, 771–802, 1972.
- [4] F. Barahona and A. Mahjoub. On the cut polytope. *Mathematical Programming*, **36**:157–173, 1986.
- [5] W.W. Barrett, C.R. Johnson, and P. Tarazaga. The real positive definite completion problem: cycle completability. *Memoirs of the American Mathematical Society*, **584**, 69 pages, 1996.
- [6] M. Belk. Realizability of graphs in three dimensions. *Disc. Comput. Geom.*, **37**:139–162, 2007.
- [7] M. Belk and R. Connelly. Realizability of graphs. *Disc. Comput. Geom.*, **37**:125–137, 2007.
- [8] J. Briët, F. de Oliveira Filho and F. Vallentin. *Grothendieck inequalities for semidefinite programs with rank constraint*, Preprint, 2011. Available at arXiv:1011.1754v1.
- [9] Y. Colin de Verdière. Multiplicities of eigenvalues and tree-width of graphs. *J. Comb. Theory B*, **74(2)**:121–146, 1998.
- [10] M.M. Deza and M. Laurent. *Geometry of Cuts and Metrics*. Springer, 1997.
- [11] S.M. Fallat and L. Hogben. The minimum rank of symmetric matrices described by a graph: A survey. *Linear Algebra Appl.*, **426**:558–582, 2007.
- [12] S.M. Fallat and L. Hogben. Variants on the minimum rank problem: A survey II. Preprint, 2011. Available at arXiv:1102.5142v1.
- [13] M.R. Garey, D. S. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W. H. Freeman & Co. 1979.

- [14] R. Grone, S. Pierce and W. Watkins. Extremal correlation matrices. *Linear Algebra Appl.*, **134**:63–70, 1990.
- [15] M. Grötschel, L. Lovász and A. Schrijver. *Geometric Algorithms and Combinatorial Optimization*. Springer, 1988.
- [16] G. Haynes, C. Park, A. Schaeffer, J. Webster and L.M, Mitchell. Orthogonal vector coloring. *Electron. J. Comb.*, 17, 2010.
- [17] M. Laurent. The real positive semidefinite completion problem for series-parallel graphs. *Linear Algebra Appl.*, **252(1-3)**:347–366, 1997.
- [18] M. Laurent. Polynomial instances of the positive semidefinite and Euclidean distance matrix completion problems. *SIAM J. Matrix Anal. A.*, **22**:874–894, 2000.
- [19] M. Laurent and A. Varvitsiotis. The Gram dimension of a graph. Preprint, 2011. Available at arXiv:1112.5960.
- [20] M. Laurent and A. Varvitsiotis. A new graph parameter related to bounded rank positive semidefinite matrix completions. Preprint, 2012.
- [21] L. Lovász. *Geometric Representations of Graphs*. Lecture Notes, 2001. <http://www.cs.elte.hu/~lovasz/geomrep.pdf>
- [22] R. Peeters. Orthogonal representations over finite fields and the chromatic number of graphs. *Combinatorica*, **16(3)**:417–431, 1996.
- [23] R. Peeters. *Ranks and Structure of Graphs*. PhD thesis, Tilburg University, 1995.
- [24] L. Porkolab and L. Khachiyan. On the complexity of semidefinite programs. *Journal of Global Optimization*, **10**:351–365, 1997.
- [25] M. Ramana. An exact duality theory for semidefinite programming and its complexity implications, *Math. Program.*, **77(1)**:129–162, 1997.
- [26] J. B. Saxe. Embeddability of weighted graphs in k-space is strongly NP-hard. *In Proc. 17-th Allerton Conf. Comm. Control Comp.*, 480–489, 1979.
- [27] L. Stockmeyer. Planar 3-colorability is polynomial complete. *SIGACT News*, **5(3)**:19–25, 1973.