Cosmic Rays used to Model Thunderclouds on Earth



How is lightning initiated in

News

Cosmic Rays used to Model Thunderclouds on Earth

Wed, 04/22/2015 - 2:17pm

- by Radboud University
- Get the latest news in High Performance Computing, Informatics, Data Analysis Software and more - Sign up now!



A particle shower initiated by a cosmic ray reaches LOFAR through a thundercloud. Courtesy of Radboud University.

thunderclouds? This is difficult to answer — how do you measure electric fields inside large, dangerously charged clouds? It was discovered, more or less by coincidence, that cosmic rays provide suitable probes to measure electric fields within thunderclouds. This surprising finding is published in *Physical Review Letters* on April 24. The measurements were performed with the LOFAR radio telescope located in the Netherlands.

Don't see your company?

Simply LOG IN / REGISTER and

GET STARTED

Once signed in, you can submit information about your company, products and boost your online presence.



Exclusives

Software and Moore's Drumbeat (Moore's Law)

May 19, 2015 2:48 pm | by James Reinders, Intel

"We used to throw away LOFAR measurements taken during thunderstorms. They were too messy." says astronomer Pim Schellart. "Well, we didn't actually throw them away of course, we just didn't analyze them."

Schellart, who completed his Ph.D. in March this year at Radboud University in Nijmegen and is supervised by Professor Heino Falcke, is interested in cosmic rays. These high-energy particles, originating from exploding stars and other astrophysical sources, continuously bombard Earth from space. High in the atmosphere these particles strike atmospheric molecules and create 'showers' of elementary particles. These showers can also be measured from the radio emission that is generated when their constituent particles are deflected by the magnetic field of the Earth. The radio emission also gives information about the original particles. These measurements are routinely conducted with LOFAR at ASTRON in Dwingeloo, but not during thunderstorms...

Modeling

That changed when the data were examined in a collaborative effort with astrophysicist Gia Trinh, Prof. Olaf Scholten from Groningen University and lightning expert Ute Ebert from the Centrum Wiskunde & Informatica in Amsterdam.

"We modeled how the electric field in thunderstorms can explain the different measurements. This worked very well. How the radio emission changes gives us a lot of information about the electric fields in thunderstorms. We could even determine the strength of the electric field at a certain height in the cloud." says Schellart.

This field can be as strong as 50 kV/m. This translates into a voltage of hundreds of millions of volts over a distance of multiple kilometers: a thundercloud contains enormous amounts of energy.

Dangerous charge

Lightning is a highly unpredictable natural phenomenon that inflicts damage to infrastructure and claims victims around the world. This new method to measure electric fields in thunderclouds will contribute to a better understanding and ultimately better predictions of lightning activity. Current measurement methods from planes, balloons or little rockets are dangerous and too localized. Most importantly, the presence of the measurement equipment influences the measurements. Cosmic rays probe the thunderclouds from top to bottom. Moving at almost the speed of light, they provide a near instantaneous 'picture' of the electric fields in the cloud. Moreover, they are created by nature and are freely available.

Cost of LIMS: True Pricing includes more than Purchase, Implementation and Annual Licensing

May 18, 2015 3:02 pm | by Siri H. Segalstad

Recap: The Week's Top Stories — May 8-14

May 15, 2015 2:34 pm | by Suzanne Tracy, Editor-in-Chief, Scientific Computing and HPC Source

Last Chance to Submit ISC Cloud & Big Data Research Papers

May 13, 2015 12:05 pm | by Suzanne Tracy, Editor-in-Chief, Scientific Computing and HPC Source

View More Exclusive Content »

Industry News

Last Chance to Submit ISC Cloud & Big Data Research Papers

May 13, 2015 12:05 pm | by Suzanne Tracy, Editor-in-Chief, Scientific Computing and HPC Source

ISC Announces the Hans Meuer Award Winning Research Paper

May 11, 2015 8:35 am | by ISC

Nominations for Three SC15 Awards due July 1

May 8, 2015 9:52 am | by SC15

"This research is an exemplary form of interdisciplinary collaboration between astronomers, particle physicists and geophysicists," says Heino Falcke. "We hope to develop the model further to ultimately answer the question: how is lightning initiated within thunderclouds?"

"Probing atmospheric electric fields in thunderstorms through radio emission from cosmic-ray induced air showers, "Schellart e.a. Physical Review Letters 24 april 2015 (highlight as editor's choice)

This research was made possible by funding from the ERC, FOM (project Cosmic Lighting), NWO en NOVA, and was a collaborative effort by researchers from Radboud University, Centrum Wiskunde & Informatica (CWI), ASTRON, KVI Rijksuniversiteit Groningen, Vrije Universiteit Brussel and many others.



View past editions of Scienctific Computing View past editions of HPC Source

Expert Contributors



Understanding Reliability Models by Mark Anawis

TOPICS MODELING DATA ACQUISITION PHYSICS

ADVERTISEMENT



Small and Medium Enterprises Enter the Limelight

Déjà Vu All Over Again:

management is

roblem, but a hallenge to the culture of an

by Steve Conway

Knowledge

not an IT p

(Ellott



Entry Deadline Approaching: May 31st, 2015

Recognizing inspired designs in science, technology, engineering, and math from the next generation of entrepreneurs, innovators, scientists, and engineers.

Evergreen Sponsor: Conference



Opening Up Performance with Open\$peedShop an Open Source Profiler by Rob Farber

Share 0 Pinit + Share



organizati

Computerized Systems in the Modern Laboratory: A **Practical Guide**

by John R. Joyce, Ph.D.

A New Dawn: Bringing HPC to Smaller Manufacturers