



A cb-Bohnenblust–Hille inequality with constant one and its applications in learning theory

Srinivasan Arunachalam^{1,2} · Arkopal Dutt¹ · Francisco Escudero Gutiérrez³ · Carlos Palazuelos⁴

Received: 22 July 2024 / Revised: 21 February 2025 / Accepted: 5 March 2025
© The Author(s) 2025

Abstract

The main result of this work shows that Bohnenblust–Hille inequality for m -homogeneous polynomials holds with constant one when the infinity norm is replaced by the completely bounded norm. Moreover, we show that this inequality finds some interesting consequences in quantum learning theory. In the second part of this paper, we broaden our investigation of the Bohnenblust–Hille inequality to other contexts. In particular, we extend recent results by Volberg and Zhang, demonstrating its applicability within a framework we have termed “Learning Low-Degree Quantum Objects”.

1 Introduction and main results

The Bohnenblust–Hille inequality states that for any $d \in \mathbb{N}$ there exists a constant C_d such that every d -homogeneous polynomial $P : \mathbb{C}^n \rightarrow \mathbb{C}$, defined as $P(z) = \sum_{|\alpha|=d} a_\alpha z^\alpha$, satisfies the following inequality:

$$\|\widehat{P}\|_{\frac{2d}{d+1}} \leq C_d \|P\|_\infty, \quad (1.1)$$

where $\|\widehat{P}\|_{\frac{2d}{d+1}}$ denotes the $\ell_{\frac{2d}{d+1}}$ sum of the coefficients $(a_\alpha)_\alpha$ and $\|P\|_\infty = \sup_{z \in \mathbb{D}^n} |P(z)|$ is the infinity norm of P [8].

We thank Jop Briët and Antonio Pérez Hernández for useful discussions. We thank Alexandros Eskenazis, Paata Ivanisvili, Alexander Volberg, and Haonan Zhang for useful comments. A.D. and S.A. thank the Institute for Pure and Applied Mathematics (IPAM) for its hospitality throughout the long program “Mathematical and Computational Challenges in Quantum Computing” in Fall 2023 during which part of this work was initiated. This research was supported by the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement no. 945045, and by the NWO Gravitation project NETWORKS under grant no. 024.002.003. F.E.G and C.P. are partially supported by the MICINN project PID2023-146758NB-I00 funded by MICIU/AEI/10.13039/501100011033, by TEC-2024/COM-84-QUITEMAD-CM, funded by Comunidad de Madrid and by Grant CEX2023-001347-S funded by MICIU/AEI/10.13039/501100011033.

Extended author information available on the last page of the article

This inequality, which generalizes the well-known Littlewood's 4/3-Inequality [24], has proven to be extremely useful in the study of the convergence of Dirichlet series and was crucial in determining the asymptotic behaviors of Bohr's radius obtained in [15]. In this regard, the authors demonstrated that the constant C_d can be taken equal to C^d , for a certain constant C . In fact, the work in [15] motivated numerous subsequent studies, where the search focused on the upper and lower bounds for C_d . The best known upper bound was given in [11], where it was proved that C_d can be actually taken to be $C^{\sqrt{d \log d}}$.

Interestingly, in recent years, the Bohnenblust–Hille inequality has proven to be useful in learning theory. This is perfectly illustrated in the striking work [19], where the authors used a version of the Bohnenblust–Hille inequality for functions defined on the hypercube $f : \{-1, 1\}^n \rightarrow \mathbb{R}$, as established in [18], to enhance the seminal Low-Degree Algorithm of Linial, Mansour, and Nisan [25]. After that, the applications of the Bohnenblust–Hille inequality to learning theory have reached the quantum computing community, motivating the study of this inequality in the non-commutative realm ([22, 32, 35]). In particular, in the work [35], a version of the Bohnenblust–Hille inequality is proven for $N \times N$ -dimensional matrices, which can be understood as a generalization of quantum Boolean functions [26].

In this work, we explore Bohnenblust–Hille inequalities from three different angles: considering the completely bounded norm instead of the infinity norm, extending the non-commutative variant proved in [35], and determining the exact constants for the case of Boolean functions.

1.1 The completely bounded Bohnenblust–Hille inequality

The completely bounded norm of a d -homogeneous polynomial P as above is defined as

$$\|P\|_{\text{cb}} = \sup \left\| \sum_{|\alpha|=d} a_\alpha Z_1^{\alpha_1} \cdots Z_n^{\alpha_n} \right\|_{\mathcal{B}(\mathcal{H})},$$

where this supremum runs over all complex Hilbert spaces \mathcal{H} and all contractions Z_1, \dots, Z_n in $\mathcal{B}(\mathcal{H})$. This norm can be understood as a non-commutative version of the infinity norm and it clearly provides an upper bound for it. Thus, one might expect that the corresponding Bohnenblust–Hille inequality involves a better constant than in the classical case. On the other hand, note that by the triangle inequality, we have that $\|P\|_{\text{cb}} \leq \sum_\alpha |a_\alpha|$. Simultaneously, the completely bounded norm has proven particularly suitable in the study of quantum algorithms, providing a notion of polynomial degree that gives a tight characterization of quantum query complexity [2]. Hence, a Bohnenblust–Hille inequality for the completely bounded norm is also motivated by its potential applications in quantum learning theory. The results of this work rigorously establish these expectations. Indeed, the main result of this work is that the Bohnenblust–Hille inequality holds with the optimal constant $C = 1$ when the infinity norm is replaced by the completely bounded norm. Additionally, we demonstrate that the exponent $2d/(d+1)$ is also optimal in the new scenario considered here, meaning that for $p < 2d/(d+1)$ there is no quantity C_d independent of n such that

$\|\widehat{P}\|_{2d/(d+1)} \leq C_d \|P\|_{\text{cb}}$ for every d -homogeneous polynomial P , as it happens with the original Bohnenblust–Hille inequality Eq. (1.1).

Theorem 1.1 *For every d -homogeneous polynomial $P : \mathbb{C}^n \rightarrow \mathbb{C}$, defined as $P(z) = \sum_{|\alpha|=d} a_\alpha z^\alpha$, the following inequality is satisfied:*

$$\|\widehat{P}\|_{\frac{2d}{d+1}} \leq \|P\|_{\text{cb}}.$$

Moreover, both the constant one in the inequalities and the exponent $\frac{2d}{d+1}$ are optimal.

In particular, our main result holds for multilinear forms. Moreover, we will also show that in the case of general (non-necessarily homogeneous) polynomials of degree d we have

$$\|\widehat{P}\|_{\frac{2d}{d+1}} \leq \sqrt{d+1} \|P\|_{\text{cb}}.$$

The optimality of Theorem 1.1 shows that the completely bounded norm fits perfectly into the study of the Bohnenblust–Hille inequality. In fact, Theorem 1.1 motivates the study of the optimality of the Bohnenblust–Hille inequality from an angle not explored to date. Rather than focusing on determining the optimal constant that satisfies the inequality (1.1), it is possible to examine the norms that satisfy the associated Bohnenblust–Hille inequality with a constant value of one. It is plausible that the second problem sheds light on the first; particularly, in the problem of finding new lower bounds for the constant C_d . Indeed, in order to find good lower bounds for the classical BH inequality, we must consider polynomials for which the infinity norm is very different from any norm for which a BH inequality with constant 1 can be proven.

Theorem 1.1 entails interesting consequences in learning theory. In particular, it allows us to improve the estimates in [19] when we restrict ourselves to certain functions arising in quantum computing. Indeed, in that work, it is proven that it is possible to learn any bounded function $f : \{-1, 1\}^n \rightarrow [-1, 1]$ of degree at most d with L_2 -accuracy ε and confidence $1 - \delta$ by using $O(\varepsilon^{-2(d+1)} C^{d^{3/2} \sqrt{\log d}} \log(n/\delta))$ random samples on the function. A particularly interesting type of these functions are those that arise from a quantum algorithm with d queries. More precisely, we consider here quantum query algorithms that prepare a state

$$|\psi_x\rangle = U_d(O_{x_d} \otimes \text{Id}_m) U_{d-1} \cdots U_1(O_{x_1} \otimes \text{Id}_m) U_0|u\rangle, \quad (1.2)$$

where m is an integer, x stands for (x_1, \dots, x_d) , O_y is the n -dimensional matrix that maps $|i\rangle$ to $y_i|i\rangle$, U_1, \dots, U_d are $(n+m)$ -dimensional unitaries and $|u\rangle$ is an $(n+m)$ -dimensional unit vector. The algorithm succeeds according to a projective measurement that measures the projection of the final state onto some fixed $(n+m)$ -dimensional unit vector $|v\rangle$. Hence, the amplitude of $|v\rangle$ is given by $T(x) = \langle v|\psi_x\rangle$, so that $|T(x)|^2$ is the acceptance probability of the algorithm. These quantum algorithms have been considered in the quantum computing literature; for example, k -fold forrelation, that witnesses the biggest possible quantum-classical separation, has this structure

[1]. As we will explain in Sect. 5, the argument in [19] alongside the Bohnenblust–Hille inequality for (bounded) multilinear forms [11] imply that the amplitudes T can be learned from $O(\varepsilon^{-2(d+1)} \text{poly}(d)^d \log(n/\delta))$ samples. Furthermore, using Theorem 1.1 instead of [11] allows us to obtain the following result for learning d -query quantum algorithms which, in particular, requires a number of samples that is polynomial on n when ε and δ are constants and $d = \log(n)$.

Corollary 1.2 *Consider a quantum algorithm that makes d queries as explained above. Then, its amplitudes can be learned with L_2^2 -accuracy ε and confidence $1 - \delta$ from $O(\varepsilon^{-2(d+1)} d^2 \log(n/\delta))$ uniform random samples.*

1.2 Extending the non-commutative Bohnenblust–Hille inequality

Motivated by the applications to learning quantum channels, we extend the non-commutative version of the BH inequality proved in [35]. This generalization concerns the Pauli coefficients of linear maps $\Phi : M_N \rightarrow M_N$, where let $N = 2^n$ and n is a natural number. These maps can be expressed as

$$\Phi(\rho) = \sum_{x, y \in \{0, 1, 2, 3\}^n} \widehat{\Phi}(x, y) \cdot \sigma_x \rho \sigma_y, \quad (1.3)$$

where $\sigma_x = \otimes_{i \in [n]} \sigma_{x_i}$ and σ_i for $i \in \{0, 1, 2, 3\}$ are the Pauli matrices

$$\sigma_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix};$$

and $\widehat{\Phi}(x, y)$ are the Pauli coefficients of the map. Given $x \in \{0, 1, 2, 3\}^n$, $|x|$ is the number of non-zero entries of x . The degree of Φ is the minimum integer d such that $\widehat{\Phi}(x, y) = 0$ if $|x| + |y| > d$.

We show that we can upper bound the $\ell_{2d/(d+1)}$ -sum of the Pauli coefficients of $\widehat{\Phi}$ in a Bohnenblust–Hille way.

Theorem 1.3 *Let $\Phi : M_N \rightarrow M_N$ be a linear map of degree d . Then,*

$$\|\widehat{\Phi}\|_{\frac{2d}{d+1}} \leq C^d \|\Phi : S_1^N \rightarrow S_\infty^N\|,$$

where C is a universal constant and S_1^N and S_∞^N denote the spaces of one and infinity Schatten classes respectively.

The proof of Theorem 1.3 follows a similar approach to the one in [35] and, in fact, extends their result. Indeed, if one considers a matrix $A \in M_N$, the main result in [35] follows from the application of Theorem 1.3 to the linear map $\Phi(X) = XA$.

We use our extension to improve the current results on learning quantum channels. From a physical perspective, quantum channels describe the transformations between quantum systems. Since quantum systems are represented by quantum states, which correspond to non-commutative probability distributions, specifically positive

semidefinite matrices with trace 1, quantum channels map one set of non-commutative probabilities to another. Mathematically, quantum channels on n -qubits are maps $\Phi : M_N \rightarrow M_N$ that are completely positive and trace-preserving. In particular, they satisfy $\|\Phi : S_1^N \rightarrow S_\infty^N\| \leq 1$ and Theorem 1.3 applies to them. Learning an n -qubit quantum channel is in general challenging and is known to require $\Theta(4^n)$ applications (queries) of the channel [21]. This exponential complexity can be drastically improved when prior information on the structure of the channel is available. For example, a recent work of Bao and Yao [13] considered k -junta quantum channels, i.e., n -qubit channels that act non-trivially only on at most k of the n (unknown) qubits leaving the rest of qubits unchanged. These channels were shown to be learnable using $\tilde{\Theta}(4^k)$ queries to the channel [13].

Using the same learning model as the recent work of Bao and Yao (see Sect. 5 for details) we prove the following result for learning low-degree channels, which contrary to the other applications of BH inequality in quantum learning theory, it has a sample complexity independent of n [22, 33–35].

Theorem 1.4 *Let Φ be a n -qubit degree- d quantum channel. Then it can be learned in L_2 -accuracy ε and confidence $\geq 1 - \delta$ by making $\exp(\tilde{O}(d^2 + d \log(1/\varepsilon))) \cdot \log(1/\delta)$ queries to Φ . Here, we use the notation \tilde{O} to hide logarithmic factors in d , $1/\varepsilon$, and $1/\delta$.*

1.3 Boolean functions

Since boolean functions $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ are particularly important in many contexts, we also analyze this case. Remember that the classical Fourier expansion in the hypercube allows one to write any function as

$$f = \sum_{s \in \{0, 1\}^n} \hat{f}(s) \chi_s, \quad (1.4)$$

where $\chi_s(x) = \prod_{i \in \text{supp}(s)} x_i$ for $s \in \{0, 1\}^n$ and $(\hat{f}(s))_s$ are the Fourier coefficients of f . Then, the degree of f is the minimum d such that $\hat{f}(s) = 0$ if $|s| > d$.

In this work, we show how the granularity property of these functions allows us to prove the corresponding optimal Bohnenblust–Hille inequality.

Proposition 1.5 *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be a function of degree at most d . Then,*

$$\left(\sum_{s \in \{0, 1\}^n} |\hat{f}(s)|^{\frac{2d}{d+1}} \right)^{\frac{d+1}{2d}} \leq 2^{\frac{d-1}{d}}.$$

The equality is witnessed by the address function.

The previous proposition might be of interest in functional analysis for two reasons. First, it is conjectured that the value of the BH constant for real d -linear forms is $2^{\frac{d-1}{d}}$ [31], so this fact proves the conjecture for the particular case of d -linear Boolean

forms. Second, the address function, that saturates the inequality, is a d -linear form that gives a lower bound for the BH constant for multilinear forms of $2^{\frac{d-1}{d}}$, which matches the best lower bound known so far for the BH inequality for real multilinear forms [17]. Together with Proposition 1.5 about Boolean functions, in this work we also study the complexity of these functions from the learning theoretical point of view and improve previous estimates in [28, Corollary 34] and [20, Corollary 4] (see Sect. 4.1 for details).

1.4 Organization

The paper is organized as follows. In Sect. 2 we introduce some preliminaries and basic results that we will use throughout the paper. Section 3 is devoted to studying the Bohnenblust–Hille inequality for the completely bounded norm and, in particular, we prove Theorem 1.1 and demonstrate its optimality. In Sect. 4 we study the BH inequality in other contexts. First, we focus on the case of boolean functions and we prove Proposition 1.5. Then, we consider the BH-inequality for linear maps $\Phi : M_N \rightarrow M_N$ and prove Theorem 1.3. Since we believe that the main interest of this article lies in its purely mathematical content, we have decided to postpone all the results on learning theory until the last section of the paper. Indeed, in Sect. 5, we explain the applications of our results to various problems and the improvements they entail compared to existing results. We believe this section will be of great interest to researchers working on mathematical problems related to those discussed in this paper, particularly those interested in potential applications. It is thus in this section where we prove Corollary 5.1, some almost-optimal results for the learnability of Boolean functions, and Theorem 1.4.

The findings outlined in this study are situated within a broader framework we termed “Learning low-degree quantum objects”. Our primary contribution lies in demonstrating novel variants of the Bohnenblust–Hille inequality, directly applicable to contextual issues. We have excluded certain additional results in learning theory that, while intriguing, do not leverage the mathematical substance of our inequalities. These omitted results can be found in [4].

2 Preliminaries

Throughout this work we will consider different constants, all of which will be denoted by C and their value will be clear from context. We will use C_d to refer to quantities that only depend on d and are constant with respect to other parameters. Given a natural number m , $[m]$ represents the set $\{1, 2, \dots, m\}$. We will often use n to refer to the number of variables or qubits, and N as 2^n . We use M_N to denote the space of complex square matrices of dimension N . Unless stated otherwise, given a matrix M , $\|M\|$ is used to denote its norm as a linear map from a Hilbert space to the same Hilbert space.

A n -qubit state ρ is an element of M_N that is positive semidefinite and has trace one. A n -qubit channel $\Phi : M_N \rightarrow M_N$ is a completely positive trace preserving

linear map. A measurement is a set $\{M_x\}_x$ of positive semidefinite matrices that sum to the identity. By the postulates of quantum mechanics, measuring a quantum state ρ with $\{M_x\}_x$ outputs x with probability $\text{Tr}[\rho M_x]$.

We will often use the Choi–Jamiołkowski isomorphism to encode a quantum channel as a quantum state. We call the resulting state as the Choi–Jamiołkowski state (or CJ state for short). The CJ representation is given by

$$J(\Phi) = \sum_{i,j \in [N]} \Phi(|i\rangle\langle j|) \otimes |i\rangle\langle j| = (\Phi \otimes I) \left(\sum_{i,j \in [N]} |i\rangle\langle j| \otimes |i\rangle\langle j| \right), \quad (2.1)$$

which is an element in $M_N \otimes M_N = M_{N^2}$. The CJ state $v(\Phi)$ is defined to be

$$v(\Phi) = \frac{J(\Phi)}{\text{Tr}[J(\Phi)]} = \frac{J(\Phi)}{N}. \quad (2.2)$$

According to (2.1), the CJ state $v(\Phi)$ can be prepared by first preparing n EPR pairs (over $2n$ qubits) and then applying Φ to the n qubits coming from the first half of each of the n EPR pairs.

In this paper, we consider the space of functions defined on the Boolean hypercube $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ endowed with the inner product $\langle f, g \rangle = \mathbb{E}_x[f(x)g(x)]$, where the expectation is taken with respect to the uniform measure of probability. For $s \in \{0, 1\}^n$, the Fourier characters, defined by $\chi_s(x) = \prod_{i:s_i=1} x_i$, constitute an orthonormal basis of this space. Hence, every f can be identified with a multilinear polynomial via the Fourier expansion

$$f = \sum_{s \in \{0, 1\}^n} \widehat{f}(s) \chi_s, \quad (2.3)$$

where $\widehat{f}(s)$ are the Fourier coefficients given by

$$\widehat{f}(s) = \langle \chi_s, f \rangle = \mathbb{E}_x[f(x) \chi_s(x)]. \quad (2.4)$$

The degree of f is the minimum d such that $\widehat{f}(s) = 0$ if $|s| > d$. We will often use Parseval's identity:

$$\|f\|_2^2 := \langle f, f \rangle = \sum_{s \in \{0, 1\}^n} \widehat{f}(s)^2. \quad (2.5)$$

We will also consider the p -norms of the Fourier spectrum, which are defined as

$$\|\widehat{f}\|_p = \left(\sum_{s \in \{0, 1\}^n} |\widehat{f}(s)|^p \right)^{1/p}.$$

We endow the space of linear maps from M_N to M_N with the inner product $\langle \Phi, \Psi \rangle = \langle J(\Phi), J(\Psi) \rangle / N^2$. An orthonormal basis is defined via the characters

$$\Phi_{x,y}(\rho) = \sigma_x \rho \sigma_y, \quad (2.6)$$

for any $x, y \in \{0, 1, 2, 3\}^n$, where for every $z \in \{0, 1, 2, 3\}^n$ we denote $\sigma_z = \sigma_{z_1} \otimes \cdots \otimes \sigma_{z_n} \in M_N = M_{2^n}$, with σ_i being the corresponding Pauli matrix. The Pauli expansion of one of these linear maps is then defined as

$$\Phi = \sum_{x,y \in \{0,1,2,3\}^n} \widehat{\Phi}(x,y) \Phi_{x,y}, \quad (2.7)$$

where $\widehat{\Phi}(x,y) = \langle \Phi_{x,y}, \Phi \rangle$ are the Pauli coefficients. As $\{\Phi_{x,y}\}_{x,y}$ is an orthonormal basis, we have a version of Parseval's identity

$$\|\Phi\|_2^2 := \langle \Phi, \Phi \rangle = \sum_{x,y \in \{0,1,2,3\}^n} |\widehat{\Phi}(x,y)|^2.$$

We will also consider the p -norms of the Pauli spectrum, which are defined as

$$\|\widehat{\Phi}\|_p = \left(\sum_{x,y \in \{0,1,2,3\}^n} |\widehat{\Phi}(x,y)|^p \right)^{1/p}.$$

We define the degree of Φ in an analogue way to the Fourier degree.

Definition 2.1 Given $\Phi : M_N \rightarrow M_N$ its *degree* is the minimum d such that $\widehat{\Phi}(x,y) = 0$ for any $x, y \in \{0, 1, 2, 3\}^n$ with $|x| + |y| > d$.

3 Bohnenblust–Hille Inequality for the completely bounded norm

In this section we will prove those results concerning the Bohnenblust–Hille Inequality for the completely bounded norm. We will first prove a general result for tensors, from where Theorem 1.1, as well as some other results will follow straightforwardly.

3.1 cb-BH inequality for d -tensors

We consider $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ and a d -tensor

$$T = \sum_{i_1, \dots, i_d=1}^n T_{i_1, \dots, i_d} e_{i_1} \otimes \cdots \otimes e_{i_d} \in \mathbb{K}^n \otimes \cdots \otimes \mathbb{K}^n.$$

In order to simplify notation, we will write $T = (T_{\mathbf{i}})_{\mathbf{i} \in [n]^d}$. Equivalently, T can be regarded as the d -linear form $T : \mathbb{K}^n \times \cdots \times \mathbb{K}^n \rightarrow \mathbb{K}$ given by

$$T(z_1, \dots, z_d) = \sum_{\mathbf{i} \in [n]^d} T_{\mathbf{i}} z_1(i_1) \cdots z_d(i_d).$$

For one such tensor, we denote

$$\|\widehat{T}\|_{\frac{2d}{d+1}} := \left(\sum_{\mathbf{i} \in [n]^d} |T_{\mathbf{i}}|^{\frac{2d}{d+1}} \right)^{\frac{d+1}{2d}} \quad \text{and} \quad \|T\|_{\text{cb}} = \sup \left\| \sum_{\mathbf{i} \in [n]^d} T_{\mathbf{i}} A_{i_1}^1 \cdots A_{i_d}^d \right\|_{\mathcal{B}(\mathcal{H})}, \quad (3.1)$$

where the supremum runs over all (real/complex) Hilbert spaces and all contractions $(A_{i_1}^1)_{i_1=1}^n, \dots, (A_{i_d}^d)_{i_d=1}^n$ in $\mathcal{B}(\mathcal{H})$. In the case $\mathbb{K} = \mathbb{C}$, our definition coincides with the completely bounded norm of T in the category of operator spaces (see for instance [30, Chapter 17]).

Remark 3.1 It is not difficult to see that one can restrict to finite dimensional Hilbert spaces in the definition of $\|T\|_{\text{cb}}$.

The main result of the section is the following cb-BH inequality for d -tensors.

Theorem 3.1 *Let $T \in \mathbb{K}^n \otimes \cdots \otimes \mathbb{K}^n$ be a d -tensor. Then,*

$$\|\widehat{T}\|_{\frac{2d}{d+1}} \leq \|T\|_{\text{cb}}.$$

We will make use of the following lemma, originally due to Blei [9]. A simple proof can be found in [11, Theorem 2.1].

Lemma 3.2 (Blei's inequality) *Given a d -tensor $T \in \mathbb{K}^n \otimes \cdots \otimes \mathbb{K}^n$, we have*

$$\|\widehat{T}\|_{\frac{2d}{d+1}} \leq \left(\prod_{s \in [d]} \sum_{i_s \in [n]} \sqrt{\sum_{i_1, \dots, i_{s-1}, i_{s+1}, \dots, i_d \in [n]} |T_{\mathbf{i}}|^2} \right)^{\frac{1}{d}}.$$

Now, we prove the key technical lemma, from where Theorem 3.1 will follow.

Lemma 3.3 *Let $T \in \mathbb{K}^n \otimes \cdots \otimes \mathbb{K}^n$ be a d -tensor and $s \in [d]$. Then,*

$$\sum_{i_s \in [n]} \sqrt{\sum_{i_1, \dots, i_{s-1}, i_{s+1}, \dots, i_d \in [n]} |T_{i_1, \dots, i_{s-1}, i_s, i_{s+1}, \dots, i_d}|^2} \leq \|T\|_{\text{cb}}.$$

Proof We fix $s \in [d]$. The proof consists of evaluating T on an explicit set of contractions. In order to define these contractions, we denote $m = \sum_{r=0}^{d-s} n^r + \sum_{r=0}^{s-1} n^r$ and let $\{e_i, f_{\mathbf{j}} : \mathbf{i} \in [n]^r, r \in \{0\} \cup [d-s], \mathbf{j} \in [n]^t, t \in \{0\} \cup [s-1]\}$ be an orthonormal

basis of $\ell_2^m(\mathbb{K})$, where we identify $[n]^0$ with \emptyset . For every $i \in [n]$ we define the matrix $Z_i \in M_m$ as:

$$\begin{aligned} Z_i e_{\mathbf{j}} &= e_{(i, \mathbf{j})}, \text{ if } \mathbf{j} \in [n]^r, r \in \{0\} \cup [d-s-1], \\ Z_i e_{\mathbf{j}} &= \frac{\sum_{\mathbf{k} \in [n]^{s-1}} \overline{T}(\mathbf{k}, i, \mathbf{j}) f_{\mathbf{k}}}{\sqrt{\sum_{k_1, \dots, k_{s-1}, k_{s+1}, \dots, k_d \in [n]} |T(k_1, \dots, k_{s-1}, i, k_{s+1}, \dots, k_d)|^2}}, \text{ if } \mathbf{j} \in [n]^{d-s}, \\ Z_i f_{\mathbf{j}} &= \delta_{i, j_t} f_{(j_1, \dots, j_{t-1})}, \text{ if } \mathbf{j} \in [n]^t, t \in \{0\} \cup [s-1], \\ Z_i f_{\emptyset} &= 0, \end{aligned}$$

where $\bar{\alpha}$ denotes the complex conjugate of α .

Assume for the moment that Z_i are contractions. One can easily check that

$$\langle f_{\emptyset}, Z_{i_1} \dots Z_{i_d} e_{\emptyset} \rangle = \frac{\overline{T}_{i_1, \dots, i_d}}{\sqrt{\sum_{k_1, \dots, k_{s-1}, k_{s+1}, \dots, k_d \in [n]} |T(k_1, \dots, k_{s-1}, i_s, k_{s+1}, \dots, k_d)|^2}}.$$

Hence, by assuming that Z_i are contractions, we can conclude

$$\begin{aligned} \|T\|_{\text{cb}} &\geq \left\| \sum_{\mathbf{i} \in [n]^d} T_{\mathbf{i}} Z_{i_1} \dots Z_{i_d} \right\|_{B(\ell_2^m(\mathbb{K}))} \geq \sum_{\mathbf{i} \in [n]^d} T_{\mathbf{i}} \langle f_{\emptyset} | Z_{i_1} \dots Z_{i_d} | e_{\emptyset} \rangle \\ &\geq \sum_{\mathbf{i} \in [n]^d} T_{\mathbf{i}} \frac{\overline{T}_{\mathbf{i}}}{\sqrt{\sum_{k_1, \dots, k_{s-1}, k_{s+1}, \dots, k_d \in [n]} |T(k_1, \dots, k_{s-1}, i_s, k_{s+1}, \dots, k_d)|^2}} \\ &= \sum_{i_s \in [n]} \sqrt{\sum_{i_1, \dots, i_{s-1}, i_{s+1}, \dots, i_d \in [n]} |T_{i_1, \dots, i_{s-1}, i_s, i_{s+1}, \dots, i_d}|^2}, \end{aligned}$$

as desired.

Thus, it remains to prove that the matrices Z_i are contractions. Given that Z_i maps the sets $\{e_{\mathbf{i}} : \mathbf{i} \in [n]^r, r \in \{0\} \cup [d-s-1]\}$, $\{e_{\mathbf{i}} : \mathbf{i} \in [n]^{d-s}\}$ and $\{f_{\mathbf{i}} : \mathbf{i} \in [n]^t, t \in \{0\} \cup [s-1]\}$ to orthogonal subspaces, it suffices to show that the Z_i are contractions when restricted to those subspaces. For the first and third sets that is clear since Z_i maps each basis vector of those sets either to a different basis vector or to 0. For the second set, just note that for every $\lambda \in \mathbb{K}^{n^{d-s}}$ we have

$$\begin{aligned} \left\| Z_i \sum_{\mathbf{j} \in [n]^{d-s}} \lambda_{\mathbf{j}} e_{\mathbf{j}} \right\|_2^2 &= \left\| \frac{\sum_{\mathbf{k} \in [n]^{s-1}} \left(\sum_{\mathbf{j} \in [n]^{d-s}} \lambda_{\mathbf{j}} \overline{T}_{\mathbf{k} \mathbf{i} \mathbf{j}} \right) f_{\mathbf{k}}}{\sqrt{\sum_{k_1, \dots, k_{s-1}, k_{s+1}, \dots, k_d \in [n]} |T(k_1, \dots, k_{s-1}, i, k_{s+1}, \dots, k_d)|^2}} \right\|_2^2 \\ &= \frac{\sum_{\mathbf{k} \in [n]^{s-1}} \left| \sum_{\mathbf{j} \in [n]^{d-s}} \lambda_{\mathbf{j}} \overline{T}_{\mathbf{k} \mathbf{i} \mathbf{j}} \right|^2}{\sum_{k_1, \dots, k_{s-1}, k_{s+1}, \dots, k_d \in [n]} |T(k_1, \dots, k_{s-1}, i, k_{s+1}, \dots, k_d)|^2} \end{aligned}$$

$$\begin{aligned} &\leq \frac{\left(\sum_{\mathbf{k} \in [n]^{s-1}} \sum_{\mathbf{j} \in [n]^{d-s}} |T_{\mathbf{k}\mathbf{j}}|^2\right) \left(\sum_{\mathbf{j} \in [n]^{d-s}} |\lambda_{\mathbf{j}}|^2\right)}{\sum_{k_1, \dots, k_{s-1}, k_{s+1}, \dots, k_d \in [n]} |T_{k_1, \dots, k_{s-1}, i, k_{s+1}, \dots, k_d}|^2} \\ &= \sum_{\mathbf{j} \in [n]^{d-s}} |\lambda_{\mathbf{j}}|^2 = \|\lambda\|_2^2, \end{aligned}$$

where we have used Cauchy-Schwarz for the sum over \mathbf{j} . \square

Proof of Theorem 3.1 According to Lemma 3.2 and Lemma 3.3 we have

$$\|\widehat{T}\|_{\frac{2d}{d+1}} \leq \left(\prod_{s \in [d]} \sum_{i_s \in [n]} \sqrt{\sum_{i_1, \dots, i_{s-1}, i_{s+1}, \dots, i_d \in [n]} |T_{\mathbf{i}}|^2} \right)^{\frac{1}{d}} \leq \left(\prod_{s \in [d]} \|T\|_{\text{cb}} \right)^{\frac{1}{d}} = \|T\|_{\text{cb}}.$$

\square

In order to extend Theorem 3.1 to homogeneous polynomials we need the following proposition, which was proven in [7, Proposition 3.2], and states that the polarization constant for the completely bounded norm is 1.

Proposition 3.4 *Let $T \in \mathbb{K}^n \otimes \cdots \otimes \mathbb{K}^n$ be a d -tensor. Then,*

$$\|T\|_{\text{cb}} = \sup \left\| \sum_{\mathbf{i} \in [n]^d} T_{\mathbf{i}} Z_{i_1} \cdots Z_{i_d} \right\|_{\mathcal{B}(\mathcal{H})}, \quad (3.2)$$

where the supremum runs over all (real/complex) Hilbert spaces and all contractions $(Z_i)_{i=1}^n$ in $\mathcal{B}(\mathcal{H})$.

3.2 cb-BH inequality for polynomials

Now we consider the case of polynomials. Given any (not necessarily homogeneous) polynomial of degree d in n variables $P : \mathbb{K}^n \rightarrow \mathbb{K}$, we can write it as

$$P = \sum_{s \in \{0\} \cup [d]} P_s, \quad (3.3)$$

where $P_s : \mathbb{K}^n \rightarrow \mathbb{K}$ is a s -homogeneous polynomial. We denote, given $s \in [d]$,

$$\mathcal{J}(s, n) = \{(j_1, \dots, j_s) \in [n]^s : j_1 \leq \dots \leq j_s\}.$$

Then, P_s can be written uniquely as

$$P_s(x) = \sum_{\mathbf{j} \in \mathcal{J}(s, n)} a_{\mathbf{j}} x_{\mathbf{j}}, \quad (3.4)$$

where we denote $x_{\mathbf{j}} = x_{j_1} \cdots x_{j_s}$. Hence, we can define the completely bounded norm of P as

$$\|P\|_{\text{cb}} = \sup \left\| \sum_{s \in \{0\} \cup [d]} \sum_{\mathbf{j} \in \mathcal{J}(s, n)} a_{\mathbf{j}} Z_{j_1} \cdots Z_{j_s} \right\|_{\mathcal{B}(\mathcal{H})},$$

where the supremum runs over all (real/complex) Hilbert spaces \mathcal{H} and all contractions Z_i in $\mathcal{B}(\mathcal{H})$ for $i = 1, \dots, n$.

Theorem 1.1, which refers to the d -homogeneous case, follows easily from Theorem 3.1.

Proof of Theorem 1.1 Given a d -homogeneous polynomial $P : \mathbb{K}^n \rightarrow \mathbb{K}$ as above, we want to prove that

$$\left(\sum_{\mathbf{j} \in \mathcal{J}(d, n)} |a_{\mathbf{j}}|^{\frac{2d}{d+1}} \right)^{\frac{d+1}{2d}} \leq \|P\|_{\text{cb}}.$$

To do that, we reduce it to the case of tensors. We define $T_{\mathbf{j}} = a_{\mathbf{j}}$ for every $\mathbf{j} \in \mathcal{J}(d, n)$ and $T_{\mathbf{j}} = 0$ for every $\mathbf{j} \in [n]^d \setminus \mathcal{J}(d, n)$. By Proposition 3.4, the tensor T satisfies

$$\left(\sum_{\mathbf{j} \in \mathcal{J}(d, n)} |a_{\mathbf{j}}|^{\frac{2d}{d+1}} \right)^{\frac{d+1}{2d}} = \left(\sum_{\mathbf{j} \in [n]^d} |T_{\mathbf{j}}|^{\frac{2d}{d+1}} \right)^{\frac{d+1}{2d}} \quad \text{and} \quad \|T\|_{\text{cb}} = \|P\|_{\text{cb}}.$$

Hence, the result follows from Theorem 3.1. \square

We will now turn our attention to the case of general polynomials. To this end, we first prove the following result:

Lemma 3.5 *Let $P : \mathbb{K}^n \rightarrow \mathbb{K}$ be a polynomial of degree d . Then,*

$$\|P\|_{\text{cb}} \geq \frac{1}{\sqrt{d+1}} \sup_{s \in [d] \cup \{0\}} \left\| u, \sum_{\substack{\alpha \in \mathbb{N}_0^n \\ \sum_i \alpha_i = s}} a_{\alpha} Z_1^{\alpha_1} \cdots Z_n^{\alpha_n} v_s \right\|,$$

where the supremum runs over all (real/complex) Hilbert spaces \mathcal{H} , all contractions Z_1, \dots, Z_n in $\mathcal{B}(\mathcal{H})$ and all vectors u, v_s in \mathcal{H} with norm less than or equal one.

Proof For $s \in \{0\} \cup [d]$, let $b_s \in \mathbb{K}$ be such that $|b_s| = 1$ and

$$\left\| u, \sum_{\substack{\alpha \in \mathbb{N}_0^n \\ \sum_i \alpha_i = s}} a_{\alpha} Z_1^{\alpha_1} \cdots Z_n^{\alpha_n} v_s \right\| = b_s \left\| u, \sum_{\substack{\alpha \in \mathbb{N}_0^n \\ \sum_i \alpha_i = s}} a_{\alpha} Z_1^{\alpha_1} \cdots Z_n^{\alpha_n} v_s \right\|.$$

Let $\{e_0, \dots, e_d\}$ be the canonical basis of \mathbb{K}^{d+1} . We define the unitary operator $B : \mathbb{K}^{d+1} \rightarrow \mathbb{K}^{d+1}$ such that $B(e_{p+1}) = e_p$ for every $p \in [d]$ and $B(e_0) = e_d$. We

also define the unit vectors

$$\xi = \frac{1}{\sqrt{d+1}} \sum_{q \in [d] \cup \{0\}} b_q v_q \otimes e_q \in \mathcal{H} \otimes \mathbb{K}^{d+1} \quad \text{and} \quad \eta = u \otimes e_0 \in \mathcal{H} \otimes \mathbb{K}^{d+1}.$$

Finally, we consider the new contractions $\tilde{Z}_i = Z_i \otimes B$ in $\mathcal{B}(\mathcal{H} \otimes \mathbb{K}^{d+1})$ for $i = 1, \dots, n$. Then, one can easily check that

$$\left\langle \tilde{u}, \sum_{s \in [d] \cup \{0\}} \sum_{\substack{\alpha \in \mathbb{N}_0^n \\ \sum_i \alpha_i = s}} a_\alpha \tilde{Z}_1^{\alpha_1} \dots \tilde{Z}_n^{\alpha_n} \xi \right\rangle = \frac{1}{\sqrt{d+1}} \left\| u, \sum_{\substack{\alpha \in \mathbb{N}_0^n \\ \sum_i \alpha_i = s}} a_\alpha Z_1^{\alpha_1} \dots Z_n^{\alpha_n} v_s \right\|,$$

from where the statement follows. \square

We can now prove a cb-BH inequality for general polynomials of degree d .

Corollary 3.6 *Let $P : \mathbb{K}^n \rightarrow \mathbb{K}$ be a polynomial of degree d . Then,*

$$\|\hat{P}\|_{\frac{2d}{d+1}} \leq \sqrt{d+1} \|P\|_{\text{cb}}.$$

Proof Let $Q : \mathbb{K}^{n+1} \rightarrow \mathbb{K}$ be the s -homogeneous polynomial defined by

$$Q(x, x_{n+1}) := \sum_{s \in \{0\} \cup [d]} P_s(x) x_{n+1}^{d-s},$$

where $x = (x_1, \dots, x_n)$ and P_s is the d -homogeneous part of P .

It is clear that $\|\hat{Q}\|_{\frac{2d}{d+1}} = \|\hat{P}\|_{\frac{2d}{d+1}}$. On the other hand, we have

$$\|Q\|_{\text{cb}} = \sup \left\langle u, \sum_{s \in \{0\} \cup [d]} \sum_{\substack{\alpha \in \mathbb{N}_0^{n+1} \\ \sum_i \alpha_i = s}} a_\alpha Z_1^{\alpha_1} \dots Z_n^{\alpha_n} Z_{n+1}^{d-s} v \right\rangle,$$

where the sup is taken over all (real/complex) Hilbert spaces \mathcal{H} , all contractions $Z_1, \dots, Z_{n+1} \in \mathcal{B}(\mathcal{H})$ and all unit vectors u and v . Then, by defining $v_s = Z_{n+1}^{d-s} |v\rangle$ we can use Lemma 3.5 to deduce

$$\|Q\|_{\text{cb}} \leq \sqrt{d+1} \|P\|_{\text{cb}}.$$

Hence, applying Theorem 1.1 to Q concludes the proof. \square

3.3 Optimality of the cb-BH inequality

We conclude this section by proving the optimality of Theorem 1.1. We will actually prove the optimality of Theorem 3.1, from where the optimality in the exponent

for the corresponding cb-BH inequality for d -multilinear forms and d -homogeneous polynomials follows.

First of all, note that constant one is the best possible in the inequality since the d -linear form $T(x_1, \dots, x_n) = x_1$ satisfies $\|\widehat{T}\|_{2d/(d+1)} = \|T\|_{\text{cb}} = 1$. Regarding the optimality in the exponent, it follows from the next statement.

Theorem 3.7 *Let $d \in \mathbb{N}$, let $\mathbb{K} \in \{\mathbb{R}, \mathbb{C}\}$ and let $q \geq 1$. For infinitely many $n \in \mathbb{N}$, there exists a d -tensor $T \in \mathbb{K}^n \otimes \dots \otimes \mathbb{K}^n$ such that $\|T\|_q = n^{\frac{d}{q}}$ and $\|T\|_{\text{cb}} \leq n^{\frac{d+1}{2}}$.*

The optimality in the exponent of Theorem 3.1 follows easily from the previous statement. Indeed, suppose that there is a constant $C_d > 0$ such that

$$\|T\|_q \leq C_d \|T\|_{\text{cb}}.$$

Then, it follows that

$$n^{\frac{d}{q}} \leq C_d n^{\frac{d+1}{2}}$$

for every $n \in \mathbb{N}$. Therefore, $q \geq 2d/(d+1)$. In order to see that this last estimate implies the optimality for the BH inequality for d -homogeneous polynomials (Theorem 1.1) just note that for any d -linear form $T : \mathbb{K}^n \times \dots \times \mathbb{K}^n \rightarrow \mathbb{K}$, we can define a d -homogeneous polynomial in $d \times n$ variables $P : (\mathbb{K}^n)^d \rightarrow \mathbb{K}$ defined as

$$P((x_1(i_1))_{i_1}, \dots, (x_d(i_d))_{i_d}) = \sum_{i_1, \dots, i_d=1}^n T_{i_1, \dots, i_d} x_1(i_1) \cdots x_d(i_d).$$

The optimality of Theorem 1.1 follows because if we consider the lexicographical order in $[n]^d$, then $\|P\|_{2d/(d+1)} = \|T\|_{2d/(d+1)}$ and $\|P\|_{\text{cb}} = \|T\|_{\text{cb}}$.

To prove 3.7 we will use the following well-known non-commutative version of the Cauchy-Schwarz inequality.

Lemma 3.8 *Let \mathcal{H}_1 and \mathcal{H}_2 be two real/complex Hilbert spaces and consider $A_i \in \mathcal{B}(\mathcal{H}_1, \mathcal{H}_2)$, $B_i \in \mathcal{B}(\mathcal{H}_2, \mathcal{H}_1)$ for $i \in [n]$. Then,*

$$\left\| \sum_{i \in [n]} A_i B_i \right\|_{\mathcal{B}(\mathcal{H}_2)}^2 \leq \left\| \sum_{i \in [n]} A_i A_i^\dagger \right\|_{\mathcal{B}(\mathcal{H}_2)} \left\| \sum_{i \in [n]} B_i^\dagger B_i \right\|_{\mathcal{B}(\mathcal{H}_2)}.$$

Our proof is based on the proof of the optimality of the exponent in the classical BH inequality (see [16, Chapter 4]).

Proof of Theorem 3.7 Let $n \in \mathbb{N}$ and let $N = 2^n$. We will identify $[N]$ with $\mathcal{P}(n)$ (the family of subsets of n elements) and $\{-1, 1\}^n$ in an arbitrary bijective way. In this sense, we define the matrix $a \in \mathbb{R}^{N \times N}$ via

$$a_{(x,S)} = \prod_{i \in S} x_i,$$

for every $x \in \{-1, 1\}^n$ and $S \subseteq [n]$. This matrix satisfies that

$$|a_{(x,S)}| = 1, \quad (3.5)$$

$$\sum_{x \in \{-1, 1\}^n} a_{(x,S)} a_{(x,S')} = N \delta_{S,S'}. \quad (3.6)$$

We define the d -tensor $T \in \mathbb{K}^n \otimes \cdots \otimes \mathbb{K}^n$ by

$$T = \sum_{\mathbf{i} \in [N]^d} a_{(i_1, i_2)} \cdots a_{(i_{d-1}, i_d)} e_{i_1} \otimes \cdots \otimes e_{i_d}.$$

According to Eq. (3.5) we immediately deduce that $\|T\|_q = N^{\frac{d}{q}}$.

In order to prove the upper bound for $\|T\|_{\text{cb}}$ we can restrict to finite-dimensional contractions, thanks to Remark 3.1. Furthermore, we can restrict supremum of Eq. (3.1) to unitary (orthogonal for the real case) matrices, because every contraction is a convex combination of unitary matrices.

Now, given arbitrary unitary matrices $U_{i_1}^1, \dots, U_{i_d}^d$, if we denote

$$R_{i_1} = \sum_{\mathbf{j} \in [N]^{d-1}} a_{(i_1, j_2)} \cdots a_{(j_{d-1}, j_d)} U_{j_2}^2 \cdots U_{j_d}^d,$$

we can apply Lemma 3.8 to write

$$\begin{aligned} \left\| \sum_{\mathbf{i} \in [N]^d} a_{(i_1, i_2)} \cdots a_{(i_{d-1}, i_d)} U_{j_1}^1 \cdots U_{j_d}^d \right\| &\leq \left\| \sum_{i_1 \in [N]} U_{i_1}^1 (U_{i_1}^1)^\dagger \right\|^{\frac{1}{2}} \left\| \sum_{i_1 \in [N]} R_{i_1}^\dagger R_{i_1} \right\|^{\frac{1}{2}} \\ &= N^{\frac{1}{2}} \left\| \sum_{i_1 \in [N]} R_{i_1}^\dagger R_{i_1} \right\|^{\frac{1}{2}}. \end{aligned}$$

Now, we note that $\sum_{i_1 \in [N]} R_{i_1}^\dagger R_{i_1}$ can be written as

$$\begin{aligned} \sum_{\mathbf{j}, \mathbf{k} \in [N]^{d-1}} \left(\sum_{i_1 \in [N]} a_{(i_1, j_2)} a_{(i_1, k_2)} \right) &a_{(j_2, j_3)} \cdots a_{(j_{d-1}, j_d)} a_{(k_2, k_3)} \\ &\cdots a_{(k_{d-1}, k_d)} (U_{j_d}^d)^\dagger \cdots (U_{j_2}^2)^\dagger U_{k_2}^2 \cdots U_{k_d}^d. \end{aligned}$$

By using Eq. (3.6) and that $U_i^2 = \text{Id}$, for $i \in [N]$, the previous expression equals

$$\begin{aligned} N \sum_{i_2 \in [N]} \sum_{\mathbf{j}, \mathbf{k} \in [N]^{d-2}} &a_{(i_2, j_3)} \cdots a_{(j_{d-1}, j_d)} a_{(i_2, k_3)} \\ &\cdots a_{(k_{d-1}, k_d)} (U_{j_d}^d)^\dagger \cdots (U_{j_3}^3)^\dagger U_{k_3}^3 \cdots U_{k_d}^d \end{aligned}$$

$$\begin{aligned}
&= N \sum_{\mathbf{j}, \mathbf{k} \in [N]^{d-2}} \left(\sum_{i_2 \in [N]} a_{(i_2, j_3)} a_{(i_2, k_3)} \right) a_{(j_3, j_4)} \\
&\quad \cdots a_{(j_{d-1}, j_d)} \cdots a_{(k_3, k_4)} a_{(k_{d-1}, k_d)} \\
&\quad (U_{j_d}^d)^\dagger \cdots (U_{j_3}^3)^\dagger U_{k_3}^3 \cdots U_{k_d}^d.
\end{aligned}$$

We see that we can iterate this process to obtain

$$\left\| \sum_{i_1 \in [N]} R_{i_1}^\dagger R_{i_1} \right\| \leq N^{d-1} \left\| \sum_{i_d \in [N]} (U_{i_d}^d)^\dagger U_{i_d}^d \right\| = N^d.$$

Therefore, we conclude that $\left\| \sum_{\mathbf{i} \in [N]^d} a_{(i_1, i_2)} \cdots a_{(i_{d-1}, i_d)} U_{i_1}^1 \cdots U_{i_d}^d \right\| \leq N^{\frac{d+1}{2}}$. \square

Remark 3.2 The d -linear form used in the proof of Theorem 3.7 also plays a central role in quantum query complexity. Indeed, it is the linear form determined by the d -forrelation problem, that optimally separates quantum and classical query complexity [1, 12]. More precisely, given d Boolean functions $f_1, \dots, f_d : \{0, 1\}^n \rightarrow \{-1, 1\}$, its d -forrelation is defined as

$$\begin{aligned}
&\text{forr}_d(f_1, \dots, f_d) \\
&= \frac{1}{2^{n \frac{d+1}{2}}} \sum_{x_1, \dots, x_d \in \{0, 1\}^n} f(x_1) (-1)^{\langle x_1, x_2 \rangle} \cdots f(x_{d-1}) (-1)^{\langle x_{d-1}, x_d \rangle} f(x_d).
\end{aligned}$$

Thus, if we consider the d -linear form T defined in the proof of Theorem 3.7 and we identify the d functions f_1, \dots, f_d with the elements of $\{-1, 1\}^{2^n}$ determined by their truth table, we have

$$T(f_1, \dots, f_d) = 2^{n \frac{d+1}{2}} \text{forr}_d(f_1, \dots, f_d).$$

4 BH-inequality in other contexts

4.1 Boolean functions

We determine the exact value of the BH constant for Boolean functions. This result follows from the well-known fact that the Fourier coefficients of Boolean functions are multiples of $2^{1-d} \mathbb{Z}$. This property is usually referred to as the granularity of Boolean functions [29, Exercise 1.11]. We sketch the proof below for the sake of completeness.

Lemma 4.1 *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ with degree at most d . Then, $\widehat{f}(s) \in 2^{1-d} \mathbb{Z}$ for every $s \in \{0, 1\}^n$.*

Proof Recall that $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$. We define $g : \{0, 1\}^n \rightarrow \{0, 1\}$ by

$$g(z) = \frac{1}{2} \left(1 - f((1 - 2z_1), \dots, (1 - 2z_n)) \right).$$

It is not difficult to see that g can be written in a unique way as

$$g(z) = \sum_{s \in \{0,1\}^n} c_s \prod_{i:s_i=1} z_i$$

for some coefficients $c_s \in \mathbb{R}$ such that $c_s = 0$ for every s with $|s| > d$. By applying induction on $|s|$, one can actually prove that $c_s \in \mathbb{Z}$ for every s . Indeed, we first note that for $s = \emptyset$, one has $c_{\emptyset^n} = g(0^n) \in \{0, 1\}$. For s with $|s| = t + 1 > 0$, assuming that $c_s \in \mathbb{Z}$ for every s with $|s| \leq t$, we have

$$c_s = g(s) - \sum_{|s'| < |s|, s'_i \leq s_i} c_{s'},$$

so c_s belongs to \mathbb{Z} . Finally, the statement for f can be obtained by just noticing that

$$f(x) = 1 - 2g\left(\frac{1-x_1}{2}, \dots, \frac{1-x_n}{2}\right) = 1 - 2 \sum_{|s| \leq d} c_s \prod_{i:s_i=1} \frac{1-x_i}{2}.$$

□

Proposition 4.2 *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ of degree at most d . Then,*

$$\left(\sum_{s \in \{0,1\}^n} |\widehat{f}(s)|^{\frac{2d}{d+1}} \right)^{\frac{d+1}{2d}} \leq 2^{\frac{d-1}{d}}.$$

The equality is witnessed by the address function.

Proof Since for Boolean functions one has $\|f\|_2 = 1$, Lemma 4.1 and Parseval's identity imply that f has at most $2^{2(d-1)}$ non-zero Fourier coefficients. Indeed, this immediately follows from the identity $\sum_s |\widehat{f}(s)|^2 = 1$ and the fact that $|\widehat{f}(s)| \geq 2^{1-d}$ for every non-zero coefficient. Hence, Hölder's inequality implies that, for $p \in [1, 2)$,

$$\sum_{s: \widehat{f}(s) \neq 0} |\widehat{f}(s)|^p \cdot 1 \leq \left(\sum_{s \in \{0,1\}^n} \widehat{f}^2(s) \right)^{\frac{p}{2}} \left(2^{2(d-1)} \right)^{\frac{2-p}{2}} = 2^{(d-1)(2-p)}.$$

Taking $p = 2d/(d+1)$ the claimed inequality follows.

The equality is witnessed by the *address function* $f : (\{-1, 1\}^n)^d \rightarrow \{-1, 1\}$ of degree d and $n = 2^{d-1}$, which is defined as

$$f(x) = \sum_{a \in \{-1,1\}^{d-1}} \underbrace{\frac{x_1(1) - a_1 x_1(2)}{2} \cdots \frac{x_{d-1}(1) - a_{d-1} x_{d-1}(2)}{2}}_{g_a(x_1, \dots, x_{d-1})} x_d(a), \quad (4.1)$$

where we identify $\{-1, 1\}^{d-1}$ with $[2^{d-1}]$ in the canonical way. The address function is Boolean because for every $(x_1, \dots, x_{d-1}) \in (\{-1, 1\}^n)^{d-1}$ there is only one $a \in$

$\{-1, 1\}^{d-1}$ such that $g_a(x_1, \dots, x_{d-1})$ is not 0, in which case it takes the value ± 1 . Given that it has $2^{2(d-1)}$ Fourier coefficients and all of them equal 2^{1-d} , we have that

$$\left(\sum_{s \in \{0, 1\}^n} |\widehat{f}(s)|^{\frac{2d}{d+1}} \right)^{\frac{d+1}{2d}} = 2^{1-d} 2^{2(d-1) \cdot \frac{d+1}{2d}} = 2^{\frac{d-1}{d}},$$

as promised. \square

4.2 A non-commutative BH inequality

In this section, we prove a Bohnenblust–Hille inequality for linear maps that are bounded in the S_1 to S_∞ norm, such as quantum channels. Recall from Eq. (2.3) that any such a function can be written as

$$f = \sum_{s \in \{0, 1\}^n} \widehat{f}(s) \chi_s,$$

and it has degree d if this is the minimal number for which $\widehat{f}(s) = 0$ if $|s| > d$. The following result was proved originally in [10], and with a better constant in [18].

Theorem 4.3 *Let $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ be a function of degree at most d . Then,*

$$\|\widehat{f}\|_{\frac{2d}{d+1}} \leq C \sqrt{d \log d} \|f\|_\infty,$$

where $C > 0$ is a constant.

In order to prove Theorem 1.3 we follow a similar argument to the one used in [35]. However, we need to modify their argument in order to consider maps from S_1 to S_∞ and not just matrices in M_N . In fact, as we explain in Remark 4.1, Theorem 1.3 generalizes the non-commutative BH inequality proved in [35].

For every $\Phi : M_N \rightarrow M_N$, we will assign it a function $f_\Phi : \{-1, 1\}^{3n} \times \{-1, 1\}^{3n} \rightarrow \mathbb{C}$ whose Fourier spectrum will be closely related to the one of $\widehat{\Phi}$, as shown in Lemma 4.4, and then we will be able to reduce to Theorem 4.3. The function f_Φ is defined as follows. For $a = (a^1, a^2, a^3)$, $b = (b^1, b^2, b^3) \in \{-1, 1\}^n \times \{-1, 1\}^n \times \{-1, 1\}^n$ and $s, t \in \{1, 2, 3\}^n$, define the following matrices (which are not necessarily states)

$$|a^s\rangle\langle b^t| = \bigotimes_{i \in [n]} |\chi_{a_i^{s(i)}}^{s(i)}\rangle\langle \chi_{b_i^{t(i)}}^{t(i)}|,$$

Here $|\chi_a^s\rangle$ is the eigenvector of σ_s with eigenvalue a . The function $f_\Phi : \{-1, 1\}^{3n} \times \{-1, 1\}^{3n} \rightarrow \mathbb{C}$ is then given by

$$f_\Phi(a, b) = \frac{1}{9^n} \sum_{s, t \in \{1, 2, 3\}^n} \text{Tr}[\Phi(|a^s\rangle\langle b^t|) |b^t\rangle\langle a^s|].$$

We recall the reader that any function $\Phi : M_N \rightarrow M_N$ can be expressed as

$$\Phi(\rho) = \sum_{x, y \in \{0, 1, 2, 3\}^n} \widehat{\Phi}(x, y) \cdot \sigma_x \rho \sigma_y, \quad (4.2)$$

where $\sigma_x = \otimes_{i \in [n]} \sigma_{x_i}$ and σ_i for $i \in \{0, 1, 2, 3\}$ are the Pauli matrices. Then, if $|x|$ denotes the number of non-zero entries of $x \in \{0, 1, 2, 3\}^n$, the degree of Φ is the minimum integer d such that $\widehat{\Phi}(x, y) = 0$ if $|x| + |y| > d$.

In the following lemma, the key properties of the function f are presented.

Lemma 4.4 *Let $\Phi : M_N \rightarrow M_N$ be a function of degree at most d . Then, f_Φ has also degree d . Moreover, $|f_\Phi(a, b)| \leq \|\Phi\|_{S_1 \rightarrow S_\infty}$ for all a, b and $\|\widehat{\Phi}\|_p \leq 3^d \|\widehat{f}_\Phi\|_p$.*

Proof We first show the bound on $|f_\Phi|$. Given that $|a^s\rangle\langle b^t|$ is a rank one operator such that $\| |a^s\rangle\|_2 = \| |b^t\rangle\|_2 = 1$, we conclude that

$$\| |a^s\rangle\langle b^t| \|_{S_1} = 1. \quad (4.3)$$

Thus, we have that:

$$\begin{aligned} |f_\Phi(a, b)| &\leq \frac{1}{9^n} \sum_{s, t \in \{1, 2, 3\}^n} |\text{Tr}[\Phi(|a^s\rangle\langle b^t|) |b^t\rangle\langle a^s|]| \\ &\leq \frac{1}{9^n} \sum_{s, t \in \{1, 2, 3\}^n} \|\Phi(|a^s\rangle\langle b^t|)\|_{S_\infty} \| |b^t\rangle\langle a^s| \|_{S_1} \\ &\leq \frac{1}{9^n} \sum_{s, t \in \{1, 2, 3\}^n} \|\Phi\|_{S_1 \rightarrow S_\infty} \| |a^s\rangle\langle b^t| \|_{S_1} \| |b^t\rangle\langle a^s| \|_{S_1} \\ &\leq \frac{1}{9^n} \sum_{s, t \in \{1, 2, 3\}^n} \|\Phi\|_{S_1 \rightarrow S_\infty} = \|\Phi\|_{S_1 \rightarrow S_\infty}, \end{aligned}$$

where in the first inequality we have used the triangle inequality, in the second inequality the duality between S_1 and S_∞ , in the third the definition of $S_1 \rightarrow S_\infty$ norm and in the fourth inequality we have used Eq. (4.3).

We now prove the estimate $\|\widehat{\Phi}\|_p \leq 3^{-d} \|\widehat{f}_\Phi\|_p$ and also that the degree of f_Φ is d . To this end, it suffices to show that

$$f_\Phi(a, b) = \sum_{x, y \in \{0, 1, 2, 3\}^n} \frac{\widehat{\Phi}(x, y)}{3^{|x|+|y|}} \prod_{i \in \text{supp}(x)} \prod_{j \in \text{supp}(y)} a_i^{x(i)} b_j^{y(j)}, \quad (4.4)$$

where $\text{supp}(x) = \{i \in [n] : x_i \neq 0\}$ and $|x|$ is the size of $\text{supp}(x)$. Indeed, this follows from the fact that for every $x, y \in \{0, 1, 2, 3\}^n$, the product $\prod_{i \in \text{supp}(x)} \prod_{j \in \text{supp}(y)} a_i^{x(i)} b_j^{y(j)}$ can be read as $\chi_{S_{x, y}}(a, b)$ for a certain $S_{x, y} \in \{-1, 1\}^{6n}$ satisfying that $S_{x, y} \neq S_{x', y'}$ whenever $(x, y) \neq (x', y')$.

To prove Eq. (4.4) the key is observing that for every $s, t \in \{1, 2, 3\}$, $x, y \in \{0, 1, 2, 3\}$ and $a, b \in \{-1, 1\}$, we have that

$$\text{Tr}[\sigma_x |\chi_a^s\rangle \langle \chi_b^t| \sigma_y |\chi_b^t\rangle \langle \chi_a^s|] = \begin{cases} 0 & \text{if } (s \neq x \text{ and } x \neq 0) \text{ or } (t \neq y \text{ and } y \neq 0), \\ 1 & \text{if } x = 0 \text{ and } y = 0, \\ a & \text{if } s = x \text{ and } y = 0, \\ b & \text{if } x = 0 \text{ and } t = y, \\ ab & \text{if } s = x \text{ and } y = t. \end{cases}$$

Hence, taking tensor products we have that for every $s, t \in \{1, 2, 3\}^n$, $x, y \in \{0, 1, 2, 3\}^n$ and $a = (a^1, a^2, a^3)$, $b = (b^1, b^2, b^3) \in \{-1, 1\}^n \times \{-1, 1\}^n \times \{-1, 1\}^n$, it holds that

$$\begin{aligned} & \text{Tr}[\sigma_x |a^s\rangle \langle b^t| \sigma_y |b^t\rangle \langle a^s|] \\ &= \langle \chi_a^s | \sigma_x | \chi_a^s \rangle \langle \chi_b^t | \sigma_y | \chi_b^t \rangle = \prod_{i \in \text{supp } x} \prod_{j \in \text{supp } y} a_i^{x(i)} b_j^{y(j)} \delta_{x(i), s(i)} \delta_{y(j), t(j)}. \end{aligned}$$

In particular, it follows that

$$\begin{aligned} f_{\Phi_{x,y}}(a, b) &\equiv \frac{1}{9^n} \sum_{s, t \in \{1, 2, 3\}^n} \text{Tr}[\sigma_x |a^s\rangle \langle b^t| \sigma_y |b^t\rangle \langle a^s|] \\ &= \frac{1}{9^n} \prod_{i \in \text{supp } x} \prod_{j \in \text{supp } y} a_i^{x(i)} b_j^{y(j)} \sum_{s \in \mathcal{X}, t \in \mathcal{Y}} 1, \end{aligned}$$

where $\mathcal{X} = \{s \in \{1, 2, 3\}^n : s(i) = x(i) \forall i \in \text{supp}(x)\}$. Since $|\mathcal{X}| = 3^{n-|x|}$, Eq. (4.4) follows for $\Phi_{x,y}$. Finally, Eq. (4.4) follows in general because

$$f_{\Phi}(a, b) = \sum_{x, y \in \{0, 1, 2, 3\}^n} \widehat{\Phi}(x, y) f_{\Phi_{x,y}}(a, b).$$

□

Proof of Theorem 1.3 Let $\Re f_{\Phi} : \{-1, 1\}^{6n} \rightarrow \mathbb{R}$ be defined as $(\Re f_{\Phi})(x) = \Re(f_{\Phi}(x))$ and $\Im f_{\Phi} : \{-1, 1\}^{6n} \rightarrow \mathbb{R}$ as $(\Im f_{\Phi})(x) = \Im(f_{\Phi}(x))$. Note that we have that $\widehat{f_{\Phi}} = \widehat{\Re f_{\Phi}} + i \widehat{\Im f_{\Phi}}$. By Lemma 4.4,

$$|(\Re f_{\Phi})(a, b)|, |(\Im f_{\Phi})(a, b)| \leq |f_{\Phi}(x)| \leq \|\Phi\|_{S_1 \rightarrow S_{\infty}},$$

and that the degree of both the real and imaginary part is at most d . Hence, by the triangle inequality and Theorem 4.3 we have

$$\|\widehat{f_{\Phi}}\|_{\frac{2d}{d+1}} \leq \|\widehat{\Re f_{\Phi}}\|_{\frac{2d}{d+1}} + \|\widehat{\Im f_{\Phi}}\|_{\frac{2d}{d+1}} \leq C^{\sqrt{d \log d}} \|\Phi\|_{S_1 \rightarrow S_{\infty}}.$$

Thus, using that $\|\widehat{\Phi}\|_{2d/(d+1)} \leq 3^d \|\widehat{f_{\Phi}}\|_{2d/(d+1)}$, it follows that $\|\widehat{\Phi}\|_{2d/(d+1)} \leq C^d \|\Phi\|_{S_1 \rightarrow S_{\infty}}$. □

Corollary 4.5 *Let $\Phi : M_N \rightarrow M_N$ be an n -qubit quantum channel of degree at most d . Then*

$$\|\widehat{\Phi}\|_{2d/(d+1)} \leq C^d,$$

Proof We just have to show that if Φ is a quantum channel, then $\|\Phi\|_{S_1 \rightarrow S_\infty} \leq 1$. This is true since $\|\Phi\|_{S_1 \rightarrow S_\infty} \leq \|\Phi\|_{S_1 \rightarrow S_1}$ and Φ^\dagger is a completely positive and unital map between C^* -algebras, so we have $\|\Phi\|_{S_1 \rightarrow S_1} = \|\Phi^\dagger\|_{S_\infty \rightarrow S_\infty} = 1$ [30, Proposition 3.2]. \square

Remark 4.1 Theorem 1.3 generalizes the non-commutative BH inequality proved by Volberg and Zhang in [35]. Indeed, given $M = \sum_x \widehat{M}(x)\sigma_x \in M_N$ the main result of [35] is recovered when one applies Theorem 1.3 to $\Phi_M(\cdot) = (\cdot)M$, which satisfies $\widehat{\Phi}_M(x, y) = \delta_{x,0^n} \widehat{M}(y)$ and $\|\Phi_M\|_{S_1 \rightarrow S_\infty} = \|M\|$.

5 Learning low-degree quantum objects

This section is devoted to explaining the applications of the results developed in the previous section to learning theory.

5.1 Why are BH inequalities useful for learning?

We start by recalling a classical problem in learning theory which includes some of the results we present next and serves as motivation for other problems that are explained further below. Consider a function $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ to which we only have access through random samples. Here, a random sample means that we have access to $(x, f(x))$ for an element x chosen uniformly at random from $\{-1, 1\}^n$. Assume that we fix $\varepsilon > 0$ and $\delta > 0$. Then, we want to devise an algorithm such that, by having access to $T(n, \varepsilon, \delta)$ random samples, produces another function $f' : \{-1, 1\}^n \rightarrow \mathbb{R}$ satisfying, with probability at least $1 - \delta$, that $\|f - f'\|_2 < \varepsilon$. In this case, we say that f can be learned within L_2 -error ε by using $T(n, \varepsilon, \delta)$ samples.¹ The goal is, of course, to minimize the number of samples needed to learn the function.

A relevant instance of the problem we just have introduced is learning a bounded function $f : \{-1, 1\}^n \rightarrow [-1, 1]$ of degree at most d . The seminal low-degree algorithm by Linial, Mansour and Nisan solves it with $O_{d,\varepsilon}(n^d)$ samples [25].² Their algorithm is based on learning the relevant part of the Fourier spectrum of the function f which, thanks to Parseval's identity, allows us to learn the function. More precisely, if f' has also degree at most d , we then have that

$$\|f - f'\|_2^2 = \sum_{s \in \{0,1\}^n, |s| \leq d} |\widehat{f}(s) - \widehat{f'}(s)|^2.$$

¹ Despite we don't mention δ explicitly, this parameter is implicit in the problem. Sometimes, one fixes $\delta = 2/3$.

² Here and below, we use $O_{d,\varepsilon}$ to hide factors that depend on d and $1/\varepsilon$ and are independent of n .

Hence, in order to learn f up to error ε , it suffices to learn each of its Fourier coefficients $\widehat{f}(s)$ with $|s| \leq d$ up to error $\varepsilon/\sqrt{n^d}$. Indeed, since there are at most $O(n^d)$ of this coefficients, this immediately implies that $\|f - f'\|_2^2 < \varepsilon^2$.

Now, we explain how to learn the Fourier coefficients $\widehat{f}(s)$ for $|s| \leq d$ with probability $\geq 1 - \delta$ and by just using $T = O(n^d \log(n^d/\delta)/\varepsilon^2)$ random samples $(x_i, f(x_i))_{i \in [T]}$. To this end, let us consider the *empirical Fourier coefficients*, defined as

$$\widehat{f'}(s) = \frac{1}{T} \sum_{i \in [T]} f(x_i) \chi_s(x_i).$$

Note that, for a fixed s , $\widehat{f'}(s)$ can be seen as the average of T independent random variables distributed identically to the random variable $h_s : \{-1, 1\}^n \rightarrow [-1, 1]$ given by $h_s(\cdot) = f(\cdot) \chi_s(\cdot)$. Fixing s , since $\mathbb{E} h_s = \widehat{f}(s)$, we can apply the Hoeffding bound to state that

$$\begin{aligned} \Pr\left(|\widehat{f'}(s) - \widehat{f}(s)| > \frac{\varepsilon}{\sqrt{n^d}}\right) &= \Pr\left(\frac{1}{T} \left| \sum_{i \in [T]} (f(x_i) \chi_s(x_i) - \widehat{f}(s)) \right| > \frac{\varepsilon}{\sqrt{n^d}}\right) \\ &\leq \exp\left(-\frac{T\varepsilon^2}{2n^d}\right). \end{aligned}$$

A union bound can then be applied to upper bound the probability that $|\widehat{f'}(s) - \widehat{f}(s)| \leq \frac{\varepsilon}{\sqrt{n^d}}$ for every $|s| \leq d$ by

$$1 - \exp\left(-\frac{T\varepsilon^2}{2n^d} + d \log n\right).$$

Hence, by choosing $T = 2n^d \log(n^d/\delta)/\varepsilon^2$, we make this upper bound equal to $1 - \delta$ as we wanted.

The algorithm by Linial et al. was the state of the art until recently, when Eskenazis and Ivanisvili showed that a function of degree d can actually be learnt by using only $O_{d,\varepsilon}(\log n)$ random samples [19]. Their key insight was to use a Bohnenblust and Hille inequality for functions defined on the hypercube $f : \{-1, 1\}^n \rightarrow \mathbb{R}$, proved in [18], which can be used to upper bound the contribution of the *small Fourier coefficients*. To illustrate this, we consider the sum of the squares of the Fourier coefficients which are smaller than a certain parameter ε' , which will be fixed later; namely

$$\sum_{s \in \{0,1\}^n, |\widehat{f}(s)| \leq \varepsilon'} |\widehat{f}(s)|^2.$$

To upper bound this quantity, one can use that $2 = 2/(d+1) + 2d/(d+1)$, so

$$\sum_{s \in \{0,1\}^n, |\widehat{f}(s)| \leq \varepsilon'} |\widehat{f}(s)|^2 \leq \varepsilon'^{\frac{2}{d+1}} \sum_{s \in \{0,1\}^n, |\widehat{f}(s)| \leq \varepsilon'} |\widehat{f}(s)|^{\frac{2d}{d+1}}.$$

Now one can use the aforementioned BH inequality, which states that $\|\widehat{f}\|_{2d/(d+1)} \leq C^{\sqrt{d \log d}} \|f\|_\infty$, to obtain

$$\sum_{s \in \{0,1\}^n, |\widehat{f}(s)| \leq \varepsilon'} |\widehat{f}(s)|^{\frac{2d}{d+1}} \leq \varepsilon'^{\frac{2}{d+1}} C^{\sqrt{d \log d}}.$$

Therefore, by setting $\varepsilon' = \varepsilon^{d+1} C^{-(d+1)\sqrt{d \log d}/2}$, it follows that

$$\sum_{s \in \{0,1\}^n, |\widehat{f}(s)| \leq \varepsilon'} |\widehat{f}(s)|^2 \leq \varepsilon^2. \quad (5.1)$$

From Eq. (5.1), Eskenazis and Ivanisvili essentially followed the ideas of Linial et al., but now they just needed to learn every low-degree Fourier coefficient up to error $\varepsilon' = \varepsilon^{d+1} C^{-(d+1)\sqrt{d \log d}/2}$, which is *much bigger* than $\varepsilon/\sqrt{n^d}$ and, in particular, independent of n . To be more precise, they proved that these functions can be learned with L_2 -error ε and confidence $1 - \delta$ by using

$$O\left(\varepsilon^{-2(d+1)} \|\widehat{f}\|_{\frac{2d}{d+1}}^2 d^2 \log\left(\frac{n}{\delta}\right)\right) \quad (5.2)$$

random samples.

5.2 Learning quantum query algorithms

In particular, the result of Eskenazis and Ivanisvili applies to the amplitudes of quantum query algorithms as in Eq. (1.2) which, since the early days of quantum query complexity, are known to be bounded d -tensors $T : (\{-1, 1\}^n)^d \rightarrow [-1, 1]$ [6, 19]. In addition, for d -tensors it is known that the BH inequality holds with a polynomial constant, $\|T\|_{2d/(d+1)} \leq \text{poly}(d) \|T\|_\infty$ [11]. Hence, it follows from Eq. (5.2) that the amplitudes of quantum query algorithms can be learned from

$$O(\varepsilon^{-2(d+1)} \text{poly}(d)^d \log(n/\delta)) \quad (5.3)$$

samples.

A key observation here, proved in [2], is that those d -linear tensors arising from quantum algorithms actually satisfy that $\|T\|_{\text{cb}} \leq 1$ [6]. Hence, Theorem 1.1 implies the following improvement with respect to Eq. (5.3).

Corollary 5.1 *Consider a quantum algorithm that makes d queries as in Eq. (1.2). Then, its amplitudes can be learned with L_2 -accuracy ε and confidence $1 - \delta$ from $O(\varepsilon^{-2(d+1)} d^2 \log(n/\delta))$ uniform random samples.*

Note that this result requires a number of samples that is polynomial on n when ε and δ are constants and $d = \log(n)$, while using (5.3) one would get $O(n^{\log \log n})$ samples as an upper bound.

5.3 Learning low-degree Boolean functions

In this section we propose almost optimal classical and quantum algorithms to learn low-degree Boolean functions. While we have already explained the classical access model (via random samples), we will also need to know what we mean by a quantum access model. The quantum counterpart of these samples are the quantum uniform samples, defined via the $(n + 1)$ -qubit states

$$|f\rangle = \frac{1}{\sqrt{2^n}} \sum_{x \in \{-1, 1\}^n} |x\rangle \otimes |f(x)\rangle \in (\mathbb{C}^2)^n \otimes \mathbb{C}^2 = (\mathbb{C}^2)^{n+1},$$

where $\{|-1\rangle, |1\rangle\}$ is the canonical (or computational) basis of \mathbb{C}^2 and we have denoted $|x\rangle = |x_1\rangle \otimes \cdots \otimes |x_n\rangle \in (\mathbb{C}^2)^n$ for every $x \in \{-1, 1\}^n$. Quantum uniform samples are at least as powerful as classical samples. Indeed, if one measures the first n qubits of $|f\rangle$ in the basis $\{|x\rangle\}_x$, then the last qubit collapses to $|f(x)\rangle$ for a uniformly random x . However, they are actually strictly more powerful, as they allow one to sample from the Fourier distribution $(|\hat{f}(s)|^2)_s$. For a proof of this well-known result, see for instance [3, Lemma 4].

Lemma 5.2 (Fourier sampling) *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be a Boolean function. There is an algorithm that inputs $|f\rangle$, succeeds with probability $1/2$ and, in this case, samples a string $s \in \{0, 1\}^n$ according to the probability distribution $(|\hat{f}(s)|^2)_s$.*

We now state the main result of this section on Boolean functions.

Proposition 5.3 *Let $f : \{-1, 1\}^n \rightarrow \{-1, 1\}$ be a degree- d function. There is a quantum algorithm that learns f exactly with probability $1 - \delta$ using $O(4^d d \log(1/\delta))$ uniform quantum samples. Also, there is a classical algorithm that only uses $O(4^d d \log(n/\delta))$ uniform samples for this task.*

Despite the simplicity of the proof of Proposition 5.3, we include it for completeness and because it seems not to be well-known. See for instance [28, Corollary 34], which proposes a quantum algorithm for the same problem that requires $O(n^d)$ samples, or [20, Corollary 4] that proposes a classical algorithm that requires $O(2^{d^2} \log n)$ samples. Proposition 5.3 highly improves those estimates. We also remark that a classical lower bound of $\Omega(2^d \log n)$ samples was recently proved, making our classical result nearly optimal [20]. Regarding the tightness of our quantum result, since learning functions $f : \{-1, 1\}^d \rightarrow \{-1, 1\}$ of degree d requires $\Omega(2^d)$ uniform quantum samples (which is folklore and follows from example from [5]), our quantum estimate is almost optimal too.

Proof For the classical upper bound we propose the following algorithm. We take $T = 2 \cdot 4^d \log(n^d/\delta)$ uniform samples $(x_i, f(x_i))$ and use them to define the empirical Fourier coefficients as

$$\hat{f}'(s) = \frac{1}{T} \sum_{i \in [T]} f(x_i) \chi_s(x_i),$$

for every $s \subseteq [n]$. Define now the event $\mathcal{E} = \{|\hat{f}(s) - \hat{f}'(s)| < 2^{-d} \forall |s| \leq d\}$. Then, one can argue exactly in the same way as in Sect. 5.1 to conclude that $\Pr[\mathcal{E}] \geq 1 - \delta$.

Once we have computed the coefficients $\hat{f}'(s)$, we round every of them to the closest number $\hat{f}''(s) \in 2^{d-1}\mathbb{Z}$. If \mathcal{E} occurs, by granularity we have that $\hat{f}''(s) = \hat{f}(s)$ for every $|s| \leq d$, so $f = \sum_s \hat{f}''(s)\chi_s$, as desired.

For the quantum upper bound we begin by sampling $N = 4^d \log(4^d/\delta)$ times from $(\hat{f}(s)^2)_{s \in \{0,1\}^n}$. This can be done, with probability $\geq 1 - \delta$, by using $T_1 = O(4^d \log(4^d/\delta))$ quantum uniform samples, thanks to Lemma 5.2 and a Hoeffding bound. Now, given s such that $\hat{f}(s) \neq 0$, the probability that a sample s' according to the distribution $(\hat{f}(s)^2)_{s \in \{0,1\}^n}$ satisfies $s' \neq s$ is given by $1 - \hat{f}(s)^2 \leq 1 - 4^{1-d}$, where we have used that $\hat{f}(s)^2 \geq 4^{1-d}$ by Lemma 4.1. Hence, if s_1, \dots, s_N are the N samples, then the probability that we have $s_i \neq s$ for every $i = 1, \dots, N$ is upper bounded by

$$(1 - 4^{1-d})^N \leq \frac{\delta}{4^d}.$$

Thus, taking a union bound over the at most 4^{d-1} non-zero Fourier coefficients (due to Lemma 4.1 and $\sum_s |\hat{f}(s)|^2 = 1$), it follows that, with probability $1 - \delta$, we will have sampled every non-zero Fourier coefficient.

In the second part of the algorithm we use $T_2 = O(4^d \log(4^d/\delta))$ quantum uniform samples and measure them in the computational basis, which generates classical uniform samples. From here, we can argue as in the classical upper bound and learn f exactly. The quantum advantage comes from Fourier sampling, that allows us to detect the non-zero Fourier coefficients, and apply the union bound only over those, that are at most 4^{d-1} . \square

5.4 Learning low-degree quantum channels

First of all, we define the access model we use. Given a channel Φ , a learning algorithm is allowed to make queries to Φ as follows: it can choose a state ρ , feed ρ to the channel to obtain $\Phi(\rho)$ and measure $\Phi(\rho)$ in any basis.

The goal here, as in the previous sections, is to produce a classical description of a map $\tilde{\Phi}$ that is close to Φ in the ℓ_2 -distance defined by the usual inner product for maps from M_N to M_N , i.e., $\langle \Phi, \tilde{\Phi} \rangle = \text{Tr}[J(\Phi)J(\tilde{\Phi})]/4^n$, where $J(\Phi)$ is the Choi–Jamiołkowski (CJ) representation of Φ .

For a reader not familiar with quantum computing, we remark that the proof of the main result of this section does not require prior knowledge of quantum computing, if one uses Lemmas 5.4 and 5.6 in a black-box manner. The reader can find in [27] an excellent reference to learn about quantum computing.

An important fact for our learning algorithm is that $\hat{\Phi} = (\hat{\Phi}(x, y))_{x, y}$ is a state that can be prepared with 1 query to Φ (see [13, Lemma 8]). This is the content of the following statement.

Lemma 5.4 *If Φ is a quantum channel, then $\widehat{\Phi}$ is a state unitarily equivalent to $v(\Phi)$. In particular, one query to Φ suffices to sample once from $(\widehat{\Phi}(x, x))_x$, which is a probability distribution.*

We will also make use of the following lemma, proved in [23, Proposition 7].

Lemma 5.5 *Let ρ, ρ' be two states. Then, one can estimate $\text{Tr}[\rho\rho']$ up to error ε with probability $1 - \delta$, by using $O((1/\varepsilon)^2 \log(1/\delta))$ copies of ρ and ρ' .*

Before proving the main theorem of the section, we show that for a given $x, y \in \{0, 1, 2, 3\}^n$, the corresponding Pauli coefficient $\widehat{\Phi}(x, y)$ can be efficiently learned.

Lemma 5.6 (Pauli coefficient estimation for channels) *Let $\Phi : M_N \rightarrow M_N$ be a quantum channel and let $x, y \in \{0, 1, 2, 3\}^n$. Then, $\widehat{\Phi}(x, y)$ can be estimated with error ε and probability $1 - \delta$ using $O((1/\varepsilon)^2 \log(1/\delta))$ queries to Φ .*

Proof If $x = y$, we just have to prepare $\widehat{\Phi}$ and apply Lemma 5.5 to $\widehat{\Phi}$ and the state $\rho = |x\rangle\langle x|$. If $x \neq y$, one first learns $\widehat{\Phi}(x, x)$ and $\widehat{\Phi}(y, y)$ with error ε as before. On the one hand, one can learn $\widehat{\Phi}(x, x) + \widehat{\Phi}(y, y) + 2\Re\widehat{\Phi}(x, y)$, with error ε by applying Lemma 5.5 to $\widehat{\Phi}$ and $|\xi\rangle\langle\xi|$, where $|\xi\rangle = 1/\sqrt{2}(|x\rangle + |y\rangle)$. Hence, one learns $\Re\widehat{\Phi}(x, y)$ with error $3\varepsilon/2$. On the other hand, one can learn $\widehat{\Phi}(x, x) + \widehat{\Phi}(y, y) + 2\Im\widehat{\Phi}(x, y)$, with error ε by applying Lemma 5.5 to $\widehat{\Phi}$ and $|\eta\rangle\langle\eta|$, where $|\eta\rangle = 1/\sqrt{2}(|x\rangle + i|y\rangle)$, and one can then learn $\Im\widehat{\Phi}(x, y)$ with error $3\varepsilon/2$. \square

We will also need the following well-known result about distribution learning theory. See [14, Theorem 9] for a proof.

Lemma 5.7 *Let $p = \{p(x)\}_x$ be a probability distribution over some set \mathcal{X} . Let $p' = (p'(x))_x$ be the empirical probability distribution obtained after sampling T times from p . Then, for $T = O((1/\varepsilon)^2 \log(1/\delta))$ with probability $\geq 1 - \delta$ we have that $|p(x) - p'(x)| \leq \varepsilon$ for every $x \in \mathcal{X}$.*

Now, we are ready to prove Theorem 1.4, which we restate for the convenience of the reader.

Theorem 1.4 *Let Φ be a n -qubit degree- d quantum channel. Then it can be learned in L_2 -accuracy ε and confidence $\geq 1 - \delta$ by making $\exp(\widetilde{O}(d^2 + d \log(1/\varepsilon))) \cdot \log(1/\delta)$ queries to Φ . Here, we use the notation \widetilde{O} to hide logarithmic factors in $d, 1/\varepsilon$, and $1/\delta$.*

Proof The algorithm consists of 2 steps. In the first one we detect the relevant Pauli coefficients, while in the second step we learn the few Pauli coefficients detected as relevant.

Step 1. Detect the big Pauli coefficients. Let $c > 0$ be a parameter to be determined later. We invoke Lemma 5.4 to sample T_1 times from $(\widehat{\Phi}(x, x))_x$ by making T_1 queries to Φ . Let $(\widehat{\Phi}'(x, x))_x$ be the empirical distribution obtained from these samples. We store the big Pauli coefficients in the set $\mathcal{X}_c = \{x : \widehat{\Phi}'(x, x) \geq c\}$. Note that, since $\sum_{x \in \mathcal{X}_c} \widehat{\Phi}'(x, x) \leq 1$, we know that

$$|\mathcal{X}_c| \leq \frac{1}{c}. \quad (5.4)$$

Step 2. Learn the big Pauli coefficients. We invoke Lemma 5.6 to state that, by querying Φ just

$$T_2 = O((1/c)^4(1/\varepsilon)^2 \log((1/c)^2(1/\delta)))$$

times, we can find approximations $\widehat{\Phi}''(x, y)$ of $\Phi(x, y)$ for the at most $(1/c)^2$ pairs $(x, y) \in \mathcal{X}_c$, such that

$$\sup_{(x,y) \in \mathcal{X}_c \times \mathcal{X}_c} |\widehat{\Phi}(x, y) - \widehat{\Phi}''(x, y)| \leq c\varepsilon. \quad (5.5)$$

happens with probability $\geq 1 - \delta$.

Output. We output $\Phi''(\cdot) = \sum_{x,y \in \mathcal{X}_c} \widehat{\Phi}''(x, y) \sigma_x(\cdot) \sigma_y$ as our approximation for Φ .

Correctness analysis. We consider the event $\mathcal{E} = \{|\widehat{\Phi}(x, x) - \widehat{\Phi}'(x, x)| \leq c \ \forall x \in \{0, 1, 2, 3\}^n\}$. By Lemma 5.7, taking $T_1 = O((1/c)^2 \log(1/\delta))$ ensures that

$$\Pr[\mathcal{E}] \geq 1 - \delta.$$

Assuming the event \mathcal{E} holds, we have that

$$x \notin \mathcal{X}_c \implies |\widehat{\Phi}(x, x)| \leq |\widehat{\Phi}'(x, x)| + ||\widehat{\Phi}(x, x) - \widehat{\Phi}'(x, x)|| \leq 2c. \quad (5.6)$$

In particular, it follows that

$$x \notin \mathcal{X}_c \implies |\widehat{\Phi}(x, y)| \leq \sqrt{|\widehat{\Phi}(x, x)| |\widehat{\Phi}(y, y)|} \leq \sqrt{2c} \ \forall y \in \{0, 1, 2, 3\}^n, \quad (5.7)$$

where in the first inequality we have used that $\widehat{\Phi}$ is positive semidefinite and in the second inequality we have used Eq. (5.6) and that $\widehat{\Phi}(y, y) \leq 1$.

Assuming that both parts of the algorithm succeed, we have that Φ'' is close to Φ . Indeed,

$$\begin{aligned} \|\Phi - \Phi''\|_2^2 &= \sum_{x,y \in \mathcal{X}_c} |\widehat{\Phi}(x, y) - \widehat{\Phi}''(x, y)|^2 + \sum_{x \vee y \notin \mathcal{X}_c} |\widehat{\Phi}(x, y)|^2 \\ &\leq \varepsilon^2 + \sum_{x \vee y \notin \mathcal{X}_c} |\widehat{\Phi}(x, y)|^{\frac{2}{d+1}} |\widehat{\Phi}(x, y)|^{\frac{2d}{d+1}} \\ &\leq \varepsilon^2 + (2c)^{\frac{1}{d+1}} \|\widehat{\Phi}\|_{\frac{2d}{d+1}}^{\frac{2d}{d+1}} \\ &\leq \varepsilon^2 + c^{\frac{1}{d+1}} C^d. \end{aligned}$$

Here, in the equality we have used Parseval's identity, in the first inequality we used Eq. (5.4), Eq. (5.5) and that $2 = 1/(d + 1/2) + 2d/(d + 1/2)$; in the second inequality

we have used Eq. (5.7) and in the third inequality we used the Bohnenblust–Hille inequality for channels (Corollary 4.5). Hence, by choosing

$$c = \varepsilon^{2d+2} C^{-d(d+1)}$$

we obtain the desired result.

Complexity analysis. Note that $T_2 > T_1$, so the complexity T_2 dominates the one of the first part of the algorithm. Hence, the total number of queries made is

$$O\left(C^{4d(d+1)}(1/\varepsilon)^{8d+10} \log(C^{2d(d+1)}(1/\varepsilon)^{4d+4}(1/\delta))\right).$$

□

Declarations

Conflict of interest On behalf of all authors, the corresponding author states that there is no conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Aaronson, S., Ambainis, A.: Forrelation: A problem that optimally separates quantum from classical computing. In *Proceedings of the Forty-Seventh Annual ACM Symposium on Theory of Computing*, STOC '15, New York, NY (2015). Association for Computing Machinery
2. Arunachalam, S., Briët, J., Palazuelos, C.: Quantum query algorithms are completely bounded forms. *SIAM J. Comput.* **48**(3), 903–925 (2019). Preliminary version in *ITCS'18*
3. Arunachalam, S., Chakraborty, S., Lee, T., Paraashar, M., De Wolf, R.: Two new results about quantum exact learning. *Quantum* **5**, 587 (2021)
4. Arunachalam, S., Dutt, A., Escudero, G., Francisco, P. C.: Learning low-degree quantum objects. [arXiv:2405.10933](https://arxiv.org/abs/2405.10933) (2024)
5. Arunachalam, S., de Wolf, R.: Optimal quantum sample complexity of learning algorithms. *J. Mach. Learn. Res.* **19**(71), 1–36 (2018)
6. Beals, R., Buhrman, H., Cleve, R., Mosca, M., De Wolf, R.: Quantum lower bounds by polynomials. *J. ACM (JACM)* **48**(4), 778–797 (2001)
7. Briët, J., Escudero Gutiérrez, F., Gribling, S.: Grothendieck inequalities characterize converses to the polynomial method. [arXiv:2212.08559](https://arxiv.org/abs/2212.08559) (2022)
8. Bohnenblust, H.F., Hille, E.: On the absolute convergence of Dirichlet series. *Ann. Math.* 600–622 (1931)
9. Blei, R.C.: Fractional cartesian products of sets. In *Annales de l'institut Fourier* **29**, 79–105 (1979)
10. Blei, R.: *Analysis in Integer and Fractional Dimensions*, vol. 71. Cambridge University Press, Cambridge (2001)
11. Bayart, F., Pellegrino, D., Seoane-Sepúlveda, J.B.: The Bohr radius of the n -dimensional polydisk is equivalent to $(\log n)/n$. *Adv. Math.* **264**, 726–746 (2014)

12. Bansal, N., Sinha, M.: k-forrelation optimally separates quantum and classical query complexity. In: Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing, STOC 2021, New York, NY (2021). Association for Computing Machinery
13. Bao, Z., Yao, P.: On Testing and Learning Quantum Junta Channels. In: Gergely, N., Lorenzo, R. (eds.), Proceedings of Thirty Sixth Conference on Learning Theory, volume 195 of *Proceedings of Machine Learning Research*, pp. 1064–1094 (2023)
14. Canonne, C.L.: A short note on learning discrete distributions. [arXiv:2002.11457](https://arxiv.org/abs/2002.11457) (2020)
15. Defant, A., Frerick, L., Ortega-Cerdà, J., Ounaïes, M., Seip, K.: The Bohnenblust–Hille inequality for homogeneous polynomials is hypercontractive. *Ann. Math.*, pp. 485–497 (2011)
16. Defant, A., García, D., Maestre, M., Peris, P.: Dirichlet series and holomorphic functions in high dimensions. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, pp 07 (2019)
17. Diniz, D., Muñoz-Fernández, G., Pellegrino, D., Seoane-Sepúlveda, J.: Lower bounds for the constants in the Bohnenblust–Hille inequality: the case of real scalars. *Proc. Am. Math. Soc.* **142**(2), 575–580 (2014)
18. Defant, A., Mastyło, M., Pérez, A.: On the Fourier spectrum of functions on boolean cubes. *Math. Ann.* **374**, 653–680 (2019)
19. Eskenazis, A., Iyannis, P.: Learning low-degree functions from a logarithmic number of random queries. In: Proceedings of the 54th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2022, New York, NY, USA (2022). Association for Computing Machinery
20. Eskenazis, A., Iyannis, P., Streck, L.: Low-degree learning and the metric entropy of polynomials. [arXiv:2203.09659](https://arxiv.org/abs/2203.09659) (2022)
21. Gutoski, G., Johnston, N.: Process tomography for unitary quantum channels. *J. Math. Phys.* **55**(3) (2014)
22. Huang, H.-Y., Chen, S., Preskill, J.: Learning to predict arbitrary quantum processes. *PRX Quant.* **4**, 040337 (2023)
23. Kobayashi, H., Matsumoto, K., Yamakami, T.: Quantum merlin-arthur proof systems: are multiple Merlins more helpful to Arthur? In: Algorithms and Computation: 14th International Symposium. ISAAC 2003, pp. 189–198. Springer, New York (2003)
24. Littlewood, J.E.: On bounded bilinear forms in an infinite number of variables. *Q. J. Math.*, pp. 164–174 (1930)
25. Linial, N., Mansour, Y., Nisan, N.: Constant depth circuits, Fourier transform, and learnability. *J. ACM (JACM)* **40**(3), 607–620 (1993)
26. Montanaro, A., Osborne, T.J.: Quantum Boolean functions. [arXiv:0810.2435](https://arxiv.org/abs/0810.2435) (2008)
27. Nielsen, M.A., Chuang, I.L.: Quantum Computation and Quantum Information, vol. 10. Anniversary Cambridge University Press, Cambridge, UK (2010)
28. Nadimpalli, S., Parham, N., Vasconcelos, F., Yuen, H.: On the Pauli spectrum of QAC0. [arXiv:2311.09631](https://arxiv.org/abs/2311.09631) (2023)
29. O’Donnell, R.: Analysis of Boolean Functions. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, p. 07 (2014)
30. Paulsen, V.: Completely Bounded Maps and Operator Algebras. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, pp. 02 (2003)
31. Pellegrino, D., Teixeira, E.V.: Towards sharp Bohnenblust–Hille constants. *Commun. Contemp. Math.* **20**(03), 1750029 (2018)
32. Rouzé, C., Wirth, M., Zhang, H.: Quantum talagrand, kkl and friedguts theorems and the learnability of quantum Boolean functions. *Commun. Math. Phys.* **405**(4), 95 (2024)
33. Slot, J., Volberg, A., Zhang, H.: Bohnenblust–Hille inequality for cyclic groups. [arXiv:2305.10560](https://arxiv.org/abs/2305.10560) (2023)
34. Slot, J., Volberg, A., Zhang, H.: Noncommutative bohnenblust–hille inequality in the Heisenberg–Weyl and gell-mann bases with applications to fast learning. [arXiv:2301.01438](https://arxiv.org/abs/2301.01438) (2023)
35. Volberg, A., Zhang, H.: Noncommutative Bohnenblust–Hille inequalities. *Math. Ann.*, pp. 1–20 (2023)

Authors and Affiliations

Srinivasan Arunachalam^{1,2} · Arkopal Dutt¹ · Francisco Escudero Gutiérrez³  · Carlos Palazuelos⁴

✉ Francisco Escudero Gutiérrez
feg@cw.nl

Srinivasan Arunachalam
Srinivasan.Arunachalam@ibm.com

Arkopal Dutt
arkopal@ibm.com

Carlos Palazuelos
cpalazue@ucm.es

¹ IBM Quantum, New York, NY, USA

² University of Illinois at Urbana-Champaign, 1409 W. Green St., Urbana, IL 61891, USA

³ Qusoft and CWI, Amsterdam, The Netherlands

⁴ Instituto de Ciencias Matemáticas (ICMAT), Facultad de Ciencias Matemáticas, Universidad Complutense de Madrid, Madrid 28040, Spain