



Approximating the Volume of a Truncated Relaxation of the Independence Polytope

Ferenc Bencs¹ · Guus Regts²

Received: 2 May 2024 / Revised: 21 January 2026 / Accepted: 22 January 2026

© The Author(s) 2026

Abstract

Answering a question of Gamarnik and Smedira [15], we give a polynomial time algorithm that approximately computes the volume of a truncation of a relaxation of the independent set polytope, improving on their quasi-polynomial time algorithm. Our algorithm is obtained by viewing the volume as an evaluation of a graph polynomial and we approximate this evaluation using Barvinok’s interpolation method.

Keywords Deterministic volume computation · Interpolation method · Independence polytope

Mathematics Subject Classification Primary: 05C31 · 52B55; Secondary: 05C30 · 68W25 · 05C69

1 Introduction

Approximately computing the volume of a polytope and more generally a convex body is one of the success stories of randomized algorithms. Indeed, Dyer, Frieze and Kannan [12] showed that with access to a membership oracle there is a randomized polynomial time algorithm to approximately compute the volume of a convex body, whereas no such deterministic algorithm can exist by a result of Bárány and Füredi [3].

Editor in Charge: Kenneth Clarkson

Funded by the Netherlands Organisation of Scientific Research (NWO): VI.Veni.222.303

Funded by the Netherlands Organisation of Scientific Research (NWO): VI.Vidi.193.068.

Ferenc Bencs
ferenc.bencs@gmail.com

Guus Regts
g.regts@uva.nl

¹ Centrum Wiskunde & Informatica, P.O. Box 94079, 1090 GB Amsterdam, The Netherlands

² Korteweg de Vries Institute for Mathematics, University of Amsterdam, P.O. Box 94248, 1090 GE Amsterdam, The Netherlands

Partly motivated by the question whether randomized algorithms are superior over deterministic algorithms in the context of approximately computing volumes of polytopes, Gamarnik and Smedira [15] considered the question of approximately computing the volume of a standard relaxation of the independent set polytope of a graph $G = (V, E)$ which is defined as

$$P_G := \{x \in [0, 1]^V \mid x_u + x_v \leq 1 \text{ for all } uv \in E\}.$$

In fact for a bipartite graph G , P_G is equal to the independent set polytope of G [28]*Theorem 19.7. Since P_G is given by inequality constraints, testing membership in P_G can be done efficiently and hence there is an efficient randomized algorithm for approximately computing its volume [12]. As far as we know, no efficient deterministic algorithm is known for this task for bounded degree graphs. For related result when the girth of G tends to infinity see [16].

The choice of polytope P_G may seem a bit arbitrary, but not much appears to be known about deterministic approximation of the volume of polytopes with high precision (see [4, 6] for deterministic approximations of the volume of an n -dimensional bounded polyhedron defined by $m > n$ linear inequalities within a multiplicative factor of $2^{O(m-n)}$) and as such it is as good a choice as any other. Moreover, since integration can be seen as an idealized form of counting [15], the polytope P_G actually presents itself as a rather natural candidate to test our ability to design efficient deterministic approximation algorithms for computing volumes. Indeed, in the ‘counting world’ it is known that for the problem of approximately computing the number of independent sets in a bounded degree graph (and more generally approximately computing evaluations of its independence polynomial) randomized algorithms do not outperform deterministic algorithms in the sense that for graphs of maximum degree at most 5 both efficient deterministic and randomized (Markov chain based) polynomial time algorithms exist for this task [1, 31], while if the maximum degree is at least 6 approximating the number of independent sets is NP-hard [14, 29]¹. The Markov chain based randomized algorithms do tend to have faster running times compared to their deterministic counterparts [1, 10, 27, 31]. Incidentally the randomized algorithm for approximating volumes of Dyer, Frieze and Kannan [12] is also based on Markov chains and as such it is natural to try to use techniques from approximate counting to design deterministic approximation algorithms in the context of volume approximation.

Gamarnik and Smedira [15] in fact considered a truncation of P_G , defined as follows. For $\delta \in [0, 1/2]$ and a graph $G = (V, E)$ let

$$P_{G,\delta} := \{(x_v)_{v \in V} \in [0, 1/2 + \delta]^V \mid x_u + x_v \leq 1 \text{ for all } uv \in E\} \quad (1.1)$$

and note that $P_{G,1/2}$ is equal to P_G . Using the successful correlation decay method from approximate counting [2, 31], Gamarnik and Smedira [15] managed to design an algorithm that approximately compute the volume of a $P_{G,\delta}$ for $\delta = O(1/\Delta(G)^2)$ in time $n^{O(\log n)}$, where n denotes the number of vertices of G .

¹ This transition in the computational complexity of approximate counting in fact extends to evaluating the independence polynomial on bounded degree graphs.

Our main result improves on their result, both in terms running time as well as in terms of the bound on δ .

Theorem 1.1 *There exists a constant $C > 0$ such that for each $\Delta > 0$ and $\delta \in [0, C/\Delta]$, there exists an algorithm that on input of an n -vertex graph of maximum degree at most Δ and $\varepsilon \in (0, 1)$ computes a number ξ such that*

$$(1 - \varepsilon)\xi \leq \text{vol}(P_{G,\delta}) \leq (1 + \varepsilon)\xi$$

in time polynomial in n/ε .

While we can provide an explicit bound on the constant C in the theorem above, we choose not to since ideally we would like to be able to take δ to be $1/2$ and this appears to be out of reach for now. Indeed, our approach to proving Theorem 1.1 consists of viewing the volume of $P_{G,\delta}$ as the evaluation of a certain graph polynomial/partition function and by applying Barvinok’s interpolation method [5, 30] to it. For this we need to show the polynomial is zero-free on some open region of the complex plane and we manage to do this for $\delta = O(1/\Delta)$. It appears to be unlikely that this can be pushed beyond $O(1/\Delta)$; we will say more about this in Section 5.

The remainder of this paper is organized as follows. In the next section we will indicate how to interpret the volume of $P_{G,\delta}$ as the evaluation of a polynomial that is naturally associated to G and indicate how this leads to a proof of Theorem 1.1 provided two conditions are met. In Sections 3 and 4 we verify these conditions. Finally, in Section 5 we conclude with some discussion and possible extensions.

2 The Volume as an Evaluation of a Graph Polynomial

In this section we will show how to view $\text{Vol}(P_{G,\delta})$ as the evaluation of a certain generating function and explain how this allows us to employ Barvinok’s interpolation method to prove Theorem 1.1. To this end we introduce some terminology.

For a total order on the edges of a graph $H = (V, E)$ and a spanning tree (V, T) , $T \subseteq E$, we call $e \in E \setminus T$ *broken* if $T \cup \{e\}$ contains a cycle in which e is the largest edge. We denote by E_T the collection of broken edges. A well known combinatorial lemma, going back to Penrose [26], states the following:

Lemma 2.1 *Let $H = (V, E)$ be a graph with a fixed total ordering of the edges. Then the collection of connected subgraphs of H can be partitioned into intervals of the form $\{F \mid T \subseteq F \subseteq T \cup E_T\}$, where T is a tree.*

Remark 2.2 A simple proof of Lemma 2.1 can be obtained by associating a connected subgraph (V, F) of H to its minimum weight spanning tree.

We start by rewriting the volume of $P_{G,\delta}$. We have

$$\text{Vol}(P_{G,\delta}) = \int \prod_{uv \in E} \mathbf{1}_{x_u+x_v \leq 1} d\mu, \tag{2.1}$$

where μ denotes the Lebesgue measure, where we integrate over $[0, 1/2 + \delta]^V$ and where $\mathbf{1}_{x_u+x_v \leq 1}$ denotes the function that on input of $\mathbf{x} \in \mathbb{R}^V$ outputs 1 if $x_u + x_v \leq 1$ and zero otherwise. In what follows it will be convenient to denote $I = I_\delta := [0, 1/2 + \delta]$. Note that we can write

$$\mathbf{1}_{x_u+x_v \leq 1} = 1 - \mathbf{1}_{x_u+x_v > 1}$$

and hence

$$\begin{aligned} \prod_{uv \in E} \mathbf{1}_{x_u+x_v \leq 1} &= \sum_{A \subseteq E} (-1)^{|A|} \prod_{uv \in A} \mathbf{1}_{x_u+x_v > 1} \\ &= \sum_{A \subseteq E} \prod_{C \in \mathcal{C}(A)} (-1)^{|C|} \prod_{uv \in C} \mathbf{1}_{x_u+x_v > 1}, \end{aligned} \tag{2.2}$$

where we denote by $\mathcal{C}(A)$ the collection of edge sets of the components of the graph (V, A) .

We can expand this further by first identifying the support of the connected components and then selecting edge sets. To this end denote for $k \in \mathbb{Z}_{\geq 0}$, by $\pi_k(V)$ the collection of unordered partitions of V into k nonempty sets. For $S \subseteq V$ we denote by $E(S)$ the set of edges of G with both endpoints in S . Then (2.2) can be written as

$$\sum_{k \geq 1} \sum_{P \in \pi_k(V)} \prod_{\substack{S \in P \\ |S| \geq 2}} \sum_{\substack{A \subseteq E(S) \\ (S,A) \text{ connected}}} (-1)^{|A|} \prod_{uv \in A} \mathbf{1}_{x_u+x_v > 1}. \tag{2.3}$$

Plugging this into the integral we obtain that (2.1) is equal to

$$\begin{aligned} &\int_{I^V} \sum_{k \geq 1} \sum_{P \in \pi_k(V)} \prod_{\substack{S \in P \\ |S| \geq 2}} \sum_{\substack{A \subseteq E(S) \\ (S,A) \text{ connected}}} (-1)^{|A|} \prod_{uv \in A} \mathbf{1}_{x_u+x_v > 1} d\mu \\ &= \sum_{k \geq 1} \sum_{P \in \pi_k(V)} \prod_{\substack{S \in P \\ |S| \geq 2}} \int_{I^S} \sum_{\substack{A \subseteq E(S) \\ (S,A) \text{ connected}}} (-1)^{|A|} \prod_{uv \in A} \mathbf{1}_{x_u+x_v > 1} d\mu \cdot \prod_{\substack{P \in \pi_k(V) \\ |P|=1}} \int_I 1 d\mu. \end{aligned} \tag{2.4}$$

Now focusing on the contribution of a single set $S \in P$ of size at least 2 and fixing some arbitrary ordering of edges of G , denoting by $E(S)_T \subseteq E(S)$ the set of broken edges associated to T , we have by Lemma 2.1,

$$\begin{aligned} &\sum_{\substack{A \subseteq E(S) \\ (S,A) \text{ connected}}} (-1)^{|A|} \prod_{uv \in A} \mathbf{1}_{x_u+x_v > 1} \\ &= \sum_{T \subseteq E(S)} \sum_{\substack{A \subseteq E(S)_T \\ \text{spanning tree}}} (-1)^{|T \cup A|} \prod_{uv \in T \cup A} \mathbf{1}_{x_u+x_v > 1} \end{aligned}$$

$$= \sum_{\substack{T \subseteq E(S) \\ \text{spanning tree}}} (-1)^{|T|} \prod_{uv \in T} \mathbf{1}_{x_u+x_v > 1} \cdot \prod_{st \in E(S)_T} \mathbf{1}_{x_s+x_t \leq 1}, \tag{2.5}$$

where for the final equality we use (2.2).

Define for an induced subgraph $H = (S, E(S))$ of G ,

$$w(H) := \frac{1}{(1/2 + \delta)^{|S|}} \sum_{\substack{T \subseteq E(S) \\ \text{spanning tree}}} (-1)^{|T|} \int_{I^S} \prod_{uv \in T} \mathbf{1}_{x_u+x_v > 1} \cdot \prod_{st \in E(S)_T} \mathbf{1}_{x_s+x_t \leq 1} d\mu.$$

We note that since the first line of (2.5) does not depend on the ordering of E this quantity really only depends on the graph H . We moreover note that in case H is not connected we have $w(H) = 0$ (since in that case H does not have any spanning trees) and if H consists of a single vertex we have $w(H) = 1$.

Let us next define the univariate polynomial $p_{G,\delta}(x)$, for an n -vertex graph G , by

$$p_{G,\delta}(x) := \sum_{P \in \pi(V)} \prod_{S \in P} w(G[S])x^{|S|-1}, \tag{2.6}$$

where we denote by $\pi(V)$ the collection of all unordered partitions of the set V .

Using that $\int_I 1 d\mu = 1/2 + \delta$, we then arrive at the following conclusion.

Proposition 2.3 *With notation as above we have*

$$(1/2 + \delta)^{-|V|} \text{Vol}(P_{G,\delta}) = p_{G,\delta}(1). \tag{2.7}$$

The upshot of this proposition is that approximating the volume of $P_{G,\delta}$ is equivalent to approximating the polynomial $p_{G,\delta}(x)$ evaluated at 1. We will do this using Barvinok’s interpolation method [5] combined with the refinement due to Patel and the second author [30]. For this it is convenient to express the coefficients of $p_{G,\delta}$ as induced subgraph counts. More precisely, for a graph H , $\text{ind}(H, G)$ denotes the number of subsets S of $V(G)$ such that $G[S]$ is isomorphic to H (i.e. for which there exists a map $\phi : V(H) \rightarrow S$ that for all $u, v \in V(H)$: $uv \in E(H)$ if and only if $\phi(u)\phi(v) \in E(G)$). We note that for $k \in \mathbb{Z}_{\geq 0}$ the coefficient of x^k in P_G is equal to

$$\sum_{\substack{H \in \mathcal{G}^* \\ |V(H)|=k+k(H)}} \text{ind}(H, G) \prod_{C \text{ comp. of } H} w(C), \tag{2.8}$$

where $k(H)$ denotes the number of components of H and where \mathcal{G}^* denotes the family of graphs in which each component has at least two vertices. This implies that we can write $p_{G,\delta}(x) = \sum_{k \geq 0} e_k x^k$, where e_k is of the form $e_k = e_k(G) = \sum_H \lambda_{H,k} \text{ind}(H, G)$ for certain numbers $\lambda_{H,k}$ that only depend on H and k and not on G . We refer to $\lambda_{H,k}$ as the coefficient of $\text{ind}(H, \cdot)$ in $p_{G,\delta}$. We note that $\lambda_{H,k} = 0$, in case $k > |V(H)|/2$ since $k(H) \leq |V(H)|/2$ for any $H \in \mathcal{G}^*$.

To be able to apply the interpolation method we then need the following ingredients. Here \mathcal{G}_Δ denotes the family of all graphs of maximum degree at most Δ .

- A region U in the complex plane that contains 0 and 1 such that for all $z \in U$ and all $G \in \mathcal{G}_\Delta$, $p_{G,\delta}(z) \neq 0$, and,
- an algorithm that on input of an m -vertex graph $H = (V, E) \in \mathcal{G}_\Delta$ computes the coefficient $\lambda_{H,k}$ of $\text{ind}(H, \cdot)$ in $p_{G,\delta}$ in time $C^{O(m)}$ for a constant $C > 0$.

In the next section we will prove the following result, taking care of the first ingredient.

Theorem 2.4 *There exist constants $C \in (0, 1)$ and $K > 1$ such that for any $\Delta > 0$ if $\delta < C/\Delta$ and $|x| < K$, then $p_{G,\delta}(x) \neq 0$ for all graphs G of maximum degree at most Δ .*

For experts in the use of statistical physics methods such as the Kotecký-Preiss conditions [22], the Dobrushin conditions [11] or Gruber-Kunz conditions [8, 19], this should not be a difficult exercise. For completeness we will however provide a detailed and self-contained proof below inspired by [21].

In the subsequent section we will prove the following result, addressing the second ingredient.

Theorem 2.5 *Let $\Delta > 0$. There exists an algorithm that on input of a m -vertex graph $G = (V, E) \in \mathcal{G}_\Delta$ computes the coefficient $\lambda_{H,k}$ of $\text{ind}(H, \cdot)$ in $p_{G,\delta}$ in time $\Delta^{O(k)}$, where the implicit constants do not depend on Δ .*

Combining these two ingredients with Theorem 3.2 from [30] we immediately obtain the desired deterministic algorithm to approximate the volume of the polytope $P_{G,\delta}$, thereby proving Theorem 1.1.

3 A Zero-Free Disk for $p_{G,\delta}$

In this section we prove the following result implying Theorem 2.4. Here e is the base of the natural logarithm.

Theorem 3.1 *Let $\Delta > 1$. Suppose $K > 1$ and $\delta \in (0, 1)$ satisfy*

$$4\delta K \leq \frac{\log(a)(1 - 1/\Delta)}{a\Delta} \tag{3.1}$$

for some $a \in (1, e)$. Then for any graph G of maximum degree at most Δ and $x \in \mathbb{C}$ such that $|x| \leq K$, $p_{G,\delta}(x) \neq 0$.

First of all it will be convenient to reformulate $p_{G,\delta}(x)$ as a certain forest generating function. To this end we introduce for a graph $G = (V, E)$ and a tree $T \subseteq E$ (not necessarily a spanning tree) its weight

$$w_T := (1/2 + \delta)^{-|V(T)|} \int_{I^{V(T)}} (-1)^{|T|} \prod_{uv \in T} \mathbf{1}_{x_u + x_v > 1} \cdot \prod_{st \in E_T} \mathbf{1}_{x_s + x_t \leq 1} d\mu, \tag{3.2}$$

where we recall from the previous section that E_T only depends on T and the graph induced by $V(T)$.

Proposition 3.2 For any graph $G = (V, E)$ and $\delta \neq -1/2$ we have

$$p_{G,\delta}(x) = \sum_{\substack{F \subseteq E(G) \\ F \text{ forest}}} x^{|F|} \prod_{T \text{ comp. of } F} w_T, \tag{3.3}$$

where we write "comp." as a shorthand for connected component.

Proof Expanding the right-hand side of (3.3) by grouping forests according to the partition of V they induce, we see that

$$\sum_{\substack{F \subseteq E(G) \\ F \text{ forest}}} x^{|F|} \prod_{T \text{ comp. of } F} w_T = \sum_{P \in \pi(V)} \prod_{S \in P} \sum_{\substack{T \subseteq E(S) \\ \text{spanning tree}}} w_T x^{|T|}.$$

Here we implicitly use that $w_\emptyset = 1$. We now recognize

$$\sum_{\substack{T \subseteq E(S) \\ \text{spanning tree}}} w_T x^{|T|} = w(G[S])$$

and hence the desired equality follows from the definition of $P_{G,\delta}(x)$. □

Now that we have the desired reformulation of $p_{G,\delta}$ as a forest generating function we require some preliminary results before we can prove Theorem 3.1. Let us denote for a graph G and a vertex $v \in V(G)$ by $T_{G,v}(x)$ the rooted tree generating function of (G, v) , defined by

$$T_{G,v}(x) = \sum_{\substack{T \subseteq E(G) \\ v \in V(T)}} x^{|T|},$$

where the sum is over trees T . We require the following simple lemma, found as Lemma 4.6 in [25].

Lemma 3.3 Let $\Delta > 0$ and let G be a graph of maximum degree at most Δ . Then, for any $a > 1$,

$$T_{G,v} \left(\frac{\log a}{a^\Delta} \right) \leq a.$$

The next lemma says we can bound the weights w_T .

Lemma 3.4 Let $G = (V, E)$ be a graph and let $\delta \in (0, 1/2)$. For any tree $T \subseteq E$ with at least one edge we have

$$|w_T| \leq \left(\frac{2\delta}{1/2 + \delta} \right)^{|V(T)|}.$$

Proof Let us denote the number of vertices of T by $k \geq 2$. The denominator in the definition of w_T (as found in (3.2)) is equal to $(1/2 + \delta)^k$. So it suffices to bound the

numerator. The numerator can simply be bounded by

$$\int_{[0, 1/2+\delta]^{V(T)}} \prod_{uv \in T} \mathbf{1}_{x_u+x_v>1} d\mu,$$

which in turn can be bounded by $(2\delta)^k$ as it is equal to the volume of a polytope contained in the box $[1/2 - \delta, 1/2 + \delta]^{V(T)}$. This proves the lemma. \square

Proof of Theorem 3.1 We will prove the theorem by proving inductively on the number of vertices of G the following two statements:

- $p_{G,\delta}(x) \neq 0$,
- for any vertex v of G , $\left| \frac{p_{G,\delta}(x)}{p_{G-v,\delta}(x)} - 1 \right| \leq 1/\Delta$.

In case that G consist of a single vertex the two statements are clearly true. Next assume that $|V(G)| > 1$. Since $1/\Delta \in (0, 1)$, the second item implies the first by induction. So we focus on proving the second item.

We have for any vertex v ,

$$p_{G,\delta}(x) = p_{G-v,\delta}(x) + \sum_{\substack{T \subseteq E \\ v \in V(T)}} x^{|T|} w_T p_{G \setminus V(T),\delta}(x),$$

where the sum is over trees T with at least one edge, containing the vertex v . This follows by considering the forests that have a nontrivial component that contains v and those that do not. Since by induction $p_{G-v,\delta}(x) \neq 0$, this implies that

$$\frac{p_{G,\delta}(x)}{p_{G-v,\delta}(x)} - 1 = \sum_{\substack{T \subseteq E \\ v \in V(T)}} x^{|T|} w_T \frac{p_{G \setminus V(T),\delta}(x)}{p_{G-v,\delta}(x)}.$$

By induction and a telescoping argument we can bound

$$\left| \frac{p_{G \setminus V(T),\delta}(x)}{p_{G-v,\delta}(x)} \right| \leq \left(\frac{1}{1 - 1/\Delta} \right)^{|E(T)|}.$$

Now simply bounding $\frac{2\delta}{1/2+\delta}$ by 4δ we have by Lemma 3.4 that $|w_T| \leq (4\delta)^{|V(T)|}$ and therefore

$$\left| \frac{p_{G,\delta}(x)}{p_{G-v,\delta}(x)} - 1 \right| \leq 4\delta(T_{G,v}(|x| \frac{4\delta}{1-1/\Delta}) - 1).$$

Next we use that $|x|\delta \leq \frac{\log(a)(1-1/\Delta)}{a4\Delta}$ by assumption, in combination with Lemma 3.3 to conclude that

$$\left| \frac{p_{G,\delta}(x)}{p_{G-v,\delta}(x)} - 1 \right| \leq 4\delta(T_{G,v}(\frac{\log a}{a\Delta}) - 1) \leq 4\delta(a - 1) < 1/\Delta,$$

as desired. \square

4 Computing the Coefficients of $\text{ind}(H, \cdot)$

The aim in this section is to give a proof of Theorem 2.5. We recall that for a graph H and $k \in \mathbb{N}$ such that $|V(H)| = k(H) + k$ we have that the coefficient $\lambda_{H,k}$ of $\text{ind}(H, G)$ is given by (cf. (2.8)),

$$\lambda_{H,k} = \prod_{C \text{ comp. of } H} w(H),$$

provided each component of H has size at least 2; otherwise $\lambda_{H,k} = 0$. We use (3.3) to obtain the following alternative expression:

$$\lambda_{H,k} = \sum_{\substack{F \subseteq E(H) \\ \text{sp. forest} \\ |F|=k}} \prod_{T \text{ comp. of } F} w_T. \tag{4.1}$$

In the next subsection we will outline how to compute the w_T , after which we will give our proof of Theorem 2.5.

4.1 Computation of w_T

Let us recall that

$$w_T = \frac{(-1)^{|T|} \int \prod_{uv \in T} \mathbf{1}_{x_u+x_v>1} \cdot \prod_{st \in E_T} \mathbf{1}_{x_u+x_v \leq 1} d\mu}{\int 1 d\mu},$$

where the integrations are over $I^{V(T)}$.

The denominator in w_T is equal to the volume of $I^{V(T)}$, that is $(1/2 + \delta)^{|V(T)|}$. Thus we only have to focus on the integral appearing in the numerator, which we will denote by $(-1)^{|V(T)|} \hat{w}_T$.

Remark 4.1 We note that \hat{w}_T can be seen as the volume of a polytope with at most $\binom{\Delta k}{k}$ many vertices and as such the geometric algorithm (based on determinants) of Lawrence [23] (see also [18]) could be used to compute \hat{w}_T in time $\Delta^{O(k)}$. We will however provide a direct/combinatorial algorithm.

Proposition 4.2 For any connected graph $G = (V, E)$ on $k \geq 2$ vertices of maximum degree Δ and any spanning tree $T \subseteq E(G)$ the quantity

$$\hat{w}_T = \int \prod_{uv \in T} \mathbf{1}_{x_u+x_v>1} \cdot \prod_{st \in E_T} \mathbf{1}_{x_u+x_v \leq 1} d\mu$$

can be computed in $\Delta^{O(k)}$ time, where the implicit constant does not depend on Δ .

The remainder of this subsection is devoted to proving Proposition 4.2. Let us denote the function appearing in the integral as $f(\mathbf{x})$. To prove the proposition, we will reduce

the problem to calculating the integral of $f(\mathbf{x})$ over some nice (but exponentially many) smaller polytopes.

First we have to compute E_T . It is clear from the definition that this can be done in polynomial time in the number of vertices, i.e. $\text{POLY}(k)$.

Since all the variables appearing in the integral have to be in the interval $[0, 1/2 + \delta]$ and T is connected, it means that any variable has to be in $J = [1/2 - \delta, 1/2 + \delta]$, thus we can restrict the integral of $f(\mathbf{x})$ over J^V . For later purposes let $J_- = [1/2 - \delta, 1/2]$.

For each $S \subseteq V$ let

$$P_S := \left\{ x \in J^V \mid \begin{array}{l} x_u \leq 1/2 \text{ if } u \in S \\ x_u \geq 1/2 \text{ if } u \notin S \end{array} \right\}.$$

Clearly, this defines a decomposition of J^V into 2^k pieces, i.e.

$$J^V = \bigcup_{S \subseteq V} P_S,$$

If $S' \neq S$, then there exists $u \in S \Delta S'$ and thus $P_S \cap P_{S'} \subseteq \{x \in J^V \mid x_u = 1/2\}$. Therefore, we have

$$\hat{w}_T = \sum_{S \subseteq V(T)} \int_{P_S} f(\mathbf{x}) d\mu.$$

Some of these integrals are actually zero, which we can rule out. If $uv \in T$ and $u, v \in S$, then $\int_{P_S} f(\mathbf{x}) d\mu = 0$, since it is the volume of a polytope with codimension at least 1 (since if $f(\mathbf{x}) \neq 0$ then $x_u = x_v = 1/2$). Similarly, if $uv \in E_T$ and $u, v \in V \setminus S$, then $\int_{P_S} f(\mathbf{x}) = 0$. We therefore call a set $S \subseteq V$ nice if S is independent in T and $V \setminus S$ is independent in E_T .

For a given nice set $S \subseteq V$ endow it with a linear ordering. Let us consider two maps $s, t : V \setminus S \rightarrow S \cup \{\oplus, \ominus\}$, such that

- if $N_T(u) \cap S = \emptyset$, then $s(u) = \oplus$, otherwise $s(u) \in N_T(u) \cap S$,
- if $N_{E_T}(u) \cap S = \emptyset$, then $t(u) = \ominus$, otherwise $t(u) \in N_{E_T}(u) \cap S$.

We denote the set of pairs (s, t) of such maps by \mathcal{F}_S . Then for each $(s, t) \in \mathcal{F}_S$ we define a polytope

$$P_{S,(s,t)} := \left\{ \mathbf{x} \in P_S \mid \forall u \in V \setminus S : \begin{array}{l} x_i \geq x_{s(u)} \text{ for all } i \in N_T(u) \cap S \\ x_i \leq x_{t(u)} \text{ for all } i \in N_{E_T}(u) \cap S \end{array} \right\},$$

where we set $x_\ominus = 1/2 - \delta$, $x_\oplus = 1/2$. Clearly, this defines a decomposition of P_S into at most Δ^{2k} parts, i.e.

$$P_S = \bigcup_{(s,t) \in \mathcal{F}_S} P_{S,(s,t)},$$

which implies that

$$\int_{P_S} f(\mathbf{x}) d\mu = \sum_{(s,t) \in \mathcal{F}_S} \int_{P_{S,(s,t)}} f(\mathbf{x}) d\mu,$$

since for distinct (s, t) and (s', t') the intersection of the polytopes $P_{S,(s,t)}$ and $P_{S,(s',t')}$ lies in an affine hyperplane.

Order the elements of $V \setminus S$ as v_1, \dots, v_ℓ . Then for a fixed $(s, t) \in \mathcal{F}_S$ we have the equality

$$\int_{P_{S,(s,t)}} f(\mathbf{x}) \, d\mu = \int_{J_-^S} \left(\int_{1-x_s(v_1)}^{1-x_t(v_1)} \dots \int_{1-x_s(v_\ell)}^{1-x_t(v_\ell)} \mathbf{1} \, d\mu(x_{v_1}) \dots d\mu(x_{v_\ell}) \right) \cdot g_{(s,t)}(\mathbf{x}) \, d\mu, \tag{4.2}$$

where $g_{(s,t)}(\mathbf{x}) : \mathbb{R}^S \rightarrow \mathbb{R}$ is defined as

$$g_{(s,t)}(\mathbf{x}) = \prod_{i=1}^{\ell} \mathbf{1}_{x_{s(v_i)} > x_{t(v_i)}} \cdot \prod_{v \in (N_T(v_i) \cap S) \cup \{\oplus\}} \mathbf{1}_{x_v > x_{s(v_i)}} \cdot \prod_{v \in (N_{E_T}(v_i) \cap S) \cup \{\ominus\}} \mathbf{1}_{x_v < x_{t(v_i)}}.$$

Indeed, this follows from the observation that for a vertex $v \in V \setminus S$ an edge uv with $u \in S$ and we have that if $uv \in T$ the condition $x_u + x_v \geq 1$ translates to $x_v \geq 1 - x_u$, while if $uv \in E_T$ the condition $x_u + x_v \leq 1$ translates to $x_v \leq 1 - x_u$ (if v has no T -neighbor in S we have $s(v) = \oplus$ and the condition $x_v \geq 1/2$ translates to $x_v \geq 1 - 1/2$ while if v has no E_T -neighbor in S we have $t(v) = \ominus$ and the condition $x_v \leq 1/2 + \delta$ translates to $x_v \leq 1 - (1/2 - \delta)$). Using that for each neighbor u of v we have by definition of the polytope $P_{S,(s,t)}$ that $x_u \geq x_{s(v)}$ and $x_u \leq x_{t(v)}$ the identity (4.2) follows.

By (4.2) we thus have

$$\int_{P_{S,(s,t)}} f(\mathbf{x}) \, d\mu = \int_{J_-^S} \underbrace{\prod_{i=1}^{\ell} (x_{s(v_i)} - x_{t(v_i)})}_{p(\mathbf{x})} \cdot g_{(s,t)}(\mathbf{x}) \, d\mu, \tag{4.3}$$

where $p(\mathbf{x})$ has total degree at most k and has at most k variables.

Let us next give an interpretation for $g_{(s,t)}(\mathbf{x})$. For a given $(s, t) \in \mathcal{F}_S$ we define the digraph $D = D_{(s,t)}$ on S where $(u, v) \in E(D)$, if there exists $v_i \in V \setminus S$ such that

- either $v = s(v_i)$ and $u = t(v_i)$,
- or $u = s(v_i)$ and $v \in N_T(v_i) \cap S$,
- or $v = t(v_i)$ and $u \in N_{E_T}(v_i) \cap S$.

Observe that each term in $g_{(s,t)}(\mathbf{x})$ corresponds to an arc of this digraph, in the sense that $0 \neq g_{(s,t)}(\mathbf{x})$ if and only if the ordering defined by \mathbf{x} on its coordinates is compatible with the digraph D , that is, for every $(u, v) \in D$ we have $x_u < x_v$. We see immediately that this integral above is 0 if D contains an oriented cycle. Therefore we may assume that $D = D_{(s,t)}$ is acyclic, i.e. it defines a poset \mathcal{P} on S . The next lemma provides an algorithm for evaluating (4.3).

Lemma 4.3 Let \mathcal{P} a poset on a set S of size k and let $a < b \in \mathbb{R}$. Let $p(\mathbf{x})$ a polynomial in variables $\{x_u\}_{u \in S}$ of total degree d with integer coefficients. Let

$$g(\mathbf{x}) = \prod_{u < v} \mathbf{1}_{x_u < x_v} \prod_{u \in S} \mathbf{1}_{b > x_u} \prod_{u \in S} \mathbf{1}_{x_u > a}.$$

Then

$$\int_{\mathbb{R}^S} p(\mathbf{x})g(\mathbf{x}) \, d\mu$$

can be computed in $O(k2^{d+3k})$ time.

Proof Let us define for each $\emptyset \neq U \subseteq S$ a function $F_U : \mathbb{R}^2 \times \mathbb{R}^{S \setminus U} \rightarrow \mathbb{R}$ by

$$F_U(m, M, \{x_v\}_{v \in S \setminus U}) := \sum_{u \in \min U} \int_m^M F_{U-u}(m, M, \{x_v\}_{v \in S \setminus U \cup \{u\}})|_{m=x_u} \, dx_u,$$

where $\min U$ denotes the set of minimal elements of the poset restricted to U , and define

$$F_{\emptyset}(m, M, \{x_v\}_{v \in S}) := p(\mathbf{x}).$$

It is clear that each F_U is polynomial in the $k + 2 - |U|$ variables $\{x_v\}_{v \notin U}$, m and M and it has total degree at most $d + |U|$. This means that among all the polynomials $\{F_U\}_{U \subseteq S}$ we see at most $\binom{d+2k+2}{k+2} < 2^{d+2k+2}$ many different monomials. This means that to calculate F_U from F_{U-u} for a minimal element u of U (by which we mean computing the coefficients of each monomial) we need at most $O(k2^{d+2k})$ time. This implies that to calculate F_S we need at most $O(k2^{d+2k} \cdot 2^k) = O(k2^{d+3k})$ time.

We next claim that

$$F_U(m, M, \{x_u\}_{u \in S \setminus U})|_{m=a, M=b} = \int_{\mathbb{R}^U} p(\mathbf{x})g_U(a, b, \mathbf{x}) \, d\mu, \tag{4.4}$$

where

$$g_U(m, M, \mathbf{x}) = \prod_{u, v \in U, u < v} \mathbf{1}_{x_u < x_v} \prod_{u \in U} \mathbf{1}_{x_u < M} \prod_{u \in U} \mathbf{1}_{x_u > m}.$$

We will prove this by induction on the size of U . If $U = \emptyset$, then it is trivial, so let us assume that $U \neq \emptyset$. Then using that we can partition the space by saying which minimal element of U is the smallest we have

$$\begin{aligned} \int_{\mathbb{R}^U} p(\mathbf{x})g_U(a, b, \mathbf{x}) \, d\mu &= \int_{\mathbb{R}^U} \sum_{u \in \min U} \left(\prod_{\substack{v \in \min U \\ v \neq u}} \mathbf{1}_{x_v > x_u} \right) p(\mathbf{x})g_U(a, b, \mathbf{x}) \, d\mu \\ &= \sum_{u \in \min U} \int_{\mathbb{R}^U} \left(\prod_{\substack{v \in \min U \\ v \neq u}} \mathbf{1}_{x_v > x_u} \right) p(\mathbf{x})g_U(a, b, \mathbf{x}) \, d\mu \end{aligned}$$

$$\begin{aligned}
 &= \sum_{u \in \min U} \int_{\mathbb{R}^U} \left(\prod_{v \in \min U - u} \mathbf{1}_{x_v > x_u} \right) p(\mathbf{x}) g_{U-u}(a, b, \mathbf{x}) \cdot \mathbf{1}_{x_u > a} \mathbf{1}_{x_u < b} d\mu \\
 &= \sum_{u \in \min U} \int_{\mathbb{R}^U} p(\mathbf{x}) g_{U-u}(x_u, b, \mathbf{x}) \cdot \mathbf{1}_{x_u > a} \mathbf{1}_{x_u < b} d\mu. \tag{4.5}
 \end{aligned}$$

Next we apply Fubini’s theorem to obtain that (4.5) is equal to

$$\begin{aligned}
 &= \sum_{u \in \min U} \int_a^b \left(\int_{\mathbb{R}^{U-u}} p(\mathbf{x}) g_{U-u}(x_u, b, \mathbf{x}) d\mu \right) dx_u \\
 &= \sum_{u \in \min U} \int_a^b F_{U-u}(x_u, b, \{x_v\}_{v \in S-U \cup \{u\}}) dx_u,
 \end{aligned}$$

where the last step follows by using induction. We thus see that $\int_{\mathbb{R}^U} p(\mathbf{x}) g_U(a, b, \mathbf{x}) d\mu$ equals $F_U(m, M, \{x_u\}_{u \in S-U})|_{m=a, M=b}$ by definition, thereby proving (4.4).

As a consequence we obtain that for any $a < b$ we have

$$\int_{\mathbb{R}^S} p(\mathbf{x}) g(\mathbf{x}) d\mu = F_S(a, b).$$

This finishes the proof, as we have already indicated how to compute the F_U for $U \subseteq S$. □

Proof of Proposition 4.2 Using the notations above we see that

$$\hat{w}_T = \sum_{\substack{S \subseteq V(T) \\ S \text{ is nice}}} \left(\sum_{\substack{(s,t) \in \mathcal{F}_S \\ D_{(s,t)} \text{ is acyclic}}} \int_{P_{S,(s,t)}} f(\mathbf{x}) d\mu \right).$$

By the previous lemma we can compute

$$\int_{P_{S,(s,t)}} f(\mathbf{x}) d\mu$$

in $O(k2^{4k})$ time, thus \hat{w}_T can be computed in $O(2^k \cdot \Delta^{2k} \cdot 2^{4k} \cdot \text{POLY}(k)) = \Delta^{O(k)}$ time. □

4.2 Computation of $\lambda_{H,k}$

We now conclude this section by giving a proof of Theorem 2.5.

Proof of Theorem 2.5 We may assume that the graph H has maximum degree at most Δ for otherwise the coefficient of $\text{ind}(H, \cdot)$ in $P_{G,\delta}$ will be zero. Since we can also assume that H has at most $2k$ vertices, it will have at most Δk edges. By (4.1) we

need to enumerate all forests F of k edges in H and compute the weights w_T of each component T of F . To enumerate all forests with k edges in H we just go over all subsets of $E(H)$ with k elements, which takes time bounded by $\binom{\Delta k}{k} \leq (e\Delta)^k$, by Stirling's formula. For each such a set we identify the components, T_1, \dots, T_ℓ and whether these components are trees with at least one edge, which takes $\text{POLY}(k)$ time.

By the algorithm of Proposition 4.2 we compute the normalized weights

$$\hat{w}_{T_i} = (-1)^{|T_i|} (1/2 + \delta)^{|V(T_i)|} w_{T_i}$$

and hence the weights w_{T_i} in $\Delta^{O(|V(T_i)|)}$ time. All together the algorithm takes $\Delta^{O(k)}$ time. □

5 Concluding Remarks, Open Problems and Discussion

We discuss here some extensions and limitations of our method and mention some open problems.

Gamarnik and Smedira [15] indicate that their method for approximating the volume of $P_{G,\delta}$ does not extend all the way to $\delta = 1/2$. Even though we improve on their bound on δ from $O(1/\Delta^2)$ to $O(1/\Delta)$, our method also comes with limitations that we will explain now. As detailed in (3.3) the volume of the polytope $P_{G,\delta}$ can be seen as the evaluation of a multivariate forest generating function, i.e. the volume can be expressed as

$$\tilde{p}_G(x; \{w_T\}) := \sum_{\substack{F \subseteq E(G) \\ F \text{ forest}}} \prod_{T \text{ comp. of } F} w_T \cdot x^{|T|},$$

where $\tilde{p}_G(x; \{w_T\})$ is the multivariate forest generating function with tree weights $\{w_T\}$. Crucial for establishing absence of zeros of $p_{G,\delta}$ is that the tree weights $|w_T|$ are exponentially small (in our setting they are bounded by $(4\delta)^{|V(T)|}$, see Lemma 3.4). This is in fact the only property that we use, in the sense that there is a $\delta' = O(1/\Delta)$ such that if for every tree T the absolute value of its weight is in $[0, \delta'^{|E(T)|}]$ then for any graph of maximum degree at most Δ the polynomial $\tilde{p}_G(x; \{\tilde{w}_T\})$ has a zero-free disk of radius bigger than 1. In particular, it means that

$$\tilde{p}_{G,\delta'}(x) := \sum_{\substack{F \subseteq E(G) \\ F \text{ forest}}} (\delta' x)^{|F|}$$

which is the forest generating function of G evaluated at $\delta'x$, has a zero-free disk of radius larger than 1 (for the variable x). A result of the first author together with Csikvári [7] implies that no zero-free disk of radius larger than 1 (for the variable x) exists when $\delta' > \frac{1}{\Delta-2}$. This suggests that to extend Theorem 2.4 beyond $\delta = O(1/\Delta)$ one needs to devise a proof that uses more information about the weights w_T than just the fact they are exponentially small. We leave it as a challenging open problem to do so, or to find another approach for deterministically approximating the volume of

$P_{G,\delta}$ for $\delta \gg 1/\Delta$, remarking that the results from [16] do suggest that this should be possible.

Now let us turn to possible extensions of our approach. Our approach relies on three ingredients: 1) express the volume of the polytope as an evaluation of a graph polynomial, 2) establish a zero-free region for this polynomial and 3) ‘efficiently’ compute the low (logarithmically small) order terms of the Taylor series.

- Volumes of polytopes.** Let P be a polytope defined by $Ax \leq b$ and $x \in [0, 1]^d$, where $A \in \mathbb{R}^{n \times d}$. If each row of A has at most 2 non-zero coefficient, then in a similar fashion we can express the volume of P as an evaluation of the multivariate forest generating polynomial. To address the case when the number of non-zero coefficients in each row of A is bounded by some $k > 2$, we obtain the volume as an evaluation of hypergraph polynomial. We expect that the ideas of [13] are applicable in this scenario to establish a zero-free region for a similarly defined polynomial when the polytope is appropriately truncated.

To compute the low order terms of the Taylor series of the logarithm one could use the algorithm of Lawrence [23]. It is not clear to us how to directly generalize our combinatorial approach.

We note that recently Guo and N [20] designed an algorithm for the volume of the fractional hypergraph matching polytope based on our approach.

- Homomorphism count.** Let H be a weighted graph on $[m] = \{1, \dots, m\}$ with adjacency matrix A . Then the number of homomorphism of G into H can be expressed as

$$\text{hom}(G, H) = \sum_{\phi: V(G) \rightarrow [m]} \prod_{uv \in E(G)} A_{\phi(u), \phi(v)}. \tag{5.1}$$

We can think about this expression as a discretized generalization of (2.1). Indeed, if we let I be the interval $[0, m]$ and let $f : I \times I \rightarrow \mathbb{R}$ be the step function representing A , that is

$$f(x, y) = \begin{cases} A_{\lfloor x/m \rfloor, \lfloor y/m \rfloor} & \text{if } x, y \in I, \\ 0 & \text{otherwise,} \end{cases}$$

then

$$\text{hom}(G, H) = \int_{I^{V(G)}} \prod_{uv \in E(G)} f(x_u, y_v) d\mu. \tag{5.2}$$

The approximation of $\text{hom}(G, H)$ has already been addressed in case all the entries of A are close to 1 [5, 30]. It is proved that for any Δ, m there is a algorithm to give ε -approximation of $\text{hom}(G, H)$ for graphs of maximum degree Δ (assuming that the entries of A are $0.35/\Delta$ close to 1) in polynomial time in $|V(G)|/\varepsilon$, however the degree of this polynomial depends on m . Our current approach allows to also treat matrices A for which not all of its entries are close to 1, see our next comment below.

- Homomorphism densities and general integrals.** One could further generalize (5.2) by simply allowing f to be an arbitrary measurable (symmetric) function on some measure space $(X, \mu)^2$. In particular, if f is $[0, 1]$ valued and (X, μ) is

a probability space, then f is a graphon and (5.2) expresses the homomorphism density of G in the graphon f (see [24]). For this setup let us remark the following observations.

Similarly as in Section 2 we have

$$\text{hom}(G, f) = p_G(x)|_{x=1} = \sum_{P \in \pi(V)} \prod_{S \in P} w(G[S])x^{|S|-1}|_{x=1},$$

where

$$w(H) = \sum_{\substack{T \subseteq E \\ (V(H), T) \text{ tree}}} (-1)^{|T|} \underbrace{\int_{X^S} \prod_{uv \in T} (1 - f(x_u, x_v)) \cdot \prod_{st \in E(S)_T} f(x_s, x_t) d\mu}_{w_T}.$$

For a given graphon f let $c(f) = \sup_{x \in X} \int_X (1 - f(x, y)) dy$. This parameter plays a crucial role in [9] to establish the convergence radius of a cluster expansion for the homomorphism count in the context of sparse graph limits. It is not hard to see that $|w_T| \leq c(f)^{|T|}$, thus one can easily obtain that if $c(f)$ is at most $O(1/\Delta)$, then $p_G(x)$ is zero-free on a disk of radius bigger than 1 (i.e. a variant of Theorem 2.4 holds). However, it is not clear how to compute the low order terms of the polynomial in an efficient way for an arbitrary function f . Our main result (Theorem 1.1) deals with functions of the form $f = \mathbf{1}_{x+y \leq 1}$ and we heavily relied on the combinatorics provided by f .

A different approach to establish an approximation algorithm for computing the homomorphism density in f would be by to approximate f with a step-function \hat{f} . This idea was used in [17] in combination with the correlation decay method to find a deterministic quasi-polynomial time approximation algorithm for computing $\text{hom}(G, f)$ (provided that f is smooth enough and close enough to a constant function). Combining this idea with our approach we can also obtain a quasi-polynomial time approximation algorithm, but we have to leave it as an open problem to improve the running time to actual polynomial time.

Acknowledgements We thank an anonymous referee for constructive feedback.

Data Availability No new data were created or analysed in this study. Data sharing is not applicable to this article.

Declarations

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Anari, N., Liu, K., Gharan, S.O.: Spectral independence in high-dimensional expanders and applications to the hardcore model. *SIAM J. Comput.* **53**(6), FOCS20–1–FOCS20–37 (2024)
2. Bandyopadhyay, A., Gamarnik, D.: Counting without sampling. New algorithms for enumeration problems using statistical physics, Proceedings of the Seventeenth Annual ACM-SIAM Symposium on Discrete Algorithms, pp. 890–899, (2006)
3. Bárány, I., Füredi, Z.: Computing the volume is difficult. *Discrete Comput. Geom.* **2**(4), 319–326 (1987)
4. Barvinok, A.: Asymptotic estimates for the number of contingency tables, integer flows, and volumes of transportation polytopes. *Int. Math. Res. Not. IMRN* **2**, 348–385 (2009)
5. Barvinok, A.: Combinatorics and complexity of partition functions, Algorithms and Combinatorics, vol. 30. Springer, Cham (2016)
6. Barvinok, A., Rudelson, M.: A quick estimate for the volume of a polyhedron. *Israel J. Math.* **262**(1), 449–473 (2024)
7. Bencs, F., Csikvári, P.: Evaluations of Tutte polynomials of regular graphs. *J. Combin. Theory Ser. B* **157**, 500–523 (2022)
8. Bissacot, R., Fernández, R., Procacci, A.: On the convergence of cluster expansions for polymer gases. *J. Stat. Phys.* **139**(4), 598–617 (2010)
9. Borgs, C., Chayes, J., Kahn, J., Lovász, L.: Left and right convergence of graphs with bounded degree. *Random Structures & Algorithms* **42**(1), 1–28 (2013)
10. Chen, Z., Liu, K., Vigoda, E.: Optimal mixing of Glauber dynamics: entropy factorization via high-dimensional expansion, Proceedings of the 53rd annual ACM SIGACT Symposium on Theory of Computing, pp. 1537–1550, (2021)
11. Dobrushin, R.L.: Estimates of semi-invariants for the Ising model at low temperatures. *Translations of the American Mathematical Society-Series* **2**(177), 59–82 (1996)
12. Dyer, M., Frieze, A., Kannan, R.: A random polynomial-time algorithm for approximating the volume of convex bodies. *J. Assoc. Comput. Mach.* **38**(1), 1–17 (1991)
13. Fadnavis, S.: On the roots of hypergraph chromatic polynomials, arXiv preprint [arXiv:1509.05950](https://arxiv.org/abs/1509.05950) (2015)
14. Galanis, A., Štefankovič, D., Vigoda, E.: Inapproximability of the partition function for the antiferromagnetic Ising and hard-core models. *Combin. Probab. Comput.* **25**(4), 500–559 (2016)
15. Gamarnik, D., Smedira, D.: Computing the volume of a restricted independent set polytope deterministically, arXiv preprint [arXiv:2312.03906](https://arxiv.org/abs/2312.03906) (2023)
16. Gamarnik, D., Ramanan, K.: Uniqueness of Gibbs measures for continuous hardcore models. *Ann. Probab.* **47**(4), 1949–1981 (2019)
17. Gamarnik, D., Smedira, D.: Integrating high-dimensional functions deterministically, arXiv preprint [arXiv:2402.08232](https://arxiv.org/abs/2402.08232) (2024)
18. Gritzmann, P., Klee, V.: Computational convexity, Handbook of discrete and computational geometry, pp. 937–964, (2017)
19. Gruber, C., Kunz, H.: General properties of polymer systems. *Comm. Math. Phys.* **22**, 133–161 (1971)
20. Guo, H., Vishvajeet N.: Deterministic approximation for the volume of the truncated fractional matching polytope, 16th Innovations in Theoretical Computer Science Conference, pp. Art. No. 57, 14, (2025)
21. Jenssen, M., Patel, V., Regts, G.: Improved bounds for the zeros of the chromatic polynomial via Whitney’s broken circuit theorem. *J. Combin. Theory Ser. B* **169**, 233–252 (2024)
22. Kotecký, R., Preiss, D.: Cluster expansion for abstract polymer models. *Comm. Math. Phys.* **103**(3), 491–498 (1986)
23. Lawrence, J.: Polytope volume computation. *Math. Comp.* **57**(195), 259–271 (1991)
24. Lovász, L.: Large networks and graph limits, American Mathematical Society Colloquium Publications, vol. 60. American Mathematical Society, Providence, RI (2012)
25. Patel, V., Regts, G.: Approximate counting using Taylor’s theorem: a survey. *Bull. Eur. Assoc. Theor. Comput. Sci. EATCS* **138**, 40–70 (2022)
26. Penrose, O.: Convergence of fugacity expansions for classical systems. *Statistical mechanics: foundations and applications* **101**, (1967)
27. Peters, H., Regts, G.: On a conjecture of Sokal concerning roots of the independence polynomial. *Michigan Math. J.* **68**(1), 33–55 (2019)

28. Schrijver, A.: Combinatorial optimization. Polyhedra and efficiency. Vol. A, Algorithms and Combinatorics, vol. 24,A, Springer-Verlag, Berlin, 2003. Paths, Flows, Matchings, Chapters 1–38
29. Sly, A., Sun, N.: Counting in two-spin models on d -regular graphs. *Ann. Probab.* **42**(6), 2383–2416 (2014)
30. Viresh Patel and Guus Regts: Deterministic polynomial-time approximation algorithms for partition functions and graph polynomials. *SIAM J. Comput.* **46**(6), 1893–1919 (2017)
31. Weitz, D.: Counting independent sets up to the tree threshold, STOC'06: Proceedings of the 38th Annual ACM Symposium on Theory of Computing, pp. 140–149, (2006)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.