



Centrum voor Wiskunde en Informatica

**REPORT**RAPPORT

**MAS**

Modelling, Analysis and Simulation



*Modelling, Analysis and Simulation*

Global existence conditions for a non-local problem  
arising in statistical mechanics

C.J. van Duijn, I.A. Guerra, M.A. Peletier

**REPORT MAS-R0119 DECEMBER 31, 2001**

CWI is the National Research Institute for Mathematics and Computer Science. It is sponsored by the Netherlands Organization for Scientific Research (NWO).

CWI is a founding member of ERCIM, the European Research Consortium for Informatics and Mathematics.

CWI's research has a theme-oriented structure and is grouped into four clusters. Listed below are the names of the clusters and in parentheses their acronyms.

Probability, Networks and Algorithms (PNA)

Software Engineering (SEN)

**Modelling, Analysis and Simulation (MAS)**

Information Systems (INS)

Copyright © 2001, Stichting Centrum voor Wiskunde en Informatica

P.O. Box 94079, 1090 GB Amsterdam (NL)

Kruislaan 413, 1098 SJ Amsterdam (NL)

Telephone +31 20 592 9333

Telefax +31 20 592 4199

ISSN 1386-3703

# Global Existence Conditions for a Non-local Problem Arising in Statistical Mechanics

C. J. van Duijn<sup>‡</sup>, I. A. Guerra<sup>†</sup>, M. A. Peletier<sup>†,‡</sup>

<sup>†</sup> *Centrum voor Wiskunde en Informatica,  
PO Box 94079 1090 GB Amsterdam, The Netherlands*

<sup>‡</sup> *Department of Mathematics and Computer Science, TU Eindhoven,  
PO Box 513 5600 MB Eindhoven, The Netherlands*

## ABSTRACT

We consider the evolution of the density and temperature of a three-dimensional cloud of self-interacting particles. This phenomenon is modelled by a parabolic equation for the density distribution combining temperature-dependent diffusion and convection driven by the gradient of the gravitational potential. This equation is coupled with Poisson's equation for the potential generated by the density distribution. The system preserves mass by imposing a zero-flux boundary condition. Finally the temperature is fixed by energy conservation, that is, the sum of kinetic energy (temperature) and gravitational energy remains constant in time. This model is thermodynamically consistent, obeying the first and the second law of thermodynamics. We prove local existence and uniqueness of weak solutions for the system, using a Schauder fixed-point theorem. In addition, we give sufficient conditions for global in time existence and blow-up for radially symmetric solutions. We do this using a comparison principle for an equation for the accumulated radial mass.

*2000 Mathematics Subject Classification:* 35K60, 35A07, 35B40, 82C21.

*Keywords and Phrases:* non-local nonlinear parabolic equation, energy relation, existence of local solutions, global and blowing up solutions.

*Note:* Work carried out under project MAS 1.2 "Partial Differential Equations in Porous Media Research".

## 1. INTRODUCTION

Let  $\Omega \subset \mathbb{R}^3$  be a bounded open set satisfying  $\sup_{x \in \Omega} |x| = 1$ . In  $\Omega$  we consider the parabolic -elliptic system

$$n_t = \operatorname{div}\{\Theta(t)\nabla n + n\nabla\phi\} \quad \text{in } \Omega \times \mathbb{R}^+, \quad (1.1)$$

$$\Delta\phi = n \quad \text{in } \Omega \times \mathbb{R}^+, \quad (1.2)$$

combined with the energy relation

$$E = \kappa\Theta(t) + \int_{\Omega} n\phi dx \quad \text{in } \mathbb{R}^+, \quad (1.3)$$

where  $E \in \mathbb{R}$  and  $\kappa > 0$  are given parameters. At the boundary  $\partial\Omega \in C^{1+\alpha}$  ( $\alpha > 0$ ) we prescribe

$$(\Theta(t)\nabla n + n\nabla\phi) \cdot \vec{\nu} = 0 \quad \text{on } \partial\Omega \times \mathbb{R}^+, \quad (1.4)$$

$$\phi = 0 \quad \text{in } \partial\Omega \times \mathbb{R}^+, \quad (1.5)$$

where  $\vec{\nu}$  denotes the exterior normal vector on  $\partial\Omega$ . At  $t = 0$  we have the initial condition

$$n(x, 0) = n_0(x) \quad \text{in } \Omega, \quad (1.6)$$

satisfying

$$\int_{\Omega} n_0 dx = 1, \quad \text{and} \quad n_0(x) \geq 0 \quad \text{in} \quad \Omega. \quad (1.7)$$

This set of equations defines Problem **P** for the unknowns  $n, \phi$  and  $\Theta$ . The underlying model is discussed in Section 2, as well as some known properties of the system.

The purpose of the paper is to demonstrate local existence for Problem **P** and to give sufficient conditions on  $E, \kappa$ , and  $n_0$  for global existence. Local existence is shown in Section 3. The proof uses a Schauder fixed-point theorem and a careful construction of an invariant set to avoid degenerate diffusion in (1.1). It requires  $n_0 \in L^p(\Omega)$  for  $p > \frac{3}{2}$ , implying that  $n \in L_{loc}^{\infty}((0, T]; L^{\infty}(\Omega))$  for some  $T \in (0, \infty)$ . Hence we can allow for certain singular initial data which result in solutions that are locally bounded in  $(0, T]$ . Let

$$T^* = \sup\{ T > 0 \mid \text{Problem } \mathbf{P} \text{ has a solution in } (0, T] \}.$$

If  $T^* = \infty$ , the solution is defined globally and if  $T^* < \infty$  we have at least  $\lim_{t \rightarrow T^*} \|n(t)\|_{L^q(\Omega)} = \infty$  for each  $q > \frac{3}{2}$ . For Problem **P** the optimal  $L^p(\Omega)$  space seems to be  $p = \frac{3}{2}$ , since there exists a singular stationary solution in the radial case belonging to  $L^{3/2}(\Omega) \setminus L^q(\Omega)$ , with  $q > \frac{3}{2}$ . This solution is given in Section 2.3. Uniqueness is proven for  $n_0 \in L^p(\Omega)$  with  $p \geq 2$ .

Problem **P** with  $\kappa = 3$  was recently studied in [17]: local existence and uniqueness was obtained for  $p > 3$ .

In Section 4 we consider an auxiliary problem in which we drop the energy relation (1.3) and treat  $\Theta(t)$  as a given function. This provides insight and bounds which we need in order to prove our main result about global existence. In Section 5, we first give the following result about blow-up:

**Theorem 1.1** *Let  $\Omega = B_1(0)$  be the unit ball in  $\mathbb{R}^3$ . If  $\kappa > 6$  and  $E < \frac{1}{4\pi}$ , then  $T^* < \infty$ .*

In Remark 5.2, we show that this choice of parameters implies  $n(x, T^*) = \delta_{x=0}$  and consequently  $\lim_{t \rightarrow T^*} \|n(t)\|_{L^p(\Omega)} = \infty$  for all  $p > 1$ . When the solution satisfies this last condition, we refer to this situation as gravitational collapse.

Before stating the global existence result we note from (1.3) at  $t = 0$ , that instead of prescribing  $E$  and  $n_0$  one could equivalently prescribe  $\Theta_0 := \Theta(0)$  and  $n_0$ . In fact it seems more natural to consider  $\Theta_0$  and  $n_0$  as initial values. In view of the physical interpretation of the model we consider  $\Theta_0 > 0$ . With this in mind we have

**Theorem 1.2** *Let  $\Omega = B_1(0)$  and assume that solutions of Problem **P** are radially symmetric. If the pair  $\langle n_0, \Theta_0 \rangle$  satisfies one of the following conditions*

- (i)  $n_0 \in L^{\infty}(\Omega)$  and  $\Theta_0$  is sufficiently large;
- (ii) there exists  $B > 0$  and  $\lambda = \lambda(n_0, \kappa) \in (0, 1]$  such that

$$\|n_0\|_{L^1(B_r(0))} \leq (1 + B) \frac{r^3}{r^2 + B} \leq r \quad \text{for} \quad r \in [0, 1], \quad \text{and} \quad \Theta_0 \geq \frac{3(1 + B)}{\lambda};$$

- (iii)  $n_0 \equiv \frac{3}{4\pi}$  and  $\Theta_0 > \frac{3}{2}$ ,

then  $T^* = \infty$  (global existence).

**Remark 1.3** (i) *Due to the parabolic regularity,  $n_0 \in L^{\infty}(\Omega)$  is not so restrictive.*

- (ii) *Condition (iii) is a special case of (ii).*

(iii) *The condition on  $n_0$  in (ii) implies a bound on the Morrey norm of exponent 3/2, since  $\|n_0\|_{M^{3/2}(\Omega)} = \sup_{x \in \mathbb{R}^3, 0 \leq r \leq 1} r^{-1} \|n_0\|_{L^1(\Omega \cap B_r(x))}$ . In [4], the space  $M^{3/2}(\Omega)$  was suggested as the natural space to prove existence.*

The proof of Theorem 1.2 contains two essential steps. To extend the local solution we first need a uniform bound from below on  $\Theta$ . To achieve this we use a Lyapunov functional associated with Problem **P**, the so-called Boltzmann entropy (2.7). This functional provides a uniform lower bound on  $\Theta$ , which only depends on the initial data and  $\kappa$ . If  $\Theta_0$  is positive, then  $\Theta$  remains positive in the whole existence interval, including the blow-up time.

In the second step we construct a control on  $n$ . Here we use the radial symmetry which allows us to transform equations (1.1) and (1.2) into a single equation, still containing  $\Theta$  as unknown. It has the crucial property that an ordered pair of given  $\Theta$ 's results in an ordered pair of solutions. As a comparison function we now use the solution of (1.1)-(1.2) with a suitably chosen fixed  $\Theta$ . Under certain hypotheses this auxiliary problem has a global solution which provides the control on  $n$ . The different conditions in Theorem 1.2 are closely related to global existence conditions for the auxiliary problem.

## 2. PRELIMINARIES

### 2.1 Model issues

Problem **P** describes the evolution of density and temperature of a self-attracting cluster of Brownian particles in a bounded three-dimensional region. During the evolution mass and energy are conserved. A detailed derivation and discussion on the physical assumptions can be found in [5, 8, 19] and the references therein. Below we present a brief summary.

Suppose a cluster of particles is contained in a bounded region  $\Omega \subset \mathbb{R}^3$ . The spatial particle density  $n$  satisfies the mass balance equation

$$n_t = \operatorname{div} \left\{ \frac{1}{\beta} (k\Theta \nabla n + n \nabla \phi) \right\} \quad \text{in } \Omega \times \mathbb{R}^+, \quad (2.1)$$

where  $\beta > 0$  is the friction coefficient,  $k$  the Boltzmann constant and  $\Theta$  the temperature of the system. To ensure that the cluster of particles preserves mass we impose zero mass flux along the boundary: i.e.

$$(k\Theta \nabla n + n \nabla \phi) \cdot \vec{\nu} = 0 \quad \text{on } \partial\Omega. \quad (2.2)$$

This implies

$$\int_{\Omega} n(x, t) dx = \text{constant} = M \quad \text{for all } t > 0,$$

where  $M$  is the total particle mass of the system, specified by the initial condition.

The function  $\phi$  in (2.1) is the gravitational potential. It satisfies

$$\Delta \phi = 4\pi G n \quad \text{in } \Omega \times \mathbb{R}^+, \quad (2.3)$$

with

$$\phi = -\frac{GM}{R} \quad \text{on } \partial\Omega \times \mathbb{R}^+. \quad (2.4)$$

Here  $G$  is the gravitational constant and  $R := \max_{x \in \Omega} |x|$ . Note that we have chosen as boundary condition the gravitational potential of a mass  $M$  centered at the origin of a ball of radius  $R$ .

In general the temperature varies in space and time. It satisfies an energy balance equation containing thermal diffusion, heat convection and a term due to gravitational effects [8, Eq. (1.4)]. This results in the so-called Streater model. However, the integrated energy balance does not contain the thermal diffusivity [8, Eq. (2.1)]. Furthermore we expect that a large thermal diffusivity will result in

a temperature which is nearly constant in space. Taking this limit in the integrated energy balance, one finds

$$E = \frac{\kappa M}{2} \Theta(t) + \frac{1}{2} \int_{\Omega} n \phi dx \quad \text{in } \mathbb{R}^+, \quad (2.5)$$

where  $E$  denotes the total energy of the system and  $\kappa$  the specific heat of the particles. If the cluster resembles an ideal gas we have  $\kappa = 3k$ .

Regarding the initial data for the system (2.1)-(2.5) there are two ways to proceed. If the energy  $E$  is given, it suffices to specify only the initial density

$$n(x, 0) = n_0(x) \geq 0 \quad \text{for } x \in \Omega. \quad (2.6)$$

Equivalently we can specify both initial density and temperature

$$\Theta(0) = \Theta_0 > 0.$$

Now  $E$  is fixed by (2.5) at  $t = 0$ .

If the temperature is constant in time as well we drop the energy balance (2.5) and obtain the isothermal model. This model also arises in the context of polytropic stars and the biological phenomena of chemotaxis. The corresponding mathematical problem has received considerable attention in the past years because of its rich structure. Blow-up in the form of singular solutions and gravitational collapse can occur, as well as global existence. To our knowledge there is no full description of these phenomena in  $\mathbb{R}^3$ . The reason is that in contrast to the two dimensional case, global existence in  $\mathbb{R}^3$  not only depends on the parameters of the problem, but also on the shape of the initial density profile. A detailed discussion and references are given in [13]. The isothermal model, however, plays a crucial role in the analysis presented in this paper.

Since Problem **P** has an additional equation, one expects that conservation of energy will act as a selection principle to favor global existence. This has been demonstrated in [18] for the two-dimensional case: the energy balance implies that temperature increases whenever density concentrates near a point. This in turn has a smoothing effect (through (2.1)) on the density profile, preventing blow-up from happening. Theorem 1.1 tells us that this general observation is not true in  $\mathbb{R}^3$ .

Problem **P** can also be derived for collisionless systems such as galaxies. The underlying argument is that rapid fluctuations of the gravitational field during the early stage of *violent relaxation* plays the same role as collisions, although the time scales involved for collisionless systems are smaller than for collisional systems (Brownian motion). The process of violent relaxation is considered in [14].

## 2.2 Non-dimensionalization

We put equations (2.1)-(2.5) in dimensionless form by setting

$$\tilde{x} = \frac{1}{R} x, \quad \tilde{n} = \frac{R^3}{M} n, \quad \tilde{\phi} = \frac{R}{4\pi GM} \left( \phi + \frac{GM}{R} \right)$$

and

$$\tilde{\Theta} = \frac{kR}{4\pi GM} \Theta, \quad \tilde{t} = \frac{4\pi GM}{\beta R^3} t.$$

Introducing  $\tilde{E} = \frac{R}{2\pi GM^2} \left( E + \frac{1}{2} \frac{GM^2}{R} \right)$  and  $\tilde{\kappa} = \frac{1}{k} \kappa$ , and dropping the tildes, results in Problem **P**.

## 2.3 Lyapunov functional and stationary solutions

If a triple  $\langle n, \phi, \Theta \rangle$  solves Problem **P**, then it is easy to check that

$$W(t) = \int_{\Omega} n \log n dx - \frac{\kappa}{2} \log \left( E - \int_{\Omega} n \phi dx \right) \quad \text{on } \mathbb{R}^+ \quad (2.7)$$

satisfies

$$\frac{d}{dt}W(t) = - \int_{\Omega} \frac{|\Theta(t)\nabla n + n\nabla\phi|^2}{\Theta(t)n} dx, \quad \text{for all } t > 0. \quad (2.8)$$

Hence  $W$  is a Lyapunov functional for Problem **P**, sometimes called the Boltzmann entropy [17].

One consequence of (2.8) is the following. Let  $\langle n_s, \phi_s, \Theta_s \rangle$  denote a stationary solution of Problem **P**. Then (2.8) implies

$$\Theta_s \nabla n_s + n_s \nabla \phi_s \equiv 0 \quad \text{in } \Omega.$$

Introducing the scaled potential  $\psi := \frac{\phi_s}{\Theta_s}$  we observe that  $n_s \equiv \frac{e^{-\psi}}{\int_{\Omega} e^{-\psi} dx}$ , where  $\psi$  satisfies

$$(S) \quad \begin{cases} \Delta \psi &= \frac{1}{\Theta_s} \frac{e^{-\psi}}{\int_{\Omega} e^{-\psi} dx} & \text{in } \Omega, \\ \psi &= 0 & \text{on } \partial\Omega. \end{cases}$$

The corresponding energy relation takes the form

$$E \frac{1}{(\Theta_s)^2} = \kappa \frac{1}{\Theta_s} - \int_{\Omega} |\nabla \psi|^2 dx. \quad (2.9)$$

Problem **S** has only one singular radially symmetric solution [12], the Chandrasekhar solution

$$n_s = U := \frac{1}{4\pi} \frac{1}{|x|^2}, \quad (2.10)$$

provided  $\Theta_s = \frac{1}{8\pi}$ . It satisfies (2.9) for  $E = \frac{(\kappa-2)}{8\pi}$ . Observe that  $U \in L^{3/2}(\Omega) \setminus L^q(\Omega)$ , with  $q > \frac{3}{2}$ . If this solution is attained by Problem **P** for  $t \uparrow T^* < \infty$ , we have a blow-up without concentration of mass at the origin.

For completeness we recall a result [5, Proposition 5.6.] for bounded radially symmetric solutions of Problem **S** and (2.9).

**Theorem 2.1** *Let  $\Omega = B_1(0)$ . For any  $\kappa > 0$ , there exists  $E_{\kappa} \in \mathbb{R}$  such that:*

- (i) *If  $E > E_{\kappa}$  there exist bounded negative solutions;*
- (ii) *If  $E < E_{\kappa}$  there are no nontrivial bounded negative solutions.*

This observation is originally due to Antonov [1] as a result of a computational approach. He also showed that stationary solutions are local maxima or saddle points of an entropy and there is no global entropy maximum.

Theorem 2.1 is still open for general domains [5]. This is related to the non-trivial nature of the set of singular solutions [12].

#### 2.4 Radially symmetric solutions

Our main theorem about global existence is stated in terms of radially symmetric solutions. Radial symmetry in Problem **P** not only reduces the spatial dimension, it also allows us to combine equations (1.1) and (1.2) into a single equation for the accumulated mass

$$Q(r, t) := \int_{B_r(0)} n(x, t) dx \quad \text{for } r \in (0, 1] \quad \text{and } t \in \mathbb{R}^+.$$

This is shown in [7]. Redefining  $t := \frac{3}{4\pi}t$  and  $\Theta := 12\pi\Theta$ , we obtain in terms of  $Q(y, t) := Q(r, t)$ , with  $y = r^3$ , the equation

$$Q_t = y^{4/3}\Theta(t)Q_{yy} + QQ_y \quad \text{for } y \in (0, 1) \quad \text{and } t \in \mathbb{R}^+. \quad (2.11)$$

To transform the energy relation (1.3), we first note that (1.2) and (1.5) give  $\int n\phi dx = -\int |\nabla\phi|^2 dx$ . Further, radial symmetry and (1.2) imply  $4\pi r^2 \partial_r \phi = Q(r, t)$ . Finally we introduce  $E := 12\pi E$ , to get in terms of  $Q(y, t)$

$$E = \kappa\Theta(t) - \int_0^1 \frac{Q^2}{y^{4/3}} dy \quad \text{for } t \in \mathbb{R}^+. \quad (2.12)$$

The boundary conditions for  $Q$  are

$$Q(0, t) = 0, \quad Q(1, t) = 1, \quad \text{for } t \in \mathbb{R}^+, \quad (2.13)$$

and the initial condition becomes

$$Q(y, 0) = Q_0(y) := \frac{4\pi}{3} \int_0^y n_0(y^{1/3}) dy \quad \text{for } 0 \leq y \leq 1. \quad (2.14)$$

Equations (2.11)-(2.14) define Problem **Q**.

Note that

$$\Theta(t) = \text{constant} = \frac{3}{2} \quad \text{and} \quad Q(y, t) = y^{1/3}$$

satisfy equation (2.11) and boundary conditions (2.13). The energy relation (2.12) is satisfied for  $E = \frac{3}{2}(\kappa - 2)$ . This is the transformed Chandrasekhar solution (2.10).

### 3. WELL-POSEDNESS FOR PROBLEM **P**

Before we give a formal solution definition for Problem **P** we observe that  $\phi$  is known in terms of  $n$  by the boundary value problem (1.2) and (1.5). Therefore we denote a solution by  $\langle n, \Theta \rangle$  instead of the triple  $\langle n, \phi, \Theta \rangle$ .

We call  $\langle n, \Theta \rangle$  a weak solution of Problem **P** if for some  $T > 0$ :

- (i)  $n \in L^2(0, T; H^1(\Omega))$  and  $n_t \in L^2(0, T; (H^1(\Omega))')$ ;
- (ii)  $\Theta \in C([0, T])$  and  $\Theta(t) > 0$  for  $t \in [0, T]$ ;
- (iii) the triple  $\langle n, \phi, \Theta \rangle$ , where  $\phi \in C([0, T]; H_0^1(\Omega))$  solves the boundary value problem (1.2) and (1.5), satisfies (1.1) in the weak sense and (1.3) for all  $t \in [0, T]$ ;
- (iv)  $n(\cdot, 0) = n_0 \geq 0$  a.e in  $\Omega$ .

**Remark 3.1** *The regularity in (i) implies  $n \in C([0, T]; L^2(\Omega))$  [20, p. 260]. Therefore  $\Theta$  and  $\phi$  are continuous in time in the sense of (ii) and (iii) respectively and the initial value of  $n$  can be prescribed.*

#### 3.1 Local existence

Let  $R_T := \Omega \times (0, T]$  for arbitrarily chosen  $T > 0$ .

The first result asserts local existence for Problem **P**.

**Theorem 3.2** *Let  $E \in \mathbb{R}$ ,  $\kappa > 0$ , and let  $n_0 \in L^2(\Omega)$  be such that  $\Theta(0) = \Theta_0 > 0$ . Then there exists a weak solution  $\langle n, \Theta \rangle$  of Problem **P** with  $T = T(\|n_0\|_{L^2(\Omega)}, \Omega, \Theta_0) > 0$ . It satisfies  $n \geq 0$  in  $R_T$  and  $n \in L_{loc}^\infty((0, T]; L^\infty(\Omega))$ .*

**Proof:** The proof uses a Schauder fixed-point theorem [21, Corollary 9.7]. For any fixed  $T > 0$ , let

$$X = \{v \in L^2(0, T; H^1(\Omega)) \quad \text{with} \quad v_t \in L^2(0, T; (H^1(\Omega))')\}$$

and let  $F: X \rightarrow C([0, T])$  be defined by

$$F(v)(t) = \frac{1}{\kappa} [\|v(t)\|_{H^{-1}(\Omega)}^2 + E] \quad \text{for any } t \in [0, T], \quad (3.1)$$



and for all  $v \in X$ . This map is clearly well-defined: observe that  $v$  and  $v_t$  belong to  $L^2(0, T; H^{-1}(\Omega))$ . Note that  $F(n)(t)$  is the temperature  $\Theta(t)$  whenever  $n$  is the solution of Problem **P**.

Next define  $N: X \rightarrow X$ , with  $u = N(v)$  satisfying

$$\left. \begin{array}{l} u_t = \operatorname{div}(F(v)(t)\nabla u + u\nabla\phi) \\ \Delta\phi = u \end{array} \right\} \quad \text{in} \quad R_T \quad (3.2)$$

$$\left. \begin{array}{l} \phi = 0 \\ (F(v)(t)\nabla u + u) \cdot \vec{\nu} = 0 \end{array} \right\} \quad \text{on} \quad \partial\Omega \times [0, T], \quad (3.3)$$

$$u(x, 0) = n_0(x) \quad \text{for} \quad x \in \Omega. \quad (3.4)$$

For given  $v \in X$ , this problem is essentially Problem **P** with prescribed temperature. As we point out in Remark 4.1, we have local existence and uniqueness provided  $F$  remains positively bounded from below. Under this condition the operator  $N$  is well-defined.

To apply the fixed point theorem, we need to prove that there exists  $\mathcal{C} \subset X$ , with  $\mathcal{C}$  convex, bounded and closed in  $(X, \|\cdot\|)$ , such that:

- (i)  $N(\mathcal{C}) \subset \mathcal{C}$ ;
- (ii)  $N$  is weakly-weakly sequentially continuous in  $X$ .

For any  $v \in \mathcal{C}$ , the operator  $N$  has to be well defined. Thus in addition to (i) and (ii) we need

- (iii) there exists  $F_0 = F_0(\mathcal{C})$  such that  $F(v)(t) \geq F_0 > 0$  for all  $t \in [0, T]$  and for all  $v \in \mathcal{C}$ .

We show below that

$$\mathcal{C} = \left\{ v \in X \mid v(0) = n_0, \quad \|v\|_{L^2(0, T; L^2(\Omega))} \leq RT^{1/2}, \quad \|\nabla v\|_{L^2(0, T; L^2(\Omega))} \leq R', \right. \\ \left. \text{and} \quad \|v_t\|_{L^2(0, T; (H^1(\Omega))')} \leq R'' \right\},$$

for suitably chosen constants  $R, R', R''$  and for  $T$  sufficiently small. In fact  $R = 2\|n_0\|_{L^2(\Omega)}$ ,  $R' = 2\|n_0\|_{L^2(\Omega)}/\Theta_0^{1/2}$  and  $R'' = 2\|n_0\|_{L^2(\Omega)}\Theta_0^{1/2} + 4C\|n_0\|_{L^2(\Omega)}^2/\Theta_0^{1/2}$ , where  $C = C(\Omega)$  is a positive constant. Clearly  $\mathcal{C}$  is convex, bounded, and closed in  $X$ . Note that  $\mathcal{C}$  is not empty: the solution of the heat equation with initial value  $n_0$  and diffusion coefficient  $\Theta_0/4$  satisfies  $\|\nabla n\|_{L^2(0, T; L^2(\Omega))} \leq \sqrt{2}\|n_0\|_{L^2(\Omega)}/\Theta_0^{1/2}$  and  $\|n_t\|_{L^2(0, T; (H^1(\Omega))')} \leq \frac{1}{2\sqrt{2}}\|n_0\|_{L^2(\Omega)}\Theta_0^{1/2}$ . Hence  $n \in \mathcal{C}$  for  $T > 0$ .

We first show (iii). Differentiating expression (3.1), applying Cauchy-Schwartz and the continuous injections  $(H^1(\Omega))' \hookrightarrow H^{-1}(\Omega)$  and  $L^2(\Omega) \hookrightarrow H^{-1}(\Omega)$ , yields the estimate

$$\kappa|F(v)_t(t)| \leq 2\|v(t)\|_{H^{-1}(\Omega)}\|v_t(t)\|_{H^{-1}(\Omega)} \leq C\|v(t)\|_{L^2(\Omega)}\|v_t(t)\|_{(H^1(\Omega))'} \quad \text{a.e. in} \quad [0, T], \quad (3.5)$$

where  $C = C(\Omega) > 0$ . Integration now gives

$$|F(v)(t) - F(n_0)| \leq \frac{C}{\kappa}R'RT^{1/2} \quad \text{for all} \quad t \in [0, T]. \quad (3.6)$$

Hence  $F(v)(t) \geq F(n_0) - \frac{C}{\kappa}R'RT^{1/2} = \Theta_0 - \frac{C}{\kappa}R'RT^{1/2}$  for  $0 < t \leq T$ . If we now choose  $F_0 = \Theta_0/2$  and  $T^*$  such that  $\frac{C}{\kappa}R'RT^{1/2} = \Theta_0/2$ , we have established (iii) for all  $0 < t \leq T \leq T^*$ .

Next we verify (i) for a suitable  $T \leq T^*$ . Starting point is inequality (4.3) with  $\Theta = F(v)(t)$  and  $v \in \mathcal{C}$ . It follows that the solution of (3.2)-(3.4) satisfies

$$\frac{1}{2} \frac{d}{dt} \|u\|_{L^2(\Omega)}^2 + F(v)(t)\|\nabla u\|_{L^2(\Omega)}^2 \leq \|u\|_{L^3(\Omega)}\|\nabla\phi\|_{L^6(\Omega)}\|\nabla n\|_{L^2(\Omega)}.$$

Since (4.4) holds for any  $\Theta > 0$ , we use it with  $\Theta = F_0$  to find

$$\frac{1}{2} \frac{d}{dt} (\|u(t)\|_{L^2(\Omega)}^2) + (F(v)(t) - \frac{F_0}{2})\|\nabla u\|_{L^2(\Omega)}^2 \leq \frac{C}{F_0^3} (\|u(t)\|_{L^2(\Omega)}^2)^3 + \frac{F_0}{2}\|u(t)\|_{L^2(\Omega)}^2 \quad (3.7)$$

for  $0 \leq t \leq T$  and for some  $C = C(\Omega) > 0$ .

Since  $v \in \mathcal{C}$  and consequently  $F(v)(t) \geq F_0$ , we obtain

$$\frac{1}{2} \frac{d}{dt} (\|u\|_{L^2(\Omega)}^2) + \frac{F_0}{2} \|\nabla u\|_{L^2(\Omega)}^2 \leq \frac{C}{F_0^3} (\|u(t)\|_{L^2(\Omega)}^2)^3 + \frac{F_0}{2} \|u(t)\|_{L^2(\Omega)}^2 \quad \text{in } [0, T]. \quad (3.8)$$

This inequality implies some useful bounds. Disregarding the gradient in the left-hand side of (3.8) gives a differential inequality in terms of  $\|u(t)\|_{L^2(\Omega)}^2$ . It follows that there exists  $T_0 = T_0(\Theta_0, \Omega, \|n_0\|_{L^2(\Omega)})$  such that  $u$  is well defined in  $R_{T_0}$  and satisfies

$$\sup_{t \in [0, T_0]} \|u(t)\|_{L^2(\Omega)}^2 \leq (2\|n_0\|_{L^2(\Omega)})^2, \quad \text{and thus} \quad \|u\|_{L^2(0, T_0; L^2(\Omega))} \leq 2\|n_0\|_{L^2(\Omega)} T_0^{1/2}. \quad (3.9)$$

Integrating (3.8) and using (3.9) gives

$$\|\nabla u\|_{L^2(0, T_0; L^2(\Omega))} \leq C T_0^{1/2} + \|n_0\|_{L^2(\Omega)} / (F_0)^{1/2} = C T_0^{1/2} + \sqrt{2} \|n_0\|_{L^2(\Omega)} / \Theta_0^{1/2}. \quad (3.10)$$

for a positive constant  $C = C(\Theta_0, \Omega, \|n_0\|_{L^2(\Omega)})$ . Note that (3.9) and (3.10) imply

$$u \in L^\infty(0, T_0; L^2(\Omega)) \quad \text{and} \quad u \in L^2(0, T_0; H^1(\Omega)).$$

To show that  $u \in \mathcal{C}$ , for sufficiently small  $T$ , it remains to prove the bound on  $u_t$ . With  $\xi \in L^2(0, T; H^1(\Omega))$ , we have from (3.2)

$$\int_0^T \langle u_t, \xi \rangle dt = - \int_0^T F(v)(t) \int_\Omega \nabla u \nabla \xi dx dt + \int_0^T \int_\Omega u \nabla \phi \nabla \xi dx dt, \quad (3.11)$$

where  $\langle \cdot, \cdot \rangle$  denotes the pairing between  $(H^1(\Omega))'$ , and  $H^1(\Omega)$ . To estimate the right hand side we first note that  $F(v) \in L^\infty(0, T^*)$ . Indeed, from (3.6) we deduce

$$F_0 < F(v)(t) \leq \frac{C}{\kappa} R R' T^{1/2} + \Theta_0, \quad \text{for } 0 \leq t \leq T \leq T^*. \quad (3.12)$$

Next we use (6.3) and interpolation inequality (6.2) from the appendix. This gives

$$\left| \int_\Omega u \nabla \phi \nabla \xi dx \right| \leq \|u\|_{L^3(\Omega)} \|\nabla \phi\|_{L^6(\Omega)} \|\nabla \xi\|_{L^2(\Omega)} \leq C_s^{1/2} C_I \|u\|_{H^1(\Omega)} \|u\|_{L^2(\Omega)} \|\nabla \xi\|_{L^2(\Omega)}.$$

Finally we combine this expression with (3.9), (3.10), and (3.12), and obtain after some manipulation

$$\int_0^T |\langle u_t, \xi \rangle| dt \leq \{C(T^{1/2} + T) + \sqrt{2} \|n_0\|_{L^2(\Omega)} \Theta_0^{1/2} + 2\sqrt{2} C_s^{1/2} C_I \|n_0\|_{L^2}^2 / \Theta_0^{1/2}\} \|\xi\|_{L^2(0, T, H^1(\Omega))}.$$

for some  $C = C(\Omega, \|n_0\|_{L^2(\Omega)}, \Theta_0)$ . Taking now  $T_1 < T_0 < T^*$  sufficiently small we obtain that  $u \in \mathcal{C}$  for  $0 \leq T \leq T_1$  and consequently  $N(\mathcal{C}) \subset \mathcal{C}$ .

Next we show (ii): i.e. we claim that  $v_k \in \mathcal{C}$ ,  $v_k \rightharpoonup v$  in  $X$  implies  $N(v_k) \rightharpoonup N(v)$  in  $X$ . For any such sequence  $v_k$ , define  $u_k := N(v_k) \in \mathcal{C}$ . Using the weak compactness of  $\mathcal{C}$  we extract a subsequence  $u_{k'} \in \mathcal{C}$  such that  $u_{k'} \rightharpoonup u^*$  in  $X$ . We show below that  $u^* = N(v)$ , which proves the assertion. Since

$$v_{k'} \rightharpoonup v \quad \text{and} \quad u_{k'} \rightharpoonup u^* \quad \text{in } \mathcal{C},$$

we obtain by Aubin's Lemma [16, pag. 58] for a subsequence, denoted again by  $k'$ ,

$$v_{k'} \rightarrow v \quad \text{and} \quad u_{k'} \rightarrow u^* \quad \text{in } L^2(0, T; L^2(\Omega)).$$

We use this in (3.11) for  $u_{k'}, v_{k'}$ , and  $\phi_{k'}$ . Since  $\Delta\phi_{k'} = u_{k'}$ , we have  $\phi_{k'} \rightarrow \phi^*$  in  $L^2(0, T; H^2(\Omega))$  satisfying  $\Delta\phi^* = u^*$ . Moreover, as  $k' \rightarrow \infty$ ,

$$\begin{aligned} u_{k'_t} &\rightharpoonup u_t^* && \text{in } L^2(0, T, (H^1(\Omega))'), \\ \nabla\phi_{k'} &\rightarrow \nabla\phi^* && \text{in } L^2(0, T, H^1(\Omega)), \\ \nabla u_{k'} &\rightharpoonup \nabla u^* && \text{in } L^2(0, T; L^2(\Omega)). \end{aligned}$$

Now suppose  $F(v_{k'}) \rightarrow F(v)$  in  $C([0, T])$ . Then letting  $k' \rightarrow \infty$  in (3.11) we obtain a solution  $u^*$  of problem (3.2)-(3.4) for the temperature  $F(v)(t)$ . By uniqueness we have  $u^* = N(v)$ .

It remains to show that  $F(v_{k'}) \rightarrow F(v)$  in  $C([0, T])$ . In view of the continuous injection  $L^2(\Omega) \hookrightarrow H^{-1}(\Omega)$ , we find from (3.1)

$$\kappa|F(v_{k'})(t) - F(v)(t)| \leq C(\Omega) [\|v_{k'}(t)\|_{L^2(\Omega)} + \|v(t)\|_{L^2(\Omega)}] \|v_{k'}(t) - v(t)\|_{L^2(\Omega)}.$$

This implies directly  $F(v_{k'}) \rightarrow F(v)$  in  $L^1([0, T])$ . Writing (3.5) for the difference  $F(v_{k'}) - F(v)$ , using the continuous injection  $(H^1(\Omega))' \hookrightarrow H^{-1}(\Omega)$ , and integrating the result gives

$$\begin{aligned} \kappa \int_0^T |F(v_{k'})_t(t) - F(v)_t(t)| dt &\leq \int_0^T |(v(t), v_{k'_t}(t) - v_t(t))_{H^{-1}}| dt \\ &+ \|v - v_{k'}\|_{L^2(0, T; L^2(\Omega))} \|(v_{k'})_t\|_{L^2(0, T; (H^1(\Omega))')}. \end{aligned}$$

Since  $v_{k'_t} \rightharpoonup v_t$ , in  $L^2(0, T, H^{-1}(\Omega))$ , we obtain  $F(v_{k'}) \rightarrow F(v)$  in  $W^{1,1}([0, T])$ . This concludes the proof of (ii) and establishes local existence for Problem **P**.

The boundedness of  $n$  follows from [6, Theorem 2] and  $n \geq 0$  a.e. in  $R_T$  is essentially demonstrated in [11].

**Remark 3.3** *Let  $n_0 \in L^p(\Omega)$  with  $p > 3$  and let  $\Theta(0) = \Theta_0 > 0$ . Then Problem **P** has a local solution satisfying  $n \in L^\infty(0, T; L^p(\Omega))$  and  $n^{p/2} \in L^2(0, T; H^1(\Omega))$ . The proof is almost identical to the proof of Theorem 3.2.*

### 3.2 Uniqueness

Uniqueness is stated for an equivalent formulation of Problem **P** in which we replace  $t$  by  $\tau = \int_0^t \Theta(t) dt$ . This transformation only affects equation (1.1), which now becomes

$$n_\tau = \operatorname{div} \left\{ \nabla n + \frac{n}{\Theta(\tau)} \nabla \phi \right\} \quad \text{in } R_{\hat{T}}. \quad (3.13)$$

for  $\hat{T} = \int_0^T \Theta(t) dt$ . The problem stated in terms of  $x$  and  $\tau$  is denoted by Problem **P<sub>e</sub>**. Without proof we remark that  $\langle n = n(x, t), \Theta = \Theta(t) \rangle$  solves Problem **P** if and only if  $\langle n = n(x, \tau), \Theta = \Theta(\tau) \rangle$  solves problem Problem **P<sub>e</sub>**. This is due to the strict positivity of  $\Theta$  in the existence interval.

**Theorem 3.4** *If  $n_0 \in L^2(\Omega)$  and  $\Theta_0 > 0$ , then Problem **P<sub>e</sub>** has at most one solution  $\langle n, \Theta \rangle$ .*

**Proof:** We use a uniqueness result of Biler & Nadzieja [11], who considered the problem

$$n_\tau = \operatorname{div}(\nabla n + nX(n)) \quad \text{in } R_T, \quad (3.14)$$

$$(\nabla n + nX(n)) \cdot \vec{\nu} = 0 \quad \text{on } \partial\Omega \times [0, T], \quad (3.15)$$

$$n(\cdot, 0) = n_0 \quad \text{in } \Omega, \quad (3.16)$$

where  $X$  is a general non-local vector field operator in  $\mathbb{R}^3$ . For this problem uniqueness in  $L^2(\Omega)$  was proved in [11, Theorem 1 (i)] under the following condition: there exists  $C > 0$  such that

$$(U) \quad \|X(u) - X(v)\|_{L^6(\Omega)} \leq C\|u - v\|_{L^2(\Omega)}$$

for all  $u, v \in L^2(\Omega)$ .

Note that the constant  $C$  in (U) does not depend on the choice of  $u, v \in L^2(\Omega)$ . In our case  $X(n) = \frac{\nabla\phi}{\Theta(\tau)}$ . Below, in Lemma 3.5, we show that again (U) holds but with  $C$  depending on both norms  $\|u\|_{L^2(\Omega)}$  and  $\|v\|_{L^2(\Omega)}$ . Now suppose that Problem  $\mathbf{P}_e$  admits two solutions  $\langle n_1, \Theta_1 \rangle$  and  $\langle n_2, \Theta_2 \rangle$  in some interval  $[0, T]$ . From the solution definition we know that both  $\|n_1(t)\|_{L^2(\Omega)}$  and  $\|n_2(t)\|_{L^2(\Omega)}$  are uniformly bounded in  $[0, T]$ . Therefore (U) is satisfied for the two solutions  $n_1(t)$  and  $n_2(t)$ , with  $0 \leq t \leq T$ , for an appropriately chosen constant  $C$ . As a consequence we can apply the result of [11]. This proves the theorem.

**Lemma 3.5** *Suppose there exist  $\delta > 0$  and  $u, v \in L^2(\Omega)$  such that*

$$\min \left\{ \Theta_u := E + \int_{\Omega} |\nabla\phi_u|^2 dx, \Theta_v := E + \int_{\Omega} |\nabla\phi_v|^2 dx \right\} \geq \delta > 0,$$

where

$$\Delta\phi_u = u, \quad \Delta\phi_v = v \quad \text{in} \quad \Omega, \quad (3.17)$$

$$\phi_u = \phi_v = 0 \quad \text{on} \quad \partial\Omega. \quad (3.18)$$

Then

$$\left\| \nabla \left( \frac{\phi_u}{\Theta_u} - \frac{\phi_v}{\Theta_v} \right) \right\|_{L^6(\Omega)} \leq C \|u - v\|_{L^2(\Omega)}$$

where  $C = C(\delta, \|u\|_{L^2(\Omega)}, \|v\|_{L^2(\Omega)})$ .

**Proof:** Using

$$\|\nabla(\phi_u - \phi_v)\|_{L^6(\Omega)} \leq C_I \|u - v\|_{L^2(\Omega)}$$

we estimate

$$\begin{aligned} \left\| \nabla \left( \frac{\phi_u}{\Theta_u} - \frac{\phi_v}{\Theta_v} \right) \right\|_{L^6(\Omega)} &= \left\| \nabla \left( \frac{\phi_u}{\Theta_u} - \frac{\phi_u}{\Theta_v} + \frac{\phi_u}{\Theta_v} - \frac{\phi_v}{\Theta_v} \right) \right\|_{L^6(\Omega)} \\ &\leq \frac{1}{\Theta_u \Theta_v} \|\nabla\phi_u\|_{L^6(\Omega)} |\Theta_u - \Theta_v| + \frac{C_I}{\Theta_v} \|u - v\|_{L^2(\Omega)} \\ &\leq \frac{C_I}{\Theta_v} \left\{ \frac{\|u\|_{L^2(\Omega)}}{\Theta_u} \left| \int_{\Omega} (|\nabla\phi_u|^2 - |\nabla\phi_v|^2) dx \right| + \|u - v\|_{L^2(\Omega)} \right\}. \end{aligned}$$

Since

$$\begin{aligned} \left| \int_{\Omega} (|\nabla\phi_u|^2 - |\nabla\phi_v|^2) dx \right| &\leq \|\nabla(\phi_u + \phi_v)\|_{L^2(\Omega)} \|\nabla(\phi_u - \phi_v)\|_{L^2(\Omega)}, \\ &\leq C(\|u\|_{L^2(\Omega)} + \|v\|_{L^2(\Omega)}) \|u - v\|_{L^2(\Omega)} \end{aligned}$$

for some  $C > 0$ , we obtain the assertion.  $\blacksquare$

**Remark 3.6** *Note that for  $n_0 \in L^p(\Omega)$  with  $p > 2$ , we can apply the above theorem and obtain uniqueness of solutions for such initial data. Thus the local solution given in Remark 3.3 of Theorem 3.2 for  $n_0 \in L^p(\Omega)$  with  $p > 3$  is unique.*

**Remark 3.7** *A slight modification of the above argument directly gives uniqueness for  $n_0 \in L^p(\Omega)$  with  $p > 3$  and  $\Theta_0 > 0$ . Again following [11, Theorem 1 (ii)], we need to show*

$$(U') \quad \|X(u) - X(v)\|_{L^\infty(\Omega)} \leq C \|u - v\|_{L^p(\Omega)} \quad (p > 3).$$

With  $X(n) = \frac{\nabla\phi}{\Theta(\tau)}$ , inequality (U') results from inequality (6.3).

The next theorem extends the local existence result for  $n_0 \in L^p(\Omega)$  with  $p > \frac{3}{2}$ . To do this we modify our definition of weak solution for  $1 < p < 2$ . We replace (i) by  $n \in L^\infty(0, T; L^p(\Omega))$  and  $n_t \in L^p(0, T; (W^{1,p'})')$  with  $\frac{1}{p} + \frac{1}{p'} = 1$ . We show that  $n \in C([0, T]; L^p(\Omega))$ , implying that  $n_0$  can be prescribed. In fact as  $n \in L^\infty(0, T; L^p(\Omega))$  and  $n \in C([0, T], (W^{1,p'}(\Omega))')$ , (since  $n \in L^p(0, T; (W^{1,p'}(\Omega))')$  and  $n_t \in L^p(0, T; (W^{1,p'}(\Omega))')$ ), we use the injection  $L^p(\Omega) \hookrightarrow (W^{1,p'}(\Omega))'$ , and follow the argument in [16, p. 23] to conclude.

**Theorem 3.8** *Let  $n_0 \in L^p(\Omega)$  with  $p > 3/2$ , and  $\Theta(0) = \Theta_0 > 0$ . Then there exists  $T = T(\Omega, \|n_0\|_{L^p(\Omega)}, \Theta_0) > 0$ , and a weak solution  $\langle n, \Theta \rangle$  of Problem **P**. It satisfies  $n^{p/2} \in L^2(0, T; H^1(\Omega))$  and furthermore  $n \in L_{loc}^\infty(0, T]; L^\infty(\Omega)$ . This solution is unique.*

**Proof:** Due to Remarks 3.3 and 3.6, we only need to demonstrate local existence for  $n_0 \in L^p(\Omega)$  with  $p \in (3/2, 3)$ . We follow the proof of [11, Theorem 1 (iii)] and approximate  $n_0 \in L^p(\Omega)$  by functions in  $L^{p^*}(\Omega)$  with  $p^* > 3$ . Thus let  $\{n_{0\epsilon}\} \subset L^{p^*}(\Omega)$  satisfy  $\|n_{0\epsilon} - n_0\|_{L^p(\Omega)} \rightarrow 0$  as  $\epsilon \rightarrow 0$ .

For each  $\epsilon > 0$  we consider Problem **P** with initial data  $\langle n_{0\epsilon}, \Theta_0 \rangle$ . By Remark 3.6 there exists a solution  $\langle n_\epsilon, \Theta_\epsilon \rangle$ , with  $\Theta_\epsilon \geq \Theta_0/2$ , in some interval  $[0, T_\epsilon]$ . Next we use the estimate [11, Eq. (10)]

$$\|n_\epsilon(t)\|_{L^p(\Omega)}^p + \int_0^t |\nabla |n_\epsilon|^{p/2}|^2 d\tau \leq \exp\left(C \int_0^t \left\| \frac{\nabla \phi_\epsilon}{\Theta_\epsilon} \right\|_{L^q(\Omega)}^{\frac{2q}{q-3}} d\tau\right) \|n_{0\epsilon}\|_{L^p(\Omega)}^p \quad (3.19)$$

for almost every  $t \in [0, T_\epsilon]$ . Here  $p \in (3/2, 3)$ ,  $1/q = 1/p - 1/3$  and  $C = C(\Omega, p)$ . Note that  $q > 3$ . Further, using  $\|\nabla \phi_\epsilon\|_{L^q(\Omega)} \leq C\|n_\epsilon\|_{L^p(\Omega)}$  and the uniform lower bound on  $\Theta_\epsilon$ , we obtain

$$\|n_\epsilon(t)\|_{L^p(\Omega)}^p \leq C \exp\left(\frac{2}{\Theta_0} \int_0^t \|n_\epsilon(\tau)\|_{L^p(\Omega)}^{\frac{2q}{q-3}} d\tau\right) \quad \text{for almost every } t \in [0, T_\epsilon],$$

where  $C = C(\bar{\epsilon}) > 0$  and  $0 < \epsilon \leq \bar{\epsilon}$ . Since  $C$  does not depend on  $\epsilon$  we have that  $T_\epsilon = T$  and

$$\|n_\epsilon(t)\|_{L^p(\Omega)} \leq C \quad \text{and} \quad \Theta_\epsilon(t) \geq \Theta_0/2 \quad \text{for almost every } t \in [0, T] \quad (3.20)$$

and for all  $0 < \epsilon \leq \bar{\epsilon}$ . Using this and (3.19), we deduce

$$\|n_\epsilon(t)^{p/2}\|_{H^1(\Omega)} \leq C \quad \text{and} \quad \Theta_\epsilon(t) \geq \Theta_0/2 \quad \text{for almost every } t \in [0, T]. \quad (3.21)$$

and for all  $0 < \epsilon \leq \bar{\epsilon}$ .

Next we separate the demonstration into two cases:  $p < 2$  and  $p > 2$ .

We begin with  $p < 2$ . Under this condition, we have

$$\|\nabla n\|_{L^p(\Omega)} \leq C(\Omega, p) \|\nabla n^{p/2}\|_{L^2(\Omega)} \|n\|_{L^p(\Omega)}^{(2-p)/2} \quad \text{for } n^{p/2} \in H^1(\Omega).$$

Combining this with (3.20) and (3.21), since  $p < 2$ , we obtain

$$\|\nabla n_\epsilon\|_{L^p(0, T; L^p(\Omega))} \leq C \quad \text{for all } 0 < \epsilon \leq \bar{\epsilon}. \quad (3.22)$$

Consequently, we can check that  $\|n_{\epsilon_t}\|_{L^p(0, T, (W^{1,p'}(\Omega))')}$  is bounded for all  $0 < \epsilon \leq \bar{\epsilon}$ . Now using a compactness theorem [16, p. 141], with  $L^p(\Omega) \hookrightarrow (W^{1,p'}(\Omega))'$ , we find for a subsequence  $\epsilon \rightarrow 0$ ,

$$n_\epsilon \rightarrow n \quad \text{in } L^p(0, T; L^p(\Omega)). \quad (3.23)$$

Now using standard arguments and above estimates, we get as  $\epsilon \rightarrow 0$

$$n_{\epsilon_t} \rightharpoonup n_t \quad \text{in } L^p(0, T, (W^{1,p'}(\Omega))'), \quad (3.24)$$

$$\nabla \phi_\epsilon \rightarrow \nabla \phi \quad \text{in } L^p(0, T, W^{1,p}(\Omega)), \quad (3.25)$$

$$\nabla n_\epsilon \rightharpoonup \nabla n \quad \text{in } L^p(0, T; L^p(\Omega)). \quad (3.26)$$

To conclude, it suffices to prove  $\Theta_\epsilon \rightarrow \Theta$  in  $C([0, T])$ . This follows from showing that

$$\kappa \int_0^T |\Theta_{\epsilon_t} - \Theta_t| dt \leq 2 \int_0^T \left| \int_\Omega n_\epsilon \phi_{\epsilon_t} - n \phi_t dx \right| dt \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0. \quad (3.27)$$

We obtain this using [21, Proposition 23.9 (d)], combining (i) strong convergence of  $n_\epsilon \rightarrow n$  in  $L^{p'}(0, T; L^p(\Omega))$  with (ii) weak convergence of  $\phi_{\epsilon_t} \rightharpoonup \phi_t$  in  $L^p(0, T; L^{p'}(\Omega))$ . In fact (i) is consequence of (3.20) and (3.23); and (ii) yields using (3.24) and the estimate  $\|\phi_{\epsilon_t}\|_{p'} \leq C \|\phi_{\epsilon_t}\|_{W^{1,p}(\Omega)} \leq C \|n_{\epsilon_t}\|_{(W^{1,p'}(\Omega))'}$ , where we have used  $p > \frac{3}{2}$ .

Now we take the limit  $\epsilon \rightarrow 0$  to conclude that  $n$  satisfies Problem **P**.

For  $p > 2$ , we use that  $n_{0_\epsilon} \in L^2(\Omega)$  and in particular (3.20) implies

$$\|n_\epsilon(t)\|_{L^2(\Omega)} \leq C \quad \text{and} \quad \Theta_\epsilon(t) \geq \Theta_0/2 \quad \text{for all } t \in [0, T].$$

We follow the proof of Theorem 3.2, to find  $\|n_{\epsilon_t}\|_{L^2(0, T, (H^1(\Omega))')} \leq C$ . With this we may apply again the compactness theorem [16, p. 141], now with  $L^p(\Omega) \hookrightarrow (H^1(\Omega))'$ , since  $p > 2$ , and obtain  $n_\epsilon \rightarrow n$  in  $L^p(0, T, L^p(\Omega))$ . Finally, we show (3.27) using  $p = p' = 2$  and obtain  $\Theta_\epsilon \rightarrow \Theta$  in  $C([0, T])$ , which concludes the proof of the theorem. ■

### 3.3 Radially symmetric solutions

In Section 2.4 we introduced Problem **Q** describing radially symmetric solutions of Problem **P** in the unit ball. In this paper we do not prove existence for Problem **Q**. Instead we shall assume that if  $\Omega = B_1(0)$  and if  $n_0$  is radially symmetric, then the corresponding weak solution is radially symmetric. By standard regularity theory weak solutions of Problem **P** satisfy equations (1.1)-(1.3) and boundary conditions (1.4)-(1.5) in a classical sense. With this in mind we introduce for Problem **Q** the following solution definition.

Let  $D_T = (0, 1) \times (0, T]$ . A pair  $\langle Q, \Theta \rangle$  solves Problem **Q**, if for some  $T > 0$ :

- (i)  $Q \in C^{2,1}(D_T) \cap C(\overline{D_T})$ , and  $\Theta \in C([0, T])$ ;
- (ii)  $\langle Q, \Theta \rangle$  satisfies equations (2.11)-(2.14);
- (iii)  $Q_y \geq 0$  in  $D_T$  and  $\Theta > 0$  in  $[0, T]$ .

Clearly radial solutions of Problem **P** with  $n_0 \in L^p(B_1(0))$ ,  $p > \frac{3}{2}$ , satisfy this definition. This follows directly from the identity

$$Q_y(y, t) = \frac{4\pi}{3} n(y^{1/3}, t) \quad \text{for } (y, t) \in D_T. \quad (3.28)$$

## 4. PRESCRIBED TEMPERATURE PROBLEM

In this section we study Problem **P** with prescribed temperature  $\Theta(t)$  satisfying

$$\Theta: [0, T] \rightarrow \mathbb{R} \quad \text{such that} \quad \Theta \in C([0, T]) \quad \text{and} \quad \Theta(t) > \delta > 0 \quad \text{for } t \in [0, T]. \quad (4.1)$$

Thus we drop the energy relation (1.3) and assume that  $\Theta$  in (1.1) and (1.4) is given and satisfies (4.1). We denote this modified problem by **P\***. Clearly, if Problem **P\*** has a radially symmetric solution, then the corresponding formulation in terms of the mass  $Q$ , which we denote by **Q\***, has a classical solution according to the definition given in Section 3.3.

We first recall some recent results of Biler & Nadzieja [3, 11], related to local existence for Problem **P\*** and global existence for Problem **Q\***.

### Remark 4.1

- (i) Let  $n_0 \in L^2(\Omega)$  and let  $\Theta$  satisfy (4.1). Then there exists  $T = T(\Omega, \|n_0\|_{L^2(\Omega)}, \delta) > 0$  so that Problem  $\mathbf{P}^*$  has a unique weak solution in  $[0, T]$  which satisfies  $n \in L_{loc}^\infty((0, T], L^\infty(\Omega))$ . Proof: see [11, Theorem 1 (i)].
- (ii) Let  $n_0 \in L^p(\Omega)$  with  $p > \frac{3}{2}$  and let  $\Theta$  satisfy (4.1). Then there exist  $T = T(\Omega, \|n_0\|_{L^p(\Omega)}, \delta) > 0$  so that Problem  $\mathbf{P}^*$  has a weak solution in  $[0, T]$  satisfying  $n \in L^\infty(0, T, L^p(\Omega))$  and  $n^{p/2} \in L^2(0, T, H^1(\Omega))$ . For  $p > 3$  the solution is unique. Proof: see [11, Theorem 1 (ii) and (iii)].
- (iii) If for some  $B > 0$ ,  $Q_0(y) \leq y^{-\frac{1+B}{y^{2/3}+B}}$  for  $0 \leq y \leq 1$  and  $\Theta(t) = \text{constant} \geq \frac{3}{2}(1+B)$  then Problem  $\mathbf{Q}^*$  has a global classical solution satisfying  $Q(y, \tau) \leq y^{-\frac{1+B}{y^{2/3}+B}}$  for  $(y, \tau) \in \overline{D_\infty}$ . Proof: see [3, Theorem 1 (iii)].

In the remainder of this section we present some new results related to Problem  $\mathbf{P}^*$  with constant temperature  $\Theta > 0$ . We first extend a global existence result of [9].

**Theorem 4.2** For a given domain  $\Omega$ , there exist positive constants,  $\alpha_1, \alpha_2, A, B$ , and  $C$  with  $\alpha_2 \geq \alpha_1$  so that if the constant temperature  $\Theta$  and the initial condition  $n_0$  satisfy

$$\Theta \geq \alpha_1 \quad \text{and} \quad \|n_0\|_{L^2(\Omega)}^2 \leq A + \frac{B}{\Theta^4}$$

or

$$\Theta \geq \alpha_2 \quad \text{and} \quad \|n_0\|_{L^2(\Omega)}^2 \leq C(\Theta^2 - \Theta),$$

then Problem  $\mathbf{P}^*$  has a global (weak) solution for which the  $L^2(\Omega)$  norm is uniformly bounded in time.

**Proof:** Integrating (1.1) we obtain the expression

$$\frac{1}{2} \frac{d}{dt} \|n\|_{L^2(\Omega)}^2 + \Theta \|\nabla n\|_{L^2(\Omega)}^2 = - \int_{\Omega} n \nabla n \nabla \phi \, dx. \quad (4.2)$$

As in the proof of [9, Theorem 2 (iii)] we estimate

$$\frac{1}{2} \frac{d}{dt} \|n\|_{L^2(\Omega)}^2 + \Theta \|\nabla n\|_{L^2(\Omega)}^2 \leq \|n\|_{L^3(\Omega)} \|\nabla \phi\|_{L^6(\Omega)} \|\nabla n\|_{L^2(\Omega)}. \quad (4.3)$$

The aim is to obtain a differential inequality for  $\|n(t)\|_{L^2(\Omega)}^2$ . From the appendix we first use (6.2) and then (6.3) with  $r = 6$  and  $p = 2$ . This gives

$$\|n\|_{L^3(\Omega)} \|\nabla \phi\|_{L^6(\Omega)} \|\nabla n\|_{L^2(\Omega)} \leq \frac{\Theta}{2} \|n\|_{H^1(\Omega)}^2 + \frac{C_1}{\Theta^3} (\|n\|_{L^2(\Omega)}^2)^3. \quad (4.4)$$

Further, we use (6.1) with  $q = 2$  and  $p = 1$  to obtain

$$2\Theta \|n\|_{L^2(\Omega)}^2 \leq \frac{\Theta}{2} \|n\|_{H^1(\Omega)}^2 + C_2 \Theta \|n\|_{L^1(\Omega)}^2. \quad (4.5)$$

The combination (4.3)-(4.5) eliminates the gradient term. Since  $\|n\|_{L^1(\Omega)} = 1$ , we are left with an inequality of the form

$$\frac{d}{dt} w \leq p_\Theta(w) := \frac{C_1}{\Theta^3} w^3 - \Theta w + C_2 \Theta \quad \text{for } t > 0 \quad (4.6)$$

with  $w(t) := \|n(t)\|_{L^2(\Omega)}^2$ . Here  $C_1$  and  $C_2$  are positive constants only depending on  $\Omega$ . The assertions of the theorem now follow from particular properties of (4.6).

First observe that if  $\Theta > \alpha_1 := \frac{3^{3/4}}{2^{1/2}} (C_2 C_1^{1/2})^{1/2}$ , then  $p_\Theta(w) = 0$  has two positive real roots  $w_* < w^*$  and  $p_\Theta(w) < 0$  for  $w_* < w < w^*$ . If  $\Theta = \alpha_1$ , these roots coincide. A simple calculation shows that

$$w_0 = C_2 + \frac{C_2^3 C_1}{\Theta^4}$$

satisfies  $0 < w_0 < w_*$  for all  $\Theta \geq \alpha_1$ . Since  $p_\Theta(w) > 0$  for  $0 \leq w < w_*$ , we deduce that  $w(t)$ , with  $w(0) \leq w_0$ , satisfies  $w(t) \leq w_*$  for all  $t \geq 0$ . This proves the first assertion.

Next consider

$$w^0 := C_1^{-1/2}(\Theta^2 - \Theta).$$

Then  $p_\Theta(w^0) \leq 0$  provided

$$\Theta \geq \alpha_2 = \left( \left( \frac{3 + C_2 C_1^{1/2}}{4} \right)^2 - \frac{1}{2} \right)^{1/2} + \frac{3 + C_2 C_1^{1/2}}{4}.$$

Clearly  $\alpha_2 \geq \alpha_1$ , since  $p_\Theta(w) > 0$  for  $\Theta < \alpha_1$  and for all  $w > 0$ . As before we have that  $w(t)$ , with  $w(0) \leq w^0$ , satisfies  $w(t) \leq w^0 \leq w^*$  for all  $t \geq 0$ . This proves the second assertion. ■

In a similar fashion global existence results are obtained in  $L^p(\Omega)$  for  $p > 3/2$ . Instead of (4.6) one now finds

$$\frac{d}{dt} w \leq \frac{C_1}{\Theta^\beta} w^\beta - \Theta w + C_2 \Theta \quad \text{for } t > 0 \quad (4.7)$$

with  $w(t) := \|n(t)\|_{L^p(\Omega)}^p$ . Here

$$\beta = \begin{cases} \frac{2p-1}{2p-3} & \text{for } 3/2 < p < 3 \\ \frac{p+2}{p} & \text{for } p > 3. \end{cases}$$

Inequality (4.7) implies the following result.

**Theorem 4.3** *For a given domain  $\Omega$ , there exist positive constants,  $\beta_1, \beta_2, \bar{A}, \bar{B}$ , and  $\bar{C}$  with  $\beta_2 \geq \beta_1$  so that if the constant temperature  $\Theta$  and the initial condition  $n_0$  satisfy*

$$\Theta \geq \beta_1 \quad \text{and} \quad \|n_0\|_{L^p(\Omega)}^p \leq \bar{A} + \frac{\bar{B}}{\Theta^{\beta+1}}$$

or

$$\Theta \geq \beta_2 \quad \text{and} \quad \|n_0\|_{L^p(\Omega)}^p \leq \bar{C}(\Theta^\gamma - \Theta), \quad \text{with } \gamma = \frac{\beta+1}{\beta-1},$$

then Problem **P\*** has a global (weak) solution for which the  $L^p(\Omega)$  norm is uniformly bounded in time.

## 5. GLOBAL EXISTENCE FOR PROBLEM **Q**

In this section we study radially symmetric solutions of Problem **P**. We will use the classical formulation in terms of Problem **Q**. Before we prove the global existence results, we first demonstrate the blow-up result Theorem 1.1.

**Theorem 5.1** *Let  $\kappa > 6$  and  $E < 3$ . Then  $T^* < \infty$ .*

**Proof:** Suppose Problem **Q** has a global solution  $Q = Q(y, t)$  and  $\Theta = \Theta(t) \in (0, \infty)$  for all  $t > 0$ . Setting

$$w_\epsilon(t) := \int_\epsilon^1 Q(y, t) y^{-1/3} dy \quad \text{for all } t > 0,$$

we find, after differentiating and using equation (2.11),

$$\begin{aligned} \frac{dw_\epsilon(t)}{dt} &= \Theta(t) Q_y(1, t) - \epsilon \Theta(t) Q_y(\epsilon, t) - \Theta(t) + \Theta(t) Q(\epsilon, t) \\ &- \frac{1}{2} \frac{Q(\epsilon, t)^2}{\epsilon^{1/3}} + \frac{1}{2} + \frac{\kappa}{6} \Theta(t) - \frac{E}{6} + \frac{1}{6} \int_0^\epsilon Q^2(y, t) y^{-4/3} dy \quad \text{for all } t > 0. \end{aligned} \quad (5.1)$$



Since  $\Theta(t) < \infty$ , we obtain from (2.12)

$$Q(y, t)^2/y^{4/3} \in L^1(0, 1) \quad \text{for all } t > 0,$$

which implies that we can choose a sequence  $\epsilon_n \downarrow 0$  along which  $\lim_{\epsilon_n \downarrow 0} \frac{Q(\epsilon_n, t)}{\epsilon_n^{1/6}} = 0$  for all  $t > 0$ . Using this and  $Q_y(1, t) \geq 0$  in (5.1), we find in the limit

$$\frac{dw_0(t)}{dt} \geq \frac{1}{2} - \Theta(t) + \frac{\kappa}{6}\Theta(t) - \frac{E}{6} \quad \text{for all } t > 0.$$

The parameter choice implies that  $\frac{dw_0(t)}{dt} \geq \delta > 0$  for all  $t > 0$ . This contradicts  $w_0 \leq \frac{3}{2}$  (implied by  $Q \leq 1$  in  $\overline{D_\infty}$ ). ■

**Remark 5.2** Using (3.28) we can express  $w_0(t)$  in terms of the second moment of the corresponding radially symmetric solution:

$$\int_{B_1(0)} n(x, t)|x|^2 dx = 1 - \frac{2}{3}w_0(t).$$

Since  $dw_0(t)/dt$  is bounded away from zero, there exists a  $T^* \in (0, \infty)$  (blow-up time) such that  $\lim_{t \uparrow T^*} w_0(t) = \frac{3}{2}$  and

$$\lim_{t \rightarrow T^*} \int_{B_1(0)} n(x, t)|x|^2 dx = 0.$$

This implies the gravitational collapse  $n(x, T^*) = \delta_{x=0}$ .

Next we turn to global existence. The proof uses a comparison principle for the  $Q$ -equation (2.11) with respect to given ordered temperatures, and the fact that temperature is positively bounded from below. The results are stated in terms of an equivalent formulation, as in (3.13), in which we replace  $t$  by  $\tau$ : i.e.

$$Q_\tau = y^{4/3}Q_{yy} + \frac{1}{\Theta(\tau)}QQ_y \quad \text{in } D_T \tag{5.2}$$

$$Q(0, \tau) = 0, \quad Q(1, \tau) = 1 \quad \text{for } \tau \in [0, T], \tag{5.3}$$

$$Q(y, 0) = Q_0(y) \quad \text{for } y \in [0, 1], \tag{5.4}$$

and the energy relation

$$E = \kappa\Theta(\tau) - \int_0^1 \frac{Q^2}{y^{4/3}} dy \quad \text{for } \tau \in [0, T]. \tag{5.5}$$

We first consider (5.2)-(5.4) for given ordered temperatures and ordered initial data.

**Proposition 5.3** Let  $i = 1, 2$ . Suppose  $Q_i$  solves (5.2)-(5.4) in  $D_{T_i}$  subject to  $Q_0 = Q_{0i}$ , and given  $\Theta = \Theta_i$ , satisfying (4.1) in  $[0, T_i]$ . Let  $T = \min\{T_1, T_2\}$ . If

$$\Theta_1 \leq \Theta_2 \quad \text{in } [0, T], \quad \text{and} \quad Q_{01} \geq Q_{02} \quad \text{in } (0, 1)$$

and if there exists  $K > 0$  such that either

$$0 \leq Q_{1y} \leq K \quad \text{or} \quad 0 \leq Q_{2y} \leq K \quad \text{in } \overline{D_T},$$

then

$$Q_1 \geq Q_2 \quad \text{in } D_T.$$

**Proof:** Suppose  $0 \leq Q_{2y} \leq K$ . Since

$$Q_{2\tau} = y^{4/3}Q_{2yy} + \frac{1}{\Theta_1(\tau)}Q_2Q_{2y} + \left(\frac{1}{\Theta_2(\tau)} - \frac{1}{\Theta_1(\tau)}\right)Q_2Q_{2y} \quad \text{in } D_T.$$

It follows from  $\Theta_1 \leq \Theta_2$  and  $Q_{2y} \geq 0$  that

$$Q_{2\tau} \leq y^{4/3}Q_{2yy} + \frac{1}{\Theta_1(\tau)}Q_2Q_{2y} \quad \text{in } D_T.$$

This inequality and the boundedness of  $Q_{2y}$  allows us to use [15, Theorem 3.2], which show that  $Q_2$  is a subsolution for the  $Q$ -equation with  $\Theta_1$ . ■

Next we use the Boltzmann entropy (2.7) in terms of  $Q = Q(y, \tau)$  to establish a positive lower bound on  $\Theta$ .

**Proposition 5.4** *Let  $\langle Q, \Theta \rangle$  be a solution of Problem (5.2)-(5.5). Suppose  $\Theta(0) = \frac{1}{\kappa} \left( E + \int_0^1 \frac{Q_0(y)}{y^{4/3}} dy \right) > 0$ . Then*

$$\Theta(\tau) \geq \lambda \Theta(0) \quad \text{for } \tau > 0, \quad \text{with } \lambda = \exp \left( -\frac{2}{\kappa} \int_0^1 Q_{0y} \log Q_{0y} dy \right). \quad (5.6)$$

**Proof:** Rewriting (2.7) results in

$$W(\tau) := \int_0^1 Q_y \log Q_y dy - \frac{\kappa}{2} \log \left( E + \int_0^1 \frac{Q^2}{y^{4/3}} dy \right) \quad \text{for } \tau > 0$$

and differentiation gives, see also (2.8),

$$\frac{dW(\tau)}{d\tau} = - \int_0^1 \frac{Q_\tau^2}{y^{4/3}Q_y} dy \leq 0 \quad \text{for } \tau > 0.$$

Hence  $W(\tau)$  is decreasing in  $\tau$ . As a consequence

$$\begin{aligned} W(\tau) &= \int_0^1 Q_y \log Q_y dy - \frac{\kappa}{2} \log \left( E + \int_0^1 \frac{Q^2}{y^{4/3}} dy \right) \leq \\ &\leq W(0) = \int_0^1 Q_{0y} \log Q_{0y} dy - \frac{\kappa}{2} \log \left( E + \int_0^1 \frac{Q_0^2}{y^{4/3}} dy \right) \quad \text{for } \tau > 0. \end{aligned}$$

Here we use Jensen's inequality to estimate

$$\int_0^1 Q_y \log Q_y dy \geq \left( \int_0^1 Q_y dy \right) \log \left( \int_0^1 Q_y dy \right) = 0,$$

from which lower bound (5.6) directly follows. ■

Note that whenever  $\Theta$  is bounded away from zero, blow-up in Problem (5.2)-(5.5) can only occur at the boundary  $y = 0$ . This is a direct consequence of classical regularity theory, which implies that  $Q$  is smooth away from  $y = 0$ . Blow-up manifests itself through singular behaviour of  $Q_y$  as  $(y, \tau)$  approaches the point  $(0, T^*)$ . This corresponds to unbounded density at the origin of the radially

symmetric solution of Problem **P**. Below we use the comparison argument (Proposition 5.3) to control the behaviour of  $Q_y(0, \tau)$ . We show that this implies a uniform bound on  $\|Q_y(\tau)\|_{L^2(0,1)}$  and thus on  $\|n(\tau)\|_{L^2(B_1(0))}$  for all  $0 \leq \tau < T^*$ . Global existence for  $Q = Q(y, \tau)$  as a consequence of Theorem 3.2. The results translate in a straightforward manner to the assertions of Theorem 1.2

**Theorem 5.5** *Let  $Q_0 : [0, 1] \mapsto [0, 1]$  be nondecreasing,  $Q_{0,y} \in L^\infty(0, 1)$  and  $Q_0(0) = 0$ ,  $Q_0(1) = 1$ . Let  $\Theta(0) = \Theta_0 > 0$ . If either*

(i)  $\Theta_0$  is sufficiently large;

or

(ii) there exists  $B > 0$  such that

$$\Theta_0 \geq \frac{3(1+B)}{2\lambda}, \quad \text{and} \quad Q_0(y) \leq \frac{y(1+B)}{y^{2/3}+B}, \quad \text{with} \quad \lambda = \exp\left(-\frac{2}{\kappa} \int_0^1 Q_{0,y} \log Q_{0,y} dy\right).$$

Then Problem (5.2)-(5.5) has a global solution  $\langle Q, \Theta \rangle$  in the sense of Section 3.3. Moreover there exist constants  $L > 0$  and  $\Theta^* > 0$  such that  $\Theta(\tau) \leq \Theta^*$  and  $\|Q_y(\tau)\|_{L^2(0,1)} \leq \|Q_{0,y}\|_{L^2(0,1)} \exp(L\tau)$  for all  $\tau > 0$ . If (ii) is satisfied we have in addition

$$Q(y, \tau) \leq \frac{y(1+B)}{y^{2/3}+B} \quad \text{for all} \quad (y, \tau) \in \overline{D_\infty}.$$

**Proof:** First we consider the auxiliary problem

$$(AP) \quad \begin{cases} \bar{Q}_\tau = y^{4/3} \bar{Q}_{yy} + \frac{1}{A} \bar{Q} \bar{Q}_y & \text{in} \quad D_\infty, \\ \bar{Q}(0, \tau) = 0, \quad \bar{Q}(1, \tau) = 1 & \text{for} \quad \tau > 0, \\ \bar{Q}(y, 0) = Q_0(y) & \text{for} \quad y \in [0, 1], \end{cases} \quad (5.7)$$

where  $A > 0$  and where  $Q_0$  satisfies the conditions of the theorem.

By Theorem 4.2 and Remark 4.1 (iii), we have: if either

$$(H_1) \quad A \geq \alpha_2 \quad \text{and} \quad \|Q_{0,y}\|_{L^2(0,1)}^2 \leq C(A^2 - A)$$

or

$$(H_2) \quad A = \frac{3}{2}(1+B) \quad \text{and} \quad Q_0(y) \leq \frac{y(1+B)}{y^{2/3}+B} \quad \text{for} \quad 0 \leq y \leq 1, \quad \text{and for some } B > 0,$$

then Problem **AP** has a global solution  $\bar{Q} : \overline{D_\infty} \mapsto [0, 1]$ . Since  $Q_{0,y} \in L^\infty(0, 1)$ , the regularity theory of [7, Theorem 2] gives

$$\|\bar{Q}_y\|_{L^\infty(0,1)} \in L_{loc}^\infty([0, \infty)). \quad (5.8)$$

The conditions on  $Q_0$  and  $\Theta_0$  guarantee that Problem (5.2)-(5.5) has a classical solution in  $D_T$  for some  $T$ . Now suppose

$$T^* = \sup\{T > 0 \mid \text{solution of Problem (5.2)-(5.5) exists in } D_T\} < \infty. \quad (5.9)$$

Fix  $A > 0$  such that  $(H_1)$  is satisfied, and choose  $\Theta_0 \geq \frac{A}{\lambda}$ . By (5.6), we have

$$\Theta(\tau) \geq A \quad \text{for all} \quad \tau \in [0, T^*) \quad (5.10)$$

and by Proposition 5.3 and (5.8), we find

$$Q(y, \tau) \leq \bar{Q}(y, \tau) \leq Ky \quad \text{for } (y, \tau) \in [0, 1] \times [0, T^*] \quad (5.11)$$

for some  $K > 0$ . Below we show that this implies a uniform bound on  $\|Q_y(\cdot)\|_{L^2(\Omega)}$  in  $[0, T^*]$ . Multiplying (5.2) by  $Q_\tau/y^{4/3}$  gives

$$\frac{1}{y^{4/3}}Q_\tau^2 = Q_\tau Q_{yy} + \frac{1}{\Theta(\tau)y^{4/3}}Q_\tau Q Q_y \quad \text{in } [0, 1] \times (0, T^*). \quad (5.12)$$

Using (5.10) and (5.11), the second term on the right can be estimated by

$$\begin{aligned} \frac{1}{\Theta(\tau)y^{4/3}}Q_\tau Q Q_y &\leq \frac{1}{y^{4/3}}Q_\tau^2 + \frac{1}{4\Theta^2(\tau)y^{4/3}}Q^2 Q_y^2 \\ &\leq \frac{1}{y^{4/3}}Q_\tau^2 + \frac{K^2}{4A^2}Q^2 Q_y^2. \end{aligned}$$

Using this in (5.12) and integrating the results gives

$$\frac{d}{d\tau}\|Q_y(\tau)\|_{L^2(0,1)}^2 \leq \frac{K^2}{2A^2}\|Q_y(\tau)\|_{L^2(0,1)}^2 \quad \text{for all } 0 \leq \tau < T^*.$$

Hence

$$\|Q_y(\tau)\|_{L^2(0,1)} \leq \|Q_{0y}\|_{L^2(0,1)} \exp\left(\left(\frac{M}{2A}\right)^2 T^*\right)$$

and

$$A \leq \Theta(\tau) \leq \frac{1}{\kappa}\left(E + \int_0^1 \frac{\bar{Q}^2(y, \tau)}{y^{4/3}} dy\right) \leq \frac{1}{\kappa}\left(E + \frac{3}{5}K^2\right)$$

for all  $0 \leq \tau < T^*$ . This allows us to use Theorem 3.2 at  $T^{*-}$ , which contradicts (5.9). The uniform upper bound in the temperature follows from the observation

$$\bar{Q}(y, \tau) = \int_0^y \bar{Q}_y(y, \tau) dy \leq y^{1/2}\|\bar{Q}_y(\tau)\|_{L^2(0,1)},$$

implying

$$\int_0^1 \frac{\bar{Q}^2(y, \tau)}{y^{4/3}} dy \leq \frac{3}{2}\|\bar{Q}_y(\tau)\|_{L^2(0,1)}^2,$$

and thus last expression is uniformly bounded if  $A$  satisfies  $(H_1)$  (Theorem 4.2).

If (ii) holds, global existence follows in an identical way. Again (5.10) and (5.11) hold, yielding the same bounds on  $\|Q_y(\tau)\|_{L^2(0,1)}$  and  $\Theta(\tau)$ . The pointwise bound on  $Q$  in  $D_\infty$  results from the fact that  $y(1+B)/(y^{2/3}+B)$  is a supersolution for Problem **AP** if  $A$  and  $Q_0$  satisfy  $(H_2)$ . Take for instance  $K = \frac{B+1}{B}$  in (5.11). The corresponding temperature bound is a direct consequence. ■

As a special case of Theorem 5.5 (ii) we have

**Corollary 5.6** *If  $Q_0(y) = y$ , and  $\Theta_0 > \frac{3}{2}$ , then Problem (5.2)-(5.5) has a global  $\langle Q, \Theta \rangle$  solution and  $\Theta_0 \leq \Theta(\tau) < \Theta_0 + \frac{12}{5\kappa}$  for all  $\tau \geq 0$ .*

**Proof:** Since  $\lambda = 1$ , we can select a sufficiently small  $B > 0$  such that Theorem 5.5 (ii) holds. The pointwise bound on  $Q$  implies  $Q(y, \tau) \leq y^{1/3}$  for all  $(y, \tau) \in \overline{D_\infty}$ . Since

$$\Theta(\tau) = \Theta_0 + \int_0^1 \frac{Q^2(y, \tau) - Q_0^2(y)}{y^{4/3}} dy,$$

the upper bound is immediate. ■

## 6. APPENDIX: INEQUALITIES

For completeness we give in this appendix some inequalities which are used at various places in the paper.

Let  $\Omega$  be a bounded open subset of  $\mathbb{R}^N$  with a  $C^{1+\alpha}$  ( $\alpha > 0$ ) boundary.

**First interpolation inequality.** Let  $N > 2$ ,  $r \leq \frac{2N}{N-2}$  and let  $p \leq q \leq r$  satisfy  $\frac{1}{q} = \frac{\alpha}{p} + \frac{(1-\alpha)}{r}$  for some  $\alpha \in (0, 1)$ . Then

$$\|n\|_{L^q(\Omega)} \leq C_s^{1-\alpha} \|n\|_{H^1(\Omega)}^{1-\alpha} \|n\|_{L^p(\Omega)}^\alpha \quad \text{for all } n \in H^1(\Omega) \cap L^p(\Omega). \quad (6.1)$$

**Proof:** Use the Sobolev inequality  $\|n\|_{L^r(\Omega)} \leq C_s \|n\|_{H^1(\Omega)}$  for  $N > 2$  and  $r \leq \frac{2N}{N-2}$ , and the interpolation inequality  $\|n\|_{L^q(\Omega)} \leq \|n\|_{L^p(\Omega)}^\alpha \|n\|_{L^r(\Omega)}^{1-\alpha}$ . ■

**Second interpolation inequality.** Let  $N = 3$ . Then

$$\|n\|_{L^3(\Omega)} \leq C_s^{1/2} \|n\|_{H^1(\Omega)}^{1/2} \|n\|_{L^2(\Omega)}^{1/2} \quad \text{for all } n \in H^1(\Omega). \quad (6.2)$$

**Proof:** Take  $p = 2$ ,  $q = 3$ ,  $r = \frac{2N}{N-2} = 6$  and  $\alpha = 1/2$  in (6.1). ■

**Poisson's equation and  $L^p$ -norms.** Let  $n \in L^p(\Omega)$ ,  $p > \frac{N}{2}$ , and let  $\phi$  satisfy (1.2) and (1.5). Then

$$\begin{cases} \|\nabla\phi\|_{L^r(\Omega)} \leq C_I \|n\|_{L^p(\Omega)} & \text{for } 1 < r \leq \frac{pN}{N-p} \quad \text{and} \quad \frac{N}{2} < p < N, \\ \|\nabla\phi\|_{L^\infty(\Omega)} \leq C_I \|n\|_{L^p(\Omega)} & \text{for } p > N. \end{cases} \quad (6.3)$$

where the constant  $C_I$  depends on  $\Omega$  and  $p$ .

**Proof:** Since  $\phi$  satisfies (1.2) with (1.5), we use the representation by the Green's function to obtain  $\|\phi\|_{L^p(\Omega)} \leq \|\Delta\phi\|_{L^p(\Omega)}$  for  $N > 2$  and  $p > N/2$ . If  $p < N$  we combine this with the Sobolev inequality  $\|\nabla\phi\|_{L^r(\Omega)} \leq C(\|\Delta\phi\|_{L^p(\Omega)} + \|\phi\|_{L^p(\Omega)})$  for  $r \leq pN/(N-p)$  to obtain the desired inequality. If  $p > N$  we proceed similarly. ■

## Acknowledgments

The authors are grateful to P.-H. Chavanis for bringing this problem to their attention and J. Dolbeault for pointing out reference [5].

## References

1. ANTONOV, V.A., Most Probable phase distribution in spherical star systems and conditions for its existence, *Vest. Leningr. gos. Univ.*, **7** (1962), 135. (English version: Hut P., ed., 1984, IAU Symp. 113, Dynamics of stars clusters. Reidel, Dordrecht)
2. BILER, P., Growth and accretion of mass in an astrophysical model, *Applicationes Math.*, **23** (1995), 179-189.
3. BILER, P. AND NADZIEJA T., Growth and accretion of mass in an astrophysical model II, *Applicationes Math.*, **23** (1995), 351-361.
4. BILER, P., Existence and nonexistence of solutions for a model of gravitational interaction of particles, III, *Colloq. Math.*, **68** (1995), 229-239.
5. BILER, P., DOLBEAULT, J., ESTEBAN, M.J., MARKOWICH, P.A., AND NADZIEJA T., Steady States for Streater's Energy-transport Models of Self-Gravitating Particles, submitted to IMA Volumes in Mathematics Series, Springer Verlag, 2000.
6. BILER, P., HEBISCH, W. AND NADZIEJA T., The Debye system: existence and long time behaviour of solutions, *Nonlinear Analysis*, **23** (1994), 1189-1209.
7. BILER, P., HILHORST D. AND NADZIEJA T., Existence and nonexistence of solutions for a model of gravitational interaction of particles, II, *Colloq. Math.*, **67** (1994), 297-308.
8. BILER, P., KRZYWICKI, A. AND NADZIEJA T., Self-interaction of Brownian particles coupled with thermodynamic processes, *Rep. Math. Phys.* **42** (1998), 359-372.
9. BILER, P. AND NADZIEJA T., Existence and nonexistence of solutions for a model of gravitational interaction of particles, I, *Colloq. Math.*, **66** (1994), 319-334.
10. BILER, P. AND NADZIEJA T., Nonlocal parabolic problems in statistical mechanics, *Nonlinear Analysis*, (8) **30** (1997), 5343-5350.
11. BILER, P. AND NADZIEJA T., A class of nonlocal parabolic problems occurring in statistical mechanics, *Colloq. Math.*, **66** (1993), 131-145.
12. BIDAUT-VÉRON, M.-F. AND VÉRON, L., Nonlinear elliptic equations on compact Riemannian manifolds and asymptotics of Emden equations, *Invent. Math.* **106** (1991), 489-539.
13. BRENNER, M.P., CONSTANTIN, P., KADANOFF, L.P., SCHNKEL, A., VENKATARAMANI, S.C., Diffusion, attraction and collapse, *Nonlinearity*, **12** (1999) 1071-1098.

14. CHAVANIS, P.H. SOMMERIA, J. AND ROBERT, R., Statistical mechanics of two-dimensional vortices and collisionless stellar systems, *Astrophys. J.* **471** (1996), 385-399.
15. HILHORST D., A nonlinear evolution problem arising in the physics of ionized gases, *SIAM J. Math. Anal.* **13** (1982), 16-39.
16. LIONS, J.L., Quelques méthodes de résolution des problèmes aux limites non linéaires, *Dunod-Gauthier Villars* (1969).
17. ROSIER C., Probleme de Cauchy pour une équation parabolique modélisant la relaxation des systèmes stellaires auto-gravitans, *C. R. Acad. Sci. Paris, Série I* **332** (2001) 903-908.
18. WOLANSKY, G., Comparison between two models of self-gravitating clusters: conditions for gravitational collapse, *Nonlinear Analysis*, (7) **24** (1995), 1119-1129.
19. WOLANSKY, G., On steady distributions of self-attracting clusters under friction and fluctuations, *Arch. Rational Mech. Anal.* **119** (1992), 355-391.
20. TEMAM, R. Navier-Stokes Equations. North Holland (1979).
21. ZEIDLER, E. Nonlinear functional analysis and its applications, Part I-IV, *Springer Verslag*, (1986)-(1988).