



Centrum voor Wiskunde en Informatica

REPORTRAPPORT

Zero and Pole Distribution of Diagonal Padé Approximants to
the Exponential Function

K.A. Driver, N.M. Temme

Modelling, Analysis and Simulation (MAS)

MAS-R9701 February 28, 1997

Report MAS-R9701
ISSN 1386-3703

CWI
P.O. Box 94079
1090 GB Amsterdam
The Netherlands

CWI is the National Research Institute for Mathematics and Computer Science. CWI is part of the Stichting Mathematisch Centrum (SMC), the Dutch foundation for promotion of mathematics and computer science and their applications.

SMC is sponsored by the Netherlands Organization for Scientific Research (NWO). CWI is a member of ERCIM, the European Research Consortium for Informatics and Mathematics.

Copyright © Stichting Mathematisch Centrum
P.O. Box 94079, 1090 GB Amsterdam (NL)
Kruislaan 413, 1098 SJ Amsterdam (NL)
Telephone +31 20 592 9333
Telefax +31 20 592 4199

Zero and Pole Distribution of Diagonal Padé Approximants to the Exponential Function

Kathy A. Driver¹ and Nico M. Temme²

¹*Mathematics Department, University of the Witwatersrand
P.O. WITS 2050, South Africa*

²*CWI, P.O. Box 94079, 1090 GB Amsterdam, The Netherlands
e-mail: 036kad@cosmos.wits.ac.za, nicot@cwi.nl*

Dedicated to the memory of Douglas Sears

This report is a preprint version of a paper that has been accepted for publication in a special issue of *Questiones Mathematicae*, to honor the South African mathematician Douglas Sears.

ABSTRACT

The polynomials P_n and Q_m having degrees n and m respectively, with P_n monic, that solve the approximation problem

$$P_n(z)e^{-z} + Q_m(z) = \mathcal{O}(z^{n+m+1})$$

will be investigated for their asymptotic behaviour, in particular in connection with the distribution of their zeros. The symbol \mathcal{O} means that the left-hand side should vanish at the origin at least to the order $n + m + 1$. This problem is discussed in great detail in a series of papers by Saff and Varga. In the present paper we show how their results can be obtained by using uniform expansions of integrals in which Airy functions are the main approximants. We shall focus on the important diagonal case when $n = m$ and the polynomials P_n and Q_n , as well as the remainder $E_{n,n}(z) = P_n(z)e^{-z} + Q_n(z)$ can be expressed in terms of Hankel and Bessel functions. The approximate location of the zeros of P_n , Q_n and $E_{n,n}$ are given in terms of the known zeros of certain Airy functions. An application is given in which the asymptotic information on the zeros is used to obtain an estimate in an approximation of the unit block function by means of the polynomials P_n , Q_n .

1991 Mathematics Subject Classification: 41A21, 30E15, 33C10, 30C15, 41A60.

Keywords & Phrases: Padé polynomials, asymptotic behaviour, uniform asymptotic methods, exponential function, Bessel functions, Airy functions, zero and pole distribution.

Note: Work carried out under project MAS2.8 Exploratory research.

1. Introduction

It is well known (cf. PERRON (1950, page 433)) that the solution of the Padé approximation problem for the exponential function, namely,

$$P_n(z)e^{-z} + Q_m(z) = \mathcal{O}(z^{n+m+1}) \quad \text{as } z \rightarrow 0, \quad (1.1)$$

where P_n and Q_m are polynomials of degree n and m respectively, with P_n monic, is given by

$$P_n(z) = \frac{1}{m!} \int_0^\infty t^m (t+z)^n e^{-t} dt,$$

$$Q_m(z) = \frac{-1}{m!} \int_0^\infty t^n (t-z)^m e^{-t} dt.$$

Explicit forms are (cf. PERRON (1950, page 436))

$$P_n(z) = n! \sum_{k=0}^n \binom{m+n-k}{m} \frac{z^k}{k!}, \quad (1.2)$$

$$Q_m(z) = -n! \sum_{k=0}^m \binom{m+n-k}{n} \frac{(-z)^k}{k!}. \quad (1.3)$$

Let the remainder $E_{n,m}$ be defined by

$$E_{n,m}(z) = P_n(z)e^{-z} + Q_m(z) \quad (1.4)$$

then

$$E_{n,m}(z) = \frac{(-1)^m z^{m+n+1}}{m!} \int_0^1 u^n (1-u)^m e^{-uz} du.$$

The quantities $P_n, Q_m, E_{n,m}$ can be expressed in terms of (confluent) hypergeometric functions. We have

$$P_n(z) = U(-n, -n-m, z) = z^{n+m+1} U(m+1, n+m+2, z)$$

$$Q_m(z) = -\frac{n!}{m!} U(-m, -n-m, -z) = -\frac{n!}{m!} (-z)^{n+m+1} U(n+1, n+m+2, -z)$$

$$E_{n,m}(z) = \frac{(-1)^{m+1} n! z^{n+m+1}}{(n+m+1)!} M(n+1, n+m+2, -z) \quad (1.5)$$

$$= \frac{(-1)^{m+1} n! z^{n+m+1}}{(n+m+1)!} e^{-z} M(m+1, n+m+2, z).$$

In a sequence of papers Saff and Varga investigated the polynomials P_n, Q_m and the remainder $E_{n,m}$, and the distribution of their zeros, for large values of n, m with fixed ratio $\sigma = m/n$ (the final paper appeared in 1978). They used saddle point methods for the integrals defining the U - and M -functions and found curves in the complex z -plane along which

the zeros are cumulating. For $m = 0$ their results agree with the earlier results obtained by Szegö on the distribution of the zeros of the exponential polynomial

$$e_n(z) = \sum_{k=0}^n \frac{z^k}{k!}.$$

In this paper, we adopt a new approach for locating the zeros of P_n, Q_m and $E_{n,m}$ by using uniform asymptotic approximations for these functions in terms of Airy functions. Approximate values of the zeros of P_n, Q_m and $E_{n,m}$ are given in terms of the (known) zeros of certain Airy functions. For all positive integers m and n , contour integral representations of P_n, Q_m and $E_{n,m}$ can be given and uniform asymptotic methods applied to these contour integrals (cf. DRIVER & TEMME (1996a,b)). Our investigations here will focus on the diagonal case $n = m$ where certain simplifications occur. For $n = m$, the quantities P_n, Q_n and $E_{n,n}$ can be expressed in terms of Hankel and Bessel functions which permit Airy-type approximations to be done via Sommerfeld contour integrals, which seem not to be available in the general non-diagonal case.

2. The relations with Bessel and Hankel functions

In the diagonal case $n = m$ we have using the representations in (1.5) (cf. also ABRAMOWITZ & STEGUN (1964, pages 509, 510))

$$P_n(z) = \pi^{-\frac{1}{2}} z^{n+\frac{1}{2}} e^{z/2} K_{n+\frac{1}{2}}(z/2), \quad E_{n,n}(z) = \frac{1}{2} (-1)^{n+1} \pi^{\frac{1}{2}} z^{n+\frac{1}{2}} e^{-z/2} I_{n+\frac{1}{2}}(z/2). \quad (2.1)$$

It follows immediately from (1.2) and (1.3) that, when $m = n$,

$$Q_n(z) = -P_n(-z). \quad (2.2)$$

Of course, since the $[m/n]$ Padé approximant for e^{-z} is $-Q_m(z)/P_n(z)$, the zeros of $P_n(z)$ are the poles of this Padé approximant and the above symmetry of P_n and Q_n means that knowledge about the zeros of either one of the polynomials suffices for both. Nevertheless, for our discussion, it is better to write P_n and Q_n in terms of different Bessel functions.

We have given in (2.1) the relations with modified Bessel functions. For obtaining the asymptotic expansions it is better to write the functions in terms of ordinary Bessel and Hankel functions. We recall the expansions (cf. WATSON (1944, page (201)) or TEMME (1996, page 239))

$$H_{n+\frac{1}{2}}^{(1)}(z) = \sqrt{2/(\pi z)} e^{-(n+1)\pi i/2+iz} \sum_{m=0}^n (+i)^m (n + \frac{1}{2}, m)(2z)^{-m}, \quad (2.3)$$

$$H_{n+\frac{1}{2}}^{(2)}(z) = \sqrt{2/(\pi z)} e^{+(n+1)\pi i/2-iz} \sum_{m=0}^n (-i)^m (n + \frac{1}{2}, m)(2z)^{-m}, \quad (2.4)$$

where Hankel's symbol (α, m) is defined by

$$(\alpha, m) = \frac{\Gamma(\frac{1}{2} + \alpha + m)}{m! \Gamma(\frac{1}{2} + \alpha - m)}, \quad m = 0, 1, 2, \dots$$

Since

$$(n + \frac{1}{2}, m) = \frac{(n + m)!}{m!(n - m)!} \quad \text{for } m = 0, 1, \dots, n,$$

a simple comparison of (2.3) and (2.4) with (1.3) and (1.2) respectively yields the relations

$$\begin{aligned} P_n(2iz) &= -i(2z)^n \sqrt{\pi z/2} e^{iz} H_{n+\frac{1}{2}}^{(2)}(z), \\ Q_n(2iz) &= -i(2z)^n \sqrt{\pi z/2} e^{-iz} H_{n+\frac{1}{2}}^{(1)}(z), \\ E_{n,n}(2iz) &= -i(2z)^n \sqrt{2\pi z} e^{-iz} J_{n+\frac{1}{2}}(z), \end{aligned} \tag{2.5}$$

where the third line in (2.5) follows from the relation

$$J_\nu(z) = \frac{1}{2} \left[H_\nu^{(1)}(z) + H_\nu^{(2)}(z) \right]. \tag{2.6}$$

It is known where the zeros of these Bessel functions occur when n is large. For $J_{n+1/2}(z)$ they are located in the interval (n, ∞) . For $E_{n,n}(z)$ this gives an infinite set of zeros along the positive imaginary axis, in the interval $(2in, i\infty)$; there is a conjugate set in $(-2in, -i\infty)$. The Hankel functions have zeros along curves in the complex plane which start also at $z = n$; see ABRAMOWITZ & STEGUN (1964, page 373). For $P_n(z)$ this gives n zeros in the half-plane $\Re z < 0$ along an arc from $2in$ to $-2in$. The arc cuts the real positive axis at $z = -2na$, where $a = 0.66274\dots$. The zeros of $Q_n(z)$ are located in the right half-plane, and they follow from $Q_n(z) = -P_n(-z)$.

We give asymptotic representations of P_n , Q_n and $E_{n,n}$ for large values of n in terms of Airy functions. In OLVER (1974, Ch. 11. §10) Airy-type expansions of the Bessel functions $J_\nu(\nu z)$ and $H_\nu^{(j)}(\nu z)$ are considered, but in the present case we need similar representations for $J_{n+\frac{1}{2}}(nz)$ and $H_{n+\frac{1}{2}}^{(j)}(nz)$.

3. Airy-type approximations for Bessel and Hankel functions

We give the main steps for deriving Airy-type approximations for Bessel and Hankel functions by using the Sommerfeld contour integrals. Since uniform asymptotic approximations of $J_\nu(\nu z)$, $H_\nu^{(1)}(\nu z)$ and $H_\nu^{(2)}(\nu z)$ can be found in OLVER (1974, p. 423–425) we can use the formulas in (2.5) to write down the corresponding approximations for $P_n[2i(n+\frac{1}{2})z]$, $Q_n[2i(n+\frac{1}{2})z]$ and $E_{n,n}[2i(n+\frac{1}{2})z]$ immediately. In Olver's book the results are derived by using the differential equation of the Bessel functions, and the results are shown to hold in large domains of complex parameters and are provided with error bounds for the remainders in the expansions.

Because in the non-diagonal case, and also in Hermite-Padé approximations to the exponential functions (both aspects are studied in our future papers DRIVER & TEMME (1996a,b))

these powerful results and methods are not available, we give in this preparatory paper the standard procedure based on the Sommerfeld integrals. A similar procedure will be used in the more general cases on more complicated integrals. In the Hermite-Padé case differential equations are not known for the corresponding polynomials.

We have (OLVER (1974, p. 58))

$$J_\nu(z) = \frac{1}{2\pi i} \int_{\mathcal{C}} e^{z \sinh t - \nu t} dt,$$

where \mathcal{C} is a contour starting at $\infty - \pi i$ and terminating at $\infty + \pi i$. It follows that

$$J_{n+\frac{1}{2}}(nz) = \frac{1}{2\pi i} \int_{\mathcal{C}} e^{n\phi(t) - \frac{1}{2}t} dt, \quad \phi(t) = z \sinh t - t.$$

Solving the equation $\phi'(t) = 0$, gives the saddle points

$$t^\pm = \pm\tau, \quad \tau = \operatorname{arccosh} \frac{1}{z}. \quad (3.1)$$

When $0 < z < 1$, these points are real, when $z > 1$ they lie on the imaginary axis. In order to obtain a standard form which can be used for obtaining a uniform expansion that is valid when the saddle points are close together, or coincide, we use the cubic transformation

$$\phi(t) = \frac{1}{3}\zeta^3 - \eta\zeta + A, \quad (3.2)$$

where the saddle points t^\pm should correspond with the saddle points $\pm\sqrt{\eta}$ in the ζ -plane. This transformation was introduced by CHESTER, FRIEDMAN & URSELL (1957). They showed that the mapping is one-to-one and analytic for all z in a neighborhood of $z = 1$, which is a local result. In fact, it can be shown (cf. DRIVER & TEMME (1996B)) that the mapping is one-to-one and analytic in a domain containing the path of integration.

The cubic transformation gives, upon substituting $t = t^\pm, \zeta = \pm\sqrt{\eta}$,

$$\frac{2}{3}\eta^{3/2} = \tau - \tanh \tau, \quad A = 0. \quad (3.3)$$

If $0 < z < 1$ then τ and η are positive. We can also write for positive z :

$$\begin{aligned} \frac{2}{3}\eta^{3/2} &= \operatorname{arctanh} \sqrt{1-z^2} - \sqrt{1-z^2}, \quad 0 < z \leq 1, \\ \frac{2}{3}(-\eta)^{3/2} &= \sqrt{z^2-1} - \arctan \sqrt{z^2-1}, \quad z \geq 1, \end{aligned} \quad (3.4)$$

which gives a better insight in the relation between η and z . We can expand the $\operatorname{arctanh}$ -function to obtain

$$\frac{2}{3}\eta^{3/2} = \frac{1}{3}(1-z^2)^{3/2} + \frac{1}{5}(1-z^2)^{5/2} + \dots,$$

which gives

$$\eta = 2^{1/3}(1-z)[1 + \mathcal{O}(1-z)], \quad z \rightarrow 1. \quad (3.5)$$

This defines the relation near $z = 1$. For complex values of z this relation should be used with analytic continuation to define which branch of the multi-valued function $\eta^{3/2}$ is used.

Using the cubic transformation we obtain

$$J_{n+\frac{1}{2}}(nz) = \frac{1}{2\pi i} \int_{\mathcal{C}} e^{n(\frac{1}{3}\zeta^3 - \eta\zeta)} f(\zeta) d\zeta, \quad f(\zeta) = e^{-\frac{1}{2}t} \frac{dt}{d\zeta}, \quad (3.6)$$

where \mathcal{C} is a contour running from $\infty \exp(-\pi i/3)$ to $\infty \exp(\pi i/3)$, through the saddle point at $\zeta = \sqrt{\eta}$. The first order Airy-type uniform approximation is obtained by replacing $f(\zeta)$ with

$$\begin{aligned} f_0 &= \frac{1}{2}[f(\sqrt{\eta}) + f(-\sqrt{\eta})] \\ &= \cosh \frac{1}{2}\tau \left. \frac{dt}{d\zeta} \right|_{\zeta=\sqrt{\eta}} \\ &= \sqrt{\frac{z+1}{2z}} \left(\frac{4\eta}{1-z^2} \right)^{1/4}. \end{aligned} \quad (3.7)$$

This gives (compare with OLVER (1974, page 425) and ABRAMOWITZ & STEGUN (1964, page 368))

$$J_{n+\frac{1}{2}}(nz) \sim n^{-1/3} f_0 \text{Ai}(\eta n^{2/3}), \quad n \rightarrow \infty \quad (3.8)$$

and, by using the third line in (2.5),

$$E_{n,n}(2izn) \sim -i\sqrt{\pi} n^{-1/3} (2zn)^{n+1/2} e^{-izn} f_0 \text{Ai}(\eta n^{2/3}). \quad (3.9)$$

In a similar way we obtain for the Hankel functions

$$\begin{aligned} H_{n+\frac{1}{2}}^{(1)}(nz) &= \frac{1}{\pi i} \int_{-\infty}^{\infty+\pi i} e^{n\phi(t) - \frac{1}{2}t} dt \\ &= \frac{1}{\pi i} \int_{\mathcal{C}^{(1)}} e^{n(\frac{1}{3}\zeta^3 - \eta\zeta)} f(\zeta) d\zeta \\ &\sim -2n^{-1/3} f_0 e^{2\pi i/3} \text{Ai}(\eta n^{2/3} e^{2\pi i/3}), \end{aligned} \quad (3.10)$$

$$Q_n(2izn) \sim i\sqrt{\pi} n^{-1/3} (2zn)^{n+1/2} e^{-izn} f_0 e^{2\pi i/3} \text{Ai}(\eta n^{2/3} e^{2\pi i/3}),$$

$$\begin{aligned} H_{n+\frac{1}{2}}^{(2)}(nz) &= \frac{-1}{\pi i} \int_{-\infty}^{\infty-\pi i} e^{n\phi(t) - \frac{1}{2}t} dt \\ &= \frac{-1}{\pi i} \int_{\mathcal{C}^{(-1)}} e^{n(\frac{1}{3}\zeta^3 - \eta\zeta)} f(\zeta) d\zeta \\ &\sim -2n^{-1/3} f_0 e^{-2\pi i/3} \text{Ai}(\eta n^{2/3} e^{-2\pi i/3}), \end{aligned} \quad (3.11)$$

$$P_n(2izn) \sim i\sqrt{\pi} n^{-1/3} (2zn)^{n+1/2} e^{izn} f_0 e^{-2\pi i/3} \text{Ai}(\eta n^{2/3} e^{-2\pi i/3}),$$

where $\mathcal{C}^{(j)}$ are contours running from $-\infty$ to $\infty \exp(j\pi i/3)$, $j = -1, 1$. The results for the Hankel functions yield the Bessel function result because of the relation in (2.6) and (cf. OLVER (1974, page 414))

$$e^{-2\pi i/3} \text{Ai}\left(ze^{-2\pi i/3}\right) + e^{2\pi i/3} \text{Ai}\left(ze^{2\pi i/3}\right) + \text{Ai}(z) = 0.$$

We can obtain more terms in the approximations given in (3.9) – (3.11) by using standard methods for obtaining uniform expansion of integrals (cf. WONG (1989)). On the other hand, as remarked at the beginning of this section, detailed information on this type of uniform expansions can be obtained from OLVER (1974).

4. Locating the zeros

Having available the asymptotic approximations in terms of the Airy functions we can obtain approximations of the locations of the zeros of $P_n, Q_n, E_{n,n}$. The only zeros of the Airy function $\text{Ai}(z)$ occur along the negative z -axis, and there are infinitely many zeros (cf. OLVER (1974, p. 415)). When the argument of the Airy function in (3.9) is negative, we have $\eta = \exp(\pm i\pi)$, and $\eta^{3/2} = \mp i$. Because the right-hand side of (3.4) is purely imaginary when $z > 1$, we conclude that an infinite string of zeros of $E_{n,n}(2izn)$ occurs near the intervals $(-\infty, -1), (1, \infty)$ (in fact all zeros are real and inside these intervals). The zeros of the polynomials $P_n(2izn), Q_n(2izn)$ are located along curves in the z -plane that are defined by $\eta \exp(-2\pi i/3) < 0, \eta \exp(+2\pi i/3) < 0$, respectively, where, again, the relation between η and z is given in (3.4).

To obtain a first approximation of these zeros, we use (3.9)–(3.11). Let a_j be the zeros of $\text{Ai}(z)$ (we have $a_1 = -2.3381\dots$; see ABRAMOWITZ & STEGUN (1964, p. 478) for more values). Then the zeros z_s of $P_n(2izn)$ follow from the single term estimate obtained from the final line in (3.11)

$$\eta_s \sim a_s e^{2\pi i/3} n^{-2/3}, \quad n \rightarrow \infty,$$

and by inverting the relation between η and z (cf. (3.4)). More precise asymptotic estimates follow from using more terms in the uniform asymptotic approximations in (3.9)–(3.11) (cf. DRIVER & TEMME (1996b)). The zeros of $P_n(2izn)$ near $z = 1$ satisfy (cf. (3.5))

$$z_j \sim 1 - a_j 2^{-1/3} e^{2\pi i/3} n^{-2/3}, \quad n \rightarrow \infty. \quad (4.1)$$

4.1. The condition for the zeros

We compare the condition for the location of the zeros of the quantities $P_n, Q_n, E_{n,n}$ as given in SAFF & VARGA (1978) with the condition that follows from the uniform Airy-type asymptotic approximation. Saff and Varga introduce the quantity

$$w_1(z) = \frac{2iz e^{\sqrt{1-z^2}}}{1 + \sqrt{1-z^2}}, \quad (4.2)$$

where we translated their notation into ours (that is, we take $\sigma = m/n = 1$ and change z into iz). According to Saff and Varga, the zeros of the three quantities $P_n, Q_m, E_{n,m}$ occur along curves in the z -plane defined by

$$|w_1(z)| = 1. \quad (4.3)$$

By using $z = 1/\cosh \tau$ (cf. (3.1)), a straightforward computation shows that

$$\ln w_1(z) = \frac{1}{2}\pi i + \tanh \tau - \tau = \frac{1}{2}\pi i - \frac{2}{3}\eta^{3/2}, \quad (4.4)$$

where $\frac{2}{3}\eta^{3/2}$ is defined in (3.3). The condition (4.3) can be read as: $\ln w_1(z)$ is purely imaginary. So an equivalent formulation of (4.3) reads: $-\eta^{3/2}$ is purely imaginary, that is, the phase of η equals $\pm\pi$ or $\pm\pi/3$. In other words, η is located on the rays where the zeros of $\text{Ai}(z)$ and $\text{Ai}(e^{\pm 2\pi/3}z)$ lie. This is in agreement with our description of the location of the zeros given earlier in this section.

5. An application: approximating the block function

As an application we consider the approximation of the block function

$$f(t) = \begin{cases} 1, & \text{if } 0 < x < 1; \\ 0, & \text{otherwise.} \end{cases} \quad (5.1)$$

This function plays an important role in, for instance, the theory of electronics. We describe a method to obtain approximations of the block function by using Padé approximants of the exponential function. By using the estimates of the zeros of the polynomials $P_n(z)$ given in §4 we derive an asymptotic estimate of the error in the approximation of the block function.

By taking the Laplace transform of f and inverting we can write

$$f(t) = \frac{1}{2\pi i} \int_{\mathcal{L}} e^{st} (1 - e^{-s}) \frac{ds}{s}, \quad (5.2)$$

where \mathcal{L} is a vertical line in the half-plane $\Re s > 0$. To obtain smooth approximations of f we can replace the exponential function $\exp(-s)$ with its Padé approximations, and we obtain

$$f_n(t) = \frac{1}{2\pi i} \int_{\mathcal{L}} e^{st} \left[1 + \frac{Q_n(s)}{P_n(s)} \right] \frac{ds}{s}, \quad n = 1, 2, \dots \quad (5.3)$$

For example,

$$f_1(t) = 2e^{-2t}, \quad f_2(t) = 4\sqrt{3} e^{-3t} \sin(\sqrt{3} t). \quad (5.4)$$

From the results in the previous sections it follows that the polynomial $P_n(s)$ in (5.3) has n simple zeros s_j in the left half-plane $\Re s < 0$. When we want to evaluate $f_n(t)$ of (5.3) numerically we can split up the rational function into partial fractions and evaluate the resulting integrals. By using residue calculus it is not difficult to prove that

$$f_n(t) = \sum_{j=1}^n e^{s_j t} \frac{Q_n(s_j)}{s_j P_n'(s_j)}. \quad (5.5)$$




Figure 5.1. The functions $f_n(t)$ of (5.3) that approximate the block function $n = 2, 3, \dots, 10$. The rate of approximation for large values of n is described in (5.7), for $t > 1$.

In this way we obtained the first elements given in (5.4) and in Figure 5.1. Observe that $s = 0$ is not a pole in (5.3).

It is of interest to know the asymptotic nature of the representation in (5.5) for large values of n . When we order the zeros s_j of $P_n(s)$ with respect to their real parts, a sum of residues as in (5.5) gives an excellent representation with respect to large values of t . Surprisingly, the situation is different in the present case for large values of n .

The zeros s_j with the largest real part (whose corresponding terms in (5.5) dominate the large t -behaviour) occur near $\pm 2in$. The behaviour of the s_j near $s = 2in$ follows from (4.1):

$$s_j \sim 2in - a_j 2^{2/3} e^{7\pi i/6} n^{1/3}, \quad n \rightarrow \infty. \quad (5.6)$$

Similar approximations hold near $s = -2in$, the real axis being an axis of symmetry. From (2.5) it follows that

$$\frac{Q_n(s_j)}{s_j P_n'(s_j)} = e^{-s_j} \frac{2i H_{n+\frac{1}{2}}^{(1)}[s_j/(2i)]}{s_j H_{n+\frac{1}{2}}^{(2)'}[s_j/(2i)]} = e^{-s_j} \mathcal{O}\left(n^{-2/3}\right),$$

where we have used well-known estimates of the Hankel functions:

$$H_\nu^{(1)}(\nu z) = \mathcal{O}\left(\nu^{-1/3}\right), \quad H_\nu^{(2)'}(\nu z) = \mathcal{O}\left(\nu^{-2/3}\right), \quad \nu \rightarrow \infty$$

with $z \sim 1$ (cf. ABRAMOWITZ & STEGUN (1964, pages 368–369)). Hence,

$$f_n(t) = e^{(t-1)s_1} \mathcal{O}\left(n^{-2/3}\right) = \mathcal{O}\left[n^{-2/3} e^{-(t-1)|a_1|2^{2/3} \cos(\pi/6)n^{1/3}}\right], \quad (5.7)$$

as $n \rightarrow \infty$ (with $a_1 = -2.3381\dots$), which estimate holds if $t > 1$.

We see that the expected estimate $f_n(t) \sim 1$ for $0 < t < 1$ cannot be obtained in this way. Apparently, if $0 < t < 1$, all exponentially large contributions in (5.5) cancel against each other. When we concentrate on the error in the approximation, that is, on

$$f(t) - f_n(t) = \frac{-1}{2\pi i} \int_{\mathcal{C}} e^{st} \frac{E_{n,n}(s)}{P_n(s)} \frac{ds}{s}, \quad n = 1, 2, \dots, \quad (5.8)$$

we can use saddle point methods. We can write this in the form

$$f(t) - f_n(t) = \frac{(-1)^{n+1}}{2i} \int_{\mathcal{C}} e^{2w(t-1)} \frac{I_{n+\frac{1}{2}}(w)}{K_{n+\frac{1}{2}}(w)} \frac{dw}{w}, \quad (5.9)$$

where we used the modified Bessel function representations given in (2.1). We are not pursuing this further, because it is outside the scope of this paper.

We conclude this section by stating an interesting property of the approximations $f_n(t)$ defined in (5.3). We have

$$\int_0^\infty f_n^2(t) dt = 1, \quad n = 1, 2, \dots, \quad (5.10)$$

which, on the one hand, says that the L^2 -norm of the approximant is the same as that of the block function, but on the other hand, it means that the electronic system, based upon these functions $f_n(t)$, is stable.

To show (5.10) we use (5.5):

$$I = \int_0^\infty f_n^2(t) dt = \sum_{j=1}^n \frac{Q_n(s_j)}{s_j P'_n(s_j)} \int_0^\infty e^{s_j t} f_n(t) dt.$$

We know from the earlier literature on Padé approximations to the exponential function (cf. SAFF & VARGA (1978) and their earlier papers) that all zeros s_j of $P_n(s)$ are in the left half-plane $\Re s < 0$. Hence, inverting (5.3), we have

$$I = - \sum_{j=1}^n \frac{Q_n(s_j)}{s_j^2 P'_n(s_j)} \left[1 + \frac{Q_n(-s_j)}{P_n(-s_j)} \right] = - \sum_{j=1}^n \frac{Q_n(s_j)}{s_j^2 P'_n(s_j)},$$

because $Q_n(-s_j) = -P_n(s_j) = 0$. It is not difficult to verify that

$$I = \frac{-1}{2\pi i} \int_{\mathcal{C}} \left[1 + \frac{Q_n(s)}{P_n(s)} \right] \frac{ds}{s^2},$$

where \mathcal{C} is a vertical line in the left half-plane such that all zeros of P_n are at left of \mathcal{C} ; on \mathcal{C} we integrate from $-i\infty$ to $i\infty$. Using $1 + Q_n(s)/P_n(s) = s + \mathcal{O}(s^2)$ as $s \rightarrow 0$ (which follows from the representations in (1.2) and (1.3) with $n = m$), we see that the integrand has a simple pole at the origin, with residue equal to 1. We shift \mathcal{C} across the pole at the origin, picking up the residue, and observe that the remaining integral over \mathcal{C} in the right half-plane vanishes. Hence, $I = 1$.

6. Concluding remarks

We have shown how to obtain the location and asymptotic approximation of zeros of the polynomials that constitute the diagonal Padé approximations to the exponential function. By using uniform Airy-type asymptotic expansions it is possible to give a clear description of the distribution of the zeros and their asymptotic approximation. We have compared our method with Saff & Varga's approach for locating the zeros of the polynomials.

We have used the asymptotic estimates of the zeros for obtaining an asymptotic estimate of functions that are smooth approximations to the block function, and we have proven a stability condition for these approximations.

In DRIVER & TEMME (1996b) more details are given on the non-diagonal case. Again we use Airy-type approximations to obtain the distribution of the zeros and the asymptotic estimates for the zeros of the polynomials.

7. Bibliography

- [1] ABRAMOWITZ, M. & I.A. STEGUN (1964). *Handbook of mathematical functions with formulas, graphs and mathematical tables*, Nat. Bur. Standards Appl. Series, **55**, U.S. Government Printing Office, Washington, D.C.
- [2] CHESTER, C., B. FRIEDMAN, B.& F. URSELL (1957). An extension of the method of steepest descent, *Proc. Cambridge Philos. Soc.* **53**, 599–611.
- [3] DRIVER, K.A. & TEMME, N.M. (1996a). On polynomials related with Hermite–Padé approximations to the exponential function, *manuscript*.
- [4] DRIVER, K.A. & TEMME, N.M. (1996b). Asymptotics and zero distribution of Padé polynomials associated with the exponential function, *manuscript*.
- [5] OLVER, F.W.J. (1974). *Asymptotics and special functions*, Academic Press, New York.
- [6] PERRON, O. (1950). *Die Lehre von den Kettenbrüchen*, Chelsea, New York.
- [7] SAFF, E.B. & R.S. VARGA, (1978). On the zeros and poles of Padé approximants to e^z . III, *Numer. Math.* **30** 241–266.
- [8] TEMME, N.M. (1996). *Special functions: An introduction to the classical functions of mathematical physics*, Wiley, New York.
- [9] WATSON, G.N. (1944). *A treatise on the theory of Bessel functions*, Cambridge University Press, London and New York.
- [10] WONG, R. (1989). *Asymptotic approximations of integrals*. Academic Press, New York.