



Self-dual morphological operators and filters

H.J.A.M. Heijmans

Department of Operations Research, Statistics, and System Theory

Report BS-R9409 February 1994

CWI is the National Research Institute for Mathematics and Computer Science. CWI is part of the Stichting Mathematisch Centrum (SMC), the Dutch foundation for promotion of mathematics and computer science and their applications.

SMC is sponsored by the Netherlands Organization for Scientific Research (NWO). CWI is a member of ERCIM, the European Research Consortium for Informatics and Mathematics.

Copyright © Stichting Mathematisch Centrum
P.O. Box 94079, 1090 GB Amsterdam (NL)
Kruislaan 413, 1098 SJ Amsterdam (NL)
Telephone +31 20 592 9333
Telefax +31 20 592 4199

Self-Dual Morphological Operators and Filters

Henk J.A.M. Heijmans

CWI

P.O. Box 94079, 1090 GB Amsterdam, The Netherlands

Abstract

The median operator is a nonlinear image transformation celebrated for its noise cleaning capacities. It treats the foreground and background of an image identically, i.e., it is self-dual. Unfortunately, the median operator has one major drawback: it is not idempotent. Even worse, subsequent iterations of a given image may lead to oscillations. This paper describes a general method for the construction of morphological operators which are self-dual. This construction is based upon the concept of a switch operator. Subsequently, the paper treats a class of operators, the so-called activity-extensive operators, which have the intriguing property that every sequence of iterates of a given image is pointwise monotone and therefore convergent. The underlying concept is that of the activity ordering. Every increasing, self-dual operator can be modified in such a way that it becomes activity-extensive. The sequence of iterates of this modification converges to a self-dual morphological filter.

AMS Subject Classification (1991): 68U10, 58F08

Keywords & Phrases: mathematical morphology, median operator, self-dual operator, idempotent operator, morphological filter, centre operator, activity ordering, activity-extensive operator, switch operator, persistent structure, iteration, cellular automata.

1. Introduction

The median operator is likely to be one of the most frequently used transformations in image processing which is nonlinear. In fact, this operator (see Section 3 for a formal definition) fits quite well in the framework of mathematical morphology [10, 13, 14, 3]; it is increasing, translation invariant, and can be decomposed as a finite union of erosions. The median operator is eminently suited for the elimination of noise [7]. Furthermore, it affects foreground and background noise in an identical manner. This is usually expressed by saying that the median operator is *self-dual*; see Section 2 for a formal definition. The major disadvantage of this operator is its non-idempotence: repeated application of the operator may change the outcome further or make previous changes undone. Even worse, iterating this operator may lead to oscillations. With regard to noise cleaning purposes, this is an undesirable property [14, 12].

Most of the operators discussed in this paper will be illustrated through their action on one particular noisy binary image X with a resolution of 128×128 pixels; see the left-most image in Figure 1. The second image in this figure shows $\mu(X)$, where μ is the median operator. The sequence of images obtained by iteration converges pixel-wise to a limit image $\mu^\infty(X)$; see the third image in Figure 1. The right-most image shows all pixels which have been flipped at

Report BS-R9409

ISSN 0924-0659

CWI

P.O. Box 94079, 1090 GB Amsterdam, The Netherlands



Fig. 1. From left to right: the original image X , the transform $\mu(X)$, where μ is the median operator, the limit image $\mu^\infty(X)$, and the set of pixels which have been flipped at at least one iteration step.

at least one iteration step; note that this set may be substantially larger than the symmetric difference of X and $\mu^\infty(X)$.

Two other important operators known from mathematical morphology, are the opening and the closing. These operators have interesting noise cleaning capabilities, too, and they are idempotent. Unfortunately, however, they lack self-duality. As a matter of fact, opening and closing are complementary in the sense that the opening affects noise particles with high intensities, whereas the closing affects noise particles with low intensities. The major question addressed in this paper is the following: “Can we find a general method to design morphological operators which are self-dual and idempotent?” A partial answer to this question was given in our previous work [3, Chapter 13], [4]. In this paper we will review this work and supplement it with a substantial number of new results and examples.

The basic idea underlying our approach is to construct self-dual operators ψ which are not necessarily idempotent, but which do satisfy the (weaker) constraint that they are *activity-extensive*. The latter means that the sequence of iterates $\psi, \psi^2, \psi^3, \dots$ is increasing with respect to the so-called activity ordering on the set of all operators; this ordering was introduced by Serra [14] and will be reviewed in Section 2. If ψ is an activity-extensive operator, then repeated application of ψ to an image yields a convergent sequence. This leaves us with the problem of finding self-dual operators which are activity-extensive. This part of the problem will constitute the main body of the present paper.

In Section 2 we recall the basic concepts which we need in this paper. As much as possible we adopt the terminology and notation of [3]. We point out that we use the terminology “operator” for an arbitrary transformation; the word “filter” is preserved for operators which are increasing and idempotent. In Section 3 we will make some general statements about self-dual operators; in particular, we will recall the notions of a (weighted) rank and median operator. We also establish a lattice isomorphism between the self-dual operators provided with the activity ordering at the one hand, and the anti-extensive operators with the pointwise ordering at the other. In Section 4 we define *switch operators* and indicate their importance in relation to our problem of finding increasing, self-dual operators. We study an important instance of a switch operator corresponding with the situation that we want to eliminate isolated noise points. Here the word “isolated” has to be interpreted in terms of some prescribed adjacency relation on the underlying point set. From Section 5 onwards, we will restrict attention to operators which are translation invariant. Under this restriction we can give a complete characterization of switch operators, and hence, of increasing, self-dual operators. Subsequently, in Section 6 we show how an arbitrary increasing, self-dual operator can be modified such that it becomes activity-extensive; then this modification can be used to construct a self-dual filter by iteration. A large number of examples

will be presented in Section 7. We end with some conclusions in Section 8; there we will also point out some open problems.

2. Terminology and basic facts

We presume that the reader is familiar with the basic notions of mathematical morphology. Throughout this paper we are mainly concerned with the space of discrete binary images $\mathcal{P}(\mathbb{Z}^2)$. If ϕ, ψ are two operators on this space, then $\phi \leq \psi$ means $\phi(X) \subseteq \psi(X)$, for every $X \subseteq \mathbb{Z}^2$. If $\psi_i, i \in I$ is a collection of operators then $\bigwedge_{i \in I} \psi_i$ and $\bigvee_{i \in I} \psi_i$ denote the infimum and supremum of this collection, respectively; see e.g. [3].

If $X \subseteq \mathbb{Z}^2$ and $h \in \mathbb{Z}^2$ then X_h denotes the translation of the set X along the vector h . Sometimes we interpret X as an indicator function. Then $X(h) = 1$ if $h \in X$ and 0 otherwise. Given a translation invariant operator ψ on $\mathcal{P}(\mathbb{Z}^2)$, we say that ψ is a *finite-window operator* if there exists a finite set $M \subseteq \mathbb{Z}^2$ such that $h \in \psi(X) \iff h \in \psi(X \cap A_h)$ for every $h \in \mathbb{Z}^2$, $X \subseteq \mathbb{Z}^2$ and $A \supseteq M$. If ψ is increasing then it suffices to consider $A = M$.

In [14] Serra introduces a partial ordering on the complete lattice of all operators on \mathcal{L} , where \mathcal{L} is a complete Boolean lattice. Here we specialize to the case $\mathcal{L} = \mathcal{P}(\mathbb{Z}^2)$. Given two operators ϕ, ψ on $\mathcal{P}(\mathbb{Z}^2)$, we say that ψ is *more active than* ϕ , written $\phi \preceq \psi$, if

$$\text{id} \wedge \psi \leq \text{id} \wedge \phi \quad \text{and} \quad \text{id} \vee \psi \geq \text{id} \vee \phi.$$

Here id is the identity operator which maps every set onto itself. The infimum of a collection of operators $\psi_i, i \in I$, with respect to the activity ordering is denoted by $\bigwedge_{i \in I} \psi_i$. It is given by the expression

$$\bigwedge_{i \in I} \psi_i = (\text{id} \wedge \bigvee_{i \in I} \psi_i) \vee \bigwedge_{i \in I} \psi_i.$$

This operator is sometimes called the *centre* of the operators ψ_i . Dually, their supremum, denoted by $\bigvee_{i \in I} \psi_i$, is given by

$$\bigvee_{i \in I} \psi_i = (\nu \wedge \bigvee_{i \in I} \psi_i) \vee \bigwedge_{i \in I} \psi_i,$$

and is called the *anti-centre*. Here ν is the complement operator given by

$$\nu(X) = X^c.$$

It is obvious that id and ν are the least and greatest operator, respectively, with respect to the activity ordering. The *negative* ψ^* of an operator ψ is given by $\psi^* = \nu\psi\nu$, i.e., $\psi^*(X) = (\psi(X^c))^c$.

The operator ψ is called *activity-extensive* if

$$\psi^n \preceq \psi^{n+1}, \quad n = 0, 1, 2, \dots$$

This means that, for a fixed set $X \subseteq \mathbb{Z}^2$, the sequence $\psi^n(X)$ is pointwise monotone: the sequence $\psi^n(X)(h)$, where $h \in \mathbb{Z}^2$ is given, is monotone, either increasing or decreasing. As such sequences consist exclusively of 0's and 1's, they must be of the form $0, 0, \dots, 0, 1, 1, \dots$ or $1, 1, \dots, 1, 0, 0, \dots$. We call $\{\psi^n(X) \mid n = 0, 1, 2, \dots\}$ the *orbit* starting at $\psi^0(X) = X$.

3. Self-dual operators

Before entering a general discussion on the construction of self-dual operators we present some examples based on Boolean functions; see also [3, §4.5]. Let $A = \{a_1, a_2, \dots, a_n\}$ be a finite structuring element, and let b be the threshold function with realization vector $(w_1, \dots, w_n \mid s)$, i.e.,

$$b(u_1, \dots, u_n) = \left[\sum_{i=1}^n w_i u_i \geq s \right];$$

here $w_1, \dots, w_n, s \in \mathbb{Z}$. Here we use the following convention. The expression $[P]$ equals 1 if P is true and 0 if P is false. Now

$$\psi_b(X) = \{x \in \mathbb{Z}^2 \mid \sum_{i=1}^n w_i X(x + a_i) \geq s\},$$

is an increasing, translation invariant, morphological operator. We call this operator *weighted rank operator*. If the weights w_i as well as the threshold s are positive integers, then this operator has the following interpretation: the point x lies in $\psi_b(X)$ if the s th value of the sequence which is obtained by putting the values $X(x + a_i)$ (counted w_i times) in decreasing order equals one. If $w_i \geq 0$ for every i , then ψ_b is an increasing operator. Its negative ψ_b^* is given by

$$\psi_b^*(X) = \{x \in \mathbb{Z}^2 \mid \sum_{i=1}^n w_i X(x + a_i) \geq \sum_{i=1}^n w_i - s + 1\}.$$

It follows immediately that ψ_b is self-dual if and only if $2s = \sum_{i=1}^n w_i + 1$. In this case ψ_b is called a *weighted median operator*. A weighted rank operator can be represented by a matrix containing the weights w_i in which the position of the origin is marked; see Example 3.1 below for an illustration.

If all weights w_i have the value 1, and if $s \in \{1, 2, \dots, n\}$, then ψ_b is called *rank operator*; this operator is denoted by $\rho_{A,s}$. It is easy to show that $h \in \rho_{A,s}(X)$ if and only if $X \cap A_h$ has at least cardinality s . In particular,

$$\rho_{A,1}(X) = X \oplus \check{A} \quad \text{and} \quad \rho_{A,n}(X) = X \ominus A.$$

Furthermore,

$$\rho_{A,n} \leq \rho_{A,n-1} \leq \dots \leq \rho_{A,1}.$$

If n is odd and $s = (n + 1)/2$ then $\rho_{A,s}$ is self-dual; we use the notation

$$\mu_A = \rho_{A,(n+1)/2},$$

if n is odd, and we call μ_A the *median operator*.

Let ψ be an operator on $\mathcal{P}(\mathbb{Z}^2)$ with negative ψ^* . The centre of ψ and ψ^* is given by

$$\gamma = (\text{id} \wedge (\psi \vee \psi^*)) \vee (\psi \wedge \psi^*).$$

The anti-centre is

$$\kappa = (\nu \wedge (\psi \vee \psi^*)) \vee (\psi \wedge \psi^*),$$

where ν is the complement operator. It is easy to see that both γ and κ are self-dual. Furthermore, γ is increasing if ψ is increasing.

3.1. Example. Let A be the 3×3 square centred at the origin, and let $\rho_{A,s}$, $1 \leq s \leq 9$, be the associated rank operators. The negative of $\rho_{A,s}$ is $\rho_{A,10-s}$, and $\rho_{A,s} \leq \rho_{A,10-s}$ if $s \geq 5$. Define θ_s as the centre of $\rho_{A,s}$ and $\rho_{A,10-s}$; hence

$$\theta_s = (\text{id} \wedge \rho_{A,10-s}) \vee \rho_{A,s}, \quad s = 5, 6, 7, 8, 9.$$

Note that $\theta_5 = \mu_A$, the median operator. Obviously, every θ_s is self-dual. Furthermore, it is easy to show that

$$\text{id} = \theta_9 \preceq \theta_8 \preceq \theta_7 \preceq \theta_6 \preceq \theta_5.$$

In Figure 2 the action of these operators is illustrated by means of an example. It is easy to show that θ_k corresponds with the weighted median operator represented by the matrix

$$w = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 2k-9 & 1 \\ 1 & 1 & 1 \end{pmatrix};$$

the centre of the matrix corresponds with the origin, and the threshold s equals k .





Fig. 2. From left to right and top to bottom: original image X and its transforms $\theta_8(X), \theta_7(X), \theta_6(X), \theta_5(X)$. This picture illustrates nicely that $\theta_8 \preceq \theta_7 \preceq \theta_6 \preceq \theta_5$.

Let us have a closer look at the median operator μ_A defined in the previous example. When a point h lies in the complement of X , then $h \in \mu_A(X)$ if it is surrounded by at least five neighbours which lie in X like in one of the following 56 configurations:



Note that there correspond eight rotations with every structuring element depicted. Dually, when h lies in X , then h lies outside $\mu_A(X)$ if it has at least five neighbours in the complement of X .

Consider an arbitrary self-dual operator ψ . We denote the points which are contained in X but not in $\psi(X)$ by $\sigma(X)$. Hence

$$\sigma(X) = X \setminus \psi(X) = X \cap (\psi(X))^c = X \cap \psi(X^c). \quad (3.1)$$

Here we have used that ψ is self-dual, hence

$$\psi(X^c) = (\psi(X))^c.$$

It is easy to check that ψ can be recovered from σ by means of the formula:

$$\psi(X) = (X \setminus \sigma(X)) \cup \sigma(X^c). \quad (3.2)$$

Motivated by the above observations we define two mappings on \mathcal{O} , the lattice of operators on $\mathcal{P}(\mathbb{Z}^2)$.

3.2. Definition. Let the mappings $\Sigma, \Psi : \mathcal{O} \rightarrow \mathcal{O}$ be defined by

$$\Sigma(\psi) = \text{id} \wedge \psi\nu, \quad \psi \in \mathcal{O}, \quad (3.3)$$

$$\Psi(\sigma) = (\text{id} \wedge \nu\sigma) \vee \sigma\nu, \quad \sigma \in \mathcal{O}. \quad (3.4)$$

Observe that (3.1) and (3.2) can be reformulated as $\sigma = \Sigma(\psi)$ and $\psi = \Psi(\sigma)$, respectively. Furthermore

$$\sigma \leq \text{id} \iff \sigma\nu \leq \nu \iff \nu \leq \nu\sigma, \quad (3.5)$$

for every operator σ . This means in particular that $\Psi(\sigma)$ is the centre of $\nu\sigma$ and $\sigma\nu$ if σ is anti-extensive.

We denote by \mathcal{O}_{sd} and \mathcal{O}_{ae} the self-dual and anti-extensive operators on $\mathcal{P}(\mathbb{Z}^2)$, respectively. It is obvious that (\mathcal{O}_{ae}, \leq) is a complete sublattice of (\mathcal{O}, \leq) . From the fact that both the centre and the anti-centre of a family of self-dual operators is self-dual (cf. [3, Prop. 3.42]), we conclude that $(\mathcal{O}_{sd}, \preceq)$ is a complete sublattice of (\mathcal{O}, \preceq) .

3.3. Proposition.

- (a) $\Sigma : (\mathcal{O}_{sd}, \preceq) \rightarrow (\mathcal{O}_{ae}, \leq)$ is increasing.
- (b) $\Psi : (\mathcal{O}_{ae}, \leq) \rightarrow (\mathcal{O}_{sd}, \preceq)$ is increasing.

PROOF. (a): It is evident that $\Sigma(\psi)$ is anti-extensive. Let ψ_1, ψ_2 be self-dual operators and $\psi_1 \preceq \psi_2$. Then $\text{id} \vee \psi_1 \leq \text{id} \vee \psi_2$, hence $\nu \vee \psi_1\nu = (\text{id} \vee \psi_1)\nu \leq (\text{id} \vee \psi_2)\nu = \nu \vee \psi_2\nu$. Taking the infimum with id we get $\text{id} \wedge (\nu \vee \psi_1\nu) \leq \text{id} \wedge (\nu \vee \psi_2\nu)$, that is, $\text{id} \wedge \psi_1\nu \leq \text{id} \wedge \psi_2\nu$.

(b): First we must show that $\Psi(\sigma)$ is self-dual if $\sigma \leq \text{id}$. This is done as follows:

$$\begin{aligned} \Psi(\sigma)^* &= \nu\Psi(\sigma)\nu \\ &= \nu((\text{id} \wedge \nu\sigma) \vee \sigma\nu)\nu \\ &= \nu((\nu \wedge \nu\sigma\nu) \vee \sigma) \\ &= \nu(\nu \wedge \nu\sigma\nu) \wedge \nu\sigma \\ &= (\text{id} \vee \sigma\nu) \wedge \nu\sigma \\ &= (\text{id} \wedge \nu\sigma) \vee \sigma\nu \\ &= \Psi(\sigma); \end{aligned}$$

here we have used that $\sigma\nu \leq \nu\sigma$ by the anti-extensivity of σ . To show that Ψ is increasing, let $\sigma_1 \leq \sigma_2 \leq \text{id}$, and let $\Psi(\sigma_1) = \psi_1$, $\Psi(\sigma_2) = \psi_2$. We show that $\psi_1 \preceq \psi_2$. First,

$$\begin{aligned} \text{id} \wedge \psi_1 &= \text{id} \wedge ((\text{id} \wedge \nu\sigma_1) \vee \sigma_1\nu) \\ &= (\text{id} \wedge (\text{id} \wedge \nu\sigma_1)) \vee (\text{id} \wedge \sigma_1\nu) \\ &= (\text{id} \wedge \nu\sigma_1) \vee (\text{id} \wedge \sigma_1\nu) \\ &= \text{id} \wedge \nu\sigma_1, \end{aligned}$$

since $\nu\sigma_1 \geq \sigma_1\nu$ by (3.5). Now $\sigma_1 \leq \sigma_2$ implies that $\text{id} \wedge \psi_1 \geq \text{id} \wedge \psi_2$. We also find

$$\text{id} \vee \psi_1 = \text{id} \vee \sigma_1\nu.$$

Clearly, $\sigma_1 \leq \sigma_2$ implies that $\text{id} \vee \psi_1 \leq \text{id} \vee \psi_2$. Thus we find that $\psi_1 \preceq \psi_2$. \blacksquare

3.4. Proposition. *The mappings $\Sigma : (\mathcal{O}_{sd}, \preceq) \rightarrow (\mathcal{O}_{ae}, \leq)$ and $\Psi : (\mathcal{O}_{ae}, \leq) \rightarrow (\mathcal{O}_{sd}, \preceq)$ are bijective (onto and one-one). Furthermore, these mappings are each other's inverses, i.e.,*

$$\Sigma(\Psi(\sigma)) = \sigma, \quad \text{for } \sigma \in \mathcal{O}_{ae}; \quad (3.6)$$

$$\Psi(\Sigma(\psi)) = \psi, \quad \text{for } \psi \in \mathcal{O}_{sd}. \quad (3.7)$$

PROOF. We prove (3.6) and leave (3.7) as an exercise for the reader. Note that the validity of these two relations implies the bijectivity of the mappings Σ and Ψ .

Let σ be anti-extensive; then

$$\begin{aligned} \Sigma(\Psi(\sigma)) &= \text{id} \wedge \Psi(\sigma)\nu = \text{id} \wedge ((\text{id} \wedge \nu\sigma) \vee \sigma\nu)\nu \\ &= \text{id} \wedge ((\nu \wedge \nu\sigma\nu) \vee \sigma) = (\text{id} \wedge \nu \wedge \nu\sigma\nu) \vee (\text{id} \wedge \sigma) \\ &= \text{id} \wedge \sigma = \sigma. \end{aligned}$$

Here we have used that $\text{id} \wedge \nu = o$, the operator which maps every set onto the empty set. \blacksquare

The following relations are easily derived from the previous result.

$$\Sigma(\bigvee_{i \in I} \psi_i) = \bigvee_{i \in I} \Sigma(\psi_i); \quad (3.8)$$

$$\Sigma(\bigwedge_{i \in I} \psi_i) = \bigwedge_{i \in I} \Sigma(\psi_i); \quad (3.9)$$

$$\Psi(\bigvee_{i \in I} \sigma_i) = \bigvee_{i \in I} \Psi(\sigma_i); \quad (3.10)$$

$$\Psi(\bigwedge_{i \in I} \sigma_i) = \bigwedge_{i \in I} \Psi(\sigma_i). \quad (3.11)$$

Here we prove (3.8). Let ψ_i be self-dual operators for $i \in I$, and let $\sigma_i = \Sigma(\psi_i)$. Since $\psi_j \preceq \bigvee_{i \in I} \psi_i$ for $j \in I$, we obtain $\sigma_j \leq \Sigma(\bigvee_{i \in I} \psi_i)$. This implies that $\bigvee_{i \in I} \sigma_i \leq \Sigma(\bigvee_{i \in I} \psi_i)$, hence that $\sigma_j \leq \bigvee_{i \in I} \sigma_i \leq \Sigma(\bigvee_{i \in I} \psi_i)$. Applying Ψ we arrive at

$$\psi_j \preceq \Psi(\bigvee_{i \in I} \sigma_i) \preceq \bigvee_{i \in I} \psi_i.$$

Taking the activity supremum over all j at the left-hand side, we get

$$\bigvee_{i \in I} \psi_i \preceq \Psi(\bigvee_{i \in I} \sigma_i) \preceq \bigvee_{i \in I} \psi_i.$$

But then all three terms must be equal, meaning that

$$\bigvee_{i \in I} \psi_i = \Psi\left(\bigvee_{i \in I} \sigma_i\right).$$

Applying Σ on both sides and using that $\Sigma\Psi = \text{id}$, gives

$$\Sigma\left(\bigvee_{i \in I} \psi_i\right) = \bigvee_{i \in I} \Sigma(\psi_i).$$

The least and greatest operator with respect to the activity ordering are id and ν , respectively. It is easy to check that $\Sigma(\text{id}) = o$ and $\Sigma(\nu) = \text{id}$, the least resp. greatest anti-extensive operator.

3.5. Remark. Let ψ be an operator, not necessarily self-dual. The centre $\psi \wedge \psi^*$ and the anti-centre $\psi \vee \psi^*$ are self-dual. A straightforward calculation shows that


$$\begin{aligned} \Sigma(\psi \wedge \psi^*) &= \Sigma(\psi \wedge \psi^*) = \text{id} \wedge (\psi \nu \wedge \nu \psi); \\ \Sigma(\psi \vee \psi^*) &= \Sigma(\psi \vee \psi^*) = \text{id} \wedge (\psi \nu \vee \nu \psi). \end{aligned}$$

The results given so far provide a general method to construct self-dual operators. We give some special attention to the class of translation invariant operators. It is a well-known fact [1], [3, Section 4.2] that every translation invariant operator ψ can be decomposed as a union of hit-or-miss operators. Recall that the hit-or-miss operator with structuring elements A, B is the map $X \mapsto X \otimes (A, B)$, where

$$X \otimes (A, B) = \{h \in \mathbb{Z}^2 \mid A_h \subseteq X, B_h \subseteq X^c\}.$$

In order that $X \otimes (A, B) \subseteq X$ for every X we must demand that $0 \in A$. Thus we may conclude that every anti-extensive operator σ can be written as a union of hit-or-miss operators $X \mapsto X \otimes (A, B)$ with $0 \in A$.

3.6. Example. Consider the following eight structuring elements (A contains the central black pixel, B contains the white pixels) respectively called $(A_1, B_1), \dots, (A_8, B_8)$. The underlined (central) pixel is located at the origin.



In [3, Example 4.6] these same patterns are used to obtain the pseudo-convex hull of a discrete set by iterations of the corresponding thickenings $X \odot (A, B) = X \cup X \otimes (A, B)$. Here we define

$$\sigma(X) = \bigcup_{k=1}^8 X \otimes (A_k, B_k).$$

Note that $\sigma \leq \text{id}$. Let $\psi = \Psi(\sigma)$ be the corresponding self-dual operator. In Figure 3 we depict an image X and its transform $\psi(X)$. If we compute the sequence of iterates $\psi^n(X)$ we find that this sequence converges to a period-2 orbit consisting of two images Y and $\psi(Y)$. In Figure 3 we depict Y and the symmetric difference $Y \Delta \psi(Y)$.



Fig. 3. The first image is X , the second $\psi(X)$, with ψ as defined in Example 3.6. The sequence $\psi^n(X)$ converges to a period-2 orbit containing two images. The third image represents one of these two images Y . The right-most image shows the symmetric difference $Y \Delta \psi(Y)$.



Fig. 4. From left to right: the original image X , the transformed image $\psi(X)$, and the limit $\psi^\infty(X)$. Here $\psi = \alpha \Upsilon \beta$, where α is the structural opening by the 3×3 square and β is the negative closing.

3.7. Example. Let α be an opening and $\sigma(X) = X \setminus \alpha(X)$; in other words, $\sigma = \text{id} \wedge \nu \alpha$. The corresponding self-dual operator $\psi = \Psi(\sigma)$ equals $\psi = (\nu \wedge \beta) \vee \alpha = (\nu \vee \alpha) \wedge \beta$, where β is the negative of α . Note that ψ is the anti-centre of α and β , i.e., $\psi = \alpha \Upsilon \beta$. In Figure 4 we have illustrated $\alpha \Upsilon \beta$, where α is the translation invariant, structural opening by the 3×3 square. The first image is the original image X , the second image is $\psi(X)$. In this particular case, the sequence $\psi^n(X)$ tends to a limit $\psi^\infty(X)$ as $n \rightarrow \infty$; this is the right-most image in Figure 4.

There do also exist images Y such that $\psi^n(Y)$ converges to a period-2 orbit. For example, if Y is such that $\alpha(Y) = \alpha(Y^c) = \emptyset$, then $\sigma(Y) = Y$ and $\sigma(Y^c) = Y^c$. Thus $\psi(Y) = Y^c$ and $\psi(Y^c) = Y$. This means that the pair $\{Y, Y^c\}$ forms a period-2 orbit.

4. Switch operators

The self-dual operators ψ in Examples 3.6–3.7 are not increasing. In this section we will answer the following question: which conditions must σ satisfy in order that the corresponding self-dual operator $\psi = \Psi(\sigma)$ is increasing. Assume that $X \subseteq Y$. From the interpretation of $\sigma(X)$ as the points in X which are deleted by ψ we make the following observations. If a point $h \in X$ is not contained in $\sigma(X)$ then it must not be contained in $\sigma(Y)$, for otherwise $h \in \psi(X)$ but $h \notin \psi(Y)$. This leads us to the following condition:

- (i) $h \in X, h \notin \sigma(X) \Rightarrow h \notin \sigma(Y)$.

A similar condition must hold for background points:

(ii) $h \in Y^c, h \notin \sigma(Y^c) \Rightarrow h \notin \sigma(X^c)$.

Suppose, finally, that a point h lies in Y but not in X . To avoid that h is mapped out of Y and at the same time into X we must assume:

(iii) $h \in \sigma(X^c) \Rightarrow h \notin \sigma(Y)$.

It is easy to see that (i) and (ii) amount to the following condition:

$$(S1) \ X \subseteq Y \text{ implies that } X \cap \sigma(Y) \subseteq \sigma(X).$$

$$(S2) \ \sigma(X \cup \{h\}) \cap \sigma(X^c \cup \{h\}) = \emptyset, \quad h \in \mathbb{Z}^2, X \subseteq \mathbb{Z}^2.$$

We show that (iii) follows from (S1)-(S2); note, however, that (iii) is not equivalent to (S2). Assume that $h \in \sigma(X^c)$ and $h \in \sigma(Y)$; we must show that this leads to a contradiction. From (S2) we conclude that $h \notin \sigma(X \cup \{h\})$. As $X \cup \{h\} \subseteq Y$ we conclude from (S1) that

$$(X \cup \{h\}) \cap \sigma(Y) \subseteq \sigma(X \cup \{h\}).$$

But then $h \in \sigma(X \cup \{h\})$, a contradiction.

4.1. Definition. An anti-extensive operator σ is called a *switch operator* if (S1) and (S2) hold.

The adjective ‘switch’ indicates that the operator σ yields all points which switch value from 1 to 0 (points in $\sigma(X)$) or from 0 to 1 (points in $\sigma(X^c)$) by application of the self-dual operator $\psi = \Psi(\sigma)$.

In the previous paragraphs we have established the following result.

4.2. Theorem. *Let σ be anti-extensive and $\psi = \Psi(\sigma)$; then ψ is increasing if and only if σ is a switch operator.*

If ψ is a self-dual increasing operator, and η is increasing and extensive, then

$$\psi' = (\text{id} \wedge \psi\eta) \vee \psi\eta^*$$

is the centre of $\psi\eta$ and its negation $\psi\eta^*$; note that $\eta^* \leq \eta$. Therefore, ψ' is self-dual and increasing too. The following result expresses the corresponding switch operator $\Sigma(\psi')$ in terms of $\Sigma(\psi)$ and η .

4.3. Proposition. *Let σ be a switch operator, $\psi = \Psi(\sigma)$, and let η be increasing and extensive; then $\sigma' = \text{id} \wedge \sigma\eta$ is a switch operator too, and $\Psi(\sigma') = (\text{id} \wedge \psi\eta) \vee \psi\eta^*$. Furthermore, $\sigma' \leq \sigma$ and $\psi' \preceq \psi$.*

PROOF. We show that $\Psi(\sigma') = (\text{id} \wedge \psi\eta) \vee \psi\eta^*$, and therefore increasing. This yields automatically that σ' is a switch operator.

$$\begin{aligned} \Psi(\sigma') &= (\text{id} \wedge \nu\sigma') \vee \sigma'\nu \\ &= (\text{id} \wedge \nu(\sigma\eta \wedge \text{id})) \vee (\sigma\eta \wedge \text{id})\nu \\ &= (\text{id} \wedge (\nu\sigma\eta \vee \nu)) \vee (\sigma\eta\nu \wedge \nu) \\ &= (\text{id} \wedge \nu\sigma\eta) \vee (\sigma\eta\nu \wedge \nu). \end{aligned}$$

Now we substitute that $\sigma = \Sigma(\psi) = \text{id} \wedge \psi\nu$, and use that $\nu\eta \leq \nu \leq \eta\nu$:

$$\begin{aligned} \Psi(\sigma') &= (\text{id} \wedge \nu(\text{id} \wedge \psi\nu)\eta) \vee ((\text{id} \wedge \psi\nu)\eta\nu \wedge \nu) \\ &= (\text{id} \wedge (\nu \vee \psi)\eta) \vee ((\eta\nu \wedge \psi\eta^*) \wedge \nu) \\ &= ((\text{id} \wedge \nu\eta) \vee (\text{id} \wedge \psi\eta)) \vee (\psi\eta^* \wedge \nu) \\ &= (\text{id} \wedge \psi\eta) \vee (\nu \wedge \psi\eta^*). \end{aligned}$$

Since $\eta^* \leq \eta$ we have $\text{id} \wedge \psi\eta^* \leq \text{id} \wedge \psi\eta$; thus

$$\begin{aligned}\Psi(\sigma') &= (\text{id} \wedge \psi\eta) \vee (\eta \wedge \psi\eta^*) \vee (\text{id} \wedge \psi\eta^*) \\ &= (\text{id} \wedge \psi\eta) \vee \psi\eta^*.\end{aligned}$$

This concludes the first part of the assertion.

Using (S1) and the extensivity of η we get that $X \cap \sigma(\eta(X)) \subseteq \sigma(X)$ for every X , hence $\sigma' \leq \sigma$. Now Proposition 3.3 gives that $\psi' \preceq \psi$. This concludes the proof. \blacksquare

From the following section onwards we will restrict ourselves to operators which are translation invariant. Before we do so we will present a class of self-dual operators (and their corresponding switch operators) based on the notion of adjacency.

Let E be an arbitrary non-empty set and let \sim be a symmetric relation on $E \times E$. Thus $x \sim y$ iff $y \sim x$, for $x, y \in E$. In this case we say that x and y are *adjacent*. The set of points adjacent to x is denoted by $\text{adj}(x)$:

$$\text{adj}(x) = \{y \in E \mid x \sim y\}.$$

N.B. It is not assumed that \sim is reflexive; hence $x \sim x$ may not be satisfied.

When the set E is finite we can supply it with a graph structure by considering the unordered pairs $\{x, y\}$ with $x \sim y$ as edges. We define the operator

$$\sigma(X) = \{h \in X \mid \text{adj}(h) \neq \emptyset \text{ and } \text{adj}(h) \cap X = \emptyset\}. \quad (4.1)$$

4.4. Proposition. σ is a switch operator.

PROOF. First we prove (S1). Let $X \subseteq Y$; we show that $X \cap \sigma(Y) \subseteq \sigma(X)$. If $x \in X \cap \sigma(Y)$, then $x \in Y$ and $\text{adj}(x) \cap Y = \emptyset$. This implies that $\text{adj}(x) \cap X = \emptyset$, too, hence $x \in \sigma(X)$.

To prove (S2), assume that $h \in \sigma(X \cup \{h\}) \cap \sigma(X^c \cup \{h\})$. Then $\text{adj}(h) \neq \emptyset$ and $\text{adj}(h) \cap (X \cup \{h\}) = \emptyset$ and $\text{adj}(h) \cap (X^c \cup \{h\}) = \emptyset$. But this yields that $\text{adj}(h) = \emptyset$, a contradiction. This concludes the proof. \blacksquare

From this point onwards we assume that

$$\text{adj}(h) \neq \emptyset, \quad \text{for every } h \in E.$$

Let ψ be the increasing, self-dual operator on $\mathcal{P}(E)$ associated with σ . The action of ψ can be described as follows: ψ switches the state of a point if and only if all its adjacent points have a different state. Or, to put it differently, ψ switches the state of the isolated points. Therefore, if x, y are adjacent and have the same state, then both points are left unaltered by ψ (and its subsequent iterates).

Define the dilation δ by

$$\delta(X) = \{h \in E \mid \text{adj}(h) \cap X \neq \emptyset\}. \quad (4.2)$$

The negative erosion $\varepsilon = \delta^*$ is then given by

$$\varepsilon(X) = \{h \in E \mid \text{adj}(h) \subseteq X\}. \quad (4.3)$$

Note that these two operators are known from the context of *graph morphology* [6].

Evidently, the pair (ε, δ) constitutes an adjunction. It is easy to check that $\nu\sigma = \nu \vee \delta$ and that $\sigma\nu = \nu \wedge \varepsilon$. Substitution in $\psi = (\text{id} \wedge \nu\sigma) \vee \sigma\nu$ gives us

$$\psi = (\text{id} \wedge \delta) \vee (\nu \wedge \varepsilon).$$

Using that $\varepsilon \leq \delta$ we get $\text{id} \wedge \varepsilon \leq \text{id} \wedge \delta$, and therefore

$$\psi = (\text{id} \wedge \delta) \vee (\nu \wedge \varepsilon) \vee (\text{id} \wedge \varepsilon),$$

that is

$$\psi = (\text{id} \wedge \delta) \vee \varepsilon = (\text{id} \vee \varepsilon) \wedge \delta. \quad (4.4)$$

We present a condition on the adjacency relation which guarantees that the operator ψ is idempotent. In general, however, ψ will not be idempotent. In Figure 5 for example, we depict a binary image on a graph whose edges represent the adjacency relation, which shows period 2 oscillations under iteration of ψ .



Fig. 5. A period-2 orbit.

4.5. Definition. Let $x, y, z \in E$; we call $\{x, y, z\}$ an *adjacency triple* if $x \sim y \sim z \sim x$.

The points x, y, z need not be distinct. In fact, if $x \sim x$, then $\{x, x, x\}$ is an adjacency triple.

The following intuitive argument shows that ψ is idempotent if every point in E is a member of at least one adjacency triple. Suppose namely that point $x \in X$ has no adjacent points inside X . Since it is a member of an adjacency triple $\{x, y, z\}$ with $y, z \in X^c$, the point x will be mapped outside X by ψ . However, the points y, z stay inside X^c because they constitute an adjacent pair. In subsequent iterations the points x, y, z will stay in X^c .

4.6. Proposition. Assume that every point in E is a member of at least one adjacency triple, then ψ is idempotent.

PROOF. We show that $\psi \leq \psi^2$. Then, taking the negative at both sides and using that $\psi^* = \psi$ we get $\psi \geq \psi^2$ as well.

Let $h \in \psi(X)$. There are two alternatives: $h \in X$ and $h \notin X$. Assume first that $h \in X$; then $h \notin \sigma(X)$, which means that $\text{adj}(h) \cap X \neq \emptyset$. Let $x \in \text{adj}(h) \cap X$, then $h \in \text{adj}(x) \cap X$, and therefore $x \notin \sigma(X)$. We conclude that $x \in \psi(X)$. But then $h \notin \sigma(\psi(X))$, meaning that $h \in \psi^2(X)$.

Assume next that $h \notin X$. Then $h \in \sigma(X^c)$ and thus $\text{adj}(h) \cap X^c = \emptyset$. By assumption, h is a member of some adjacency triple: $h \sim x \sim y \sim h$. Necessarily, $x, y \in X$. But then $h, x, y \in \psi(X)$ and it follows that $h \in \psi^2(X)$ (as well as $x, y \in \psi^2(X)$). This concludes the proof. ■

Apparently, the adjacency triple condition is not satisfied in the example of Figure 5.

Let us apply this result to the case that $E = \mathbb{Z}^2$ (or \mathbb{Z}^d), where \sim is translation invariant, i.e., $x \sim y$ implies that $x + h \sim y + h$ for all $x, y, h \in \mathbb{Z}^2$. Putting $A = \text{adj}(0)$, we have $\text{adj}(h) = A_h$. The fact that the adjacency relation is symmetric implies that A is symmetric;

i.e., $A = \check{A}$, where $\check{A} = \{-a \mid a \in A\}$. The adjacency triple condition can be formulated as follows:

$$0 \in A \oplus A \oplus A.$$

Furthermore

$$\delta(X) = X \oplus A, \quad \varepsilon(X) = X \ominus A.$$

We arrive at the following result.

4.7. Theorem. *Let $A \subseteq \mathbb{Z}^2$ be a symmetric structuring element ($A = \check{A}$) which satisfies*

$$0 \in A \oplus A \oplus A. \quad (4.5)$$

Then the increasing, translation invariant operator ψ given by

$$\psi(X) = (X \cap (X \oplus A)) \cup (X \ominus A) \quad (4.6)$$

is self-dual and idempotent.

Note that the conditions on A are satisfied by

$$A = \begin{array}{|c|} \hline \bullet & \bullet & \bullet \\ \hline \bullet & \vdots & \bullet \\ \hline \bullet & \bullet & \bullet \\ \hline \end{array}$$

The operator ψ in Theorem 4.7 switches all pixels whose state is different from that of all their 8-neighbours (i.e., the isolated pixels); see also example 10 in Section 7.

5. Switch operators which are translation invariant

Arriving at this point, the most obvious question is: how to design switch operators? Below we shall give a complete characterization of translation invariant switch operators. But first we give a characterization of the kernel of a translation invariant self-dual operator. Recall that the kernel of a translation invariant operator ψ on $\mathcal{P}(\mathbb{Z}^2)$ is defined as [13, 3]:

$$\mathcal{V}(\psi) = \{A \subseteq \mathbb{Z}^2 \mid 0 \in \psi(A)\}.$$

Every increasing, translation invariant operator ψ can be written as a union of erosions:

$$\psi(X) = \bigcup_{A \in \mathcal{V}(\psi)} X \ominus A. \quad (5.1)$$

Recall also that [3]

$$\mathcal{V}\left(\bigwedge_{i \in I} \psi_i\right) = \bigcap_{i \in I} \mathcal{V}(\psi_i) \quad \text{and} \quad \mathcal{V}\left(\bigvee_{i \in I} \psi_i\right) = \bigcup_{i \in I} \mathcal{V}(\psi_i). \quad (5.2)$$

5.1. Proposition. *Let ψ be a translation invariant operator; then ψ is self-dual if and only if*

$$A \in \mathcal{V}(\psi) \iff A^c \notin \mathcal{V}(\psi).$$

If, moreover, ψ is increasing, then

$$A, B \in \mathcal{V}(\psi) \Rightarrow A \cap B \neq \emptyset.$$

PROOF. $A \in \mathcal{V}(\psi)$ iff $0 \in \psi(A) = \psi^*(A) = (\psi(A^c))^c$. This implies the first assertion.

Assume that ψ is also increasing. If $A, B \in \mathcal{V}(\psi)$ and $A \cap B = \emptyset$, then $A \subseteq B^c$. This implies that $0 \in \psi(A) \subseteq \psi(B^c)$, a contradiction. \blacksquare

Assume that ψ is an increasing, translation invariant, self-dual operator, and let $\sigma = \Sigma(\psi)$. Using (5.1) we get

$$\sigma(X) = \bigcup_{A \in \mathcal{V}(\psi)} X \cap (X^c \ominus A).$$

If $0 \in A$ then $X \cap (X^c \ominus A) = \emptyset$, and thus we find

$$\sigma(X) = \bigcup_{A \in \mathcal{V}(\psi), 0 \notin A} X \cap (X^c \ominus A).$$

Using $\sigma = \text{id} \wedge \psi \nu$ and (5.2) we obtain $\mathcal{V}(\sigma) = \mathcal{V}(\text{id}) \cap \mathcal{V}(\psi \nu)$. This is equivalent to

$$A \in \mathcal{V}(\psi), 0 \notin A \iff 0 \in \sigma(A^c) \iff A^c \in \mathcal{V}(\sigma).$$

Define

$$a(\sigma) = \{A \subseteq \mathbb{Z}^2 \mid A^c \in \mathcal{V}(\sigma)\}. \quad (5.3)$$

Proposition 5.1 gives that $A \cap B \neq \emptyset$ for $A, B \in a(\sigma)$. We have thus shown the following result.

5.2. Proposition. *Let σ be a switch operator and let $a(\sigma)$ be given by (5.3); then*

$$0 \notin A \quad \text{and} \quad A \cap B \neq \emptyset, \quad \text{for } A, B \in a(\sigma). \quad (5.4)$$

Furthermore, σ is given by

$$\sigma(X) = X \cap \bigcup_{A \in a(\sigma)} X^c \ominus A. \quad (5.5)$$

The following question arises. If ψ is an increasing, translation invariant, self-dual operator, then ψ can be represented as a supremum of erosions, $\psi = \bigvee_{A \in \mathcal{V}(\psi)} \varepsilon_A$, where ε_A is the erosion $\varepsilon_A(X) = X \ominus A$. The corresponding switch operator $\sigma = \Sigma(\psi)$ is characterized completely by a subset of $\mathcal{V}(\psi)$, namely those structuring elements which do not contain the origin. It is easy to see that $\psi \geq \bigvee_{A \in a(\sigma)} \varepsilon_A$; in general, equality does not hold. Yet the collection $a(\sigma)$ is sufficiently large to recover ψ . How can we express ψ in terms of this collection?

More generally, let \mathcal{A} be a collection of structuring elements which obey

$$0 \notin A \quad \text{and} \quad A \cap B \neq \emptyset, \quad \text{for } A, B \in \mathcal{A}. \quad (5.6)$$

Let

$$\sigma(X) = \sigma_{\mathcal{A}}(X) = X \cap \bigcup_{A \in \mathcal{A}} X^c \ominus A, \quad (5.7)$$

that is,

$$\sigma_{\mathcal{A}} = \text{id} \wedge \bigvee_{A \in \mathcal{A}} \varepsilon_A \nu. \quad (5.8)$$

To show that σ is a switch operator, we compute $\psi_{\mathcal{A}} = \Psi(\sigma_{\mathcal{A}})$ and show that it is an increasing, self-dual operator.

$$\begin{aligned} \psi_{\mathcal{A}} &= (\text{id} \wedge \nu \sigma_{\mathcal{A}}) \vee \sigma_{\mathcal{A}} \nu \\ &= (\text{id} \wedge \bigwedge_{A \in \mathcal{A}} \delta_{\bar{A}}) \vee (\nu \wedge \bigvee_{A \in \mathcal{A}} \varepsilon_A). \end{aligned}$$

Since $A \cap B \neq \emptyset$ for $A, B \in \mathcal{A}$ we get that $\delta_{\bar{A}} \geq \varepsilon_B$ for $A, B \in \mathcal{A}$. This implies that $\text{id} \wedge \bigwedge_{A \in \mathcal{A}} \delta_{\bar{A}} \geq \text{id} \wedge \bigvee_{A \in \mathcal{A}} \varepsilon_A$, which leads us to

$$\begin{aligned} \psi_{\mathcal{A}} &= (\text{id} \wedge \bigwedge_{A \in \mathcal{A}} \delta_{\bar{A}}) \vee (\nu \wedge \bigvee_{A \in \mathcal{A}} \varepsilon_A) \vee (\text{id} \wedge \bigvee_{A \in \mathcal{A}} \varepsilon_A) \\ &= (\text{id} \wedge \bigwedge_{A \in \mathcal{A}} \delta_{\bar{A}}) \vee \bigvee_{A \in \mathcal{A}} \varepsilon_A. \end{aligned}$$

In other words, $\psi_{\mathcal{A}}$ is the centre of the increasing operator $\bigvee_{A \in \mathcal{A}} \varepsilon_A$ and its negation $\bigwedge_{A \in \mathcal{A}} \delta_{\bar{A}}$.

5.3. Theorem. *Every translation invariant switch operator σ is of the form $\sigma = \sigma_{\mathcal{A}}$, where \mathcal{A} is a collection of structuring elements satisfying (5.6). The corresponding self-dual operator $\psi_{\mathcal{A}} = \Psi(\sigma_{\mathcal{A}})$ is given by*

$$\psi_{\mathcal{A}}(X) = (X \cap \bigcap_{A \in \mathcal{A}} X \oplus \check{A}) \cup \bigcup_{A \in \mathcal{A}} X \ominus A. \quad (5.9)$$

If σ is a switch operator and \mathcal{A} a collection of structuring elements satisfying (5.6) such that $\sigma = \sigma_{\mathcal{A}}$, then we say that σ *generated by* \mathcal{A} . Note that, in general, \mathcal{A} is a subcollection of $\mathcal{a}(\sigma)$. To compute the set $\mathcal{a}(\sigma_{\mathcal{A}})$ observe that

$$B \in \mathcal{V}(\psi_{\mathcal{A}}) \iff B \in \mathcal{V}(\text{id} \wedge \bigwedge_{A \in \mathcal{A}} \delta_{\check{A}}) \text{ or } B \in \mathcal{V}(\bigvee_{A \in \mathcal{A}} \varepsilon_A).$$

Elements $B \in \mathcal{V}(\text{id} \wedge \bigwedge_{A \in \mathcal{A}} \delta_{\check{A}})$ contain the origin, and as such they are not contained in $\mathcal{a}(\sigma_{\mathcal{A}})$. Furthermore, $B \in \mathcal{V}(\bigvee_{A \in \mathcal{A}} \varepsilon_A)$ yields that $A \subseteq B$ for some $A \in \mathcal{A}$. Thus we conclude that

$$\mathcal{a}(\sigma_{\mathcal{A}}) = \{B \mid 0 \notin B \text{ and } A \subseteq B \text{ for some } A \in \mathcal{A}\}.$$

A structuring element $B \in \mathcal{A}$ is redundant if there exists another structuring element $A \in \mathcal{A}$ which is a strict subset of B .

Note that $\sigma_{\mathcal{A}}$ can also be written as a union of hit-or-miss operators, namely

$$\sigma_{\mathcal{A}}(X) = \bigcup_{A \in \mathcal{A}} X \otimes (\{0\}, A).$$

We conclude this section with the following result.

5.4. Proposition.

(a) *Let σ, σ' be translation invariant switch operators; then*

$$\sigma \leq \sigma' \Rightarrow \mathcal{a}(\sigma) \subseteq \mathcal{a}(\sigma').$$

(b) *Let \mathcal{A}, \mathcal{B} be subsets of $\mathcal{P}(\mathbb{Z}^2)$. If $\mathcal{A} \subseteq \mathcal{B}$ then $\sigma_{\mathcal{A}} \leq \sigma_{\mathcal{B}}$.*

(c) *The following two assertions are equivalent:*

(i) $\sigma_{\mathcal{A}} \leq \sigma_{\mathcal{B}}$;

(ii) *for every $A \in \mathcal{A}$ there exists a $B \in \mathcal{B}$ such that $B \subseteq A$.*

PROOF. (a) and (b) are obvious (in fact, (b) is a consequence of (c)). We prove (c). First assume (i) and let $A \in \mathcal{A}$. Then $0 \in A^c$ and $0 \in \sigma_{\mathcal{A}}(A^c)$. As $\sigma_{\mathcal{A}} \leq \sigma_{\mathcal{B}}$, we conclude that $0 \in \sigma_{\mathcal{B}}(A^c)$; so there exists a $B \in \mathcal{B}$ such that $B \subseteq A$. This proves (ii)

The proof that (ii) \Rightarrow (i) is straightforward. ■

5.5. Corollary. *If \mathcal{A}, \mathcal{B} are subsets of $\mathcal{P}(\mathbb{Z}^2)$ such that*

(i) $\mathcal{B} \subseteq \mathcal{A}$;

(ii) *for every $A \in \mathcal{A}$ there exists a $B \in \mathcal{B}$ such that $B \subseteq A$;*

then $\sigma_{\mathcal{A}} = \sigma_{\mathcal{B}}$.

5.6. Example. Consider the operators $\theta_5, \theta_6, \dots, \theta_9$ defined in Example 3.1, and define $\sigma_k = \Sigma(\theta_k)$, $k = 5, \dots, 9$. Then σ_k is generated by the collection \mathcal{A}_k of subsets A of the 3×3 square centred at the origin with $0 \notin A$ and $\text{card}(A) = k$. This collection \mathcal{A}_k contains $\binom{8}{k}$ subsets (which equals 0 if $k = 9$). For example, \mathcal{A}_6 consists of 28 structuring elements:

Refer to Section 7, Examples 7 and 8, for some additional results concerning this example.

6. Construction of self-dual filters

In [12, 5, 3] it has been shown how to construct morphological filters by iteration. We recall the following result.

6.1. Proposition. *Let ψ be an increasing, finite-window operator, let ψ^∞ be an increasing operator, and assume that $\psi^n \rightarrow \psi^\infty$ (pointwise) as $n \rightarrow \infty$. Then ψ^∞ is a morphological filter.*

It is obvious that the sequence ψ^n converges to a limit if ψ is extensive or anti-extensive. In Section 2 we have seen that an activity-extensive operator ψ has the property that every sequence $\psi^n(X)$ is pointwise monotone, and hence convergent. Therefore, Proposition 6.1 applies to operators which are activity-extensive.

Suppose that ψ is an increasing, translation invariant, self-dual operator. Is there a simple way to verify whether or not ψ is activity-extensive? A simple, but very useful, criterion is given by the following result.

6.2. Proposition. *Let ψ be an arbitrary operator, and assume that there exists a set X such that $\psi(X) \neq X$ and $\psi^p(X) = X$ for some integer $p > 1$. Then ψ is not activity-extensive.*

In this case we call X *periodic with respect to ψ* . If $p > 1$ is the smallest integer for which $\psi^p(X) = X$ then we say that X is p -periodic, or alternatively, that X has period p . For example, the chess board is 2-periodic with respect to the median operator which uses the rhombus (origin and its four 4-neighbours) as structuring element.

Let ψ be an increasing, translation invariant, self-dual operator with $\Sigma(\psi) = \sigma_{\mathcal{A}}$. If $A \in \mathcal{A}$, then $0 \notin A$ and $0 \in \psi(A)$. If there exists an integer $p > 1$ such that $0 \notin \psi^p(A)$, then the sequence $\psi^n(A)$ is not pointwise monotone, and therefore ψ is not activity-extensive. On the other hand, suppose that $0 \in \psi^n(A)$ for every $n \geq 1$ and $A \in \mathcal{A}$. We show that ψ is activity-extensive. Suppose, namely, that ψ is not activity-extensive. Then there exists a set X such that $\psi^n(X)$ is not pointwise monotone. Without loss of generality we can assume that $0 \notin X$, $0 \in \psi(X)$, and $0 \notin \psi^p(X)$ for some integer $p > 1$. Thus $0 \in \sigma(X^c)$, which means that $A \subseteq X$ for some $A \in \mathcal{A}$. But then, by assumption, $0 \in \psi^n(A) \subseteq \psi^n(X)$ for every $n \geq 1$, a contradiction. We have proved the following result.

6.3. Proposition. *Let ψ be an increasing, translation invariant, self-dual operator with switch operator $\sigma_{\mathcal{A}}$. Then ψ is activity-extensive if and only if $0 \in \psi^n(A)$, $n \geq 1$, for every $A \in \mathcal{A}$.*

In practice, the structuring elements in \mathcal{A} are finite and the sequence $\psi^n(A)$ becomes stationary (i.e., constant) after finitely many iterations. In such cases, verification of the condition in Proposition 6.3 requires only finitely many computations.


6.4. Example. (a) The operator θ_6 defined in Example 3.1 is not activity-extensive since

$$\begin{array}{|c|c|c|} \hline \bullet & \bullet & \bullet \\ \hline \bullet & \cdot & \bullet \\ \hline \cdot & \cdot & \bullet \\ \hline \end{array} \xrightarrow{\theta_6} \begin{array}{|c|c|c|} \hline \cdot & \bullet & \cdot \\ \hline \cdot & \bullet & \cdot \\ \hline \cdot & \cdot & \cdot \\ \hline \end{array} \xrightarrow{\theta_6} \begin{array}{|c|c|c|} \hline \cdot & \cdot & \cdot \\ \hline \cdot & \cdot & \cdot \\ \hline \cdot & \cdot & \cdot \\ \hline \end{array}$$

(b) The operator θ_7 is activity-extensive, since

$$\begin{array}{|c|c|c|} \hline \bullet & \bullet & \bullet \\ \hline \bullet & \cdot & \bullet \\ \hline \cdot & \cdot & \bullet \\ \hline \end{array} \xrightarrow{\theta_7} \begin{array}{|c|c|c|} \hline \bullet & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet \\ \hline \cdot & \bullet & \bullet \\ \hline \end{array} \xrightarrow{\theta_7} \text{idem}$$

and



The other structuring elements in \mathcal{A}_7 are 90° -rotations of these two elements.

In [4] and [3] we have explained how one can modify an increasing, self-dual operator which is not activity-extensive in such a way that it becomes activity-extensive. In fact this approach is based on the following two results.

6.5. Proposition. (See [14] and [3, Chapter 13].) *Assume that ϕ, ψ are finite-window operators, $\phi \leq \psi$, and that ϕ is an inf-overfilter and ψ a sup-underfilter. Then the centre $\gamma = (\text{id} \wedge \psi) \vee \phi$ is activity-extensive and the sequence γ^n converges to the strong middle filter of ϕ and ψ .*

6.6. Proposition. (See [3, Prop. 6.28].) *Let ψ be an increasing operator.*

- (a) *If α is an opening with $\alpha \leq \psi$ then $\psi\alpha$ is an inf-overfilter.*
- (b) *If β is a closing with $\beta \geq \psi$ then $\psi\beta$ is a sup-underfilter.*

If we combine the previous two results with Proposition 4.3 we arrive at the following construction method for self-dual morphological filters.

6.7. Proposition (Construction of self-dual filters).

Let ψ be an increasing, self-dual operator with switch operator $\sigma = \Sigma(\psi)$. Let α be an opening $\leq \psi$, and let $\beta = \alpha^$ be the negative closing. Then*

$$\pi = (\text{id} \wedge \psi\beta) \vee \psi\alpha$$

is an increasing self-dual operator with $\Sigma(\pi) = \text{id} \wedge \sigma\beta$. The operator π is activity-extensive and satisfies $\pi \preceq \psi$.

If the operators ψ, α, β are finite-window operators, then π is a finite-window operator too, and π^n converges towards a strong self-dual morphological filter π^∞ . (In fact, π^∞ is the middle filter of $\psi\alpha$ and $\psi\beta$.)

This result makes clear that we must look for openings α which are $\leq \psi$. Recall that the structural opening α_B is given by $\alpha_B(X) = X \circ B$. It is a well-known fact [3] that

$$\alpha_B \leq \psi \text{ iff } \psi(B) \geq B.$$

The following result characterizes all structuring elements B with this property.

6.8. Proposition. *Assume that ψ is increasing and self-dual, and that the switch operator $\sigma = \Sigma(\psi)$ can be represented as $\sigma = \sigma_{\mathcal{A}}$, for some collection $\mathcal{A} \subseteq \mathcal{P}(\mathbb{Z}^2)$. The following assertions are equivalent:*

- (i) $\psi(B) \geq B$;
- (ii) $\sigma(B) = \emptyset$;
- (iii) $\forall b \in B \forall A \in \mathcal{A} : A_b \cap B \neq \emptyset$;
- (iv) $\forall A \in \mathcal{A} : 0 \in \check{A} \bullet B$.

PROOF.

(i) \iff (ii): using (3.1) we get

$$\begin{aligned} \sigma(B) = \emptyset &\iff B \cap \psi(B^c) = \emptyset \\ &\iff \psi(B^c) \subseteq B^c \\ &\iff B \subseteq (\psi(B^c))^c = \psi^*(B) = \psi(B). \end{aligned}$$

(ii) \iff (iii): using the expression for $\sigma_{\mathcal{A}}$ in (5.7) we get

$$\begin{aligned}
\sigma_{\mathcal{A}}(B) = \emptyset &\iff B \cap \bigcup_{A \in \mathcal{A}} B^c \ominus A = \emptyset \\
&\iff \forall b \in B : b \notin \bigcup_{A \in \mathcal{A}} B^c \ominus A \\
&\iff \forall b \in B \forall A \in \mathcal{A} : b \notin B^c \ominus A \\
&\iff \forall b \in B \forall A \in \mathcal{A} : A_b \not\subseteq B^c \\
&\iff \forall b \in B \forall A \in \mathcal{A} : A_b \cap B \neq \emptyset.
\end{aligned}$$

(iii) \iff (iv): above we have seen that (iii) is equivalent with

$$\forall b \in B \forall A \in \mathcal{A} : b \notin B^c \ominus A.$$

This implies

$$\begin{aligned}
&\forall b \in B \forall A \in \mathcal{A} : b \in (B^c \ominus A)^c = B \oplus \check{A} = \check{A} \oplus B \\
&\iff \forall A \in \mathcal{A} : B \subseteq \check{A} \oplus B \\
&\iff \forall A \in \mathcal{A} : 0 \in (\check{A} \oplus B) \ominus B = \check{A} \bullet B.
\end{aligned}$$

This concludes the proof. \blacksquare

A structuring element B which satisfies the equivalent properties in Proposition 6.8 is called *persistent with respect to ψ* . Note that, unless $\psi = \text{id}$, singletons are not persistent. It is obvious that the collection of persistent sets is closed under unions: if B_i is persistent for every i in some index set I , then $\bigcup_{i \in I} B_i$ is persistent, too.

Assume that ψ, ψ' are increasing, self-dual operators, and that $\psi' \preceq \psi$. If B is persistent with respect to ψ then it is also persistent with respect to ψ' .

7. Examples

In this section we present several increasing, translation invariant, self-dual operators ψ on $\mathcal{P}(\mathbb{Z}^2)$. All examples contain the following ingredients:

- characterization of a (minimal) collection of structuring elements \mathcal{A} such that $\sigma_{\mathcal{A}}$ is the switch operator associated with ψ ;
- an examination of the activity-extensivity of ψ ;
- (in case that ψ is not activity-extensive) characterization of a collection of structuring elements which are persistent with respect to ψ .

In the latter case one can use the opening $\alpha(X) = \bigvee_{i \in I} X \circ B_i$, where $B_i, i \in I$, are persistent structuring elements, to obtain a self-dual operator

$$\pi = (\text{id} \wedge \psi \alpha^*) \vee \psi \alpha \tag{7.1}$$

which is activity-extensive (see Proposition 6.7). It should be evident that our interest goes to “small” persistent structuring elements, as these cause the smallest “loss of activity” if one replaces ψ with π . The operators ψ, π in Example i are denoted by ψ_i, π_i , respectively ($i = 1, 2, \dots, 13$)

1. Let A be the rhombus comprising the origin and its four horizontal and vertical neighbours. Let $\rho_{A,s}$ be the rank operators defined in Section 3, and let ψ_1 be the centre of the operator

$\rho_{A,2}$ and its dual $\rho_{A,4}$, i.e., $\psi_1 = (\text{id} \wedge \rho_{A,2}) \vee \rho_{A,4}$ (cf. Example 3.1); then ψ_1 is self-dual. It is easy to verify that $\psi_1 = \psi_A$, where A contains one structuring element:

$$A = \begin{array}{|c|} \hline \bullet \\ \hline \bullet \quad \vdots \quad \bullet \\ \hline \bullet \\ \hline \end{array}$$

The operator ψ_1 is not activity-extensive for the chess board is a periodic structure. The elements

$$\begin{array}{|c|} \hline \bullet \quad \bullet \\ \hline \end{array} \quad \text{and} \quad \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array}$$

are persistent. The operator ψ_1 switches the state of all pixels which are isolated in the 4-connected sense. Let A be the structuring element depicted above, then ψ_1 is given by (4.6). However, the adjacency triple condition governed by (4.5) is not satisfied.

2. Again, let A be the rhombus, and let ψ_2 be the corresponding median operator. Then $\psi_2 = \psi_A$ with

$$A = \begin{array}{|c|} \hline \bullet \\ \hline \bullet \quad \vdots \quad \bullet \\ \hline \bullet \\ \hline \end{array} + 90^\circ\text{-rotations}$$

Like in the previous example, ψ_2 is not activity-extensive since the chess board is 2-periodic with respect to ψ_2 . The operator ψ_2 has an infinite number of persistent elements which cannot be decomposed into smaller parts which are persistent, for example:

$$\begin{array}{|c|} \hline \bullet \quad \bullet \\ \hline \bullet \quad \bullet \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}$$

In fact, every structuring element in which every pixel has at least two 4-neighbours is persistent.

3. Consider the median operator ψ_3 associated with the 3×3 square. In Section 3 we have depicted the structuring elements in A such that $\psi_3 = \psi_A$; we recall them here for the sake of convenience.

$$\begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \vdots \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \vdots \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \vdots \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \vdots \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \vdots \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \vdots \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \quad \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \vdots \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array} + \text{rotations.}$$

ψ_3 is not activity-extensive, because the following structure is 2-periodic:

$$\begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \quad \text{etc.}$$

The set

$$\begin{array}{|c|} \hline \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \bullet \quad \bullet \\ \hline \end{array}$$


is a persistent structuring element. Let α be the corresponding structural opening, and let π_3 be the modification of ψ_3 resulting from (7.1). The median operator ψ_3 , its modification π_3 , and the filter π_3^∞ obtained by infinite iteration are illustrated in the second row of Figure 6.

4. Consider the subcollection \mathcal{B} of \mathcal{A} in Example 3 which lacks the first structuring element along with its three 90° -rotations. The corresponding self-dual operator ψ_4 is less active than ψ_3 . However, it is not activity-extensive for the structure depicted in **3** is 2-periodic. Of course, the persistent element in **3** is also persistent with respect to ψ_4 , but there also exists a smaller element which does the job, namely




Denote by π_4 the modification of ψ_4 resulting from (7.1), where α is the opening by this structuring element. Refer to the third row in Figure 6 for an illustration.

5. Consider the subcollection \mathcal{C} of \mathcal{A} in **3** which lacks the 45° , 135° , 225° and 315° -rotations of the first structuring element, i.e., the elements



Then $\psi_5 := \psi_{\mathcal{C}}$ is less active than ψ_3 but it is not activity-extensive for, again, the structure depicted in **3** is 2-periodic with respect to ψ_5 . In this case the rhombus




is persistent with respect to ψ_5 .

Denote by π_5 the modification of ψ_5 resulting from (7.1), where α is the opening by this structuring element. The operators in this example are illustrated in the fourth row of Figure 6.

6. We can combine 4 and 5 and consider the subcollection \mathcal{D} of \mathcal{A} which lacks the eight rotations of the first structuring element in \mathcal{A} , i.e., $\mathcal{D} = \mathcal{B} \cap \mathcal{C}$. It is relatively easy to show that $\sigma_{\mathcal{D}} = \sigma_{\mathcal{B}} \wedge \sigma_{\mathcal{C}}$ (here one must use that $B \in \mathcal{B}$ and $C \in \mathcal{C}$ implies that $D \subseteq B \cup C$ for some $D \in \mathcal{D}$). Now we conclude from (3.11) that $\psi_6 := \psi_{\mathcal{D}}$ is the centre of $\psi_4 = \psi_{\mathcal{B}}$ and $\psi_5 = \psi_{\mathcal{C}}$:


$$\psi_6 = \psi_4 \wedge \psi_5.$$

In this case



are persistent with respect to ψ_6 . Let π_6 be the modification of ψ_6 resulting from (7.1) if we take for α the union of the structural openings by these two elements. The action of the operators in this example are depicted in the bottom row of Figure 6.

7. Consider again the operator θ_6 introduced in Example 3.1; see also Example 5.6. Here we shall denote this operator by ψ_7 . The collection of structuring elements associated with $\Sigma(\psi_7)$ is given by




+ rotations




Fig. 6. The top-right image is the original image X . The four subsequent rows show (from left to right) the images $\psi_i(X), \pi_i(X), \pi_i^\infty(X)$ defined in examples $i = 3, 4, 5, 6$, respectively. For a better understanding of the usefulness of the modifications π_i and their iterates one should compare these images with the ones in Figure 1.

The operator ψ_7 is not activity-extensive since the pattern



is 2-periodic. The elements




are persistent. Let π_7 be the modification of ψ_7 obtained from (7.1) if one takes for α the opening by these two structuring elements. The second row in Figure 7 depicts the action of ψ_7, π_7 , and π_7^∞ .




Fig. 7. The top-right image is the original image X . The second row illustrates the operators ψ_7, π_7 , and π_7^∞ discussed in Example 7. The bottom row depicts $\psi_8(X)$ and $\psi_8^\infty(X)$ examined in Example 8.


8. Consider the subcollection \mathcal{B} of \mathcal{A}_6 which is obtained by deleting the elements



We use Proposition 6.3 to show that $\psi_8 = \psi_{\mathcal{B}}$ is activity-extensive. Thereto we must consider the sequences $\psi_8^n(B)$ for $B \in \mathcal{B}$. First we note that the elements



are invariant under ψ_8 . Now




This implies that ψ_8 is activity-extensive. The bottom row in Figure 7 depicts the transformed sets $\psi_8(X)$ (left) and $\psi_8^\infty(X)$ (right). These figures show clearly the invariance of the triangles depicted above.

9. Consider the operator θ_7 introduced in Example 3.1. We define $\psi_9 := \theta_7$. The associated collection \mathcal{A}_7 consists of eight structuring elements:

$$\mathcal{A}_7 = \begin{bmatrix} \bullet & \bullet & \bullet \\ \bullet & \cdot & \bullet \\ \bullet & \bullet & \cdot \end{bmatrix} + \text{rotations.}$$

Using Proposition 6.3 it is easy to show that ψ_9 is activity-extensive. Use namely that



In Figure 8 we depict $X, \psi_9(X)$ and $\psi_9^\infty(X)$.



Fig. 8. From left to right: $X, \psi_9(X)$ and $\psi_9^\infty(X)$; see Example 9.

10. Consider the operator $\psi_{10} := \theta_8$; here \mathcal{A}_8 contains one structuring element, namely

$$\mathcal{A}_8 = \begin{bmatrix} \bullet & \bullet & \bullet \\ \bullet & \cdot & \bullet \\ \bullet & \bullet & \bullet \end{bmatrix}$$

The structuring element in \mathcal{A}_8 satisfies condition (4.5) in Theorem 4.7, meaning that the resulting operator ψ_{10} is idempotent. Its action consists hereof that it removes isolated foreground and background pixels (isolated in the sense of 8-neighbours). An illustration can be found in Figure 2.

11. Consider the weighted median operator ψ_{11} with matrix

$$\begin{pmatrix} 1 & 3 & 1 \\ 3 & 5 & 3 \\ 1 & 3 & 1 \end{pmatrix}$$

and threshold 11. The collection of structuring elements \mathcal{A} yielding the corresponding switch operator is

If \mathcal{B} is the collection of structuring elements associated with the rhombus median operator (see Example 2) then condition (ii) in Proposition 5.4(c) is satisfied, hence the operator ψ_{11} is less active than the rhombus median operator. The chess board is periodic with respect to ψ_{11} and therefore ψ_{11} is not activity-extensive. The pattern

is persistent. Let π_{11} be the resulting activity-extensive modification of ψ_{11} . In Figure 9, second row, we depict $\psi_{11}(X), \pi_{11}(X)$, and $\pi_{11}^\infty(X)$.

12. Let ρ be the weighted rank operator obtained by changing the threshold in the previous example to 9. Let ψ_{12} be the centre of ρ and its negative ρ^* . It is evident that this operator is less active than the one in 11. The switch operator $\Sigma(\psi_{12})$ is generated by the structuring elements:

$$\mathcal{A} = \begin{bmatrix} \bullet & \bullet & \cdot \\ \bullet & \cdot & \bullet \\ \cdot & \bullet & \cdot \end{bmatrix} \begin{bmatrix} \cdot & \bullet & \bullet \\ \bullet & \cdot & \bullet \\ \cdot & \bullet & \cdot \end{bmatrix} \begin{bmatrix} \bullet & \bullet & \bullet \\ \bullet & \cdot & \bullet \\ \bullet & \cdot & \bullet \end{bmatrix} + 90^\circ\text{-rotations}$$


We use Proposition 6.3 to show that ψ_{12} is activity-extensive:



Fig. 9. The top-right image is the original image X . The second row illustrates the operators ψ_{11} , π_{11} , and π_{11}^∞ . The third row illustrates ψ_{12} and ψ_{12}^∞ , and the bottom row depicts the operators ψ_{13} , π_{13} , and π_{13}^∞ .

$$\begin{array}{ccc}
 \begin{array}{|c|} \hline \bullet & \bullet & \bullet \\ \hline \bullet & \cdot & \bullet \\ \hline \bullet & \cdot & \bullet \\ \hline \end{array} & \xrightarrow{\psi_{12}} & \begin{array}{|c|} \hline \bullet & \bullet & \bullet \\ \hline \bullet & \cdot & \bullet \\ \hline \bullet & \cdot & \bullet \\ \hline \end{array} & \xrightarrow{\psi_{12}} & \text{idem}
 \end{array}$$

This implies that ψ_{12} is activity-extensive. Observe that the elementary triangles



are persistent. In the third row in Figure 9 we depict $\psi_{12}(X)$ (left) and $\psi_{12}^\infty(X)$ (right). These images show clearly the invariance of the elementary triangles. In this respect this example resembles the one in Example 8.

13. If we add

$$\begin{array}{|c|c|c|} \hline \bullet & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet \\ \hline \bullet & \bullet & \bullet \\ \hline \end{array} + 90^\circ\text{-rotations}$$

to the collection \mathcal{A} in 11 we obtain an operator ψ_{13} which is more active than the one in 11. Note that this operator corresponds with the weighted median with matrix

$$\begin{pmatrix} 1 & 2 & 1 \\ 2 & 3 & 2 \\ 1 & 2 & 1 \end{pmatrix}$$

and threshold 8. The chess board is periodic and therefore ψ_{13} is not activity-extensive. Again

$$\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \bullet & \bullet \\ \hline \end{array}$$

is persistent. Let π_{13} be the corresponding activity-extensive modification of ψ_{13} . In the bottom row of Figure 9 we depict $\psi_{13}(X)$, $\pi_{13}(X)$, and $\pi_{13}^\infty(X)$.

8. Conclusions and open problems

The basic morphological operators like dilations, erosions, openings, closings, and alternating sequential filters are not self-dual. The best known morphological operator which is self-dual is the median operator μ . Unfortunately, this operator has a property which is undesirable from the view point of noise filtering: the sequence of iterates $\mu^n(X)$ of an input image X needs not be convergent. In fact, it is easy to find images X such that $\{X, \mu(X)\}$ constitutes a period-2 orbit.

In this paper two goals have been accomplished. First, we have presented a general method, based on the concept of a switch operator, for the construction of increasing operators which are self-dual. Under the extra assumption of translation invariance we have derived a general formula for self-dual operators:

$$\psi_{\mathcal{A}}(X) = \left(X \cap \bigcap_{A \in \mathcal{A}} X \oplus \check{A} \right) \cup \bigcup_{A \in \mathcal{A}} X \ominus A, \quad (8.1)$$

where \mathcal{A} is an arbitrary collection of structuring elements satisfying

$$0 \notin A \quad \text{and} \quad A \cap B \neq \emptyset, \text{ for } A, B \in \mathcal{A}.$$

In this paper we have restricted ourselves to binary images for simplicity. Yet, it is obvious that formula (8.1) carries over immediately to grey-scale images, and in fact, this remark applies to most of the results established in this paper.

A second accomplishment is that we have derived a general method for the modification of an arbitrary increasing, self-dual operator ψ yielding another increasing, self-dual operator π which is less active than ψ , but which has the desirable property that it is activity-extensive. This means in particular that π^n converges to a self-dual morphological filter π^∞ (presumed that it is a finite-window operator).

We have illustrated our theoretical results with a large number of examples. We have limited ourselves to the square grid, and we have only considered structuring elements which fit inside a 3×3 window. Furthermore, the structuring elements have been chosen in such a

way that the resulting operator, besides being translation invariant, is invariant under rotations about 90 degrees. From a theoretical point of view there is no need to make such restrictions. In fact, in an earlier paper [2] we have investigated modifications of the median operator on the hexagonal grid, such as the Maisonneuve filter [9]; see also [3, Ex. 13.59].

We have resisted the temptation to make quantitative statements about the noise cleaning capacities of the operators and filters discussed in our examples, others than those in terms of the activity ordering. For example the illustration of Example 8 in Figure 7 and Example 12 in Figure 9 show that $\psi_8^\infty(X)$ is very similar to $\psi_{12}^\infty(X)$, though both images differ at a few points. The only theoretical support for this resemblance is that both operators keep elementary triangles unchanged.

The research described here also raises a number of intriguing theoretical questions which fall outside the scope of this paper, but which, nevertheless deserve to be mentioned in this final section.

Question 1. *Do there exist operators which have orbits with period greater than 2?*

Period-3 orbits may be of special interest because of the possible analogy with the famous result obtained by Li and Yorke [8] in the context of dynamical systems theory which says that “period three implies chaos”.

The following question is related to the previous one.

Question 2. *Given an increasing, translation invariant (self-dual) operator ψ (e.g. the median operator). What kind of behaviour can the orbits $\{X, \psi(X), \psi^2(X), \dots\}$ exhibit? Do they necessarily “converge” to a periodic orbit? Can such morphological dynamical systems exhibit “chaotic behaviour”?*

These appear to be difficult questions which require a lot of fundamental research. Maybe one can find (partial) answers from the literature on *cellular automata* [15]. To our knowledge however, the overlap between these two fields, that is, mathematical morphology on the one hand and cellular automata on the other, is almost completely unexplored up till today; refer to [11] for some related discussions. Hopefully, the theory developed in this paper will serve as a stimulus to fill this gap.

References.

- [1] BANON, G. J. F., AND BARRERA, J. Minimal representation for translation-invariant set mappings by mathematical morphology. *SIAM J. Appl. Math.* 51 (1991), 1782–1798.
- [2] HEIJMANS, H. J. A. M. Morphological filtering and iteration. In *Visual Communications and Image Processing '90* (Lausanne, 1990), vol. 1360 of *SPIE Proceedings*, pp. 166–175.
- [3] HEIJMANS, H. J. A. M. *Morphological Image Operators*. Academic Press, Boston, 1994.
- [4] HEIJMANS, H. J. A. M. On the construction of morphological operators which are selfdual and activity-extensive. *Signal Processing* (to appear).
- [5] HEIJMANS, H. J. A. M., AND SERRA, J. Convergence, continuity and iteration in mathematical morphology. *Journal of Visual Communication and Image Representation* 3 (1992), 84–102.
- [6] HEIJMANS, H. J. A. M., AND VINCENT, L. Graph morphology in image analysis. In *Mathematical Morphology in Image Processing*, E. R. Dougherty, Ed. Marcel Dekker, New York, 1993, ch. 6, pp. 171–203.

- [7] JUSTUSSON, B. I. Median filtering: statistical properties. In *Two-Dimensional Digital Signal Processing II: Transforms and Median Filters*, T. S. Huang, Ed. Springer-Verlag, Berlin, 1981, pp. 161–196.
- [8] LI, T., AND YORKE, J. Period three implies chaos. *Amer. Math. Monthly* 82 (1975), 985–992.
- [9] MAISONNEUVE, F. Ordinaux transfinis et sur- (ou sous-) potentes. Technical report N780, Ecole Nationale Supérieure des Mines de Paris, 1982.
- [10] MATHERON, G. *Random Sets and Integral Geometry*. John Wiley and Sons, New York, 1975.
- [11] PRESTON JR., K., DUFF, M., LEVIALDI, S., NORGRÉN, P., AND TORIWAKI, J.-I. Basics of cellular logic with some applications in medical image processing. *Proc. IEEE* 67 (1979), 826–856.
- [12] RONSE, C., AND HEIJMANS, H. J. A. M. The algebraic basis of mathematical morphology – part II: Openings and closings. *Computer Vision, Graphics and Image Processing: Image Understanding* 54 (1991), 74–97.
- [13] SERRA, J. *Image Analysis and Mathematical Morphology*. Academic Press, London, 1982.
- [14] SERRA, J., Ed. *Image Analysis and Mathematical Morphology. II: Theoretical Advances*. Academic Press, London, 1988.
- [15] WOLFRAM, S., Ed. *Theory and applications of cellular automata : including selected papers 1983-1986*. World Scientific, Singapore, 1986.