



**Centrum voor Wiskunde en Informatica**  
Centre for Mathematics and Computer Science

---

A.M.H. Gerards

On Tutte's characterization of graphic matroids - a graphic proof

Department of Operations Research, Statistics, and System Theory    Report BS-R9028    October

---

The Centre for Mathematics and Computer Science is a research institute of the Stichting Mathematisch Centrum, which was founded on February 11, 1946, as a nonprofit institution aiming at the promotion of mathematics, computer science, and their applications. It is sponsored by the Dutch Government through the Netherlands Organization for the Advancement of Research (N.W.O.).

# On Tutte's Characterization of Graphic Matroids – a graphic proof

A.M.H. Gerards  
Centre for Mathematics and Computer Science  
P.O. Box 4079  
1009 AB Amsterdam, the Netherlands

## Abstract

In this paper we present a relatively simple proof of Tutte's characterization of graphic matroids. The proof uses the notion of 'signed graph' and it is 'graphic' in the sense that it can be presented almost entirely by drawing (signed) graphs.

1980 Mathematics Subject Classification: 05B35.

Key Words and Phrases: binary matroid, graphic matroid, minor, signed graph.

## 1 Introduction—Tutte's Characterization of Graphic Matroids

The aim of this paper is to present a proof of the following result.

**Theorem 1 (Tutte, 1959 [6])** *Let  $\mathcal{M}$  be a binary matroid. Then  $\mathcal{M}$  is graphic if and only if  $\mathcal{M}$  has no  $F_7$ ,  $F_7^*$ ,  $\mathcal{M}^*(K_5)$  or  $\mathcal{M}^*(K_{3,3})$  minor.*

We assume the reader to be familiar with matroid terminology. In fact what we need are the following notions (cf. Welsh [9]): *binary matroid*; *representation (matrix) (over  $GF(2)$ )* of a binary matroid; *contraction*, *deletion* and *minor*; and *graphic matroid*, i.e. the *cycle matroid* (= *polygon matroid*) of a graph. Representation matrices (over  $GF(2)$ ) of the four matroids mentioned in Theorem 1 are

$$\begin{bmatrix} 1 & 0 & 0 & 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 1 \end{bmatrix} (F_7), \quad \begin{bmatrix} 1 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 & 1 \end{bmatrix} (F_7^*),$$

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 \end{bmatrix} (\mathcal{M}^*(K_5)), \quad \begin{bmatrix} 1 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 1 & 1 \end{bmatrix} (\mathcal{M}^*(K_{3,3})).$$

Besides Tutte's original proof in [6] there exist proofs of Theorem 1 derived by: Ghouila-Houri [1]; Inukai and Weinberg [2], using Tutte's "wheels and whirls theorem" (Tutte [7]);

Seymour [4], using results from [3]; Wagner [8] and Truemper [5]. Our proof of Theorem 1 is in Section 3. The notions we need for it are given in Section 2. To motivate them and to guide the reader into the proof we have included the key ideas of our proof as observations in Section 2.

## 2 Preliminaries—the key ideas

### Signed Graphs

A *signed graph* is a pair  $(G, \Sigma)$  where  $G = (V(G), E(G))$  is an undirected graph, possibly with loops and parallel edges, and  $\Sigma$  is a subset  $E(G)$ . Edges in  $\Sigma$  are called *odd*. The other edges are called *even*. A circuit in  $G$  is called *odd (even)* in  $(G, \Sigma)$ , if it contains an odd (even) number of odd edges.

The *extended even cycle matroid*,  $\mathcal{S}(G, \Sigma)$ , of  $(G, \Sigma)$  is the binary matroid represented over  $\text{GF}(2)$  by the matrix

$$\begin{bmatrix} 1 & \chi_\Sigma \\ 0 & M_G \end{bmatrix}.$$

Here  $\chi_\Sigma$  denotes the characteristic vector of  $\Sigma$  as a subset of  $E(G)$ .  $M_G$  denotes the *node-edge incidence matrix* of  $G$ , i.e. the matrix in  $\text{GF}(2)^{V(G) \times E(G)}$  defined by

$$(M_G)_{u,e} = \begin{cases} 1 & \text{if } e = uv, \text{ for some node } v \neq u; \\ 0 & \text{else.} \end{cases}$$

The element of  $\mathcal{S}(G, \Sigma)$  which is not in  $E(G)$  will be denoted by  $\sigma$ . The circuits of  $\mathcal{S}(G, \Sigma)$  are: even circuits in  $(G, \Sigma)$ , the union of two odd circuits in  $(G, \Sigma)$  when they have at most one node in common, and the union of an odd circuit in  $(G, \Sigma)$  with  $\{\sigma\}$ .

*Resigning*  $(G, \Sigma)$  (on  $U \subseteq V(G)$ ), means replacing  $\Sigma$  by  $\Sigma \Delta \delta(U)$  ( $\Delta :=$  symmetric difference,  $\delta(U) := \{uv \in E(G) \mid u \in U, v \notin U\}$ ). As resigning leaves odd (even) circuits odd (even), it does not affect  $\mathcal{S}(G, \Sigma)$ . A signed graph that comes from  $(G, \Sigma)$  by a series of the following operations: resigning, deletion of an edge, and contraction of an *even* edge, is called a *minor* of  $(G, \Sigma)$ . Obviously the extended even cycle matroid of a minor of  $(G, \Sigma)$  is a minor of  $\mathcal{S}(G, \Sigma)$ . Moreover contracting  $\sigma$  in  $\mathcal{S}(G, \Sigma)$  yields the cycle matroid,  $\mathcal{M}(G)$ , of  $G$ .

The following observation makes clear why we are interested in signed graphs.

**Observation 1** *Let  $\mathcal{M}$  be a binary matroid, with representation  $M$  over  $\text{GF}(2)$ . If all its proper minors are graphic and  $M \neq 0$ , then  $\mathcal{M}$  is the extended even cycle matroid of a signed graph.*

**Proof:** We may assume that (eventually after applying row operations over  $\text{GF}(2)$ )

$$M = \begin{bmatrix} 1 & a^\top \\ 0 & N \end{bmatrix}.$$

Matrix  $N$  represents a minor of  $\mathcal{M}$ . Hence  $N$  represents the cycle matroid of a graph,  $G$  say. If  $\Sigma := \{e \in E(G) \mid a_e = 1\}$ , then  $\mathcal{M} = \mathcal{S}(G, \Sigma)$ .  $\square$




Figure 1:

### Examples—the four forbidden minors

In view of Observation 1 one might expect that the four matroids in Theorem 1 are extended even cycle matroids of signed graphs. Indeed, the four signed graphs, depicted in Figure 1 satisfy

**Observation 2**  $S(\overline{K}_4) = F_7^*$ ,  $S(\overline{K}_3^2) = F_7$ ,  $S(\overline{P}_3) = \mathcal{M}^*(K_5)$ , and  $S(\overline{K}_4^2) = \mathcal{M}^*(K_{3,3})$ .

(In Figure 1 bold lines denote odd edges and thin lines denote even edges.)

### Almost Bipartite Signed Graphs

A signed graph  $(G, \Sigma)$  is *bipartite* if  $\Sigma = \delta(U)$  for some  $U \subseteq V(G)$ , in other words, if one can resign  $(G, \Sigma)$  to  $(G, \emptyset)$ . So a signed graph is bipartite if and only if it contains no odd circuits. A signed graph  $(G, \Sigma)$  is called *almost bipartite* if there exists a node  $u \in V(G)$ , called a *blocknode*, such that  $u \in V(C)$  for each odd circuit  $C$  in  $(G, \Sigma)$ .

**Observation 3** Let  $(G, \Sigma)$  be a signed graph. If  $(G, \Sigma)$  is almost bipartite, then  $S(G, \Sigma)$  is graphic.

**Proof:** Let  $u$  be a blocknode in  $(G, \Sigma)$ . Resign such that  $\Sigma \subseteq \delta(u)$ . We construct a new graph  $H$  as follows (see Figure 2). We split node  $u$  into two new nodes  $u_1$  and  $u_2$ . We split  $\delta(u)$  into  $\delta(u_1)$  and  $\delta(u_2)$  such that  $\delta(u_1) = \delta(u) \setminus \Sigma$  and  $\delta(u_2) = \Sigma$ . (One should understand this in such a way that an odd loop with endpoint  $u$  becomes an edge from  $u_1$  to  $u_2$ .) Finally we add an edge  $e_\sigma$  from  $u_1$  to  $u_2$ . It is not hard to see that  $S(G, \Sigma) = \mathcal{M}(H)$ .  $\square$




Figure 2:

## Switching—essentially almost bipartite signed graphs

Let  $(G, \Sigma)$  be a signed graph. Moreover let  $G_1$  and  $G_2$  be two subgraphs of  $G$ , and let  $\Sigma_i := \Sigma \cap E(G_i)$  for  $i = 1, 2$ . We say that  $[(G_1, \Sigma_1), (G_2, \Sigma_2)]$  is a  $k$ -partition if the following holds:

$$\begin{aligned} E(G_1) \cup E(G_2) &= E(G), & E(G_1) \cap E(G_2) &= \emptyset, \\ V(G_1) \cup V(G_2) &= V(G), & |V(G_1) \cap V(G_2)| &= k. \end{aligned}$$

We call the  $k$ -partition *strong* if both  $(G_1, \Sigma_1)$  and  $(G_2, \Sigma_2)$  are not bipartite, and  $|E(G_1)|, |E(G_2)| \geq 3$ .

If  $(G, \Sigma)$  has a 1-partition  $[(G_1, \Sigma_1), (G_2, \Sigma_2)]$ , and the signed graph  $(H, \Theta)$  consists of two disjoint copies of  $(G_1, \Sigma_1)$  and  $(G_2, \Sigma_2)$ , then we say that  $(H, \Theta)$  comes from  $(G, \Sigma)$  by *breaking (at a cutnode)*. Conversely we say that  $(G, \Sigma)$  comes from  $(H, \Theta)$  by *glueing (at a node)*.

Let  $(G, \Sigma)$  have a 2-partition  $[(G_1, \Sigma_1), (G_2, \Sigma_2)]$ . Let  $u$  and  $v$  be the two nodes that  $G_1$  and  $G_2$  have in common. Consider disjoint copies  $(H_1, \Theta_1)$  and  $(H_2, \Theta_2)$  of  $(G_1, \Sigma_1)$  and  $(G_2, \Sigma_2)$  respectively. For  $i = 1, 2$ , let  $u_i$  and  $v_i$  be the copies of  $u$  and  $v$  in  $H_i$ . Construct  $(H, \Theta)$  from  $(H_1, \Theta_1)$  and  $(H_2, \Theta_2)$  by identifying  $u_1$  with  $v_2$  and  $u_2$  with  $v_1$ . The operation carrying  $(G, \Sigma)$  into  $(H, \Theta)$  is called *switching*. (See Figure 3a.)




Figure 3:

We call a signed graph *essentially almost bipartite* if it can be turned into an almost bipartite graph by a series of breakings, glueings and switchings. It is obvious that these operations leave  $S(G, \Sigma)$  invariant. Hence Observation 3 yields

**Observation 4** *Let  $(G, \Sigma)$  be a signed graph. If  $(G, \Sigma)$  is essentially almost bipartite, then  $S(G, \Sigma)$  is graphic.*

## 3 The Proof

Let  $\mathcal{M}$  be a binary matroid with no  $F_7$ ,  $F_7^*$ ,  $\mathcal{M}^*(K_5)$ , or  $\mathcal{M}^*(K_{3,3})$  minor. We have to prove that  $\mathcal{M}$  is graphic. We may assume that all its proper minors are graphic. Hence, by Observation 1,  $\mathcal{M}$  is the extended even cycle matroid of a signed graph,  $(G, \Sigma)$  say. The graphicness of  $\mathcal{M}$  follows from Observations 2 and 4 and the result below (which is the special case of Tutte's Theorem 1 when restricted to extended even cycle matroids of signed graphs, see the remark below).

**Theorem 2** Let  $(G, \Sigma)$  be a signed graph with no  $\overline{K}_4$ ,  $\overline{K}_3^2$ ,  $\overline{P}_3$ , or  $\overline{K}_4^2$  minor. Then  $(G, \Sigma)$  is essentially almost bipartite.

**Proof:** Let  $(G, \Sigma)$  be a signed graph with no  $\overline{K}_4$ ,  $\overline{K}_3^2$ ,  $\overline{P}_3$ , or  $\overline{K}_4^2$  minor. Assume that all the proper minors of  $(G, \Sigma)$  are essentially almost bipartite. We consider three cases.

**Case 1**  $(G, \Sigma)$  has a 0- or 1-partition.

We may assume that in fact  $(G, \Sigma)$  has a 0-partition. (If not, break.) As all components are proper minors of  $(G, \Sigma)$ , we may assume that they are almost bipartite. (If not, switch.) Glueing the blocknodes of the components to one new node yields an almost bipartite graph. So  $(G, \Sigma)$  is essentially almost bipartite.

**Case 2**  $(G, \Sigma)$  has no 0- or 1-partition, but has a strong 2-partition.

Let  $[(G_1, \Sigma_1), (G_2, \Sigma_2)]$  be a strong 2-partition of  $(G, \Sigma)$ . Let  $u$  and  $v$  be the two nodes that  $G_1$  and  $G_2$  have in common. For  $i = 1, 2$ , let  $(G_i^+, \Sigma_i^+)$  denote the signed graph obtained from  $(G_i, \Sigma_i)$  by adding two edges,  $e_i$  and  $f_i$  say, from  $u$  to  $v$ , with  $e_i$  odd, and  $f_i$  even.  $(G_1^+, \Sigma_1^+)$  and  $(G_2^+, \Sigma_2^+)$  are minors of  $(G, \Sigma)$ . Indeed, as the partition is strong,  $(G_2, \Sigma_2)$  contains an odd circuit,  $C$  say. As  $(G, \Sigma)$  has no 0- or 1-partition, by Menger's Theorem, in  $G_2$  there exist two node disjoint paths  $P$  and  $Q$  from  $C$  to  $\{u, v\}$  (see Figure 3b). From this it is easy to see that  $(G_1^+, \Sigma_1^+)$  is a minor of  $(G, \Sigma)$ . By symmetry, so is  $(G_2^+, \Sigma_2^+)$ .

It is not hard to see that every switching in  $(G_1^+, \Sigma_1^+)$  or in  $(G_2^+, \Sigma_2^+)$  can be carried out in  $(G, \Sigma)$ . As  $(G_1^+, \Sigma_1^+)$  and  $(G_2^+, \Sigma_2^+)$  are proper minors of  $(G, \Sigma)$ , they are essentially almost bipartite. Carry out all the switchings needed to make  $(G_1^+, \Sigma_1^+)$  and  $(G_2^+, \Sigma_2^+)$  almost bipartite in  $(G, \Sigma)$ . Then all odd circuits in  $(G_1^+, \Sigma_1^+)$  contain  $u$ , or all these odd circuits contain  $v$  (since  $e_1$  and  $f_1$  form an odd circuit). The same is true for  $(G_2^+, \Sigma_2^+)$ . So  $(G, \Sigma)$  is almost bipartite or can be made almost bipartite by just one switching.

**Case 3**  $(G, \Sigma)$  has no 0-, 1- or strong 2-partition.

An *odd*- $K_4$  and an *odd*- $K_3^2$  are as depicted in Figure 4. In that figure wriggled and dotted lines stand for pairwise openly disjoint paths. Wriggled lines correspond to paths with at least one edge, dotted lines may have length zero. The word *odd* in a region denotes that the circuit bounding that region is odd.




Figure 4:




Figure 5:

**Claim 1**  $(G, \Sigma)$  contains no odd- $K_4$  and no odd- $K_3^2$ . Moreover  $(G, \Sigma)$  contains no pair of node disjoint odd circuits.

**Proof of claim 1:** It is an easy exercise to show that if  $(G, \Sigma)$  would contain an odd- $K_4$  or an odd- $K_3^2$ , then it has a  $\overline{K}_4$  or a  $\overline{K}_3^2$  minor. Suppose that  $C_1$  and  $C_2$  are two node disjoint odd circuits. Assume that  $|V(C_1)| + |V(C_2)|$  is as large as possible. First we show that  $C_1$  or  $C_2$  has two nodes only. Indeed, if this is not the case then  $(G, \Sigma)$  contains a signed subgraph as in Figure 5a. (This follows from Menger's Theorem and the fact that  $(G, \Sigma)$  has no 0-, 1- or strong 2-partition.) As  $(G, \Sigma)$  contains no odd- $K_4$ , each of the faces marked with “\*” in Figure 5a is bounded by an even circuit. But this is impossible, as  $(G, \Sigma)$  has no  $\overline{P}_3$  minor. So we may assume that  $V(C_1) := \{u_1, u_2\}$ . Again by Menger's Theorem and the fact that  $(G, \Sigma)$  has no 0-, 1- or strong 2-partition, it follows that there exist two paths  $P_1^1$  and  $P_1^2$  from  $u_1$  to  $C_2$  which have only  $u_1$  in common and do not contain  $u_2$ . Similarly we have  $P_2^1$  and  $P_2^2$  from  $u_2$  to  $C_2$ . Obviously  $P_1^i$  and  $P_2^j$  are node disjoint. (If not, we could find an odd circuit disjoint from  $C_2$  and with more edges than  $C_1$ .) Hence  $(G, \Sigma)$  contains one of the signed graphs in Figure 5b and c. One easily sees now that  $(G, \Sigma)$  has  $\overline{K}_4^2$  or  $\hat{K}_4^2$  (see Figure 5d) as minor. As  $\overline{K}_4$  is a minor of  $\hat{K}_4^2$ , this contradicts our assumptions on  $(G, \Sigma)$ .

**end of proof of claim 1**

Let  $C_1$  and  $C_2$  be two odd circuits such that their *intersection graph*  $P$ , i.e.  $V(P) = V(C_1) \cap V(C_2)$ ,  $E(P) = E(C_1) \cap E(C_2)$  is a path. It is easy to see that such two odd circuits exist. Assume they are chosen such that  $|E(P)|$  is as small as possible. Let  $u$  be one of the endpoints of  $P$ . For  $i = 1, 2$ , let  $e_i \in (E(C_i) \cap \delta(u)) \setminus E(P)$ . (See Figure 6 (a) and (b).) Let  $H$  denote the *union graph* of  $C_1$  and  $C_2$ , i.e.  $V(H) = V(C_1) \cup V(C_2)$ ,  $E(H) = E(C_1) \cup E(C_2)$ . We resign such that  $\Sigma \cap E(H) = \{e_1, e_2\}$ . An *(st-) arc* is a path from  $s \in V(H) \setminus \{u\}$  to  $t \in V(H) \setminus \{u\}$  which is internally node disjoint with  $H$ . The arc is *odd (even)* if it contains an odd (even) number of odd edges.

**Claim 2** If  $Q$  is an odd *st-arc* then  $s \in V(C_1) \setminus V(C_2)$  and  $t \in V(C_2) \setminus V(C_1)$  (or conversely). Moreover in that case  $E(P) = \emptyset$ .

**Proof of claim 2:** Let  $Q_H$  be the unique *st-path* with edges only in  $E(H) \setminus \{e_1, e_2\}$ . Then  $Q_H$  and  $Q$  form an odd circuit. Hence  $l_i := |E(Q_H) \cap E(C_i)| \geq |E(P)|$  for  $i = 1, 2$ . If






Figure 6:

$l_1 = 0$  (or if  $l_2 = 0$ ), then  $|E(P)| = 0$ , and one of the endpoints of  $Q$  is  $u$ , which contradicts the definition of an arc. So  $l_1, l_2 > 0$ . Hence  $s \in V(C_1) \setminus V(C_2)$  and  $t \in V(C_2) \setminus V(C_1)$  (or conversely). If  $|E(P)| > 0$ , then  $Q$  and  $H$  form an odd- $K_4$  (see Figure 6(c)).

**end of proof of claim 2**

**Claim 3** *If  $E(P) = \emptyset$ , then there exists no pair of internally node disjoint paths of different parity from  $V(C_1) \setminus V(C_2)$  to  $V(C_2) \setminus V(C_1)$*

**Proof of claim 3:** If such paths would exist then  $\overline{K}_3^2$  is a minor of  $(G, \Sigma)$  (see Figure 6(d)).

**end of proof of claim 3**

We now prove that  $u$  is a blocknode. Suppose this is not the case. Let  $C$  be an odd circuit not containing  $u$ . As it meets both  $C_1$  and  $C_2$  it is the disjoint union of paths on  $H$  (disjoint from  $u$ ) and of arcs. As  $C$  is odd an odd number of the arcs is odd. By Claims 2 and 3 it follows that  $E(P) = \emptyset$  and that  $C$  contains an odd number of arcs from  $V(C_1) \setminus V(C_2)$  to  $V(C_2) \setminus V(C_1)$ . This is absurd.  $\square$

**Remark:** That Theorem 2 is a special case of Theorem 1 is quite easy to see. First observe that the only signed graphs with an extended even cycle matroid among  $F_7$ ,  $F_7^*$ ,  $\mathcal{M}^*(K_5)$  and  $\mathcal{M}^*(K_{3,3})$  are  $\mathcal{S}(\overline{K}_4)$ ,  $\mathcal{S}(\overline{K}_3^2)$ ,  $\mathcal{S}(\overline{P}_3)$ , and  $\mathcal{S}(\overline{K}_4^2)$ . So if a signed graph  $(G, \Sigma)$  has none of these four signed graphs as a minor, then, by Theorem 1,  $\mathcal{S}(G, \Sigma)$  is graphic. Let  $H$  be a graph such that  $\mathcal{S}(G, \Sigma)$  is isomorphic to  $\mathcal{M}(H)$ . Let  $e_\sigma$  be the edge in  $H$  corresponding to the special element  $\sigma$  in  $\mathcal{S}(G, \Sigma)$ . Let  $\tilde{H}$  be obtained from  $H$  by contracting  $e_\sigma$ . Then  $G$  and  $\tilde{H}$  have isomorphic cycle matroids. Hence, by Whitney's switching theorem (Whitney [10]), it follows that  $G$  can be turned into  $\tilde{H}$  by breaking, glueing and switching. From this it easily follows that  $(G, \Sigma)$  is essentially almost bipartite. This shows how Theorem 2 follows from Theorem 1.

## References

- [1] A. GHOUILA-HOURI, *Chapter III, in: Flots et tensions dans un graphe*, Annales Scientifique de l'Ecole Normale Supérieure, 81 (1964), pp. 267–339.
- [2] T. INUKAI AND L. WEINBERG, *Graph-realizability of matroids*, Annals of the New York Academy of Sciences, 319 (1979), pp. 289–305.
- [3] P. D. SEYMOUR, *Decomposition of regular matroids*, Journal of Combinatorial Theory, Series B, 28 (1980), pp. 305–354.
- [4] —, *On Tutte's characterization of graphic matroids*, Annals of Discrete Mathematics, 8 (1980), pp. 83–90.
- [5] K. TRUEMPER, *A decomposition theory for matroids II, minimal violation matroids*, Journal of Combinatorial Theory, Series B, 39 (1985), pp. 282–297.
- [6] W. T. TUTTE, *Matroids and graphs*, Transactions of the American Mathematical Society, 90 (1959), pp. 527–552.
- [7] —, *Connectivity in matroids*, Canadian Journal of Mathematics, 18 (1966), pp. 1301–1324.
- [8] D. K. WAGNER, *On theorems of Whitney and Tutte*, Discrete Mathematics, 57 (1985), pp. 147–154.
- [9] D. J. A. WELSH, *Matroid Theory*, Academic Press, New York, 1976.
- [10] H. WHITNEY, *2-isomorphic graphs*, American Journal of Mathematics, 21 (1933), pp. 73–84.