



**Centrum voor Wiskunde en Informatica**  
Centre for Mathematics and Computer Science

---

H.C.P. Berbee, W.Th.F. den Hollander

Tail triviality for sums of stationary random variables

Department of Mathematical Statistics    Report MS-R8810    June

---

*Bibliotheek*  
Centrum voor Wiskunde en Informatica  
Amsterdam

The Centre for Mathematics and Computer Science is a research institute of the Stichting Mathematisch Centrum, which was founded on February 11, 1946, as a nonprofit institution aiming at the promotion of mathematics, computer science, and their applications. It is sponsored by the Dutch Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O.).

# Tail Triviality for Sums of Stationary Random Variables

H.C.P. Berbee

*Centre for Mathematics and Computer Science  
P.O. Box 4079, 1009 AB Amsterdam, The Netherlands*

W.Th.F. den Hollander

*Delft University of Technology  
Faculty of Mathematics and Informatics  
Julianalaan 132, 2628 BL Delft, The Netherlands*

We study tail  $\sigma$ -fields and loss of memory associated with sums of stationary integer-valued random variables. An application concerns convergence in distribution of interarrival times in 0-1 sequences.

*AMS 1980 Subject Classification: Primary 60F20, secondary 60G10, 28D20, 28D05, 60J15.*

*Keywords and phrases: Tail  $\sigma$ -field for sums, entropy, strongly aperiodic, random walk with stationary increments, loss of memory, zero-two theorem, coupling, interarrival times.*

## 1. Statement of results

Let  $X_1, X_2, \dots$  be a *strictly stationary* sequence of *integer-valued* random variables and let  $S_1, S_2, \dots$  be the sums

$$S_n = X_1 + \dots + X_n \quad (n \geq 1).$$

If the increments are independent, then according to the Hewitt-Savage zero-one law (Breiman (1968)) the tail  $\sigma$ -field

$$\bigcap_{N>0} \sigma(S_n : n \geq N)$$

is trivial. Without the independence this need no longer be true. The main question that will be addressed in this paper is what can be said about the tail behavior of the sums in this more general setting.

It is natural to extend  $(X_n)_{n \geq 1}$  to a double-sided process  $X := (X_n)_{n \in \mathbb{Z}}$  and to extend also the sums to a double-sided process  $(S_n)_{n \in \mathbb{Z}}$  by requiring that

$$(1.1) \quad S_0 = 0; \quad S_n - S_{n-1} = X_n \quad (n \in \mathbb{Z}).$$

The process  $X$  is defined on the probability space  $(\Omega, \mathcal{F}, P)$  with  $\Omega = \mathbb{Z}^{\mathbb{Z}}$ ,  $X$  is the identity on  $\Omega$ ,  $\mathcal{F}$  is the product  $\sigma$ -field generated by the discrete topology on  $\mathbb{Z}$ , and  $P$  is a  $T$ -invariant probability measure with  $T$  the shift defined by  $(TX)_n = X_{n+1}$  ( $n \in \mathbb{Z}$ ). We shall be interested in the following tail  $\sigma$ -fields associated with  $(S_n)$ :

$$\begin{aligned} \mathcal{G}_\infty &:= \bigcap_{M, N \geq 0} \sigma((S_m, S_n) : m \leq -M, n \geq N), \\ \mathcal{G}_\infty^{inv} &:= \bigcap_{M, N \geq 0} \sigma(S_n - S_m : m \leq -M, n \geq N). \end{aligned}$$

Note that  $\mathcal{G}_\infty^{inv} = \bigcap_N T^{-N} \mathcal{G}_\infty$ , so  $\mathcal{G}_\infty^{inv}$  is a  $T$ -invariant  $\sigma$ -field included in the double tail  $\sigma$ -field  $\mathcal{G}_\infty$ . The following example shows that  $\mathcal{G}_\infty^{inv}$  and  $\mathcal{G}_\infty$  may be different.

**EXAMPLE (1.2):** Let  $(Y_n)$  be i.i.d. with  $P(Y_n = -1) = P(Y_n = 1) = \frac{1}{2}$  and let  $X_n = Y_n - Y_{n-1}$  ( $n \in \mathbb{Z}$ ). Then  $S_n - S_m = X_{m+1} + \dots + X_n = Y_n - Y_m$  and so  $\mathcal{G}_\infty^{inv}$  is trivial by the Kolmogorov zero-one law (Breiman (1968)). On the other hand,  $S_n = X_1 + \dots + X_n = Y_n - Y_0$  and hence  $\lim_{N \rightarrow \infty} N^{-1} \sum_{n=1}^N S_n = -Y_0$  a.s. Thus  $Y_0$  is  $\mathcal{G}_\infty$ -measurable, so  $\mathcal{G}_\infty$  is not trivial. In fact,  $\mathcal{G}_\infty = \sigma(Y_0)$  a.s.

Let

$$\mathcal{F}_\infty := \bigcap_{M, N \geq 0} \sigma(X_n : m \leq -M, n > N)$$

be the double tail  $\sigma$ -field associated with  $X$ . Since

$$\sigma(S_n - S_m : m \leq -M, n \geq N) = \sigma(S_N - S_{-M}) \vee \sigma(X_n : m \leq -M, n > N)$$

for all  $M, N \geq 0$ , it is clear that  $\mathcal{G}_\infty^{inv} \supset \mathcal{F}_\infty$ . In section 2 we prove:

**THEOREM (1.3).** *Let  $(X_n)$  be stationary integer-valued and let  $(S_n)$  be given by (1.1). If  $X_0$  has finite entropy, then  $\mathcal{G}_\infty^{inv} = \mathcal{F}_\infty$  a.s.*

**COROLLARY(1.4).** *If  $X_0$  has finite entropy, then  $\mathcal{G}_\infty^{inv}$  is trivial iff  $\mathcal{F}_\infty$  is trivial.*

As example (1.2) shows, to obtain triviality of the larger tail  $\sigma$ -field  $\mathcal{G}_\infty$  we need to assume more than triviality of  $\mathcal{F}_\infty$ . Also, it will not be enough to impose stronger mixing conditions on  $X$ . In section 3 we prove the following zero-two theorem which will be seen to be the key to our study of  $\mathcal{G}_\infty$ .

**THEOREM(1.5).** *Let  $(X_n)$  be stationary, ergodic and real-valued and let  $(S_n)$  be given by (1.1). For every real  $h$ , either*

$$\| P(S_N \in \cdot | \mathcal{F}_N^*) - P(S_N + h \in \cdot | \mathcal{F}_N^*) \| = 2 \quad \text{for all } N \geq 1 \quad \text{a.s.}$$

or else

$$\lim_{N \rightarrow \infty} \| P(S_N \in \cdot | \mathcal{F}_N^*) - P(S_N + h \in \cdot | \mathcal{F}_N^*) \| = 0 \quad \text{a.s.}$$

Here  $\mathcal{F}_N^* := \sigma(X_n : n \notin \{1, \dots, N\})$  and  $\| \cdot \|$  denotes total variation.

For i.i.d. sequences the corresponding zero-two theorem was proved by Stam (1967) and by Ornstein (1969), while Berbee (1979) gave a proof for mixing sequences. Most zero-two theorems in the literature relate to Markov processes.

Theorem(1.5) leads us in a natural way to associate a group  $H$  with  $(S_n)$  as follows:

$$H := \{h \in \mathbb{R} : \text{there exists } N \geq 1 \text{ such that with positive probability} \\ P(S_N \in \cdot | \mathcal{F}_N^*) \text{ and } P(S_N + h \in \cdot | \mathcal{F}_N^*) \text{ have mass in common}\}.$$

By theorem (1.5), the set  $H$  is a group. For  $(X_n)$  integer-valued, we say that  $(S_n)$  is *strongly aperiodic* if  $H = \mathbb{Z}$ .

Theorem (1.6) below gives sufficient conditions for triviality of  $\mathcal{G}_\infty$  as well as of the right tail  $\sigma$ -field

$$\mathcal{G}_\infty^+ := \bigcap_{N \geq 0} \sigma(S_n : n \geq N).$$

Let

$$\mathcal{F}_\infty^+ := \bigcap_{N \geq 0} \sigma(X_n : n > N)$$

and note that  $\mathcal{G}_\infty^+ \supset \mathcal{F}_\infty^+$ . In section 4 we prove:

**THEOREM(1.6).** *Let  $(X_n)$  be stationary integer-valued and let  $(S_n)$  be given by (1.1). Assume that  $(S_n)$  is strongly aperiodic. If  $\mathcal{F}_\infty^+$  is trivial then  $\mathcal{G}_\infty^+$  is trivial. If  $\mathcal{F}_\infty$  is trivial then  $\mathcal{G}_\infty$  is trivial.*

Note that in example (1.2) triviality of  $\mathcal{F}_\infty$  holds because  $(X_n)$  is one-dependent but strong aperiodicity fails because  $S_N = Y_N - Y_0$  while  $Y_0$  and  $Y_N$  are  $\mathcal{F}_N^*$ -measurable, so  $H = \{0\}$ .

As an application of theorem (1.6) we consider a stationary ergodic 0-1 sequence  $(X_n)$ . Let  $T_1, T_2, \dots$  be the random positive times at which  $X_n$  assumes the value 1:

$$(1.7) \quad \begin{aligned} T_1 &= \inf\{n \geq 1 : X_n = 1\}, \\ T_k &= \inf\{n > T_{k-1} : X_n = 1\} \quad (k > 1). \end{aligned}$$

By stationarity and ergodicity, if  $P(X_1 = 1) > 0$  then  $T_k < \infty$  a.s. for all  $k \geq 1$ . We call  $T_{k+1} - T_k$  the  $k^{\text{th}}$  interarrival time. In section 5 we prove:

**COROLLARY(1.7).** *Let  $(X_n)$  be a stationary 0-1 sequence with  $P(X_1 = 1) > 0$  and let  $(S_n)$  be given by (1.1) and  $(T_k)$  by (1.7). If  $\mathcal{G}_\infty^+$  is trivial, then for any integer  $t \geq 0$*

$$\lim_{k \rightarrow \infty} P(T_{k+1} - T_k > t) = P(T_2 - T_1 > t | X_1 = 1).$$

Corollary (1.7), together with theorem (1.6), extends a result by Janson (1984) for  $m$ -dependent sequences. Janson (1984) contains an example of a 1-dependent sequence with non-converging interarrival times. Corollary (1.4) has recently been applied by Den Hollander (1987) in a paper on mixing properties for random walk in random scenery.

## 2. Proof of theorem (1.3)

We start by recalling a few properties of entropy (see e.g. Smorodinsky (1971) or Parry (1981)). Let  $Z$  be a discrete random variable on our probability space  $(\Omega, \mathcal{F}, P)$  taking values in a countable set. The entropy of  $Z$  is defined as

$$H(Z) := - \sum_z P(Z = z) \log P(Z = z)$$

( $0 \log 0 = 0$ ). For a sub- $\sigma$ -field  $\mathcal{A} \subset \mathcal{F}$  the conditional entropy of  $Z$  given  $\mathcal{A}$  is defined as

$$H(Z|\mathcal{A}) := -E \sum_z P(Z = z|\mathcal{A}) \log P(Z = z|\mathcal{A}).$$

For two sub- $\sigma$ -fields  $\mathcal{A}, \mathcal{B} \subset \mathcal{F}$  we denote by  $\mathcal{A} \vee \mathcal{B}$  the smallest  $\sigma$ -field containing  $\mathcal{A}$  and  $\mathcal{B}$ . The following properties will be used below:

$$(2.1) \quad H(Z|\mathcal{A}) \leq H(Z|\mathcal{B}) \text{ if } \mathcal{B} \subset \mathcal{A},$$

$$(2.2) \quad H(Z|\mathcal{A}) \leq H(Z),$$

$$(2.3) \quad \begin{aligned} H(Z_1, Z_2|\mathcal{A}) &= H(Z_1|\mathcal{A}) + H(Z_2|\mathcal{A} \vee \sigma(Z_1)) \\ &= H(Z_2|\mathcal{A}) + H(Z_1|\mathcal{A} \vee \sigma(Z_2)), \end{aligned}$$

$$(2.4) \quad H(Z_1 + Z_2) \leq H(Z_1) + H(Z_2),$$

$$(2.5) \quad H(Z) \leq \log |\{z : P(Z = z) > 0\}|.$$

PROOF OF THEOREM(1.3): We introduce the notation

$$\begin{aligned} S_{m,n} &:= S_n - S_m = X_{m+1} + \cdots + X_n \quad (m, n \in \mathbb{Z}; m < n), \\ \mathcal{F}_N &:= \sigma(X_n : n \leq -N, n > N), \\ \mathcal{G}_N &:= \sigma(S_{m,n} : m \leq -N, n \geq N) = \sigma(S_{-N,N}) \vee \mathcal{F}_N. \end{aligned}$$

Note that  $\mathcal{F}_\infty = \bigcap_{N \geq 0} \mathcal{F}_N$  and  $\mathcal{G}_\infty^{inv} = \bigcap_{N \geq 0} \mathcal{G}_N$ .

By the martingale convergence theorem (Smorodinsky (1971))

$$\begin{aligned} H(X_0 | \mathcal{F}_\infty) &= \lim_{N \rightarrow \infty} H(X_0 | \mathcal{F}_N), \\ H(X_0 | \mathcal{G}_\infty^{inv}) &= \lim_{N \rightarrow \infty} H(X_0 | \mathcal{G}_N). \end{aligned}$$

Here the monotonicity of  $\mathcal{F}_N$  and  $\mathcal{G}_N$  is used together with (2.2) and the assumption  $H(X_0) < \infty$ . As a first step towards proving  $\mathcal{G}_\infty^{inv} = \mathcal{F}_\infty$  a.s. we claim that

$$(2.6) \quad H(X_0 | \mathcal{F}_\infty) = H(X_0 | \mathcal{G}_\infty^{inv}).$$

To get (2.6) we note that

$$(2.7) \quad H(X_m | \mathcal{G}_N) \leq H(X_0 | \mathcal{G}_{N+|m|}) \quad \text{for all } m \text{ and } N.$$

This follows from stationarity, (2.1) and the inclusion  $T^{-m} \mathcal{G}_{N+|m|} \subset \mathcal{G}_N$ . Now fix two positive integers  $K, L$  and consider the following quantity:

$$H((X_{jK})_{|j| \leq L-1}, S_{-LK, LK} | \mathcal{F}_{LK}),$$

i.e. the conditional entropy given  $\mathcal{F}_{LK}$  of the sum  $S_{-LK, LK}$  and of the  $X_n$  with  $|n| \leq (L-1)K$  but *sampled over gaps* of size  $K$ . The trick is to write this out in two different ways, using (2.3),

$$\begin{aligned} \text{I. } & H((X_{jK})_{|j| \leq L-1} | \mathcal{F}_{LK}) + H(S_{-LK, LK} | \mathcal{F}_{LK} \vee \sigma((X_{jK})_{|j| \leq L-1})), \\ \text{II. } & H(S_{-LK, LK} | \mathcal{F}_{LK}) + H((X_{jK})_{|j| \leq L-1} | \mathcal{F}_{LK} \vee \sigma(S_{-LK, LK})), \end{aligned}$$

to derive a lower bound for I and an upper bound for II, and thus to get an inequality between these two bounds which can then be exploited to prove (2.6).

Indeed, if we iterate the first term of I using (2.3), then we get  $2L-1$  terms each of which is bounded below by  $H(X_0 | \mathcal{F}_K)$  because of stationarity and (2.1). Hence, by ignoring the second term of I we get

$$\text{I} \geq (2L-1)H(X_0 | \mathcal{F}_K).$$

In II, on the other hand, the first term is bounded above by  $H(S_{-LK, LK})$  because of (2.2), while the second term equals  $H((X_{jK})_{|j| \leq L-1} | \mathcal{G}_{LK})$  and is bounded above by  $\sum_{|j| \leq L-1} H(X_{jK} | \mathcal{G}_{LK})$  via (2.1) and (2.3). Together with (2.7) this gives

$$\text{II} \leq H(S_{-LK, LK}) + (2L-1)H(X_0 | \mathcal{G}_{2LK}).$$

Since I=II we thus have

$$H(X_0 | \mathcal{F}_K) \leq (2L-1)^{-1} H(S_{-LK, LK}) + H(X_0 | \mathcal{G}_{2LK}).$$

Next let  $L \rightarrow \infty$  while keeping  $K$  fixed, and use the following lemma which we shall prove below.

LEMMA (2.8). If  $H(X_0) < \infty$ , then  $H(S_{-N,N}) = o(N)$ .

Finally we let  $K \rightarrow \infty$  and we end up with  $H(X_0|\mathcal{F}_\infty) \leq H(X_0|\mathcal{G}_\infty^{inv})$ . But this implies (2.6) since the reverse inequality holds by (2.1) and  $\mathcal{F}_\infty \subset \mathcal{G}_\infty^{inv}$ .

Now, (2.6) is only the first step in proving  $\mathcal{G}_\infty^{inv} = \mathcal{F}_\infty$  a.s. The next step is to prove that for any positive integer  $M$

$$(2.9) \quad H((X_n)_{|n| \leq M} | \mathcal{F}_\infty) = H((X_n)_{|n| \leq M} | \mathcal{G}_\infty^{inv}).$$

This can be done in exactly the same way as above. Instead of sampling  $2L - 1$  single variables  $X_n$  over gaps of size  $K$ , we must now sample  $2L - 1$  blocks of variables of fixed length  $2M + 1$  separated by gaps of size  $K$  and then again take the limit  $L \rightarrow \infty$  followed by  $K \rightarrow \infty$ . The key inequality is the obvious modification of (2.7). These steps are left for the reader to verify.

The assertion now follows from (2.9) and the following lemma which we shall prove below.

LEMMA(2.10). Assume  $H(X_0) < \infty$ . Let  $\mathcal{A}, \mathcal{B} \subset \mathcal{F}$  be two sub- $\sigma$ -fields such that  $\mathcal{A} \subset \mathcal{B}$ . If

$$H((X_n)_{|n| \leq m} | \mathcal{A}) = H((X_n)_{|n| \leq m} | \mathcal{B}) \quad \text{for all } m \geq 0,$$

then  $\mathcal{A} = \mathcal{B}$  a.s.

This completes the proof of theorem (1.3). ■

PROOF OF LEMMA(2.8): We follow a standard truncation argument. Fix a positive integer  $M$  and define

$$Y_n = \begin{cases} 0 & \text{if } |X_n| \leq M \\ X_n & \text{if } |X_n| > M \end{cases} \quad (n \in \mathbb{Z}).$$

Write  $S_{-N,N} = \sum_{n=-N+1}^N Y_n + \sum_{n=-N+1}^N (X_n - Y_n)$ . By stationarity and (2.4),

$$H(S_{-N,N}) \leq 2NH(Y_0) + H(\sum_{n=-N+1}^N (X_n - Y_n)).$$

Since  $X_n - Y_n$  can at most take  $2M + 1$  distinct values, the last term is at most  $\log(2N(2M + 1))$  by (2.5). Letting  $N \rightarrow \infty$  while keeping  $M$  fixed, we get

$$\limsup_{N \rightarrow \infty} (2N)^{-1} H(S_{-N,N}) \leq H(Y_0).$$

But  $H(Y_0)$  can be made arbitrarily small by letting  $M \rightarrow \infty$ , because  $H(X_0) < \infty$ . ■

PROOF OF LEMMA (2.10): Let  $E \in \mathcal{B}$ . Select  $E_m \in \sigma((X_n)_{|n| \leq m})$  such that

$$(2.11) \quad \lim_{m \rightarrow \infty} P(E_m \Delta E) = 0.$$

From (2.3), for all  $m \geq 0$

$$H(1_{E_m} | \mathcal{A}) + H((X_n)_{|n| \leq m} | \mathcal{A} \vee \sigma(1_{E_m})) = H((X_n)_{|n| \leq m} | \mathcal{A})$$

and by assumption this equals

$$H(1_{E_m} | \mathcal{B}) + H((X_n)_{|n| \leq m} | \mathcal{B} \vee \sigma(1_{E_m})) = H((X_n)_{|n| \leq m} | \mathcal{B}).$$

Because  $\mathcal{A} \subset \mathcal{B}$  there must be term by term equality, so  $H(1_{E_m} | \mathcal{A}) = H(1_{E_m} | \mathcal{B})$ . But (2.11) implies  $\lim_{m \rightarrow \infty} H(1_{E_m} | \mathcal{B}) = H(1_E | \mathcal{B}) = 0$ , hence  $\lim_{m \rightarrow \infty} H(1_{E_m} | \mathcal{A}) = H(1_E | \mathcal{A}) = 0$ . This in turn implies  $P(E | \mathcal{A}) = 0$  or 1 a.s., and so  $E \in \mathcal{A}$  a.s. Thus  $\mathcal{B} \subset \mathcal{A}$ . ■

### 3. Proof of theorem(1.5)

We follow the proof of the contraction lemma in Berbee (1986), which goes back to an idea of Bradley (1983). Berbee (1979) contains a proof for mixing sequences using coupling techniques. The ergodic theoretic proof below is considerably shorter but also less transparent.

Let us write  $P(X \in \cdot | Z)$  for the conditional distribution of  $X$  given  $\sigma(Z)$ . The following facts will be used below:

$$(3.1) \quad E \| P(X \in \cdot | Z) - P(Y \in \cdot | Z) \| = \| P((X, Z) \in \cdot) - P((Y, Z) \in \cdot) \|,$$

$$(3.2) \quad \| P(f(X) \in \cdot) - P(f(Y) \in \cdot) \| \leq \| P(X \in \cdot) - P(Y \in \cdot) \|,$$

where  $X$  and  $Y$  are random variables taking values in the same measurable space,  $Z$  is any other random variable and  $f$  is any measurable function.

PROOF OF THEOREM (1.5): Let  $P_1$  and  $P_2$  be the two probability measures on  $\mathcal{F}$  defined by

$$\begin{aligned} P_1 &= P \\ P_2 &= P \circ U_h \end{aligned}$$

where  $U_h : \Omega \rightarrow \Omega$  is the map

$$(U_h \omega)_n = \begin{cases} \omega_n & n \neq 1 \\ \omega_n + h & n = 1 \end{cases} (\omega \in \Omega = \mathbb{R}^{\mathbb{Z}}).$$

We compare the restrictions of  $P_1$  and  $P_2$  to the nested sequence of  $\sigma$ -fields

$$\mathcal{G}_{M,N} := \sigma(S_{m,n} : m \leq -M, n \geq N), \quad M, N \geq 0.$$

Since  $\mathcal{G}_\infty^{inv} = \bigcap_{M, N \geq 0} \mathcal{G}_{M, N}$ , it follows via the martingale convergence theorem that

$$(3.3) \quad \lim_{M, N \rightarrow \infty} \|P_1 - P_2\|_{\mathcal{G}_{M, N}} = \|P_1 - P_2\|_{\mathcal{G}_\infty^{inv}}.$$

Because  $\mathcal{G}_\infty^{inv}$  is  $T$ -invariant and  $P$  is  $T$ -invariant and ergodic on  $\mathcal{F}$ , it is immediate that  $P_1$  is  $T$ -invariant and ergodic on  $\mathcal{G}_\infty^{inv}$ . The same is true, however, for  $P_2$ . This follows from the commutation rule  $U_h T^{-1} = T^{-1} U_h$  which holds on  $\mathcal{G}_{M, N}$  for any  $M, N \geq 1$ . Indeed, by this rule we have

$$P_2(T^{-1}A) = P(U_h T^{-1}A) = P(T^{-1}U_h A) = P(U_h A) = P_2(A),$$

while ergodicity follows because if  $A$  is  $T$ -invariant so is  $U_h A$  by this rule. By an elementary application of the ergodic theorem we get from these ergodic properties that  $P_1$  and  $P_2$  on  $\mathcal{G}_\infty^{inv}$  are either identical or mutually singular and thus the right hand side of (3.3) equals 0 or 2.

The left hand side of (3.3) depends only on  $N + M$  because of stationarity. If we take  $M = 0$ , note that  $\mathcal{G}_{0, N} = \mathcal{F}_N^* \vee \sigma(S_N)$  and use (3.1), then (3.3) becomes equivalent to  $\lim_{N \rightarrow \infty} EZ_N = 0$  or 2 with

$$Z_N = \|P(S_N \in \cdot | \mathcal{F}_N^*) - P(S_N + h \in \cdot | \mathcal{F}_N^*)\|.$$

Because  $S_{N+1}$  is  $(S_N, X_{N+1})$ -measurable, we get from (3.1) and (3.2) that  $Z_N$  is a reverse submartingale, i.e.  $Z_{N+1} \leq E(Z_N | \mathcal{F}_{N+1}^*)$ . Since  $0 \leq Z_N \leq 2$ , it follows that  $Z_N$  converges a.s. with limit either 0 or 2. In the latter case we must have  $Z_N = 2$  a.s. for all  $N \geq 1$  because  $EZ_N$  is decreasing in  $N$ . ■

#### 4. Proof of theorem (1.6)

We need the following technical corollary of theorem (1.5). Once this has been proved theorem (1.6) will follow easily.

**COROLLARY(4.1).** *Let  $(X_n)$  be stationary, ergodic and integer-valued and let  $(S_n)$  be given by (1.1). If  $(S_n)$  is strongly aperiodic and if  $U$  and  $V$  are integer-valued  $(S_n)$ -measurable random variables, then for every  $K \geq 0$*

$$\lim_{N \rightarrow \infty} \|P((S_n + U, S_{-n} + V)_{n \geq N} \in \cdot | (S_n)_{|n| \leq K}) - P((S_n, S_{-n})_{n \geq N} \in \cdot | (S_n)_{|n| \leq K})\| = 0 \quad a.s.$$

**PROOF OF COROLLARY (4.1):** Because  $U$  is integer-valued we can approximate  $U$  by random variables  $U_m$  which are  $(S_n)_{|n| \leq m}$ -measurable such that  $\lim_{m \rightarrow \infty} P(U_m \neq U) = 0$ . Hence it is enough to prove the assertion for  $U$  measurable with respect to  $(S_n)_{|n| \leq m}$  for every fixed  $m \geq 0$ , and the same for  $V$ .

By theorem (1.5) and property (3.1) we have that for any integer  $h$

$$\lim_{N \rightarrow \infty} \|P((S_n + h)_{n \geq N} \in \cdot | (S_n)_{n \leq 0}) - P((S_n)_{n \geq N} \in \cdot | (S_n)_{n \leq 0})\| = 0 \quad a.s.$$

By stationarity this is the same as

$$\lim_{N \rightarrow \infty} \|P((S_{n+m} - S_m + h)_{n \geq N} \in \cdot | (S_{n+m} - S_m)_{n \leq 0}) - P((S_{n+m} - S_m)_{n \geq N} \in \cdot | (S_{n+m} - S_m)_{n \leq 0})\| = 0 \quad \text{a.s. for every } m.$$

Now,  $(S_{n+m} - S_m)_{n \leq 0}$  and  $(S_n)_{n \leq m}$  determine each other when  $m \geq 0$  (because  $S_0 = 0$ ). Hence we may replace  $h$  by any  $(S_n)_{n \leq m}$ -measurable random variable, and since moreover for all  $N \geq -m$  also  $(S_{-n} + V)_{n \geq N}$  is  $(S_n)_{n \leq m}$ -measurable we obtain

$$\lim_{N \rightarrow \infty} \|P((S_n + U, S_{-n} + V)_{n \geq N} \in \cdot | (S_n)_{n \leq m}) - P((S_n, S_{-n} + V)_{n \geq N} \in \cdot | (S_n)_{n \leq m})\| = 0 \quad \text{a.s.}$$

By (3.1) and (3.2) it follows that for all  $K \leq m$

$$\lim_{N \rightarrow \infty} \|P((S_n + U, S_{-n} + V)_{n \geq N} \in \cdot | (S_n)_{|n| \leq K}) - P((S_n, S_{-n} + V)_{n \geq N} \in \cdot | (S_n)_{|n| \leq K})\| = 0 \quad \text{a.s.}$$

The same type of argument shows that  $V$  may be replaced by 0 and that the role of  $U$  and  $V$  may be interchanged. The assertion now follows by using the triangle inequality. ■

In the proof of theorem (1.6) below we use the notation  $\beta(X, Y)$  to denote the *dependence coefficient* of random variables  $X$  and  $Y$  defined as

$$\beta(X, Y) := E \|P(Y \in \cdot | X) - P(Y \in \cdot)\|.$$

We shall need the inequality

$$(4.2) \quad \beta(X, Y) \leq 2E \|P(Y \in \cdot | X) - P(Z \in \cdot)\|$$

which holds for any random variable  $Z$  taking values in the same measurable space as  $Y$ . This follows from the inequality

$$E \|P(Y \in \cdot) - P(Z \in \cdot)\| \leq E \|P(Y \in \cdot | X) - P(Z \in \cdot)\|$$

which is easily deduced from (3.1) and (3.2).

PROOF OF THEOREM (1.6): Let  $K \leq m$ . Apply corollary (4.1) with  $U = -S_m$  and  $V = -S_{-m}$ , and combine this with

$$\begin{aligned} & E \|P((S_n - S_m, S_{-n} - S_{-m})_{n \geq N} \in \cdot | (S_n)_{|n| \leq K}) \\ & \quad - P((S_n - S_m, S_{-n} - S_{-m})_{n \geq N} \in \cdot)\| \\ & \leq \beta((X_n)_{|n| \leq K}, (X_n)_{|n| \geq m}) \\ & =: \varepsilon_{m, K}, \end{aligned}$$

where we use (3.1) and (3.2). This yields

$$(4.3) \quad \limsup_{N \rightarrow \infty} E \| P((S_n - S_m, S_{-n} - S_{-m})_{n \geq N} \in \cdot) - P((S_n, S_{-n})_{n \geq N} \in \cdot | (S_n)_{|n| \leq K}) \| \leq \varepsilon_{m,K}.$$

Now use (4.2) and (4.3) to obtain

$$(4.4) \quad \limsup_{N \rightarrow \infty} \beta((S_n)_{|n| \leq K}, (S_n)_{|n| \geq N}) \leq 2\varepsilon_{m,K}.$$

But if  $\mathcal{F}_\infty$  is trivial then  $\lim_{m \rightarrow \infty} \varepsilon_{m,K} = 0$  for every fixed  $K$  (by a standard martingale argument) and so

$$\lim_{N \rightarrow \infty} \beta((S_n)_{|n| \leq K}, (S_n)_{|n| \geq N}) = 0 \text{ for all } K \geq 0.$$

By the same token, this is equivalent to triviality of  $\mathcal{G}_\infty$ . Thus the second assertion is proved. The proof of triviality of  $\mathcal{G}_\infty^+$  from triviality of  $\mathcal{F}_\infty^+$  is an easy adaptation of the above proof. ■

## 5. Proof of Corollary (1.7)

The proof of Corollary (1.7) uses coupling. Let  $(Z_n^1)_{n \geq 1}$  and  $(Z_n^2)_{n \geq 1}$  be two arbitrary processes defined on probability spaces  $(\Omega^1, \mathcal{F}^1, P^1)$  and  $(\Omega^2, \mathcal{F}^2, P^2)$ , respectively. A *coupling* is a construction of these processes on a common probability space  $(\Omega^1 \times \Omega^2, \mathcal{F}^1 \times \mathcal{F}^2, P^{1,2})$  such that  $P^{1,2}|_{\mathcal{F}^1} = P^1$  and  $P^{1,2}|_{\mathcal{F}^2} = P^2$ . Given two coupled processes taking values in the same measurable space  $(\Omega^1, \mathcal{F}^1) = (\Omega^2, \mathcal{F}^2)$ , we say that the coupling is *successful* if  $P^{1,2}(Z_n^1 = Z_n^2 \text{ for all } n \text{ sufficiently large}) = 1$ .

The key to corollary (1.7) is the following maximal coupling theorem of Goldstein (1979).

**MAXIMAL COUPLING THEOREM.** *Let  $(Z_n^1)_{n \geq 1}$  and  $(Z_n^2)_{n \geq 1}$  be two arbitrary processes taking values in the same Borel space. There exists a successful coupling  $P^{1,2}$  iff  $P^1$  and  $P^2$  agree on the tail  $\sigma$ -field.*

The idea is to apply this theorem to two copies  $(S_n^1)$  and  $(S_n^2)$  of our sum process  $(S_n)$  obtained by conditioning on  $\{X_1 = 0\}$  and  $\{X_1 = 1\}$ , respectively. That is

$$\begin{aligned} P^1((S_n^1) \in \cdot) &= P((S_n) \in \cdot | X_1 = 0), \\ P^2((S_n^2) \in \cdot) &= P((S_n) \in \cdot | X_1 = 1). \end{aligned}$$

If  $\mathcal{G}_\infty^+$  is trivial, then obviously

$$P^1((S_n^1)_{n \geq 1} \in \cdot) = P^2((S_n^2)_{n \geq 1} \in \cdot) \text{ on } \mathcal{G}_\infty^+$$

and so by Goldstein's theorem there exists a successful coupling such that

$$P^{1,2}(S_n^1 = S_n^2 \text{ for all } n \text{ sufficiently large}) = 1.$$

Now let  $(T_k^1)_{k \geq 1}$  and  $(T_k^2)_{k \geq 1}$  denote the corresponding sequences of random positive times at which  $X_n^1$  and  $X_n^2$  assume the value 1. Then we have for any  $t \geq 0$

$$(5.1) \quad \begin{aligned} & |P(T_{k+1} - T_k > t | X_1 = 0) - P(T_{k+1} - T_k > t | X_1 = 1)| \\ &= |P^1(T_{k+1}^1 - T_k^1 > t) - P^2(T_{k+1}^2 - T_k^2 > t)| \\ &\leq P^{1,2}(T_k^1 < \tau^{1,2} \text{ or } T_k^2 < \tau^{1,2}) \end{aligned}$$

where

$$\tau^{1,2} = \inf\{m \geq 1 : S_n^1 = S_n^2 \text{ for all } n \geq m\}$$

denotes the coupling time. Since  $P(\lim_{k \rightarrow \infty} T_k = \infty) = 1$ , we have  $P^{1,2}(\lim_{k \rightarrow \infty} T_k^1 = \lim_{k \rightarrow \infty} T_k^2 = \infty) = 1$  and so

$$(5.2) \quad \lim_{k \rightarrow \infty} P^{1,2}(T_k^1 < \tau^{1,2} \text{ or } T_k^2 < \tau^{1,2}) = P^{1,2}(\tau^{1,2} = \infty) = 0.$$

Conditioning on  $\{X_1 = 1\}$  is natural because the process  $(T_{k+1} - T_k)_{k \geq 1}$  is stationary given  $\{X_1 = 1\}$ , i.e.

$$(5.3) \quad P(T_{k+1} - T_k > t | X_1 = 1) \text{ is independent of } k.$$

For this well-known fact and its implications we refer to Kakutani (1943) and Kac (1947). If we now combine (5.1-3), then the assertion follows. ■

**ACKNOWLEDGMENT:** The authors would like to acknowledge discussions with J. Aaronson and M. Smorodinsky (Tel Aviv) and B. Weiss (Jerusalem). The proof of theorem (1.3) as given in section 2 was suggested by B. Weiss and improves an original proof by the authors.

## References

- Berbee, H.C.P. (1979). Random walks with stationary increments and renewal theory. *Math. Centre Tract* **112**.
- Berbee, H.C.P. (1986). Periodicity and absolute regularity. *Israel J. Math.* **55**, 289-304.
- Bradley, R.C. (1983). Absolute regularity and functions of Markov chains. *Stoch. Process. Appl.* **14**, 67-77.
- Breiman, L. (1968) *Probability*. Addison-Wesley, Reading.
- Goldstein, S. (1979). Maximal coupling. *Z. Wahrscheinlichkeitstheorie verw. Gebiete* **46**, 193-204.
- Hollander, W.Th.F. den (1988). Mixing properties for random walk in random scenery. *Ann. Probab.*
- Janson, S. (1984). Runs in  $m$ -dependent sequences. *Ann. Probab.* **12**, 805-818.
- Kakutani, S. (1943). Induced measure preserving transformations. *Proc. Japan Acad. Ser. A Math. Sci.* **19**, 635-641.
- Kac, M. (1947). On the notion of recurrence in discrete stochastic processes. *Bull. Amer. Math. Soc.* **53**, 1002-1010.
- Ornstein, D.S. (1969). Random walks I. *Trans. Amer. Math. Soc.* **138**, 1-43.
- Parry, W. (1981). *Topics in Ergodic Theory*. Cambridge Tracts in Mathematics **75**. Cambridge University Press.
- Smorodinsky, M. (1971). *Ergodic Theory, Entropy*. Lecture Notes in Mathematics **214**. Springer.
- Stam, A.J. (1967). On shifting iterated convolutions. *Compositio Mathematica* **17**, 268-280, and **18**, 201-228.

Henry Berbee  
Center for Mathematics and Computer Science  
Kruislaan 413, 1098 SJ Amsterdam  
The Netherlands.

Frank den Hollander  
Delft University of Technology  
Faculty of Mathematics and Informatics  
Julianalaan 132, 2628 BL Delft  
The Netherlands.