



Centrum voor Wiskunde en Informatica
Centre for Mathematics and Computer Science

A. Blokhuis, A.E. Brouwer

Locally 4-by-4 grid graphs

Department of Pure Mathematics

Report PM-R8401 January

The Centre for Mathematics and Computer Science is a research institute of the Stichting Mathematisch Centrum, which was founded on February 11, 1946, as a nonprofit institution aiming at the promotion of mathematics, computer science, and their applications. It is sponsored by the Dutch Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O.).

LOCALLY 4-BY-4 GRID GRAPHS

A. BLOKHUIS, A.E. BROUWER

Centre for Mathematics and Computer Science, Amsterdam

We investigate locally grid graphs. Main results are: (i) a characterization of the Johnson graphs (and certain quotients of these) as locally grid graphs such that two points at distance two have precisely four common neighbours, and (ii) a complete determination of all graphs that are locally a 4×4 grid (it turns out that there are four such graphs, with 35, 40, 40 and 70 vertices).

1980 MATHEMATICS SUBJECT CLASSIFICATION: 05C25, 05C30

KEY WORDS & PHRASES: Johnson scheme, locally grid graph.

NOTE: This paper will be submitted for publication elsewhere.

Report PM-R8401

Centre for Mathematics and Computer Science

P.O. Box 4079, 1009 AB Amsterdam, The Netherlands



0. INTRODUCTION

Let us denote by $\binom{n}{m}$ the graph with as vertices the m -subsets of an n -set, where two m -sets are adjacent when they have an $(m-1)$ -set in common. [These graphs are commonly known as Johnson graphs; in the cases $m = 2$ and $m = 3$ also as triangular and tetrahedral graphs.]

Let us denote by $p \times q$ the graph with vertex set $P \times Q$ where $|P| = p$, $|Q| = q$ and (x_1, y_1) adjacent to (x_2, y_2) if and only if $x_1 = x_2$ or $y_1 = y_2$ (but not both). The graphs $p \times q$ are called *grids*. A graph Γ is called *locally G* if for each vertex x of Γ the graph induced by Γ on the set of neighbours of x is isomorphic to the graph G .

With this terminology the graph $\binom{n}{m}$ is locally $m \times (n-m)$ and one might want to characterize it as such; without additional hypotheses this seems impossible (J.I. Hall has indicated large classes of locally grid graphs), but adding the hypothesis that the graph induced on the set of common neighbours of two points at distance two is a 4-cycle (which is true in $\binom{n}{m}$) we do indeed obtain a characterization of the Johnson graphs and certain

quotients.

Let us denote by $\mu(x,y)$ (where x and y are two vertices at distance 2 in a graph Γ) the graph induced on the set of common neighbours of x and y ; we shall call subgraphs of Γ of this form μ -graphs. Characterizing locally $p \times q$ graphs (without hypothesis on the μ -graphs) is trivial for $p \leq 2$ and has been done by J.I. Hall [7,8] for $p = 3$. Here we settle the first unsolved case by finding all locally 4×4 graphs. The result is surprising in that there are besides the expected graphs $\binom{8}{4}$ and $\frac{1}{2}\binom{8}{4}$ (the quotient of $\binom{8}{4}$ obtained by identifying complementary 4-sets of the 8-set), two more examples on 40 vertices, one very regular graph (its automorphism group is transitive on points, edges, triangles, 4-cliques and 5-cliques) described in section 3, and a twisted version of the previous, described in section 4.

It will be clear to those who know the language of Buekenhout-Tits diagrams that what we consider here are geometries belonging to the diagram



Related work (characterizing certain classes of locally polar graphs) was done by Buekenhout & Hubaut [4]. There are two possible ways for a generalized quadrangle to be thin (and thus difficult to handle geometrically) - one is the case with two lines on each point, the case we consider here, and the other is the case with two points on each line. But in the latter case we have a locally $K_{p,q}$ graph, and it is trivial to check that such graphs do not exist for $p \neq q$ while the only locally $K_{p,p}$ graph is $K_{p,p,p}$.

[In contrast to the difficulty of characterizing locally $p \times q$ graphs, it is completely trivial to characterize locally $\overline{p \times q}$ graphs (where the bar denotes complementation) in all interesting cases- cf. Buset [5].]

1. CHARACTERIZATION OF THE JOHNSON GRAPHS

Let us first state the results we shall prove in this section.

THEOREM 1. *Let Γ be connected and locally grid and assume that the connected components of each μ -graph are 4-cycles. Then there are integers m, n such that Γ is locally $m \times (n-m)$, and either $\Gamma \cong \binom{n}{m}$ or $n = 2m$ and Γ is obtained*

from $\binom{2m}{m}$ by identifying each m -set with the image of its complement under σ , where σ is a permutation of the $2m$ symbols satisfying $\sigma^2 = \tau$ and having at least 8 fix points.

REMARK. For $m \leq 4$ we must have $\sigma = 1$ and $m = 4$ so that the only possible quotient is $\frac{1}{2}\binom{8}{4}$. (The quotients $\frac{1}{2}\binom{4}{2}$ and $\frac{1}{2}\binom{6}{3}$ are complete graphs K_3 and K_{10} .) In general the number of transpositions in the cycle representation of σ is an invariant of the quotient so that one obtains several non-isomorphic quotients.

When σ has at least 10 fix points then each μ -graph of the quotient is a 4-cycle; when σ has precisely 8 fix points then each μ -graph is the union of two 4-cycles.

REMARK. Some version of the above theorem was communicated by the second author to J.I. Hall at the Pullman conference in honour of T.G. Ostrom. Hall subsequently showed (in a letter dated May 21, 1981) that this theorem is essentially equivalent to theorem 2 in Sprague [10].

Under the weaker assumption that Γ is locally a graph on $m(n-m)$ vertices with valency $(m-1) + (n-m-1)$ and each μ -graph has at most 4 vertices one can prove that $\Gamma \cong \binom{n}{m}$ provided n is large enough. (T.A. Dowling [6] proved this for $n > 2m(m-1) + 4$ and A.E. Brouwer [2] for $n \geq \max(6m-1, m^2+2m+1)$.)

A. Moon [9] obtains a characterization for $n > 4m$, characterizing $\binom{n}{m}$ not as a graph but as an association scheme, where all the parameters p_{jk}^i of the association scheme are given.

In all cases the difficult part of the proof is to find cliques of the right size - thus (the analogue of) theorem 1 is trivial in these contexts; only when $n = 2m$ one has to be a little bit careful.

COROLLARY. Let Γ be connected locally $2 \times q$. Then Γ is the triangular graph $\binom{q+2}{2}$.

LEMMA. Let Γ be connected and locally grid. Then there are integers p, q such that Γ is locally $p \times q$. Each μ -graph is a union of cycles of even length. If C is a maximal clique and $x \notin C$ then x is adjacent to either 0 or 2


points of C . Each edge is in precisely two maximal cliques: one of size $p+1$ and one of size $q+1$. Each triangle is in a unique maximal clique.

PROOF OF THE LEMMA. Suppose that the neighbourhood of x is isomorphic to $p \times q$ and that $x \sim y$. Then the edge xy is in cliques of size $(p+1)$ and $(q+1)$ and hence the neighbourhood of y is also isomorphic to $p \times q$. By connectedness of Γ the graph is locally $p \times q$.

If $d(u,v) = 2$ and $a \in \mu(u,v)$ then looking at the neighbours of a we find that u,v,a have two common neighbours, so $\mu(u,v)$ is regular of valency 2. Since $\mu(u,v) \subset \Gamma(u)$ the edges of a component of $\mu(u,v)$ are alternately 'horizontal' and 'vertical', i.e., each component of $\mu(u,v)$ is an even cycle. □

PROOF OF THE COROLLARY. The only union of cycles of even length that occurs as subgraph of $2 \times q$ is a 4-cycle. Thus the theorem applies. □

PROOF OF THE THEOREM. By the lemma, Γ is locally $m \times (n-m)$ and we may assume $m \leq n-m$. The proof is by induction on m . For $m = 0$, Γ has valency zero and hence is a single point, indeed $\Gamma \cong \binom{n}{0}$ for all $n \geq 0$. Consider the graph Γ^* that has the $(n-m+1)$ -cliques of Γ as vertices, and where two such cliques are adjacent when they have precisely one point in common. We want to show that each connected component of Γ^* satisfies the hypotheses of the theorem and is locally $(m-1) \times (n-m+1)$.



To this end first observe that if $C = \{x_0, \dots, x_{n-m}\}$ is a maximal clique of Γ then $\Gamma^*(C)$ has a partition into $n-m+1$ cliques of size $m-1$, namely the sets of points in $\Gamma^*(C)$ containing some fixed x_i ($0 \leq i \leq n-m$).

Next assume that C_1 and C_2 are points of Γ^* such that $C_1 \cap C_2 = \{x_0\}$, say

$$C_1 = \{x_0, x_1, \dots, x_{n-m}\}, \quad C_2 = \{x_0, y_1, \dots, y_{n-m}\} \text{ where } x_i \sim y_i \quad (1 \leq i \leq n-m).$$

Let D_i be the maximal clique containing x_i, y_i but not x_0 ($1 \leq i \leq n-m$). Then $\{C_1, C_2, D_1, \dots, D_{n-m}\}$ is a clique in Γ^* . [For: let $i \neq j$. The point x_i has two neighbours in D_j , one is x_j and the other is a point not in $\Gamma(x_0)$, say a . Now $\mu(x_0, a)$ contains x_i, x_j, y_j and hence by our hypothesis on μ -graphs also y_i . Thus D_i is the clique determined by a, x_i, y_i and D_i is

adjacent to D_j in Γ^* .] Thus Γ^* is locally $(m-1) \times (n-m+1)$.

Next we check that the μ -graphs in Γ^* are unions of 4-cycles. Let C_1 and C_2 be as before and let D meet C_2 in the point y_1 , where $|C_1 \cap D| \neq 1$. Then $|C_1 \cap D| = 0$. The point x_0 has two neighbours on D , y_1 and, say z_1 . Looking at $\Gamma(x_0)$ we see that $x_1 \sim z_1$, and $\mu(C_1, D)$ contains the 4-cycle $(C_2, D(x_1, y_1), D(x_1, z_1), C_3)$, where $D(p, q)$ is the maximal clique on p, q not containing x_0 , and $C_3 = \{x_0, z_1, \dots, z_{n-m}\}$ is the maximal clique on x_0, z_1 not containing x_1, y_1 .


Thus, by induction, each component of Γ^* is the Johnson graph $J(n, m-1)$. First assume that $n > 2m$; then each edge is in a unique $(n-m+1)$ -clique and Γ^* is connected. Label the points of Γ^* with $(m-1)$ -sets; next label each point x of Γ with the union of the labels of the $(n-m+1)$ -cliques on x . Then x is labeled with an m -set (since the cliques incident with x form a small maximal clique in Γ^*) and clearly two points of Γ are adjacent iff they are together in a clique, i.e., iff their labels have an $(m-1)$ -set in common. Thus $\Gamma \cong J(n, m)$. [Note Γ has the right number of vertices and no label can be repeated.] Next assume $n = 2m$. Now each edge is in precisely two $(m+1)$ -cliques, and Γ^* has either one or two components. If Γ^* has two components then use the labels of one of these components to label Γ and we find $\Gamma \cong J(n, m)$ again. So assume that Γ^* is connected. Now each point x of Γ determines two m -cliques in Γ^* , and, taking the union of the labels of the members of each m -clique we find two labels for x . Both labels are m -sets, and if $x \sim y$ then each of the labels of x has an $(m-1)$ -set in common with a unique label of y . Conversely, if a label of x and a label of y have an $(m-1)$ -set in common then $x \sim y$ and also the other labels have an $(m-1)$ -set in common. Thus Γ is a quotient of $J(2m, m)$ under an automorphism σ of order 2. Now the automorphism group of $J(2m, m)$ is the direct product $\text{Sym}(2m) \times \mathbb{Z}_2$ where the \mathbb{Z}_2 sends an m -set to its complement. Also, if σ interchanges the m -sets M, N then the Johnson distance between M and N is at least 4 (otherwise the quotient will not be locally $m \times m$), i.e. $|\sigma(M) \cap M| \leq m-4$. If σ is an involution in $\text{Sym}(2m)$ then there are m -sets M such that $|\sigma(M) \cap M| \geq m-1$. Hence σ must be complementation followed by an involution σ_0 in $\text{Sym}(2m)$. Now the requirement $|\sigma(M) \cap M| \leq m-4$ means $|\sigma_0(M) \cap M| \geq 4$, i.e., σ_0 has at least 8 fix points. \square

2. LOCALLY $p \times q$ GRAPHS WITH MAXIMAL μ

LEMMA. Let Γ be locally $p \times q$ and suppose $|\mu(x,y)| = 2p$ for vertices x,y of Γ with $d(x,y) = 2$. Then no neighbour of y has distance 3 to x . In particular, if $|\mu(x,y)| = 2p$ for all pairs x,y with $d(x,y) = 2$ then Γ has diameter (at most) two.

PROOF. Obvious. □

Assume that Γ is locally $m \times n$ with $n \geq m \geq 2$, where each μ -graph has $\mu = 2m$ vertices. Then Γ is strongly regular with diagram



and parameters

$$v = 1 + mn + (m-1) \binom{n}{2} = (n+1) \left(1 + \frac{1}{2}n(m-1)\right)$$

vertices, eigenvalues

$$k = mn, \quad r = n-2, \quad s = -m$$

with multiplicities

$$1, \quad f = \frac{m(m-1)n(n+1)}{2(n+m-2)}, \quad g = v - 1 - f.$$

(For definition and properties of strongly regular graphs and partial geometries see e.g. [3].)

Since f is an integer, we see that for fixed $m > 3$ only finitely many n are possible. For example, when $m = 4$ then $(n+2) \mid 12$ so that $n = 4$ or $n = 10$.

The Hoffman bound for the maximum size of a clique in a strongly regular graph with these parameters yields $|C| \leq n+1$ and tells us that when $|C| = n+1$ and $x \notin C$ then x is adjacent to precisely two points of C .

(It is also easy to see this directly.)

This means that when $n > m$ we have a partial geometry $pg(K=n+1, R=m, T=2)$, while if $n = m$ we have a Zara graph with parameters $K = n + 1, e = 2$.

(For definition and properties of Zara graphs see F. Zara [11], A. Blokhuis [1].) For $n > m$ one easily sees that the line graph of the partial geometry has the property that each of its μ -graphs is the disjoint union of $\frac{1}{2}(n+1)$ 4-cycles. It follows that n must be odd. (But it is possible that $n = m$ for even n .)

This observation immediately kills $(m, n) = (4, 10)$. (Another way to dispose of this parameter set is to observe that the line graph of $pg(11, 4, 2)$ violates the absolute bound and hence does not exist; in general the absolute bound for the line graph states

$$V \leq \frac{1}{2} G(G+3) \text{ for } V = m(1+(m-1) \cdot \frac{1}{2}n) \text{ and } G = m-1 + \frac{m(m-1)(m-3)n}{2(n+m-2)}$$


which yields a restriction of the form $n \leq \frac{1}{2}m^4$.)

Thus, when $m=4$ then necessarily $n=4$, and there is a unique graph in this case:

PROPOSITION. *There is a unique Zara graph with $K = 5, e = 2$ on $v = 35$ vertices. It is $\frac{1}{2}\binom{8}{4}$.*

PROOF. Clearly such a graph must be locally 4×4 . In order to apply the theorem of the previous section we have to prove that the μ -graphs are unions of 4-cycles. To this end, consider two disjoint 5-cliques C, D . The edges joining a point from C to a point from D form a bipartite graph of valency 2, i.e., a union of (even) cycles; the only possibilities are (i) a union of a 4-cycle and a 6-cycle, and (ii) a 10-cycle. We shall show that the former always is the case. Suppose $C = \{a_1, a_2, a_3, a_4, a_5\}$ and let C_{ij} be the 5-clique on a_i and a_j distinct from C ($1 \leq i < j \leq 5$). Then C_{12} and C_{34} are disjoint and we see a 4-cycle (1324) between C_{12} and C_{34} so the pair (C_{12}, C_{34}) is of type (i). Counting all such pairs we find 56.10.3 ordered pairs (C', C'') of type (i). On the other hand each clique C meets itself in 5 points, 10 others in 2 points, 15 others in 1 point and the remaining $(56-1-10-15) = 30$ in 0 points. Thus the number of pairs

of disjoint cliques equals the number of such pairs of type (i) and type (ii) does not occur.




Now let x, y be two points at distance 2, and let $r \sim p \sim q$ be a path of length 3 in $\mu(x, y)$. Let C be the 5-clique on x, r, s , where s is the point completing the 4-cycle $(p q s r)$ in $\Gamma(x)$. Let D be the 5-clique on y, p, q .

Between (the disjoint 5-cliques) C and D we see the path $s \sim q \sim x \sim p \sim r \sim y$ and this must complete to a 6-cycle. Hence $y \sim s$ so that $\mu(x, y)$ contains the 4-cycle $(p q s r)$. □


3. AN INTERESTING GRAPH ON 40 VERTICES

There exists a unique locally 4×4 graph Γ with diagram



with respect to each of its points.

It has 40 vertices and 64 blocks (5-cliques). Its group of automorphisms is $2^6 \cdot S_5$ of order 7680 acting transitively on j -cliques for $0 \leq j \leq 5$. Switching antipodes is an automorphism. If we call two blocks adjacent when they meet in a unique point then the graph Γ^* on the blocks has 2 components of size 32. Each of the components has diagram




and is locally $GQ(2, 2)$.

(Unicity will be the subject of section 5. Here we are concerned with the existence.) The nicest description we know is the following:


Let Δ_{01} be the graph with as vertices the 2×2 matrices with entries in \mathbb{F}_4 and trace 0 or 1 where two matrices are adjacent when their difference has rank 1.

Let $\bar{\Delta}_{01}$ be the quotient of Δ_{01} under identification of A and $A + I$ for all matrices A .

Then $\bar{\Delta}_{01}$ has 64 vertices and diagram




(The subgraph $\bar{\Delta}_0$ of $\bar{\Delta}_{01}$ consisting of the points with trace 0 has diagram



Let Γ be the graph with as vertices the 8-cliques of $\bar{\Delta}_{01}$, two cliques being adjacent when they have nonempty intersection, then Γ has the properties stated above (and the vertices of $\bar{\Delta}_{01}$ are the 5-cliques in Γ).


[From the description one immediately sees the group of automorphisms $2^6 \cdot \text{PFL}(2,4)$ but $\text{PFL}(2,4) \cong S_5$. The 8-cliques of $\bar{\Delta}_{01}$ have 4 points with trace 0 and 4 points with trace 1; in fact the graph Δ on all 256 2×2 matrices over \mathbb{F}_4 is strongly regular with diagram




and the maximal cliques meeting the Hoffman bound are the two-dimensional affine subspaces such that any two matrices in such a subspace differ by a rank 1 matrix. From this it is not difficult to see that Γ will be locally 4×4 . The automorphism switching antipodes is $A \mapsto A + \begin{pmatrix} \omega & 0 \\ 0 & \omega \end{pmatrix}$. The partition of Γ into 5 cocliques of size 8 corresponds to the parallelism on Δ .]

4. ANOTHER GRAPH ON 40 VERTICES

Unfortunately, there is another graph on 40 points which is locally 4×4 . It has diagram



with respect to 8 of its points, while it looks as follows around each of the remaining 32 points:



(thus, its diameter is 3, but for 32 points x one has $\Gamma_3(x) = \emptyset$).

A direct description can be given as follows.

Let $X = \Omega \cup Y \cup Z$, where $\Omega = \{\infty_i, \bar{\infty}_i \mid i \in \mathbb{Z}_4\}$, Y and Z are both copies of $\mathbb{Z}_4 \times \mathbb{Z}_4$ - write (i,j) for elements of Y and $[i,j]$ for elements of Z . The graph will have $d(\infty_i, \bar{\infty}_i) = 3$ ($i \in \mathbb{Z}_4$), while $\Gamma_3(x) = \emptyset$ for $x \in X \setminus \Omega$.

The induced subgraph on Ω is a coclique, and on Y, Z we have the natural 4×4 grid. Each point in $Y \cup Z$ that is non-adjacent to ∞_i is adjacent to $\bar{\infty}_i$, so we need only specify the adjacencies of ∞_i ($i \in \mathbb{Z}_4$).

Let $\Gamma(\infty_0) = Y$ (so that $\Gamma(\bar{\infty}_0) = Z$), and

$$\Gamma(\infty_i) = \{(a,a), (a,a+i), [a,a], [a,a-i] \mid a \in \mathbb{Z}_4\} \text{ for } i = \pm 1, \text{ and}$$

$$\Gamma(\infty_2) = \{(a,a), (a,a+2), [a,a+1], [a,a-1] \mid a \in \mathbb{Z}_4\}.$$

Let the point $(a,b) \in Y$ be adjacent to the 6 points $[a-\delta+i, b+\delta+j]$ for $i, j \in \mathbb{Z}_4, |\{0, i, -j\}| = 3, \delta = b - a$.

This defines a graph Γ , and by inspection it is locally 4×4 . Its automorphism group has order 2^9 , is transitive on the 64 5-cliques and has two orbits of sizes 8, 32 on the 40 points.

The graph Γ^* with as vertices the 5-cliques in Γ , where $C \sim C'$ iff $|C \cap C'| = 1$ has two components of size 32.

5. THE LOCALLY 4×4 GRAPHS

THEOREM. *There are up to isomorphism precisely four locally 4×4 graphs, namely*

- (i) $\binom{8}{4}$ (with $v = 70$ vertices, all μ -graphs are 4-cycles),
- (ii) $\frac{1}{2} \binom{8}{4}$ (with $v = 70$ vertices, all μ -graphs are unions of two 4-cycles),
- (iii) G_{40} , the graph described in section 3 (with $v = 40$ vertices, each μ -graph is either a 6-cycle or a union of two 4-cycles).
- (iv) G'_{40} , the graph described in section 4 (with $v = 40$ vertices, each μ -graph is either a 4-cycle, a 6-cycle, an 8-cycle or the union of two 4-cycles).

The proof is split up in a series of lemmas. Let Γ be a locally 4×4 graph. Write $\Gamma_i(x)$ for the set of vertices at distance i from a vertex x .

LEMMA 1. (i) Γ has valency 16, each edge is in 6 triangles and the graph $\lambda(p, q)$ on the common neighbours of two adjacent points p and q is the union of two triangles.

- (ii) Each edge is in two 5-cliques; each triangle is in a unique 5-clique.
- (iii) If Γ has v vertices and b 5-cliques then $b = \frac{8}{5}v$. In particular $5|v$.
- (iv) For $d(p, q) = 2$ the graph $\mu(p, q)$ is either a 4-cycle or a 6-cycle or an 8-cycle or the union of two 4-cycles. If $|\mu(p, q)| = 4, 6, 8$, then q has at most 4, 1, 0 neighbours in $\Gamma_3(p)$ (respectively). \square

LEMMA 2. *If C is a 5-clique and $d(x, C) = 2$ then x has distance two to at least 3 points of C . Consequently, if $d(x, y) = 3$ then the number of neighbours of y at distance two to x is either 9 or at least 12.*

PROOF. (i) Let $x \sim p$ with $d(p,C) = 1$. Then p has two neighbours on C , say y and z . $\mu(x,y)$ is not a clique so cannot be contained in $\lambda(y,z)$. It follows that $x \sim q$ where q has two neighbours on C but $q \not\sim z$.

(ii) Now assume $d(x,y) = 3$. Looking at the 4×4 grid that is the neighbourhood of y we see on each line 0, 3 or 4 points at distance 2 to x . \square

LEMMA 3. *Equivalent are:*

(i) $v \geq 70$, (ii) $\text{diam } \Gamma \geq 4$, (iii) $\Gamma \cong \binom{8}{4}$.

PROOF. Estimating the number of points k_i at distance i from a given point x we find

$$k_0 = 1, k_1 = 16, k_2 \leq \frac{16 \cdot 9}{4} = 36, k_3 \leq \frac{36 \cdot 4}{9} = 16, k_4 \leq \frac{16 \cdot 1}{9} = 1, k_5 \leq \frac{1 \cdot 1}{9} = 0.$$


Hence $v \leq 1 + 16 + 36 + 16 + 1 = 70$ with equality iff $|\mu(x,y)| = 4$ for all pairs x,y with $d(x,y) = 2$. Also, if $\text{diam } \Gamma = 4$ then $k_4 = 1$ and the 16 neighbours of a point at distance 4 of x must have distance to x so that $k_3 = 16$ and we have equality everywhere so that $v = 70$. Now apply the theorem in section 1. \square

From now on we may assume that $\text{diam } \Gamma \leq 3$.

LEMMA 4. *Let $d(x,C) = 2$ and $|\Gamma_2(x) \cap C| = 3$. Then $|\mu(x,c)| = 4$ for all $c \in \Gamma_2(x) \cap C$.*

PROOF. Follows immediately from lemma 1 (iv). \square

REMARK. In the case of this lemma one necessarily has the following situation:



(Here the points in $\Gamma(x)$ labeled ij are adjacent to c_i and c_j .)

I.e., the points in $\Gamma(x)$ at distance 1 from C lie on a 2×3 grid.

LEMMA 5. *For no point x and 5-clique C do we have $C \subset \Gamma_3(x)$.*

PROOF. Let $A = \{a \mid a \sim x, d(a,C) = 2\}$, $B = \{b \mid b \in \Gamma_2(x), d(b,C) = 1\}$.

Let $C = \{c_1, c_2, c_3, c_4, c_5\}$.


(i) The neighbours of a point $c_i \in C$ lie on the four cliques C_{ij} on c_i, c_j ($j \neq i$) with $C_{ij} \neq C$. Each c_i has at least 9 neighbours in B , i.e., there is at most one clique C_{ij} on c_i not meeting B . But $|C|$ is odd, so there is a point $c \in C$, say $c = c_1$ such that all the cliques C_{ij} ($j \neq 1$) meet B .

(ii) For $b \in B$ we have $|\mu(b,x)| = 4$ (cf. Lemma 1 (iv)), so $\sum_{b \sim c} |\mu(b,x)| = 48$.

$$\text{But } 48 = \sum_{b \sim c} |\mu(b,x)| = \sum_{\substack{a \sim x \\ d(a,c)=2}} |\mu(a,c)|, \text{ and } |\mu(a,c)| \in \{4,6\} \text{ for } a \in A \cap \Gamma_2(c).$$


If for some $a \in A \cap \Gamma_2(c)$ we have $|\mu(a,c)| = 6$, then $|A \cap \Gamma_2(c)| < 12$ so $|A \cap \Gamma_2(c)| = 9$ and $A \cap \Gamma_2(c)$ forms a 3×3 grid in $\Gamma(x)$. If D is a 5-clique on a and x then $|D \cap \Gamma_2(c)| = 3$ so $|\mu(a,c)| = 4$ by Lemma 4, contradiction. Hence always $|\mu(a,c)| = 4$ and $|A \cap \Gamma_2(c)| = 12$.

Considering the adjacencies between $\Gamma(c)$ and A we see that one necessarily has



(where points in $A \cap \Gamma_2(c)$ are labeled with the 4 points of $B \cap \Gamma(c)$ they are adjacent to).

Let a be the point labeled "12pq" and consider the neighbourhood of a :



Let D be the clique $\{7,8,9,c,d\}$. We see that $|\mu(a,7)| = |\mu(a,8)| = 6$, $|\mu(a,9)| = 0$, (since it is impossible to find a cycle in $\Gamma(a)$ avoiding the non-neighbours of the point 9) $|\mu(a,c)| = 4$, $|\mu(a,d)| = 0$. But this contradicts the previous lemma. \square


COROLLARY. If $d(x,y) = 3$ then $|\Gamma(y) \cap \Gamma_2(x)| \geq 12$. □

LEMMA 6. Let C be a 5-clique, $d(x,C) = 2$ and $|\Gamma_2(x) \cap C| = 4$.
Then there is a $\mu \in \{4,6\}$ so that for all $c \in \Gamma_2(x) \cap C$ we have $|\mu(x,c)| = \mu$.

PROOF. If $c, c' \in \Gamma_2(x) \cap C$ then $\mu(x,c) \cap \mu(x,c')$ is either empty or an edge (for if $d \in \mu(x,c) \cap \mu(x,c')$ then x has two neighbours on the 5-clique on c, c', d). The edges of this form are pairwise disjoint and partition $\frac{1}{2} \sum_{c \in \Gamma_2(x) \cap C} \mu(x,c)$ points, so this latter number is even. This means that if $|\mu(x,c)|$ is not constant then it is twice 4 and twice 6. But one quickly checks that it is impossible to find two 4-cycles in $\Gamma(x)$, each having one edge in common with a 6-cycle in $\Gamma(x)$ (where these two edges are parallel). □

LEMMA 7. Suppose $d(x,y) = 3$, $d(x,z) = 2$, $y \sim z$, $|\mu(x,z)| = 6$.
Then $|\Gamma_3(x)| = 1$ and $v = 40$ and $|\mu(x,p)| \in \{6,8\}$ for all $p \in \Gamma_2(x)$.

PROOF. Look at the neighbourhood of y . Lemma's 4 and 6 state that if for one point z of a line in this 4×4 grid we have $d(x,z) = 2$ and $|\mu(x,z)| = 6$ then this holds for all points of this line. Thus $\Gamma(y) \subset \Gamma_2(x)$ and for each neighbour z of y we have $|\mu(x,z)| = 6$. This leads to the situation



: from $\Gamma(x)$ there are 144 edges to $\Gamma_2(x)$; 96 end in $\Gamma(y)$, 48 in $\Gamma_2(x) \setminus \Gamma(y) =: Z$. [Clearly $6 = \frac{48}{8} \leq |Z| \leq \frac{48}{4} = 12$.] By Lemma 1(iv), the points in $\Gamma(y)$ have no neighbours in $\Gamma_3(x) \setminus \{y\}$, so any additional point in $\Gamma_3(x)$ has all its neighbours in $Z \cup \Gamma_3(x)$. We show that Z is a coclique: in fact looking at the neighbours of a point $a \in \Gamma(x) \cup \Gamma(y)$ we see that

the three neighbours of a in Z form a coclique. Now if we have an edge pq in Z then pq is contained in a 5-clique $C \subset Z \cup \Gamma_3(x)$. Let $x \sim r \not\sim p$, then r has two neighbours in $C \cap Z$, contradiction.

Now it follows that $\Gamma_3(x) = \{y\}$, since any additional point has at least 12 neighbours in Z , and Z could not be a coclique.

Since there emerge precisely 96 edges from Z we have $|Z| = 6$, $v = 40$. \square

REMARK. Of course the situation of this lemma will yield conclusion (iii) of the Theorem, but we shall first prove that when we know that the situation is the same with respect to any point x of Γ .

LEMMA 8. *Suppose $C \subset \Gamma_2(x)$ for some 5-clique C .*

Then $|\mu(c,x)| = 4$ for a unique point $c \in C$ and $|\mu(c,x)| = 6$ for all other points $c \in C$.

PROOF. As we saw in the proof of lemma 6, the sets $\mu(c,x) \cap \mu(c',x)$ are either empty or an edge (for $c, c' \in C$), and the edges of this form partition the $\frac{1}{2} \sum_{c \in C} |\mu(c,x)|$ points in $\Gamma(x)$ at distance 1 to C . Thus the vector $\underline{\mu} = (|\mu(c,x)|)_{c \in C}$ takes one of the values $(4,4,4,4,4)$, $(4,4,4,6,6)$ or $(4,6,6,6,6)$. In the first case we have five 4-cycles in $\Gamma(x)$, covering five parallel edges, impossible. In the second case we find a 6-cycle having an edge in common with each of two disjoint 4-cycles (these edges being parallel), and we saw already in the proof of lemma 6 that this is impossible. So the third case $\underline{\mu} = (4,6,6,6,6)$ holds. \square

LEMMA 9. *If for some pair of points x, z with $d(x, z) = 2$ we have $|\mu(x, z)| = 4$, then $\Gamma_3(x) = \emptyset$.*

PROOF. Let $A = \{a \in \Gamma_2(x) \mid \Gamma(a) \cap \Gamma_3(x) = \emptyset\}$ and $B = \Gamma_2(x) \setminus A$. If for some $b \in B$ we have $|\mu(b,x)| = 6$ then we are in the situation of lemma 7 and $|\mu(x,z)| = 4$ is impossible. Hence we have $|\mu(b,x)| = 4$ for each $b \in B$. Considering the neighbourhood of a point $b \in B$ we see at least one point in $\Gamma_3(x)$, and for each b' in $\Gamma(b) \cap \Gamma(y)$ we have $|\mu(x, b')| = 4$, so by lemma 8 no 5-clique on b is contained in $\Gamma_2(x)$: 4 of the 5-cliques on b meet $\Gamma(x)$ and the remaining 4 meet $\Gamma_3(x)$. But this means that no point from B is adjacent to a point in A . Let C be a 5-clique meeting A .

Since $A \subset \Gamma_3(y)$ for $y \in \Gamma_3(x)$ we have $|C \cap A| \leq 2$. Also $|C \cap \Gamma(x)| \leq 2$. But there is nowhere the fifth point of C could go, so there is no such C and $A = \emptyset$. (Now $|\Gamma_2(x)| = 36$.)

Let $y \in \Gamma_3(x)$. Counting paths of length 3 between X and Y we see that $p := |\Gamma(y) \cap \Gamma_2(x)| = |\Gamma(x) \cap \Gamma_2(y)|$. If $p = 12$ then there is no room for a 2×3 grid in $\Gamma(x) \cap \Gamma_2(y)$ so each 5-clique on Y has 4 points in $\Gamma_2(x)$ and $p = 16$, contradiction. Thus $p > 12$, and estimating $\Gamma_3(x)$ we find $5 = \frac{\lceil 2 \cdot 36 \rceil}{16} \leq |\Gamma_3(x)| \leq \lfloor \frac{4 \cdot 36}{13} \rfloor = 11$ so that $58 \leq v \leq 64$ and since $5|v$ we must have $v = 60$, $|\Gamma_3(x)| = 7$.

Now a counting argument will kill this situation: let s be the number of edges between $\Gamma_2(x)$ and $\Gamma_3(x)$, t the number of edges in $\Gamma_3(x)$ and a_i ($2 \leq i \leq 4$) the number of points $b \in \Gamma_2(x)$ with precisely i neighbours in $\Gamma_3(x)$. Then we have

$$\begin{aligned} a_2 + a_3 + a_4 &= 36 \\ 2a_2 + 3a_3 + 4a_4 &= s \\ a_3 + 2a_4 &= t \\ s + 2t &= 7 \cdot 16 = 112 \end{aligned}$$

so that $s = t + 72$, $3t = 40$ and t is not integral. \square

LEMMA 10. *If $\Gamma_3(x) = \emptyset$ for some x then $\Gamma \cong \frac{1}{2} \binom{8}{4}$ or $\Gamma \cong G'_{40}$.*

PROOF. $18 = \frac{144}{8} \leq |\Gamma_2(x)| \leq \frac{144}{4} = 36$. If $|\Gamma_2(x)| = 18$ then $v = 35$ and each μ -graph has 8 points, so that Γ is strongly regular. By Lemma 8 Γ is a Zara graph with $K = 5$, $e = 2$, and now the proposition in section 2 shows that $\Gamma \cong \frac{1}{2} \binom{8}{4}$. Now let $v > 35$, and count cliques. There are 8 5-cliques on x , 48 cliques not on x meeting $\Gamma(x)$ and hence $\frac{8}{5}(v-35)$ cliques contained in $\Gamma_2(x)$. Let $M_\mu = \{b \in \Gamma_2(x) \mid |\mu(x,b)| = \mu\}$. By lemma 8 each clique C in $\Gamma_2(x)$ has 4 points in M_6 and 1 point in M_4 , so $|M_6| = \frac{16}{5}(v-35)$, $|M_4| = \frac{2}{5}(v-35)$ and $v = 1 + 16 + \frac{18}{5}(v-35) + |M_8|$, $144 = 4|M_4| + 6|M_6| + 8|M_8| = \frac{104}{5}(v-35) + 8|M_8|$. It follows that $v < 45$, and since $5|v$ we have $v = 40$, $|M_8| = 5$, $|M_6| = 16$, $|M_4| = 2$. Now enough information has been gathered to define all adjacencies without loss of generality (a rather tedious task). One finds a unique solution, namely the graph described in section 4. \square


LEMMA 11. If $\Gamma_3(x) \neq \emptyset$ for all x , then $\Gamma \cong G_{40}$, the graph described in section 3.

PROOF. Clearly it suffices to show that Γ is determined up to isomorphism.


By lemma's 1(iv), 7, 9, 10 we have that for each $x \in \Gamma$ there is a unique antipode $\bar{x} \in \Gamma$ with $d(x, \bar{x}) = 3$. Clearly the antipodes of the points in $\Gamma(x)$ are in $\Gamma(\bar{x})$, so $x \mapsto \bar{x}$ is an automorphism of Γ .

Fix $x \in \Gamma$ and let $Z = \Gamma_2(x) \cap \Gamma_2(\bar{x}) (=M_8$, see lemma 7). Then $Z = \{p, q, r, \bar{p}, \bar{q}, \bar{r}\}$. Thus given x we find a clique of size 8 $O_x = Z \cup \{x, \bar{x}\}$, and starting with another point, say $p \in O_x$, we find $O_p = O_x$. (Note that $\mu(x, p)$ and $\mu(x, \bar{p})$ partition $\Gamma(x)$.) Consequently, we have a partition of Γ in five 8-cocliques.

Consider the quotient $\bar{\Gamma}$ of Γ under identifying a with \bar{a} for each $a \in \Gamma$. It has 20 points, a partition into five 4-cocliques and valency 16, i.e. is the complete multipartite graph $K_{5 \times 4}$. There are two ways of distributing four 4-cocliques over $\Gamma(x)$, corresponding to the two latin squares of order 4, namely



In the first case, label the 4 rows of $\Gamma(x)$ with a, b, c, d and let $a1$ be the point in row a and coclique 1, etc. Let $A1 = \bar{a1}$, etc. Now we see locally at $a1$:



Since $p \neq q$, $\mu(p, q) \cap \Gamma(x)$ is a coclique (and idem with \bar{x} instead of x) so $|\mu(p, q) \cap \Gamma(x)| = 4$. Also, this remains a coclique in $\bar{\Gamma}$, so $\mu(p, q) \cap \Gamma(x)$ consists of four points labeled with the same digit. Since there are 4 points in Z distinct from p, \bar{p} and each point in $\Gamma(x)$ has three neighbours in Z we see that w.l.o.g. the points $p, \bar{p}, q, \bar{q}, r, \bar{r}$ are adjacent to the

points in $\Gamma(x)$ labeled 1 or 2, 3 or 4, 1 or 3, 2 or 4, 2 or 3, 1 or 4 respectively. Taking antipodes we find the edges between Z and $\Gamma(\bar{x})$. Also the edges between $\Gamma(x)$ and $\Gamma(\bar{x})$ are determined: e.g. the point A_1 is not adjacent to a point in the same row or column as a_1 and not adjacent to a point in coclique 1 so its neighbours in $\Gamma(x)$ must be $b_4, b_3, c_4, c_2, d_3, d_2$.

□

Thus we showed that for $\text{diam } \Gamma = 2, 3, 4$ we have $\Gamma \cong \frac{1}{2} \binom{8}{4}, G_{40}$ or $G'_{40}, \binom{8}{4}$, respectively. Clearly this proves the Theorem.

Acknowledgement. We thank prof. Timmesfeld for suggesting the way S_5 could act on a module 2^6 and thus providing us with the description of the graph G_{40} given in section 3.

REFERENCES

- [1] BLOKHUIS, A., *Few-distance sets*, thesis, Techn. Univ. Eindhoven, 1983.
- [2] BROUWER, A.E., *A note on the uniqueness of the Johnson scheme*, MC report ZN 105, Amsterdam, Sept. 1983.
- [3] BROUWER, A.E. & J.H. van LINT, *Strongly regular graphs and partial geometries*, in: *Proceedings Silver Jubilee Conference*, S.A. Vanstone (ed.), to appear.
- [4] BUEKENHOUT, F. & X. HUBAUT, *Locally polar spaces and related rank 3-groups*, J. Algebra 45 (1977) 391-434.
- [5] BUSET, D., *Graphs which are locally a cube*, Discrete Math. 46, (1983) 221-226.
- [6] DOWLING, T.A., *A characterization of the T_m graph*, J. Combinatorial Theory 6 (1969) 251-263.
- [7] HALL, J.I., *Classifying copolar spaces and graphs*, Quarterly J. Math. 33 (1982) 421-449.
- [8] HALL, J.I., *Personal communication*.

- [9] MOON, A., *On the uniqueness of the graphs $G(n,k)$ of the Johnson schemes*, *J. Combinatorial Theory (B)* 33 (1982) 256-264.
- [10] SPRAGUE, A., *Pasch's axiom and projective spaces*, *Discrete Math.* 33 (1981) 79-87.
- [11] ZARA, F., *Graphes liés aux espaces polaires*, preprint.

ONTVANGEN 1 5 MAART 1984