

**stichting
mathematisch
centrum**



AFDELING ZUIVERE WISKUNDE
(DEPARTMENT OF PURE MATHEMATICS)

ZW 158/81

MAART

A.E. BROUWER

POLARITIES OF G. HIGMAN'S SYMMETRIC DESIGN AND
A STRONGLY REGULAR GRAPH ON 176 VERTICES

Preprint

kruislaan 413 1098 SJ amsterdam

Printed at the Mathematical Centre, 413 Kruislaan, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O.).

Polarities of G. Higman's symmetric design and a strongly regular graph on 176 vertices *)

by

A.E. Brouwer

ABSTRACT

We investigate the polarities of G. Higman's symmetric 2 - (176,50,14) design and find that there are two of them (up to conjugacy), one having 80 and the other 176 absolute points. From the latter we can derive a strongly regular graph with parameters $(v,k,\lambda,\mu) = (176,49,12,14)$. Its group of automorphisms is $\text{Sym}(8)$ with orbits of size 8 and 168 on the vertices. It does not carry a partial geometry or a delta space.

KEY WORDS & PHRASES: *Higman-Sims simple group, strongly regular graph*

*) This report will be submitted for publication elsewhere.

INTRODUCTION

It is well-known that given a symmetric $2 - (V, K, \Lambda)$ design with a polarity, we can find a strongly regular graph with parameters $(v, k, \lambda, \mu) = (V, K, \Lambda, \Lambda)$ if the polarity has no absolute points, and one with parameters $(V, K-1, \Lambda-2, \Lambda)$ if all points are absolute. (In the first case take the symmetric incidence matrix of the design as adjacency matrix of the graph; in the second case first delete the diagonal.)

In [4] G. HIGMAN constructs a symmetric $2 - (176, 50, 14)$ design with the Higman-Sims group as group of automorphisms, and indicates a polarity. Using the description given in SIMS [6] it is trivial to check that the polarity has only absolute points (but note that Sims's statement about the orbit lengths is inaccurate) so that we get a $(176, 49, 12, 14)$ graph.

Computing the multiplicities of the eigenvalues shows that no $(176, 50, 14, 14)$ graph can exist, i.e., any polarity of the design must have absolute points. In fact the number of nonabsolute points is divisible by 12, so that the number of absolute points is congruent $8 \pmod{12}$. Below we study this graph in some more detail. Looking at the possible polarities of the symmetric design we find that up to conjugation there is only one other polarity - one with 80 absolute points. Consequently our graph is the only one that can be derived in this way from the symmetric design.

1. G. HIGMAN'S SYMMETRIC DESIGN

C.C. SIMS [6] gives the following description of a symmetric $2 - (176, 50, 14)$ design: There are six ways to make a set of six elements into a projective line, or, equivalently there are six transitive representations of $\text{PGL}(2, 5)$ on six given points. Now fix a set B of eight elements and let C be the set of $168 = 6 \cdot \binom{8}{2}$ groups $\text{PGL}(2, 5)$ acting on B with orbits of sizes 1, 1, 6. Both the points and the blocks of the design will be indexed by $B \cup C$. If $x \in B \cup C$ then let x denote a point of the design and L_x a block of the design; $\pi: x \mapsto L_x$ will be a polarity of the design. Incidences are defined as follows:

$$\text{for } b \in B: L_b = B \cup \{H \in C \mid H \text{ fixes } b\},$$

for $H \in C$: $L_H = \{b \in B \mid H \text{ fixes } b\} \cup$

$\cup \{G \in C \mid H \text{ has less than 60 conjugates under } N_{\text{sym}(8)}(G)\}.$

Since the 168 elements of C fall into orbits of sizes $1 + 5 + (6+6) + 30 + (30+30) + 60$ under conjugation with elements of G (where the bracketed orbits fuse under $N(G)$) we see that $|L_b| = 8 + 42 = 50$ and $|L_H| = 2 + 1 + 5 + 12 + 30 = 50$ so that blocks have size 50. It is an easy (but somewhat tedious) exercise to check that $\lambda = 14$. {It might be useful to notice that the orbits can be described as follows:

- * 1: $G = H$;
- * 5: G and H have same fixed points but are not identical;
- * 6+6: G and H have one fixed point in common and $|G \cap H| = 20$ (i.e., $G \cap H$ contains an element of order 5);
- 30+30: G and H have one fixed point in common and $|G \cap H| = 4$;
- * 30: G and H have no fixed points in common and $|G \cap H| = 4$ (i.e., $G \cap H$ contains an element of order 4);
- 60: G and H have no fixed points in common and $|G \cap H| = 2$.

Here the asterisks mark the cases where $G \in L_H$ or, equivalently $H \in L_G$. So, obviously, $x \mapsto L_x$ indeed is a polarity.}

The automorphism group of this design is the simple Higman-Sims group HS . It has index 2 in the group of correlations $\langle HS, \pi \rangle \cong \text{Aut } HS$. From the description given it is clear that $\text{Sym}(8)$ centralizes π ; in fact $C_{HS}(\pi) \cong \text{Sym}(8)$. Consequently, if Γ is the graph derived from the design with polarity π then $\text{Aut } \Gamma \cong \text{Sym}(8)$, and $\text{Aut } \Gamma$ has orbits B and C of sizes $8 + 168$ on the vertices of Γ .

2. POLARITIES OF THE DESIGN

From the character table (see FRAME [2]) we see that there are two conjugacy classes of involutions in $(\text{Aut } HS) \setminus HS$. One is the above polarity with 176 absolute points; we want to find the number of absolute points under the

other polarity. To this end, consider the permutation representation of Aut HS on the unordered flags, and compute the corresponding character χ . For the classes of involutions C'_1 and C'_{10} (in Frame's notation) we find $\chi(C'_1) = 176$ and $\chi(C'_{10}) = 80$ so that the other polarity has only 80 absolute points and cannot be used to construct a strongly regular graph. (As follows: first identify the elements of $\text{Sym}(7)$, the stabilizer of a flag in HS, among the conjugacy classes of HS. In fact we can identify the elements of $\text{Sym}(8)$ as follows:

$$\begin{aligned} 1^8 &\subset C_1, & 17 &\subset C_4, & 1^5 3 \cup 1^2 3^2 &\subset C_6, \\ 12^2 3 \cup 26 &\subset C_7, & 1^4 2^2 \cup 2^4 &\subset C_8, & 2^2 4 \cup 4^2 &\subset C_9, \\ 1^2 24 &\subset C_{10}, & 8 &\subset C_{11}, & 35 &\subset C_{14}, & 1^3 5 &\subset C_{15}, \\ 125 &\subset C_{16}, & 1^6 2 \cup 1^2 2^3 &\subset C_{17}, & 1^3 23 \cup 1^2 6 \cup 23^2 &\subset C_{18}, \\ 34 &\subset C_{19}, & 1^4 4 &\subset C_{20}. \end{aligned}$$

Next observe that χ is induced by 1 on $\text{Sym}(7)$, so that we find the decomposition of χ in HS by Frobenius reciprocity. This yields (in HS)

$$\chi = 1 + 77 + 3 \cdot 175 + 231 + 2 \cdot 693 + 154 + 1386 + 2 \cdot 2520$$

so that χ has values

$$(8800, 0, 1, 0, 25, 1, 160, 0, 8, 0, 0, 0, 30, 2, 72, 9, 1, 160, 0, 0, 0)$$

on the classes of HS (in the order given by Frame). The requirement that χ has nonnegative values now determines χ on the classes of $(\text{Aut HS}) \setminus \text{HS}$. We find the decomposition (in Aut HS)

$$\chi = 1 + 77 + 2 \cdot 175 + \overline{175} + \overline{231} + 2 \cdot 693 + 154 + \overline{1386} + 2520 + \overline{2520}$$

and the values

$$(176, 1, 5, 1, 16, 0, 0, 0, 0, 80, 5, 5, 0, 6, 0, 0, 0, 0)$$

on the classes of $(\text{Aut HS}) \backslash \text{HS}$. Here the bar above a character denotes the associate character (i.e., the negative of the column given by Frame). The computations were checked by explicit computation of χ on the classes of $\text{Sym}(8)$.}

We can describe the polarities explicitly in several ways.

A. In Sims' representation the elements 1^8 and $1^6 2$ of $\text{Sym}(8)$ give a polarity with 176 absolute points, while $1^4 2^2$ and $1^2 2^3$ give polarities with 80 absolute points.

B. MARGARET SMITH [7] has given a beautiful description of the symmetric design, as follows: Let u, v be two fixed points of the Steiner system $S(5,8,24)$. Let the points of the design be the blocks of the Steiner system containing u but not v . Let the blocks of the design be the blocks of the Steiner system containing v but not u . Define incidence between points and blocks by $B \sim B'$ iff $|B \cap B'| \in \{0,4\}$. Using the parameters of the near-hexagon associated with $S(5,8,24)$ one immediately sees that this defines a design with the right parameters.

From CONWAY [1] we learn an explicit representation of $S(5,8,24)$ showing large parts of its group of automorphisms. In fact consider 4×6 matrices with entries 0 or 1 satisfying the following restraints:

0	1		1	1		
1		1				
ω				1	1	
$\bar{\omega}$				1		1
	(0	1	0	1	ω	$\bar{\omega}$)

- (i) *the six column sums and the sum of the first row have the same parity.*
- (ii) *if r_i denotes the i -th row ($1 \leq i \leq 4$) and $\mathbb{F}_4 = \{0, 1, \omega, \bar{\omega}\}$ and C is the linear $[n, k, d] = [6, 3, 4]$ code over \mathbb{F}_4 generated by the rows of the matrix*

$$\begin{pmatrix} 1 & 0 & 0 & 1 & \omega & \bar{\omega} \\ 0 & 1 & 0 & 1 & \bar{\omega} & \omega \\ 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}$$

then $r_2 + \omega r_3 + \bar{\omega} r_4 \in C$.


It is almost trivial to verify that these matrices form a linear $[24,12,8]$ code over \mathbb{F}_2 , i.e., the extended Golay code. The Steiner system $S(5,8,24)$ is obtained by taking the matrices with weight 8.

Now if σ is an involution in M_{24} interchanging u and v (say $u = (1,5)$, $v = (1,6)$) then σ defines a polarity of the design. The involutions with 8 fixed points (such as: interchange the last two rows and the last two columns) have 176 absolute points, and those without fixed points (such as: interchange 1st and 2nd, 3rd and 4th, 5th and 6th column, and 2nd and 3rd row) have 80 absolute points.

3. THE STRUCTURE OF THE GRAPH Γ


Let Γ be the $(176,49,12,14)$ strongly regular graph constructed above. By choosing one fixed polarity we destroyed much of the symmetry in the design - the graph Γ is a rather ugly-looking animal. As we saw above, its group is $\text{Sym}(8)$ and has orbits of sizes 8 and 168 on the vertices. Since the graph has the right parameters for the point graph of a partial geometry $\text{pg}(8,7,2)$ we investigate its structure somewhat more closely. Let a *line* be an 8-clique.

Each point $b \in B$ is in 8 lines, forming the 'comb' configuration in Γ_b :



In fact Γ_b can be obtained from the Hoffman-Singleton graph by deleting a vertex x_0 , making its set of neighbours $\{x_1, \dots, x_7\}$ into a 7-clique, and then making the set of neighbours of x_i together with x_i itself into a 7-clique for $1 \leq i \leq 7$.


Each point $c \in C$ is in 6 lines, forming a comb with two teeth missing:



Γ_c contains the following maximal cliques: $18 \cdot K_2$, $50 \cdot K_3$ and $6 \cdot K_7$. It can be obtained from the Hoffman-Singleton graph by deleting a vertex x_0 , making its set of neighbours $\{x_1, \dots, x_7\}$ into a 7-clique, next making the set of neighbours of x_i together with x_i itself into a 7-clique for $1 \leq i \leq 5$, and finally putting a $K_{6,6}$ between the neighbours of x_6 and those of x_7 . We see that 12 of the neighbours of c are not in 7-cliques, i.e., 12 of the edges incident with c are not in a line, and the graph does not carry a partial geometry.

The complementary graph $\bar{\Gamma}$ satisfies the parameter restrictions for delta-spaces $\Delta(3,63,35,45)$ or $\Delta(4,42,35,30)$. But if ab is an edge of Γ contained in two lines then the subgraph of $\bar{\Gamma}$ consisting of the points non-adjacent to both a and b is the complete bipartite graph $K_{6,6}$ and does not contain cliques of size 3 or 4. Hence $\bar{\Gamma}$ does not carry a nontrivial delta-space.

It is not inconceivable that a distance regular graph with diagram



exists. The 'distance 3' graph has the parameters of $\bar{\Gamma}$, and so such a distance regular graph might be obtained by choosing 7 of the 49 edges incident with each vertex of Γ . However, I cannot see a reasonable way to do this.

REFERENCES

- [1] CONWAY, J.H., Oral communication by J.H. van Lint, Oct. 1980.
- [2] FRAME, J.S., *Computation of characters of the Higman-Sims group and its Automorphism group*, J. Algebra 20 (1972) 320-349.

- [3] HIGMAN, D.G. & C.C. SIMS, *A simple group of order 44,352,000*, Math. Z. 91 (1966) 70-86.
- [4] HIGMAN, G., *On the simple group of D.G. Higman and C.C. Sims*, Illinois J. Math. 13 (1969) 74-80. MR 39 # 1545.
- [5] MAGLIVERAS, S.S., *The subgroup structure of the Higman-Sims simple group*, Bull. Amer. Math. Soc. 77 (1971) 535-539.
- [6] SIMS, C.C., *On the isomorphism of two groups of order 44,352,000*, in: *Theory of Finite Groups - A Symposium*, Brauer and Sah (eds), Benjamin, New York, 1969, 101-108.
- [7] SMITH, M.S., *On the isomorphism of two simple groups of order 44,352,000*, J. Algebra 41 (1976) 172-174.
- [8] SMITH, M.S., *A combinatorial configuration associated with the Higman-Sims simple group*, J. Algebra 41 (1976) 175-195.

ONTVANGEN 27 APR. 1981