

**stichting
mathematisch
centrum**



AFDELING ZUIVERE WISKUNDE
(DEPARTMENT OF PURE MATHEMATICS)

ZW 133/79 DECEMBER

M. VOORHOEVE

ANGULAR VARIATION AND THE ZEROS
OF CERTAIN FUNCTIONS

Preprint

2e boerhaavestraat 49 amsterdam

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O).

Angular variation and the zeros of certain functions^{*)}

by

M. Voorhoeve

ABSTRACT

In this paper we study the zeros of functions

$$f(z) = \int_{\Gamma} e^{\zeta z} d\mu(\zeta),$$

where Γ is a Jordan arc and μ a complex measure on Γ .

Under certain conditions (i.e. if Γ is "almost a straight line"), we give explicit upper bounds for the number of zeros of f in a disk. The most important tool for proving this result is the concept of angular variation of a sequence developed in this paper.

KEY WORDS & PHRASES: *Zeros of analytic functions, Laplace transform, power sums*

^{*)} This report will be submitted for publication elsewhere.

1. INTRODUCTION

In this paper we investigate the zeros of the Laplace transform of a measure μ , i.e. the function

$$(1.1) \quad f(z) = \int_{\Gamma} e^{\zeta z} d\mu(\zeta),$$

where $\Gamma \subset \mathcal{C}$ is a Jordan arc and μ is a complex measure on Γ . Under certain specific conditions upon Γ and μ , we find an upper bound for the number of zeros of f in a disk around the origin. The resulting theorem and the method of its proof are best understood by following the order of the paper.

In Section 2 we define the *angle* of the complex numbers a, b as

$$A(a, b) = \begin{cases} |\text{Arg}(b/a)| & \text{if } ab \neq 0 \\ \frac{1}{2}\pi & \text{if } ab = 0, a \neq b, \\ 0 & \text{if } a = b = 0 \end{cases}.$$

We define the *angular variation* of a sequence $\{x_1, x_2, \dots, x_n\}$ of complex numbers as

$$A(x_1, x_2, \dots, x_n) = A(x_1, x_2) + A(x_2, x_3) + \dots + A(x_{n-1}, x_n).$$

We prove that the angular variation of a sequence is never inferior to the angular variation of the sequence of its partial sums.

In Section 3 we apply the results of the previous section to derive, by induction, a theorem on the moduli of the last terms of sequences that are obtained by iterated partial summation of a given sequence.

This result is then applied, in Sections 4 and 5, to power sums

$$s(k) = \sum_{\ell=1}^N C_{\ell} \omega_{\ell}^k,$$

where C_1, \dots, C_N and $\omega_1, \dots, \omega_N$ are given complex numbers. Under the condition that

$$A(\omega_1^{-\omega_2}, \omega_2^{-\omega_3}, \dots, \omega_{N-1}^{-\omega_N}, \omega_N) < \frac{1}{2}\pi,$$

we derive constants n and $u(t,k)$ such that for $t = 1, 2, \dots$

$$|s(n+t)| \leq \sum_{k=0}^n u(t,k) |s(k)|.$$

By limit transition, we apply this last result in Section 6 to the functions (1.1). Let $P = \{x_1, x_2, \dots, x_m\}$ be a partition of Γ . We define Γ_k as the section of Γ between x_{k-1} and x_k and set

$$\rho = \rho(\Gamma) = \sup_P A(x_1 - x_2, x_2 - x_3, \dots, x_{m-1} - x_m),$$

$$V(\Gamma, \mu) = \sup_P A(\mu(\Gamma_1), \dots, \mu(\Gamma_m)),$$

$$\lambda = \lambda(\Gamma) = \text{length of } \Gamma = \sup_P (|x_1 - x_2| + \dots + |x_{m-1} - x_m|),$$

the suprema being taken over all partitions P of Γ . Under the conditions that $\rho(\Gamma) < \frac{1}{2}\pi$ and $V(\Gamma, \mu) < \infty$ we arrive at an estimate for the Taylor coefficients of f .

By standard techniques (see e.g. TIJDEMAN [1]) we then derive the following upper bound for the number $N_T(f)$ of zeros of f in a disk around the origin with radius T :

$$(1.2) \quad N_T(f) \leq 3n + 4T\lambda,$$

where n is the upper integer part of $\frac{V(\Gamma, \mu) - \frac{1}{2}\pi}{\frac{1}{2}\pi - \rho(\Gamma)}$.

2. ANGULAR VARIATION

A nonzero complex number ζ can be represented uniquely as

$$\zeta = re^{i\phi}; \quad r \neq 0, \quad -\pi < \phi \leq \pi.$$

For such a $\zeta = re^{i\phi}$, we define $\text{Arg} \zeta := \phi$. Let $a, b \in \mathcal{C}$. Then we define their angle $A(a, b)$ by

$$A(a,b) = \begin{cases} |\operatorname{Arg}(b/a)| & \text{if } ab \neq 0 \\ \frac{1}{2}\pi & \text{if } ab = 0, a \neq b, \\ 0 & \text{if } a = b = 0 \end{cases} .$$

Note that $A(a,b)$ is a kind of distance function, satisfying

$$A(a,b) = A(b,a); \quad A(a,a) = 0; \quad A(a,b) + A(b,c) \geq A(a,c).$$

We also have

$$A(ac,bc) = A(a,b) \text{ if } c \neq 0; \quad A(ac,bd) \leq A(a,b) + A(c,d).$$

LEMMA 2.1. For $a,b \in \mathbb{C}$

$$A(a,b) = A(a,a+b) + A(a+b,b).$$

PROOF. If a or $b = 0$, the proof follows immediately. The desired equation is symmetric in a and b , so we may assume without loss of generality that $\operatorname{Arg}(b/a) \geq 0$. For $x,y \in \mathbb{R}$ such that $x^2 + y^2 \neq 0$,

$$|\operatorname{Arg}(x+iy)| = \left| \int_0^{y/x} \frac{1}{1+t^2} dt \right|$$

decreases with increasing x if y is constant. Hence, since $\operatorname{Im}(b/a) \geq 0$,

$$0 \leq \operatorname{Arg}\left(1 + \frac{b}{a}\right) \leq \operatorname{Arg}\left(\frac{b}{a}\right),$$

so

$$\operatorname{Arg}\left(\frac{b}{a}\right) = \operatorname{Arg}\left(\frac{b+a}{a}\right) + \operatorname{Arg}\left(\frac{b}{b+a}\right),$$

proving the lemma. \square

LEMMA 2.2. Let $a,b \in \mathbb{C}$, such that $|a| > |b|$. Then

$$A(a,b-a) > \frac{1}{2}\pi.$$

PROOF. Since $|b| < |a|$,

$$\operatorname{Re}\left(\frac{b}{a} - 1\right) = \operatorname{Re}\left(\frac{b}{a}\right) - 1 \leq \left|\frac{b}{a}\right| - 1 < 0,$$

so

$$A(a, b-a) = \left|\operatorname{Arg}\left(\frac{b}{a} - 1\right)\right| > \frac{1}{2}\pi. \quad \square$$

For a sequence $S = \{x_1, x_2, \dots, x_n\}$ of complex numbers we define its angular variation

$$A(S) = A(x_1, x_2, \dots, x_n) = \sum_{k=1}^{n-1} A(x_k, x_{k+1}).$$

We have trivially

$$(2.1) \quad A(x_1 y_1, x_2 y_2, \dots, x_n y_n) \leq A(x_1, \dots, x_n) + A(y_1, \dots, y_n).$$

THEOREM 2.3. Let $S = \{x_1, x_2, \dots, x_n\}$ be a sequence of complex numbers and let T be the sequence of its partial sums $x_1, x_1+x_2, \dots, x_1+x_2+\dots+x_n$. Then

$$A(T) \leq A(S) - A(x_1+x_2+\dots+x_n, x_n).$$

PROOF. We apply induction on n . If $n = 1$, there is nothing to prove, so let $n > 1$. Let S' and T' be the sequences obtained from S and T respectively by deleting their last term. By the induction hypothesis

$$\begin{aligned} A(T) &= A(T') + A(x_1+\dots+x_{n-1}, x_1+\dots+x_n) \leq \\ &\leq A(S') - A(x_1+\dots+x_{n-1}, x_{n-1}) + A(x_1+\dots+x_{n-1}, x_1+\dots+x_n) = \\ &= A(S) - A(x_{n-1}, x_n) - A(x_1+\dots+x_{n-1}, x_{n-1}) + A(x_1+\dots+x_{n-1}, x_1+\dots+x_n) \leq \\ &\leq A(S) - A(x_1+\dots+x_{n-1}, x_n) + A(x_1+\dots+x_{n-1}, x_1+\dots+x_n) = \\ &= A(S) - A(x_1+x_2+\dots+x_n, x_n), \end{aligned}$$

by applying Lemma 2.1 with $a = x_1 + \dots + x_{n-1}$, $b = x_n$. This proves our theorem. \square

3. THE FUNDAMENTAL THEOREM

THEOREM 3.1. For $k = 1, 2, \dots$ and $j = 1, 2, \dots, N$, let $f_{k,j}$ and C_j be complex numbers and denote $M_k = \max_{j=1, \dots, N} f_{k,j}$. Define for $m = 1, 2, \dots, N$,

$$s_0(m) = \sum_{j=1}^m C_j,$$

$$s_k(m) = \sum_{j=1}^m f_{k,j} s_{k-1}(j), \quad k = 1, 2, \dots$$

For any $n \geq 0$ such that

$$(3.1) \quad A(C_1, \dots, C_N) \leq \frac{1}{2}(n+1)\pi - \sum_{k=1}^n A(f_{k,1}, \dots, f_{k,N}),$$

we have for $t = 1, 2, \dots$

$$(3.2) \quad |s_{n+t}(N)| \leq \sum_{k=0}^n \binom{n+t-k-1}{n-k} \binom{N+n+t-k-1}{n+t-k} M_{n+t} \dots M_{k+1} |s_k(N)|.$$

PROOF. We write

$$u_n(p, k, N) = \begin{cases} \binom{p-k-1}{n-k} \binom{N+p-k-1}{p-k} & \text{if } p > n, 0 \leq k \leq n, \\ 1 & \text{if } 0 \leq p = k \leq n, \\ 0 & \text{otherwise} \end{cases}$$

and we shall prove by induction on N and p that - under the Condition

(3.1) -

$$(3.3) \quad |s_p(N)| \leq \sum_{k=0}^n u_n(p, k, N) M_p \dots M_{k+1} |s_k(N)|.$$

If $N = 1$ or if $p \leq n$, then (3.3) is trivial. So let $N > 1$, $p > n$.

We have

$$|s_p(N)| = |s_p(N-1) + f_{p,N} s_{p-1}(N)| \leq |s_p(N-1)| + M_p |s_{p-1}(N)|.$$

By the induction hypothesis on p

$$(3.4) \quad |s_{p-1}(N)| \leq \sum_{k=0}^n u_n(p-1, k, n) M_{p-1} \dots M_{k+1} |s_k(N)|.$$

Now suppose that $|s_0(N)| \geq |s_0(N-1)|$. Since (3.1) also holds if we replace N by $N-1$, we have by the induction hypothesis on N

$$\begin{aligned} |s_p(N-1)| &\leq \sum_{k=0}^n u_n(p, k, N-1) M_p \dots M_{k+1} |s_k(N-1)| \leq \\ &\leq \sum_{k=1}^n u_n(p, k, N-1) M_p \dots M_{k+1} |s_k(N-1)| + \\ &\quad + u_n(p, 0, N-1) M_p \dots M_1 |s_0(N)|. \end{aligned}$$

If $|s_0(N)| < |s_0(N-1)|$, then by Lemma 2.2

$$A(s_0(N-1), s_0(N) - s_0(N-1)) = A(s_0(N-1), C_N) > \frac{1}{2}\pi.$$

Hence by Theorem 2.3

$$\begin{aligned} A(s_0(1), \dots, s_0(N-1)) &\leq A(C_1, \dots, C_{N-1}) - A(s_0(N-1), C_{N-1}) = \\ (3.5) \quad &= A(C_1, \dots, C_N) - A(C_{N-1}, C_N) - A(s_0(N-1), C_{N-1}) \\ &< A(C_1, \dots, C_N) - \frac{1}{2}\pi. \end{aligned}$$

For $m = 1, 2, \dots, N-1$, set

$$D_m = f_{1,m} s_0(m); \quad t_0(m) = \sum_{j=1}^m D_j; \quad t_k(m) = \sum_{j=1}^m f_{k-1,j} t_{k-1}(j).$$

Then by (2.1) and (3.5), applying (3.1),

$$\begin{aligned} A(D_1, \dots, D_{N-1}) &\leq A(C_1, \dots, C_N) - \frac{1}{2}\pi + A(f_{1,1}, \dots, f_{1,N-1}) \leq \\ &\leq \frac{1}{2}n\pi - \sum_{k=2}^n A(f_{k,1}, \dots, f_{k,N}). \end{aligned}$$

So the conditions of the theorem apply to the $t_k(m)$'s, with n replaced by $n-1$ and N by $N-1$. By the induction hypothesis on N we thus have

$$\begin{aligned} |s_p^{(N-1)}| &= |t_{p-1}^{(N-1)}| \leq \sum_{k=0}^{n-1} u_{n-1}(p-1, k, N-1) M_p \dots M_{k+2} |t_k^{(N-1)}| = \\ &= \sum_{k=1}^n u_n(p, k, N-1) M_p \dots M_{k+1} |s_k^{(N-1)}|. \end{aligned}$$

So either way we have

$$\begin{aligned} |s_p^{(N-1)}| &\leq \sum_{k=1}^n u_n(p, k, N-1) M_p \dots M_{k+1} |s_k^{(N-1)}| + \\ &\quad + u_n(p, 0, N-1) M_p \dots M_1 |s_0^{(N)}|. \end{aligned}$$

Since for $k \geq 1$

$$|s_k^{(N-1)}| \leq |s_k^{(N)}| + M_k |s_{k-1}^{(N)}|,$$

we find

$$|s_p^{(N-1)}| \leq \sum_{k=0}^n (u_n(p, k, N-1) + u_n(p, k+1, N-1)) M_p \dots M_{k+1} |s_k^{(N)}|.$$

Hence, by (3.4),

$$\begin{aligned} |s_p^{(N)}| &\leq \sum_{k=0}^n (u_n(p-1, k, N) + u_n(p, k, N-1) + u_n(p, k+1, N-1)) M_p \dots \\ &\quad \dots M_{k+1} |s_k^{(N)}|. \end{aligned}$$

An easy calculation shows that

$$u_n(p-1, k, N) + u_n(p, k, N-1) + u_n(p, k+1, N-1) \leq u_n(p, k, N). \quad \square$$

4. POWER SUMS

THEOREM 4.1. Let $\omega_1, \omega_2, \dots, \omega_N$ and C_1, \dots, C_N be given complex numbers. We define inductively

$$s_0(m) = \sum_{k=1}^m C_k$$

$$s_\ell(m) = \sum_{k=1}^m s_{\ell-1}(k) (\omega_k - \omega_{k+\ell}) \quad (\ell > 1),$$

where

$$\omega_{N+j} := 0 \text{ if } j > 0.$$

Then

$$s_\ell(m) = \sum_{k=1}^m C_k (\omega_k - \omega_{m+1}) \dots (\omega_k - \omega_{m+\ell}).$$

PROOF. We apply induction on ℓ . The theorem is trivial for $\ell = 0$. If $\ell > 0$, then by the induction hypothesis

$$s_\ell(m) = \sum_{k=1}^m (\omega_k - \omega_{k+\ell}) \sum_{\lambda=1}^k C_\lambda (\omega_\lambda - \omega_{k+1}) \dots (\omega_\lambda - \omega_{k+\ell-1}) =$$

$$= \sum_{\lambda=1}^m C_\lambda \sum_{k=\lambda}^m (\omega_\lambda - \omega_{k+1}) (\omega_\lambda - \omega_{k+2}) \dots (\omega_\lambda - \omega_{k+\ell-1}) (\omega_k - \omega_{k+\ell}).$$

It is thus sufficient to prove that

$$(4.1) \quad \sum_{k=\lambda}^m (\omega_\lambda - \omega_{k+1}) \dots (\omega_\lambda - \omega_{k+\ell-1}) (\omega_k - \omega_{k+\ell}) = (\omega_\lambda - \omega_{m+1}) \dots (\omega_\lambda - \omega_{m+\ell}).$$

This we prove by induction on m . If $m = \lambda$, then (4.1) follows immediately.

If $m > \lambda$, then by the induction hypothesis

$$\sum_{k=\lambda}^m (\omega_\lambda - \omega_{k+1}) \dots (\omega_\lambda - \omega_{k+\ell-1}) (\omega_k - \omega_{k+\ell}) =$$

$$= (\omega_\lambda - \omega_m) (\omega_\lambda - \omega_{m+1}) \dots (\omega_\lambda - \omega_{m+\ell-1}) + (\omega_\lambda - \omega_{m+1}) \dots (\omega_\lambda - \omega_{m+\ell-1}) (\omega_m - \omega_{m+\ell}) =$$

$$= (\omega_\lambda - \omega_{m+1}) \dots (\omega_\lambda - \omega_{m+\ell-1}) (\omega_\lambda - \omega_{m+\ell}),$$

proving (4.1) and the theorem. \square

COROLLARY 4.2. In the notation of Theorem 4.1, $s_\ell(N) = \sum_{k=1}^N C_k \omega_k^\ell$.

Corollary 4.2 shows that Theorem 3.1 can be applied to power sums, with $f_{\ell,k} = \omega_k^{-\omega_{k+\ell}}$. So we shall investigate the angular variation of the sequences $\{\omega_1^{-\omega_{\ell+1}}, \omega_2^{-\omega_{\ell+2}}, \dots, \omega_N^{-\omega_{\ell+N}}\}$.

LEMMA 4.3. *Let $a, b, c, d \in \mathcal{C}$ such that $A(a-b, b-c, c-d) \leq \pi$. Then*

$$(4.2) \quad A(a-c, b-c, b-d) \geq A(a-c, a-d, b-d).$$

PROOF. If $a = b$, then (4.2) is trivial. If $b = c$, then by Lemma 2.1,

$$A(a-c, a-d, b-d) = A(a-b, a-d, b-d) = A(a-b, b-d) \leq A(a-c, b-c, b-d)$$

by the triangle inequality, proving (4.2). If $a = c$, then $A(a-b, b-c, c-d) \leq \pi$ implies that $A(b-c, c-d) = A(b-d, c-d) = 0$, so (4.2) holds. If $a = d$, by Lemma 2.1

$$\pi \geq A(a-b, b-c, c-d) = A(a-b, a-c, b-c, c-a),$$

so $A(a-b, a-c) = 0$. Hence

$$A(a-c, b-c, b-a) \geq A(a-c, b-a) = \pi = A(a-c, 0, b-a),$$

if $b \neq a$ and $c \neq a$, proving (4.2). Since the lemma is not altered by the transformation $T: (a, b, c, d) \rightarrow (d, c, b, a)$, we may thus assume that a, b, c and d are all different.

If $A(a-b, b-c, c-d) = A(a-b, c-d)$, then by repeated application of Lemma 2.1

$$\begin{aligned} A(a-b, c-d) &= A(a-b, a-c, b-c, c-d) = A(a-b, a-c, c-d) = \\ &= A(a-b, a-c, a-d, c-d). \end{aligned}$$

Likewise, $A(a-b, c-d) = A(a-b, a-d, b-d, c-d)$, so

$$\begin{aligned} A(a-b, c-d) &= A(a-b, a-c, a-d, b-d, c-d) = A(a-b, b-c, c-d) = \\ &= A(a-b, a-c, b-c, b-d, c-d). \end{aligned}$$

Hence $A(a-c, a-d, b-d) = A(a-c, b-c, b-d)$, proving the lemma. So we may assume that $A(a-c, b-c, c-d) > A(a-b, c-d)$. Without loss of generality we may assume that


$$A(b-c, c-d) \geq A(a-b, b-c); \operatorname{Arg}(c-d) = 0; \operatorname{Arg}(b-c) \geq 0,$$

since we can apply T , multiply by a nonzero constant and take conjugates everywhere without altering the lemma. So, again applying Lemma 2.1,

$$0 \leq \operatorname{Arg}(a-b) \leq \operatorname{Arg}(a-c) \leq \operatorname{Arg}(b-c),$$

$$0 = \operatorname{Arg}(c-d) \leq \operatorname{Arg}(a-d) \leq \operatorname{Arg}(a-c) \leq \operatorname{Arg}(b-c).$$

If $A(a-c, a-d, b-d) = A(a-c, b-d)$, then (4.2) immediately follows, so we may assume that $\operatorname{Arg}(b-d) > \operatorname{Arg}(a-d)$. The points d , c , a and b then form a quadrangle as in the figure.



Since $A(b-d, c-d) + A(a-b, a-c) \leq A(a-b, b-c, c-d) \leq \pi$, the point a is exterior to or on the circle through b , c and d . Hence $A(a-c, a-d) \leq A(b-c, b-d)$. Similarly $A(a-d, b-d) \leq A(a-c, b-c)$. This proves the lemma. \square

THEOREM 4.4. Let $\omega_1, \omega_2, \dots, \omega_N$ be given complex numbers, and for $k = 1, 2, \dots$ let T_k be the sequence $T_k = \{\omega_1^{-\omega_{k+1}}, \omega_1^{-\omega_{k+2}}, \omega_2^{-\omega_{k+2}}, \omega_2^{-\omega_{k+3}}, \dots, \omega_{N-1}^{-\omega_{k+N}}, \omega_N^{-\omega_{k+N}}\}$, where $\omega_j := 0$ if $j > N$. If $A(T_1) \leq \pi$, then for $k = 1, 2, \dots$

$$A(T_{k+1}) \leq A(T_k) - A(\omega_1^{-\omega_{k+1}}, \omega_1^{-\omega_{k+2}}).$$

PROOF. Let S_k be the sequence obtained from T_k by omitting the first term $\omega_1^{-\omega_{k+1}}$ and adding a final term $\omega_N^{-\omega_{k+N+1}}$. Since $A(\omega_N^{-\omega_{k+N}}, \omega_N^{-\omega_{k+N+1}}) = 0$ if $k \geq 1$, we have

$$A(S_k) = A(T_k) - A(\omega_1^{-\omega_{k+1}}, \omega_1^{-\omega_{k+2}}).$$

Now T_{k+1} is obtained from S_k by changing each term $\omega_{\ell}^{-\omega_{k+\ell}}$ in S_k into $\omega_{\ell-1}^{-\omega_{k+\ell+1}}$ for $\ell = 2, 3, \dots, N$. Since for $\ell = 2, 3, \dots, N-k$, by repeated application of Lemma 2.1,

$$\begin{aligned} \pi &\geq A(T_1) \geq A(\omega_{\ell-1}^{-\omega_{\ell}}, \omega_{\ell}^{-\omega_{\ell+1}}, \omega_{\ell+1}^{-\omega_{\ell+2}}, \dots, \omega_{\ell+k}^{-\omega_{\ell+k+1}}) \geq \\ &\geq A(\omega_{\ell-1}^{-\omega_{\ell}}, \omega_{\ell}^{-\omega_{\ell+2}}, \omega_{\ell+2}^{-\omega_{\ell+3}}, \dots, \omega_{\ell+k}^{-\omega_{\ell+k+1}}) \geq \dots \geq \\ &\geq A(\omega_{\ell-1}^{-\omega_{\ell}}, \omega_{\ell}^{-\omega_{\ell+k}}, \omega_{\ell+k}^{-\omega_{\ell+k+1}}), \end{aligned}$$

we have for $\ell = 2, 3, \dots, N-k$ by Lemma 4.3,

$$\begin{aligned} &A(\omega_{\ell-1}^{-\omega_{k+\ell}}, \omega_{\ell}^{-\omega_{k+\ell}}, \omega_{\ell}^{-\omega_{k+\ell+1}}) \geq \\ &\geq A(\omega_{\ell-1}^{-\omega_{k+\ell}}, \omega_{\ell-1}^{-\omega_{k+\ell+1}}, \omega_{\ell}^{-\omega_{k+\ell+1}}). \end{aligned}$$

For $\ell > N-k$ the same inequality trivially holds. So

$$A(T_{k+1}) \leq A(S_k) = A(T_k) - A(\omega_1^{-\omega_{k+1}}, \omega_1^{-\omega_{k+2}}),$$

proving the theorem. \square

COROLLARY 4.5. *If $A(\omega_1^{-\omega_2}, \omega_2^{-\omega_3}, \dots, \omega_N^{-\omega_{N-1}}) \leq \pi$, then for $k = 1, 2, \dots$,*

$$A(\omega_1^{-\omega_{k+1}}, \omega_2^{-\omega_{k+2}}, \dots, \omega_N^{-\omega_{k+N}}) \leq A(\omega_1^{-\omega_2}, \omega_2^{-\omega_3}, \dots, \omega_N^{-\omega_{N+1}}).$$

PROOF. The sequence $S'_k = \{\omega_1^{-\omega_{k+1}}, \omega_2^{-\omega_{k+2}}, \dots, \omega_N^{-\omega_{k+N}}\}$ is a subsequence of T_k , so by Theorem 4.4, if $A(T_1) \leq \pi$, then

$$A(S'_k) \leq A(T_k) \leq A(T_1).$$

By Lemma 2.2

$$\begin{aligned} A(T_1) &= A(\omega_1^{-\omega_2}, \omega_1^{-\omega_3}, \omega_2^{-\omega_3}, \dots, \omega_{N-1}^{-\omega_{N+1}}, \omega_N^{-\omega_{N+1}}) = \\ &= A(\omega_1^{-\omega_2}, \omega_2^{-\omega_3}, \dots, \omega_N^{-\omega_{N+1}}) \leq \pi. \end{aligned}$$

This proves the corollary. \square

5. THE MAIN THEOREM

THEOREM 5.1. Let C_1, \dots, C_N and $\omega_1, \dots, \omega_N$ be given complex numbers, such that

$$A(\omega_1^{-\omega_2}, \omega_2^{-\omega_3}, \dots, \omega_{N-1}^{-\omega_N}, \omega_N) = \rho < \frac{1}{2}\pi.$$

Define

$$\lambda = |\omega_1^{-\omega_2}| + \dots + |\omega_{N-1}^{-\omega_N}| + |\omega_N|;$$

$$n = \max\left(0, \left\lceil \frac{A(C_1, \dots, C_N) - \frac{1}{2}\pi}{\frac{1}{2}\pi - \rho} \right\rceil\right),$$

where $\lceil x \rceil$ denotes the smallest integer m such that $m \geq x$. Then for $t = 1, 2, \dots$, the power sums $s(k) = \sum_{\ell=1}^N C_\ell \omega_\ell^k$ satisfy

$$|s(n+t)| \leq \sum_{k=0}^n \binom{n+t-k-1}{n-k} \binom{n+t}{k} \lambda^{n+t-k} |s(k)|.$$

PROOF. Let $\mu := \max_{k=1, \dots, N} |\omega_k^{-\omega_{k+1}}|$, where we define $\omega_{N+1} := 0$. Set $f_{\ell, k} = \omega_k^{-\omega_{k+\ell}}$. By Corollary 4.2, $s(k) = s_k(N)$ in the notation of Theorem 4.1. By Corollary 4.5

$$A(f_{\ell, 1}, \dots, f_{\ell, N}) \leq \rho,$$

thus

$$A(C_1, \dots, C_N) \leq \frac{1}{2}\pi + \sum_{\ell=1}^n \{\frac{1}{2}\pi - A(f_{\ell, 1}, \dots, f_{\ell, N})\},$$

so we can apply Theorem 3.1. Since for $\ell = 1, 2, \dots$

$$M_\ell = \max_{k=1, \dots, N} |\omega_k^{-\omega_{k+\ell}}| \leq \ell\mu,$$

we have by Theorem 3.1

$$(5.1) \quad |s(n+t)| \leq \sum_{k=0}^n (n+t) \dots (k+1) \binom{n+t-k-1}{n-k} \binom{N+n+t-k-1}{n+t-k} \mu^{n+t-k} |s(k)|.$$

This estimate can be improved by choosing equidistant points $\alpha_{i, 1}, \dots, \alpha_{i, t(i)}$ on the line segment between ω_i and ω_{i+1} in such a way that the total number M of points is very large and

$$\begin{aligned} & \max_{i=1, \dots, n} \left\{ \max_{\lambda=1, \dots, t(i)} (|\alpha_{i,1}^{-\omega_i}|, |\alpha_{i,\lambda+1}^{-\alpha_{i,\lambda}}|, |\omega_{i+1}^{-\alpha_{i,t(i)}}|) \right\} \leq \\ & \leq \frac{\lambda(1+\varepsilon)}{M} \end{aligned}$$

for a small $\varepsilon > 0$. We now form for $\delta > 0$ the power sums

$$s_\delta(k) = \sum_{i=1}^N c_i \left(\omega_i^k + \sum_{\lambda=1}^{t(i)} \delta \alpha_{i,\lambda}^k \right).$$

Since $A(\omega_i^{-\alpha_{i,1}}, \dots, \alpha_{i,t(i)}^{-\omega_{i+1}}) = A(\omega_i, \omega_{i+1})$ and $A(C_i, C_{i+1}) = A(C_i, \delta C_i, \dots, \delta C_i, C_{i+1})$, we have by (5.1)

$$|s_\delta(n+t)| \leq \sum_{k=0}^n \binom{n+t-k-1}{n-k} \cdot \frac{(n+t)!}{k!} \cdot \binom{M+n+t-k-1}{n+t-k} \left(\frac{\lambda(1+\varepsilon)}{M} \right)^{n+t-k} |s_\delta(k)|.$$

Now we let $M \rightarrow \infty$ and $\varepsilon \downarrow 0$, giving

$$|s_\delta(n+t)| \leq \sum_{k=0}^n \binom{n+t-k-1}{n-k} \binom{n+t}{k} \lambda^{n+t-k} |s_\delta(k)|.$$

By letting $\delta \downarrow 0$ the theorem is proved. \square

6. LAPLACE TRANSFORMS

Let Γ be a Jordan arc in \mathcal{C} of finite length running from a to b . Let μ be a complex measure on Γ , that is a combination $\mu_1 + i\mu_2 - \mu_3 - i\mu_4$ of four positive measures on Γ . We consider not identically vanishing functions of the form (1.1):

$$f(z) = \int_{\Gamma} e^{\zeta z} d\mu(\zeta).$$

In studying the zeros of f , we may assume without loss of generality that $b = 0$, since we can multiply f by e^{-bz} without altering its zeros. Now let

$$P = \{a = x_1, x_2, \dots, x_m = b\}$$

be a partition of Γ and, for $k = 1, 2, \dots, m-1$, let Γ_k be the segment of Γ

running from x_k to x_{k+1} . We set

$$M(P) = \max_{k=1, \dots, m-1} |x_k - x_{k-1}|.$$

We now construct the exponential polynomials

$$F_P(z) = \sum_{k=1}^{m-1} \exp(x_k z) \mu(\Gamma_k).$$

LEMMA 6.1. Let $D \subset \mathcal{T}$ be bounded and let P_1, P_2, \dots be a sequence of partitions such that $\lim_{k \rightarrow \infty} M(P_k) = 0$. Then

$$\lim_{k \rightarrow \infty} F_{P_k}(z) = f(z),$$

uniformly for $z \in D$.

PROOF. Observe that for a partition $P = \{x_1, x_2, \dots, x_m\}$

$$\begin{aligned} |F_P(z) - f(z)| &= \left| \sum_{k=1}^{m-1} \int_{\Gamma_k} (\exp(x_k z) - \exp(\zeta z)) d\mu(\zeta) \right| \leq \\ &\leq \sum_{j=1}^4 \mu_j(\Gamma) \cdot \max_{k=1, \dots, m-1} \max_{\zeta \in \Gamma_k} |\exp(x_k z) - \exp(\zeta z)|. \end{aligned}$$

For $z \in D$,

$$|\exp(x_k z) - \exp(\zeta z)| \rightarrow 0$$

uniformly if $\max |x_k - \zeta| \rightarrow 0$. \square

We consider curves Γ with the property that $b = 0$ and

$$\rho(\Gamma) := \sup_P \{A(x_1 - x_2, x_2 - x_3, \dots, x_{m-1} - x_m)\} < \frac{1}{2}\pi,$$

the supremum being taken over all partitions $P = \{a = x_1, x_2, \dots, x_m = 0\}$ of Γ . We define

$$V(\Gamma, \mu) = \sup_P A(\mu(\Gamma_1), \dots, \mu(\Gamma_m))$$

and set

$$(6.1) \quad n = \left\lceil \frac{V(\Gamma, \mu) - \frac{1}{2}\pi}{\frac{1}{2}\pi - \rho(\Gamma)} \right\rceil.$$

Write $\lambda = \lambda(\Gamma) = \sup_P (|x_1 - x_2| + \dots + |x_{m-1} - x_m|)$, taken over all partitions $P = \{x_1, \dots, x_m\}$ of Γ . This is the length of Γ . By Theorem 5.1, since $x_m = b = 0$ and since for $l = 0, 1, \dots$,

$$F_P^{(l)}(0) = \sum_{k=1}^{m-1} \mu(\Gamma_k) x_k^l,$$

we now have

$$(6.2) \quad |F_P^{(n+t)}(0)| \leq \sum_{\ell=0}^n \binom{n+t-\ell-1}{n-\ell} \binom{n+t}{\ell} \lambda^{n+t-\ell} |F_P^{(\ell)}(0)|.$$

Writing $f(z) = \sum_{k=0}^{\infty} \phi_k z^k$, and observing that, by Lemma 6.1, $\phi_k = f^{(k)}(0)/k!$ is approximated by $F_P^{(k)}(0)/k!$, we find from (6.2) for $t = 1, 2, \dots$,

$$(6.3) \quad |\phi_{n+t}| \leq \sum_{\ell=0}^n \frac{\lambda^{n+t-\ell}}{(n-\ell)!(t-1)!(n+t-\ell)} |\phi_{\ell}|.$$

Hence

$$\begin{aligned} \sum_{t=1}^{\infty} |\phi_{n+t}| R^{n+t} &\leq (R/r)^n \sum_{t=1}^{\infty} (R\lambda)^t / t! \sum_{\ell=0}^n \frac{(\lambda r)^{n-\ell}}{(n-\ell)!} |\phi_{\ell}| r^{\ell} \leq \\ &\leq (R/r)^n (e^{\lambda R} - 1) e^{\lambda r} \max_{|z| \leq r} |f(z)|. \end{aligned}$$

Moreover,

$$\begin{aligned} \sum_{\ell=0}^n |\phi_{\ell}| R^{\ell} &\leq (R/r)^n (1 + (r/R) + \dots + (r/R)^n) \max_{\ell=0, \dots, n} (|\phi_{\ell}| r^{\ell}) \leq \\ &\leq (R/r)^n \cdot (R/(R-r)) \cdot \max_{|z| \leq r} |f(z)|. \end{aligned}$$

Hence,

$$(6.4) \quad \max_{|z| \leq R} |f(z)| < (R/r)^n e^{\lambda R} \max\{e^{\lambda R}, R/(R-r)\} \max_{z \leq r} |f(z)|. \quad \square$$

The following lemma is more or less standard.

LEMMA 6.2. *Let r , R and T be real numbers satisfying $R > r > 0$; $R \geq T > 0$. If $f \neq 0$ is analytic in a neighbourhood of the disk $|z| \leq R$, then the number $N_T(f)$ of zeros of f in the disk $\{z \in \mathcal{D}; |z| \leq T\}$, counted according to multiplicity, satisfies*

$$N_T(f) \log\left(\frac{R^2 - rT}{R(T+r)}\right) \leq \log\left\{ \frac{\max_{|z| \leq R} |f(z)|}{\max_{|z| \leq r} |f(z)|} \right\}.$$

PROOF. See WALDSCHMIDT [4], page 6.4, Lemma 6.2.1. \square

For other versions of Lemma 6.2, compare TIJDEMAN [1], Lemma 1 and VOORHOEVE [2], Lemma 4.4. We can now prove our final theorem on the zeros of the Laplace transform (1.1):

THEOREM 6.3. *For functions $f \neq 0$ defined by (1.1) we have*

$$(6.5) \quad N_T(f) \leq 3n + 4T\lambda,$$

where n is defined by (6.1) and λ is the length of Γ .

PROOF. Apply Lemma 6.2 with $R = 4T$ and $r = T/10$. Inserting (6.4), we obtain

$$N_T(f) \log(159/44) \leq n \log 40 + 4\lambda T + \max\{\lambda T/10, \log(40/39)\}.$$

This yields (6.5) if $n > 0$ or if $\lambda T > 1/5$. If $n = 0$ and $\lambda T \leq 1/5$, observe that $N_T(f) \leq 4/5 + \log(40/39) < 1$, so $N_T(f) = 0$, proving (6.5) completely. \square

We can use Theorem 6.3 to estimate the number of zeros of f in a disk with centre $c \neq 0$, either directly by estimating $N_{T+|c|}(f)$, or indirectly by observing that

$$\tilde{f}(z) = f(z+c) = \int_{\Gamma} e^{\zeta z} d\tilde{\mu}(\zeta),$$

where $\tilde{\mu}$ is defined by $\tilde{\mu}(\gamma) = \int_{\gamma} e^{c\zeta} d\mu(\zeta)$ for a measurable $\gamma \subset \Gamma$. So

$$(6.6) \quad N_T(\tilde{f}) \leq 3\tilde{n} + 4\lambda T,$$

where

$$\tilde{n} = \left\lceil \frac{V(\Gamma, \tilde{\mu}) - \frac{1}{2}\pi}{\frac{1}{2}\pi - \rho} \right\rceil.$$

One can estimate $V(\Gamma, \tilde{\mu})$ by

$$V(\Gamma, \tilde{\mu}) \leq V(\Gamma, \mu) + |c|\lambda,$$

so $\tilde{n} \leq n + \lceil |c|\lambda/(\frac{1}{2}\pi - \rho) \rceil$. This rough estimate for \tilde{n} , inserted in (6.4), does not give an essentially better result than the direct approach. In some cases, though, one can give better estimates for \tilde{n} .

7. REMARKS

1. If Γ is a straight line segment (i.e. $\rho(\Gamma) = 0$), then a more direct approach in estimating the zeros of f is possible; see VOORHOEVE [3]. Also if $V(\Gamma, \mu) < \pi$, it is easy to show that $f(0) \neq 0$ and it is not hard to give estimates of the form $|f^{(t)}(0)| \leq u(t)|f(0)|$ for $t = 1, 2, \dots$.
2. It is clear that the dependency on T and λ in the estimate (6.5) is right - that is, up to a constant factor. For instance we can take

$$f(z) = 1 + e^{\lambda z} = \int_{\Gamma} e^{\zeta z} d\mu(\zeta),$$

where $a = \lambda$, $b = 0$ and $\mu = 0$ except for two point masses at a and b . Then $N_T(f) \sim \lambda T/(2\pi)$. However, it may be possible to improve the constant n and the condition that $\rho(\Gamma) < \frac{1}{2}\pi$.

3. By this method, one cannot derive an analogue of Theorem 6.3 for all curves Γ with $\rho(\Gamma) > \pi$, because such curves can be closed. For closed curves Γ , we can derive counterexamples to such a theorem, using Cauchy's theorem

$$f(z) = \int_{\Gamma} e^{\zeta z} d\mu(\zeta) = \int_{\Gamma} e^{\zeta z} d(\mu(\zeta) + C\mu_0(\zeta)),$$

where μ_0 is the Lebesgue measure. By letting $C \rightarrow \infty$, the $V(\Gamma, \mu + C\mu_0)$ approximate $V(\Gamma, \mu_0)$. So this would imply that there is an absolute

upper bound for $N_T(f)$, independent of μ . By a suitable choice of μ , however, $f(z)$ can have zeros of arbitrary multiplicity. This is a contradiction.

REFERENCES

- [1] TIJDEMAN, R., *On the number of zeros of general exponential polynomials*, Nederl. Akad. Wet. Proc. Ser. A 74 = Indag. Math. 33 (1971) 1-7.
- [2] VOORHOEVE, M., *Zeros of exponential polynomials*, Ph.D. Thesis, Leiden State University 1977.
- [3] VOORHOEVE, M., *A generalization of Descartes' rule*, to appear in J. London Math. Soc.
- [4] WALDSCHMIDT, M., *Nombres Transcendants*, Lecture Notes in Mathematics 402, Springer-Verlag, Berlin, 1974.

