

**stichting  
mathematisch  
centrum**



---

AFDELING ZUIVERE WISKUNDE  
(DEPARTMENT OF PURE MATHEMATICS)

ZW 127/79

SEPTEMBER

H.A. WILBRINK

FINITE MINKOWSKI PLANES

Preprint

---

**2e boerhaavestraat 49 amsterdam**

BIBLIOTHEEK MATHEMATISCH CENTRUM  
AMSTERDAM

*Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.*

*The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O).*

Finite Minkowski planes<sup>\*)</sup>

by

H.A. Wilbrink

ABSTRACT

In this paper we give second characterizations of a certain class of finite Minkowski planes.

KEY WORDS & PHRASES: *Minkowski plane, affine plane, nearaffine plane.*

---

<sup>\*)</sup> This report will be submitted for publication elsewhere



## 1. INTRODUCTION

It is well known, see e.g. [5], that with each point of a Minkowski plane there is associated an affine plane, its so-called derived plane. It is the purpose of this paper to show that, under certain additional hypotheses, with each point of a Minkowski plane there is also associated a nearaffine plane, its *residual* plane. In addition we show that the "known" Minkowski plane are characterized by the fact that these nearaffine planes are nearaffine translation planes (see [9]). Using this result a configurational condition is obtained in a completely natural way which characterizes the known Minkowski planes.

## 2. BASIC CONCEPTS

Let  $M$  be a set of *points* and  $L^+$ ,  $L^-$ ,  $C$  three collections of subsets of  $M$ . The elements of  $L := L^+ \cup L^-$  are called *lines* or *generators*, the elements of  $C$  are called *circles*. We say that  $M = (M, L^+, L^-, C)$  is a *Minkowski plane* if the following axioms are satisfied (cf. [5]):

- (M1):  $L^+$  and  $L^-$  are partitions of  $M$ .
- (M2):  $|\ell^+ \cap \ell^-| = 1$  for all  $\ell^+ \in L^+$ ,  $\ell^- \in L^-$ .
- (M3): Given any three points no two on a line, there is a unique circle passing through these three points.
- (M4):  $|\ell \cap c| = 1$  for all  $\ell \in L$ ,  $c \in C$ .
- (M5): There exist three points no two of which are on one line.
- (M6): Given a circle  $c$ , a point  $P \in c$  and a point  $Q \notin c$ ,  $P$  and  $Q$  not on one line, there is a unique circle  $d$  such that  $P, Q \in d$  and  $c \cap d = \{P\}$ .

Two points  $P$  and  $Q$  are called *plus-parallel* (notation  $P \parallel_+ Q$ ) if  $P$  and  $Q$  are on a line of  $L^+$ , *minus-parallel* ( $P \parallel_- Q$ ) if  $P$  and  $Q$  are on a line of  $L^-$ . *Parallel* ( $P \parallel Q$ ) means either  $P \parallel_+ Q$  or  $P \parallel_- Q$ . For  $P \in M$ ,  $\epsilon = +, -$  we denote by  $[P]_\epsilon$  the unique line in  $L^\epsilon$  incident with  $P$ . If  $P, Q$  and  $R$  are (distinct) nonparallel points, then we denote by  $(P, Q, R)$  the unique circle containing  $P, Q$  and  $R$ . Two circles  $c$  and  $d$  *touch* in a point  $P$  if  $c \cap d = \{P\}$ .

Fix a point  $Z$  and put

$$M_Z := M \setminus ([Z]_+ \cup [Z]_-),$$

$$L_Z := \{c^* \mid c \in C, z \in c\} \cup \{\ell^* \mid \ell \in L \setminus \{[Z]_+, [Z]_-\}\},$$

where the  $*$  indicates that we have removed the point that the circle or line has in common with  $[Z]_+ \cup [Z]_-$ . Then  $M_Z := (M_Z, L_Z)$  is an affine plane with pointset  $M_Z$  and lineset  $L_Z$  (see e.g. [5]). We call  $M_Z$  the *derived plane* with respect to the point  $Z$ . We shall only consider finite Minkowski planes, i.e. Minkowski planes with a finite number of points. For finite Minkowski planes (M6) is a consequence of the other axioms (see [5]). It is easily seen that  $|L^+| = |L^-| = |\ell| = |c| =: n+1$  for all  $\ell \in L, c \in C$ . The integer  $n$  is called the *order* of the Minkowski plane. Notice that  $n$  is also the order of the derived planes  $M_Z$ .

Following BENZ [1] we sketch the close relationship between (finite) Minkowski planes and sharply 3-transitive sets of permutations. Let  $\Omega$  be a finite set,  $|\Omega| = n+1 \geq 3$ , and  $G$  a subset of  $S^\Omega$ , the symmetric group on  $\Omega$ , acting sharply triply transitively on  $\Omega$ .

Define

$$M := \Omega \times \Omega,$$

$$L^+ := \{(\alpha, \beta) \mid \alpha \in \Omega \mid \beta \in \Omega\},$$

$$L^- := \{(\alpha, \beta) \mid \beta \in \Omega \mid \alpha \in \Omega\},$$

$$C := \{(\alpha, \alpha^g) \mid \alpha \in \Omega \mid g \in G\}.$$

Then  $M := (\Omega, G) := (M, L^+, L^-, C)$  is a Minkowski plane of order  $n$ . Conversely, every Minkowski plane can be obtained in this way.

Two Minkowski planes  $M = (\Omega, G) = (M, L^+, L^-, C)$  and  $M' = (\Omega', G') = (M', L'^+, L'^-, C')$  are said to be *isomorphic* if there is a bijection  $s: M \rightarrow M'$  such that

$$L^s = L' \quad \text{and} \quad C^s = C'.$$

since  $s$  maps the disjoint lines of  $L^+$  onto disjoint lines there are only two possibilities, either  $(L^\varepsilon)^s = L^\varepsilon$  or  $(L^\varepsilon)^s = L^{-\varepsilon}$ ,  $\varepsilon = +, -$ . In the first case  $s$  is called a *positive isomorphism* in the second case a *negative*

*isomorphism*. If  $s$  is a positive isomorphism then there exist bijections  $a, b: \Omega \rightarrow \Omega'$  such that  $(\alpha, \beta)^s = (\alpha^a, \beta^b)$  for all  $\alpha, \beta \in \Omega$ , and  $G' = a^{-1}Gb$ . If  $s$  is a negative isomorphism then there exist bijections  $a, b: \Omega \rightarrow \Omega'$  such that  $(\alpha, \beta)^s = (\beta^b, \alpha^a)$ , and  $G' = b^{-1}G^{-1}a$ . It follows that we may assume w.l.o.g that  $\text{id} \in G$ .

A (positive, negative) automorphism of a Minkowski plane  $M$  is a (positive, negative) isomorphism of  $M$  onto itself. The automorphism group  $\text{Aut}(\Omega, G) \leq S^{\Omega \times \Omega}$  of the Minkowski plane  $(\Omega, G)$  is given by

$$\text{Aut}(\Omega, G) = \{(a, b) \mid a^{-1}Gb = G\} \cup \{(a, b) \mid a^{-1}Gb = G^{-1}\} \tau$$

where  $\tau$  is the permutation which sends  $(\alpha, \beta)$  to  $(\beta, \alpha)$ .

### 3. THE RESIDUAL PLANE

Let  $M = (M, L^+, L^-, C)$  be a Minkowski plane. Fix a point  $Z \in M$  and define  $M_Z = M \setminus ([Z]_+ \cup [Z]_-)$ . We have already remarked that the lines  $\neq [Z]_+, [Z]_-$  together with the circles which are incident with  $Z$  are the lines of an affine plane with pointset  $M_Z$ . We shall show that the lines  $\neq [Z]_+, [Z]_-$  together with the circles not incident with  $Z$  are the lines of a nearaffine plane with the same pointset if suitable conditions are assumed to hold in  $M$ .

For each point  $P \in M_Z$  we let the points  $P^+$  and  $P^-$  be defined by  $P^+ := [Z]_+ \cap [P]_-$ ,  $P^- := [Z]_- \cap [P]_+$ . The restriction of a line  $\ell$  or circle  $c$  to  $M_Z$  is denoted by  $\ell^* := \ell \cap M_Z$  resp.  $c^* := c \cap M_Z$ . For any two distinct points  $P, Q \in M_Z$  we define

$$P \sqcup Q := \begin{cases} \ell^* & \text{iff } P, Q \in \ell \in L, \\ \{P\} \cup (P^+, P^-, Q)^* & \text{iff } P \text{ and } Q \text{ are nonparallel.} \end{cases}$$

Since two circles can have at most two points in common it follows that  $P \sqcup Q = Q \sqcup P$  if and only if  $P \sqcup Q = \ell^*$  for some  $\ell \in L$ , provided the order  $n$  of  $M$  is at least 5. The verification of the axioms (L1), (L2) and L(3) (see [9]) is now straightforward. In order to define parallelism we have to require that the following condition holds in  $M$  for every point  $Z$ .

(A): Let  $P_1, Q_1, P_2, Q_2 \in M_Z$  and suppose that  $P_1$  and  $Q_1, P_2$  and  $Q_2, P_1$  and  $P_2$  are nonparallel. If there exists a circle  $c$  touching  $(P_1^+, P_1^-, Q_1)$  in  $P_1^-$  and touching  $(P_2^+, P_2^-, Q_2)$  in  $P_2^+$ , then there also exists a circle  $d$  touching  $(P_1^+, P_1^-, Q_1)$  in  $P_1^+$  and touching  $(P_2^+, P_2^-, Q_2)$  in  $P_2^-$  (see figure 1).




Fig. 1.

In the definition of  $P_1 \sqcup Q_1 \parallel P_2 \sqcup Q_2$  we have to distinguish several cases.

Case 1:  $P_1$  and  $Q_1$  parallel, say  $P_1 \sqcup Q_1 = \ell_1^*$  for some  $\ell_1 \in L^\mathcal{E}$ .

$$P_1 \sqcup Q_1 \parallel P_2 \sqcup Q_2 : \iff P_2 \sqcup Q_2 = \ell_2^* \quad \text{for some } \ell_2 \in L^\mathcal{E}.$$

Case 2:  $P_1$  and  $Q_1$  nonparallel,  $P_1, P_2$  parallel, say  $P_1, P_2 \in \ell \in L^\mathcal{E}$ . From [9], proposition 3.1, it is clear that we have to define

$$P_1 \sqsubset Q_1 \parallel P_2 \sqsubset Q_2 : \Leftrightarrow P_1 \sqsubset Q_1 = P_2 \sqsubset Q_2 \quad \text{or} \quad (P_1 \sqsubset Q_1) \cap (P_2 \sqsubset Q_2) = \emptyset.$$

Case 3:  $P_1$  and  $Q_1$  nonparallel and  $P_1, P_2$  nonparallel. Put  $P_3 = [P_1]_+ \cap [P_2]_-$  and  $P_4 := [P_1]_- \cap [P_2]_+$  (see fig. 1).  $P_1 \sqsubset Q_1 \parallel P_2 \sqsubset Q_2 : \Leftrightarrow$  There exists  $P_3 \sqsubset Q_3$  such that

$$(P_3 \sqsubset Q_3) \cap (P_1 \cap Q_1) = \emptyset = (P_3 \cap Q_3) \cap (P_2 \sqsubset Q_2).$$

Notice that condition (A) is equivalent to:  $P_1 \sqsubset Q_1 \parallel P_2 \sqsubset Q_2$  implies  $P_2 \sqsubset Q_2 \parallel P_1 \sqsubset Q_1$ , i.e. parallelism is a symmetric relation. We prove that parallelism is a transitive relation. Suppose  $P_1 \sqsubset Q_1 \parallel P_2 \sqsubset Q_2$  and  $P_2 \sqsubset Q_2 \parallel P_3 \sqsubset Q_3$  (with distinct  $P_1, P_2, P_3$ ). We prove that  $P_1 \sqsubset Q_1 \parallel P_3 \sqsubset Q_3$ .

Case a):  $P_1 \parallel Q_1$ . Trivial

Case b):  $P_1 \not\parallel Q_1, P_1, P_2, P_3 \in \ell$  for some  $\ell \in L$ . The transitivity follows at once from the following observation. If  $c, d, e, \in C$  and  $c$  and  $d$  touch in a point  $P$ ,  $d$  and  $e$  touch in the same point  $P$ , then  $c$  and  $e$  touch in  $P$ . To show this suppose  $Q \in c \cap e, Q \neq P$ , then there are two circles through  $Q$ , namely  $c$  and  $e$ , touching  $d$  in  $P$ . This contradicts (M6).

Case c):  $P_1 \not\parallel Q_1, P_1 \in [P_2]_\varepsilon, P_3 \in [P_2]_{-\varepsilon}$  for some  $\varepsilon = +, -$ . By definition  $P_1 \sqsubset Q_1 \parallel P_3 \sqsubset Q_3$ .

Case d):  $P_1 \not\parallel Q_1, P_1 \parallel_\varepsilon P_2$  for some  $\varepsilon = +, -$ ,  $P_3 \not\parallel P_1, P_3 \not\parallel P_2$ . Put  $P_4 := [P_2]_\varepsilon \cap [P_3]_{-\varepsilon}$ . Since  $P_2 \sqsubset Q_2 \parallel P_3 \sqsubset Q_3$  there exists  $Q_4$  such that  $P_2 \sqsubset Q_2 \parallel P_4 \sqsubset Q_4 \parallel P_3 \sqsubset Q_3$ . Apply case b) to find  $P_1 \sqsubset Q_1 \parallel P_4 \sqsubset Q_4$  and case c) to find  $P_1 \sqsubset Q_1 \parallel P_3 \sqsubset Q_3$ .

Case e):  $P_1 \not\parallel Q_1, P_1 \parallel_\varepsilon P_3$  for some  $\varepsilon = +, -$ ,  $P_2 \not\parallel P_1, P_2 \not\parallel P_3$ . Put  $P_4 := [P_1]_\varepsilon \cap [P_2]_{-\varepsilon}$ . There exists  $Q_4$  such that  $P_1 \sqsubset Q_1 \parallel P_4 \sqsubset Q_4$  and  $P_4 \sqsubset Q_4 \parallel P_3 \sqsubset Q_3$ . Apply case b).

Case f):  $P_1 \not\parallel Q_1, P_1, P_2, P_3$  mutually nonparallel. Put  $P_4 := [P_1]_+ \cap [P_2]_-$ . There exists  $Q_4$  and that  $P_1 \sqsubset Q_1 \parallel P_4 \sqsubset Q_4$  and  $P_4 \sqsubset Q_4 \parallel P_2 \sqsubset Q_2$ . Apply case d) to find  $P_4 \sqsubset Q_4 \parallel P_3 \sqsubset Q_3$  and so  $P_1 \sqsubset Q_1 \parallel P_3 \sqsubset Q_3$ .

Let  $L^Z$  be the set of all  $P \sqsubset Q, P, Q \in M_Z, P \neq Q$ . It is not hard to show that  $M^Z := (M_Z, L^Z, \sqsubset, \parallel)$  satisfies all the axioms of a nearaffine plane

except possibly (P2) or (P2'). For (P2) to hold we have to require:

(B): Let  $P_1, Q_1, P_2, Q_2$  be points as in (A). If  $P_1 \in (P_2^+, P_2^-, Q_2)$  and  $P_2 \in (P_1^+, P_1^-, Q_1)$ . Then circles  $c$  and  $d$  as described in (A) exist.

If we content ourself with the weaker (P2') we have to require:

(C): Let  $\varepsilon$  be + or -,  $A$  and  $B$  two distinct points on  $[Z]_\varepsilon$ ,  $A \neq Z \neq B$  and  $c_1$  and  $c_2$  two circles touching in  $A$ . Put (see figure 2)

$$\begin{aligned} C_i &:= [Z]_{-\varepsilon} \cap c_i, & i = 1, 2, \\ P_i &:= [A]_{-\varepsilon} \cap [C_i]_\varepsilon, & i = 1, 2, \\ Q_i &:= [B]_\varepsilon \cap c_i, & i = 1, 2, \\ D_i &:= [Q_i]_\varepsilon \cap [Z]_{-\varepsilon}, & i = 1, 2, \\ d_i &:= (P_i, D_i, B), & i = 1, 2. \end{aligned}$$

Then  $d_1$  and  $d_2$  touch in  $B$ .




Fig. 2.

If  $M$  is a Minkowski plane satisfying the conditions (A) and (B) or (A) and (C) and  $Z$  a point of  $M$ , then the nearaffine plane  $M^Z$  is called the *residual plane* with respect to  $Z$ .

For the remainder of this section let  $M = (M, L^+, L^-, C)$  be a Minkowski plane satisfying the conditions (A) and (C). Since  $\perp$  and  $\parallel$  are defined strictly in terms of the incidence in  $M$  it follows at once that an automorphism of  $M$  fixing a point  $Z$ , induces an automorphism of  $M^Z$ , i.e.  $\text{Aut}(M)_Z \simeq \text{Aut}(M^Z)$ . In fact,  $\text{Aut}(M)_Z \simeq \text{Aut}(M^Z)$  as we shall see in a moment. The crucial observation is the following lemma.

3.1. LEMMA. *Let  $Z$  be a point of  $M$ . For any two nonparallel points  $A$  and  $B$  of  $M_Z$  let  $[A, B]$  be the set of points consisting of  $A, B, Z$  and the points  $C \in M_Z$ , nonparallel to  $A$  and  $B$ , for which there is no set  $P \perp Q \setminus \{P\}$  containing  $A, B$  and  $C$ . Then*

$$[A, B] = (A, B, Z).$$

PROOF. Clearly both  $[A, B]$  and  $(A, B, Z)$  contain  $A, B$  and  $Z$ . Let  $C \in (A, B, Z)$ ,  $C \neq A, B, Z$  then  $(A, A, C) = (A, B, Z)$ . Suppose for some  $P, Q \in M_Z$  we have  $A, B, C \in P \perp Q \setminus \{P\}$ . Then  $A, B, C \in (P^+, P^-, Q) \setminus \{P^+, P^-\}$ , so  $(A, B, C) = (P^+, P^-, C)$  a circle not passing through  $Z$ , a contradiction. Conversely, let  $C \in [A, B]$ ,  $C \neq A, B, Z$  and suppose  $C \in (A, B, Z)$ . Then  $Z \notin (A, B, C)$  and so  $(A, B, C)$  intersects  $[Z]_+$  and  $[Z]_-$  in points  $P^+$  and  $P^-$  respectively, different from  $Z$ . So, with  $P$  defined by  $P = [P^+]_- \cap [P^-]_+$ ,  $A, B, C$  are on  $P \perp Q \setminus \{P\}$ , a contradiction.  $\square$

The lemma just proved shows that the residual plane  $M^Z$  completely determines the Minkowski plane  $M$ . The lines of  $M$  can be recovered from the straight lines of  $M^Z$ , the circles not containing  $Z$  from the proper lines of  $M^Z$ , and the circles containing  $Z$  from the sets  $[A, B]$ . This proves the following theorem.

3.2. THEOREM. *Let  $Y$  and  $Z$  be the points of  $M$ . Then*

- a)  $M^Y \simeq M^Z$  iff there exists  $\phi \in \text{Aut}(M)$  such that  $Y^\phi = Z$ .
- b) Any automorphism of  $M^Z$  can be extended to an automorphism of  $M$  fixing  $Z$ .
- c)  $\text{Aut}(M)_Z \simeq \text{Aut}(M^Z)$ .

It is not hard to show that for any point  $Z$  of  $M$  the residual plane  $M^Z$  satisfies the Veblen-condition (V'). In fact we can prove somewhat more.

**3.3. THEOREM.** *Let  $Z \in M$ ,  $\ell \in L$ ,  $\ell \neq [Z]_+, [Z]_-$  and let  $Y$  be defined by  $Y = \ell \cap ([Z]_+ \cup [Z]_-)$ . Then*

$$\begin{array}{c} M^Z \\ \ell^* \end{array} \simeq M_Y,$$

where  $\ell^*$  is the straight line  $\ell \setminus \{Y\}$  of  $M^Z$  (notation as in [9]).

**PROOF.** Define an isomorphism  $\phi: M_Z \rightarrow M_Y$  of  $M_{\ell^*}^Z$  onto  $M_Y$  as follows. For  $P \in M_Z$ ,  $P \notin \ell^*$  we define  $P^\phi := P$ , and for  $P \in M_Z$ ,  $P \in \ell^*$ ,  $P^\phi := [P]_{-\varepsilon} \cap [Z]_\varepsilon$ , where  $\varepsilon$  is determined by  $\ell \in L^\varepsilon$ .  $\square$

As a direct consequence of this theorem we have the following result.

**3.4. THEOREM.** *If the derived plane  $M_Z$  is a translation plane for every  $Z \in M$ , then the residual plane  $M^Z$  is a nearaffine translation plane for every  $Z \in M$ .*

**PROOF.** Apply 3.3 and 5.2 of [9].  $\square$

As a converse to this theorem we mention the following theorem.

**3.5. THEOREM.** *Let  $Z$  be a point of  $M$ . If  $M^Z$  is a nearaffine translation plane, then  $M_Z$  is a translation plane and  $M^Z$  and  $M_Z$  have the same translation group.*

**PROOF.** By 3.2 every automorphism of  $M^Z$  is also an automorphism of  $M_Z$ , and it is not hard to show that a straight translation of  $M^Z$  with a direction corresponding to  $L^\varepsilon$  is also a translation of  $M$ . Let  $T_+$  and  $T_-$  be the translation groups of  $M^Z$  with directions  $L^+$  and  $L^-$  respectively. Since  $T_+$  and  $T_-$  are also translation groups of  $M_Z$  it follows that  $T_+$  and  $T_-$  are elementary abelian. Hence, by 4.12 of [9], the set  $T$  of all translation of  $M^Z$  is a group and  $T = T_+ T_-$  = the full translation group of  $M_Z$ .  $\square$

## 4. CHARACTERIZATIONS OF THE KNOWN FINITE MODELS

Using the correspondence with sharply triply transitive sets of permutations all known (finite) Minkowski planes can be described as follows. Let  $p$  be a prime,  $h$  a positive integer,  $q := p^h$  and  $\phi$  an automorphism of  $GF(q)$ . Let  $G(\phi)$  be the set of permutations acting on the projective line  $\Omega := PG(1, q) = GF(q) \cup \{\infty\}$  given by

$$x \rightarrow \frac{ax+b}{cx+d}, \quad a, b, c, d \in GF(q), \quad ad-bc = (\text{nonzero}) \text{ square in } GF(q),$$

$$x \rightarrow \frac{ax^\phi+b}{cx^\phi+d}, \quad a, b, c, d \in GF(q), \quad ad-bc = \text{nonsquare in } GF(q),$$

i.e.  $G(\phi) = G_1 \cup \phi G_2$ , where  $G_1 := PSL(2, q)$  and  $G_2 := PG(2, q) \setminus PSL(2, q)$ . Then  $G(\phi)$  is sharply triply transitive on  $\Omega$  (cf. [7], [8], [10]). The residual planes of  $(\Omega, G(\phi))$  are easily seen to be the nearaffine translation planes described in [9], section 8. We shall show that a Minkowski plane whose residual planes are nearaffine translation planes, is isomorphic to an  $(\Omega, G(\phi))$ .

Let  $c$  be a circle of a Minkowski plane  $M$  of order  $n$  and  $Z$  a point of  $M$ ,  $Z \notin c$ . If  $M_Z$  is augmented to a projective plane, then the points of  $c^* = c \setminus ([Z]_+ \cup [Z]_-)$  together with the two ideal points corresponding to  $L^+$  and  $L^-$  constitute an oval in this projective plane. If  $n$  is even, there exists a point (the *nucleus* of the oval) in the projective plane such that the  $n+1$  lines incident with this point are the  $n+1$  tangents of the oval. If  $n$  is odd, each point of the projective plane is incident with 0 or 2 tangents (see [3]). From this observation we deduce the following lemma.

**4.1. LEMMA.** *Let  $M$  be a Minkowski plane of order  $n$ . If  $n$  is even, there cannot exist 3 distinct circles  $c_1, c_2, d$  such that  $c_1$  and  $c_2$  touch in a point  $Z$  and  $c_i$  touches  $d$  in  $P_i \neq Z$ ,  $i = 1, 2$ . In any case there cannot exist 4 distinct circles  $c_1, c_2, c_3$  and  $d$  such that  $c_1, c_2, c_3$  touch in a point  $Z$  and such that  $c_i$  touches  $d$  in a point  $P_i \neq Z$ ,  $i = 1, 2, 3$ .*

**PROOF.** Case  $n$  is even. Suppose circles  $c_1, c_2$  and  $d$  as described exist. The lines  $[[Z]_+ \cap d]_-$  and  $[[Z]_- \cap d]_+$  are tangents to the oval corresponding with

$d$  in the projective plane associated with  $M_Z$ . They intersect in a point of  $M_Z$ . Also  $c_1$  and  $c_2$  are tangents to the oval. They intersect in an ideal point of the projective plane, a contradiction.

Case  $n$  is odd. Now  $c_1$ ,  $c_2$  and  $c_3$  correspond to tangents of the oval  $d$  in the projective plane associated with  $M_Z$ . They intersect in one (ideal) point, a contradiction.  $\square$

**4.2. THEOREM.** *Let  $M = (\Omega, G) = (M, L^+, L^-, C)$  be a Minkowski plane of order  $n \geq 5$ . Suppose conditions (A) and (C) hold in  $M$  and that  $M^Z$  is a nearaffine translation plane for every point  $Z$ . Then  $M \simeq (\Omega, G(\phi))$ .*

**PROOF.** Fix  $\alpha_1 \in \Omega$ . For each point  $(\alpha_1, \beta) \in M$  there is an elementary abelian group  $T_{-}(\alpha_1, \beta)$  of translations of  $M^{(\alpha_1, \beta)}$  and  $M_{(\alpha_1, \beta)}$ , and  $T_{-}(\alpha_1, \beta) \lesssim \text{Aut}(M)$  (3.2, 3.4, 3.5). Each  $T_{-}(\alpha_1, \beta)$  fixes all lines of  $L^{-}$  and one line of  $L^{+}$  (namely the line  $\{(\alpha, \beta) \mid \alpha \in \Omega\}$ ). Using the notation of section 2, each  $T_{-}(\alpha_1, \beta)$  consists of positive automorphisms of the form  $(1, b)$ , where  $b \in S^{\Omega}$  fixes  $\beta$  and  $Gb = G$ , i.e. for each  $\beta \in \Omega$  there is an elementary abelian group  $B(\beta)$  which fixes  $\beta$ , acts regularly on  $\Omega \setminus \{\beta\}$ , and for which  $GB(\beta) = G$ . Define  $B := \langle B(\beta) \mid \beta \in \Omega \rangle$ , then  $B$  is doubly transitive on  $\Omega$  and  $GB = G$ . Therefore  $G$  is a union of cosets of  $B$  and in particular  $B \subseteq G$ . Hence, no nontrivial permutation in  $B$  leaves 3 letters fixed. By a theorem of FEIT ([4]),  $B$  contains a normal subgroup of order  $n+1$  or there exists an exactly triply transitive permutation group  $B_0$  containing  $B$  such that  $[B_0 : B] \leq 2$ . Suppose  $B$  contains a normal subgroup of order  $n+1$ , then  $B$  also contains a sharply doubly transitive subgroup  $B^*$ . The circles  $\{(\alpha, \alpha^g) \mid \alpha \in \Omega\}$ ,  $g \in B^*$  together with the lines  $\ell \in L$  now constitute an affine plane of order  $n+1$  and hence configuration as described in 4.1 exist, a contradiction. Therefore  $B \leq B_0$ , where  $B_0$  is sharply 3-transitive, and  $[B_0 : B] \leq 2$ . All sharply triply transitive groups are known (see [6]). If  $n$  is even, then  $B_0 \simeq \text{PSL}(2, n)$  and so  $B = G = \text{PSL}(2, n)$ , i.e.  $M$  is the classical Minkowski plane of order  $n = 2^h$ . If  $n$  is odd, there are at most two sharply 3-transitive groups of degree  $n+1$  and such a group certainly contains  $\text{PSL}(2, n)$ . The Sylow  $p$ -subgroups  $B(\beta)$  of  $B$  are the Sylow  $p$ -subgroups of  $\text{PSL}(2, n)$ . Therefore  $B \leq \text{PSL}(2, n)$  and since  $|B| \geq \frac{1}{2}(n+1)(n)(n-1)$  it follows that  $B \simeq \text{PSL}(2, n)$ . Thus, with  $G_1 := \text{PSL}(2, n)$  and  $G_2 := \text{PSL}(2, n) \setminus \text{PSL}(2, n)$ ,

$$G = G_1 \cup \phi G_2$$

for some  $\phi \in S^\Omega$ . It remains to show that  $\phi$  is an automorphism of  $GF(n)$ .

If  $x, y$  and  $z$  are three distinct points of  $\Omega$ , then there is a  $g \in G_1$  such that  $x^\phi = x^g$ ,  $y^\phi = y^g$ ,  $z^\phi = z^g$  for otherwise there exists  $h \in G_2$  such that  $x^\phi = x^{\phi h}$ ,  $y^\phi = y^{\phi h}$ ,  $z^\phi = z^{\phi h}$ , i.e.  $h = 1$ , contradicting  $h \in G_2$ . It follows that we may assume w.l.o.g. that  $\phi$  fixes  $0, 1$  and  $\infty$ . If we do so it also follows that

$$\frac{x^\phi - y^\phi}{x - y} = \text{square in } GF(n) \text{ for all } x, y \in GF(n), \quad x \neq y,$$

for  $g \in G_1$  determined by  $x^\phi = x^g$ ,  $y^\phi = y^g$ ,  $\infty = \infty^\phi = \infty^g$  has determinant  $\frac{x^\phi - y^\phi}{x - y}$ . By a theorem of BRUEN and LEVINGER (see [2]) it follows that  $\phi$  is an automorphism of  $GF(n)$ .  $\square$

Using the previous theorem it is possible to give a geometric characterization of the Minkowski planes  $(\Omega, G(\phi))$ . Consider the following configurational condition:

(D): Let  $\varepsilon$  be + or -,  $\ell \in L^\varepsilon$  and  $V, W$  two distinct points on  $\ell$ . Suppose  $c$  and  $c'$  are two distinct circles touching in  $V$ . Let  $Y$  and  $Q$  be two distinct points on  $c$ ,  $Y \setminus W$ ,  $Q \setminus W$ . Define

$$Y' := c' \cap [Y]_{-\varepsilon},$$

$$Q' := c' \cap [Q]_{-\varepsilon},$$

$$d := (Y, Q, W),$$

$$d' := (Y', Q', W).$$

Then  $d$  and  $d'$  touch in  $W$  (see figure 3).




Fig. 3.

Notice that (D) is nothing but a special case of the Desargues configuration (D1) in  $M^Z$  on the points  $P, Q, R, P', Q', R'$ .

**4.3. THEOREM.** *Let  $M$  be a Minkowski plane of order  $n \geq 5$ , and suppose (D) holds in  $M$ . Then  $M$  is isomorphic to one of the planes  $(\Omega, G(\phi))$ .*

Of course the proof of 4.3 is based on 4.2 and it is clear that (D) implies (A). Also (C) is a consequence of (D).

**4.4. LEMMA.** *Let  $M$  be a Minkowski plane of order  $n$*

- a) *If  $n$  is even then (A) implies (B) (hence (C)).*
- b) *In any case (D) implies (C).*

**PROOF.** a) The following statement is easily seen to be equivalent to (B): If the circles  $c$  and  $d$  as described in (A) exist, then  $P_1 \in (P_2^+, P_2^-, Q_2) \iff P_2 \in (P_1^+, P_1^-, Q_1)$ . To prove this last statement, consider the configuration of condition (A) and suppose  $c$  and  $d$  exist,  $P_2 \in (P_1^+, P_1^-, Q_1)$  but  $P_1 \notin (P_2^+, P_2^-, Q_2)$ . Let  $e$  be the circle through  $P_1$  touching  $(P_2^+, P_2^-, Q_2)$  and  $c$  in  $P_2^+$ ,  $f$  the circle through  $P_1$  touching  $(P_1^+, P_1^-, Q_1)$  in  $P_2^-$ . By (A)  $e$  and  $f$  touch in  $P_1$ . Similarly it follows that the circle  $g$  through  $P_1$  touching  $(P_2^+, P_2^-, Q_2)$  in  $P_2^-$  touches  $f$  in  $P_1$ . Therefore  $g$  and  $e$  touch in  $P_1$  and so the circles  $g, e, (P_2^+, P_2^-, Q_2)$  touch each

other in  $P_2^+, P_2^-, P_1$ . This contradicts 4.1 since  $n$  is even.

b) Consider the configuration of condition (C). We claim that  $(P_1, Q_1, Z)$  and  $(P_2, Q_2, Z)$  touch in  $Z$ . If  $(P_i, Q_i, Z)$  touches  $c_i$  in  $Q_i$  for  $i = 1, 2$ , this follows from (A). Suppose therefore that  $(P_1, Q_1, Z)$  does not touch  $c_1$  in  $Q_1$ , i.e. suppose that  $(P_1, Q_1, Z)$  has another point  $E_1 \neq Q_1$  in common with  $c_1$ . Put  $E_2 := [E_1]_{-\varepsilon} \cap c_2$ . By (D) the circles  $(E_2, Q_2, Z)$  and  $(E_1, Q_1, Z) = (P_1, Q_1, Z)$  touch in  $Z$ . Suppose  $(E_2, Q_2, Z)$  intersects  $[A]_{-\varepsilon}$  in a point  $P_2' \neq P_2$ . Let  $Y$  be the point of intersection of  $[Z]_{\varepsilon}$  and  $(E_2, P_2', C_2)$ . If we apply (D) twice it follows that  $(E_1, P_1, Y)$  and  $(E_1, C_1, Y)$  both touch  $(E_2, P_2', C_2)$  in  $Y$ . Hence  $(E_1, P_1, Y) = (E_1, C_1, Y)$  and impossibility because  $P_1 \parallel C_1$ . We have proved  $P_2 \in (E_2, Q_2, Z)$ , i.e.  $(P_1, Q_1, Z)$  and  $(P_2, Q_2, Z)$  touch in  $Z$ . So:  $c_1$  and  $c_2$  touch in  $A$  implies  $(P_1, Q_1, Z)$  and  $(P_2, Q_2, Z)$  touch in  $Z$ . It is easily seen that the converse also holds. If we replace  $c_i$  by  $d_i$ ,  $i = 1, 2$ , it follows that  $d_1$  and  $d_2$  touch in  $B$ .  $\square$

To finish the proof of 4.3 we have to show that all residual planes  $M^Z$  are nearaffine translation planes. By 3.4 it suffices to show that all derived planes  $M_Z$  are translation planes.

**4.5. LEMMA.** *Let  $M$  be a Minkowski plane satisfying (D), then  $M_Z$  is a translation plane for every point  $Z$ .*

**PROOF.** Let  $Z \in M$  and  $P, Q, R, P', Q', R' \in M_Z$  such that  $P \parallel P', Q \parallel Q', R \parallel R'$ , the line  $PQ$  (in  $M_Z$ ) is parallel to  $P'Q'$  and  $PR$  is parallel to  $P'R'$ . We have to show that  $QR$  is parallel to  $Q'R'$ , i.e. we have to show that the circles  $(Z, Q, R)$  and  $(Z, Q', R')$  touch in  $Z$ . We assume here that  $P, Q, R$  (and also  $P', Q', R'$ ) are mutually nonparallel. The other cases follow from the cases we do consider. Put  $Y = (P, Q, R) \cap [Z]_+$ . If we apply (D) to  $(P, Q, Z)$ ,  $(P', Q', Z)$ ,  $(P, Q, Y) = (P, Q, R)$  and  $(P', Q', Y)$ , it follows that  $(P, Q, R)$  and  $(P', Q', Y)$  touch in  $Y$ . Application of (D) to  $(P, R, Z)$ ,  $(P', R', Z)$ ,  $(P, R, Y) = (P, Q, R)$  and  $(P', R', Y)$  yields  $(P, Q, R)$  and  $(P', R', Y)$  touch in  $Y$ . Hence  $(P', Q', Y) = (P', R', Y) = (P', Q', R')$ . Finally we apply (D) to  $(Q, R, Y)$ ,  $(Q', R', Y)$ ,  $(Q, R, Z)$  and  $(Q', R', Z)$  and obtain the desired result.  $\square$

Notice that it is possible to give a proof of 4.3 without using the theory of nearaffine planes. Show directly, using (D), that any translation

of a desired plane  $M_{\mathbb{Z}}$  extends to an automorphism of  $M$ . Then argue as we did in 4.2.

## REFERENCES

1. BENZ, W., *Vorlesungen über Geometrie der Algebren*, Springer-Verlag, Berlin - New York (1973).
2. BRUEN, A. & LEVINGER, B., A theorem on permutations of a finite field, *Can. J. Math.*, 25 (1973), 1060-1065.
3. DEMBOWSKI, P., *Finite geometries*, Springer-Verlag, Berlin - Heidelberg - New York (1968).
4. FEIT, W., On a class of doubly transitive permutation groups, *Ill. J. Math.*, 4 (1960), 170-186.
5. HEISE, W. & KARSEL, H., Symmetrische Minkowski-Ebenen, *J. Geometry*, 3 (1973), 5-20.
6. HUPPERT, B., Scharf dreifach transitive Permutationsgruppen, *Arc. Math.*, 13 (1962), 61-72.
7. PEDRINI, C., 3-reti (non immergibili) aventi dei piani duali di quelli di Moulton quali sotto piani, *Atti Accad. Naz. Lincei Rend. Sc. fis. mat. e nat.*, 40 (1966), 385-392.
8. QUATTROCCHI, P., Sugli insiemi di sostituzioni strettamente 3- transitivi-finiti, *Atti Sem. mat. fis. Univ. Modena*, 24 (1975), 279-289 (1976).
9. WILBRINK, H., Nearaffine planes, To appear.
10. WILBRINK, H., Nearaffine planes and Minkowski planes, Master's thesis, Tech Univ. Eindhoven (1978).

H. Wilbrink  
 Mathematisch Centrum  
 2e Boerhaavestraat 49  
 1091 AL Amsterdam  
 The Netherlands



ONTVANGEN 1 9 OKT 1979