

PREPRINT
NOT FOR REVIEW

**stichting
mathematisch
centrum**



DEPARTMENT OF PURE MATHEMATICS

ZW 67/76 JANUARY

K.R. APT

RECURSIVE EMBEDDINGS OF PARTIAL ORDERINGS

Prepublication

2e boerhaavestraat 49 amsterdam

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O), by the Municipality of Amsterdam, by the University of Amsterdam, by the Free University at Amsterdam, and by industries.

Recursive embeddings of partial orderings ^{*)}

by

K.R. Apt

ABSTRACT

Let A be a countable atomless Boolean algebra and let X be a countable partial ordering. We prove that there exists an embedding of X into A which is recursive in X, A and which destroys all suprema and infima of X which can be destroyed. We show that the above theorem is false when we try to preserve all suprema and infima of X instead of destroying them. Finally we indicate that if A and B are countable Boolean algebras and B is atomless then A can be embedded into B by a function which is recursive in A, B . If A is also atomless, then there is an isomorphism from A into B which is recursive in A, B .

KEY WORDS & PHRASES: *countable atomless Boolean algebras, partial orderings, recursive embeddings*

^{*)} This paper is not for review; it is meant for publication elsewhere.

§1. PRELIMINARIES*

Throughout the paper ω denotes the set of natural numbers, and ϕ the empty set. If X is a set and n a natural number then X^n denotes the set of all n -tuples of elements of X . We say that X is a partial ordering on a set A (p.o. on A) if for some $B \subset A$ $X \subset B^2$ and for all $x, y, z \in B$

1. $(x, x) \in X$,
2. $((x, y) \in X \wedge (y, x) \in X) \longrightarrow x = y$,
3. $((x, y) \in X \wedge (y, z) \in X) \longrightarrow (x, z) \in X$.

If $(x, y) \in X$, we write $x \leq_X y$. If $(x, y) \in X \wedge x \neq y$, we write $x <_X y$.

If $(x, y) \notin X$ and $(y, x) \notin X$, we say that x and y are X -incomparable and we write $x \parallel_X y$.

z is called the supremum of x and y in X ($x \cup y = z$), if

$$x \leq_X z \wedge y \leq_X z \wedge \forall t[(x \leq_X t \wedge y \leq_X t) \longrightarrow z \leq_X t]$$

z is called the infimum of x and y in X ($x \cap y = z$), if

$$z \leq_X x \wedge z \leq_X y \wedge \forall t[(t \leq_X x \wedge t \leq_X y) \longrightarrow t \leq_X z]$$

By $\text{Fld}(X)$ we denote the set $\{x : (x, x) \in X\}$.

For the definition of a Boolean algebra we refer the reader to SIKORSKI [4].

If A is a Boolean algebra then 0 denotes its smallest element and 1 the greatest one.

If x and y are elements of A , then we write $x \leq y$ if $x \cup y = y$ and $x < y$ if $x \leq y$ and $x \neq y$. We write $x \parallel y$ if $\neg(x \leq y)$ and $\neg(y \leq x)$.

We say that A is a Boolean algebra on a set A , if every element of A is an element of A .

In this paper we are interested in partial orderings on ω and Boolean algebras on ω .

*Some of the results of this paper were obtained in 1971 when the author was a student at Wrocław University, Poland.

DEFINITION 1. Let X be a p.o. on a set A and \bar{A} a Boolean algebra. f is called an embedding of X into \bar{A} if f is an injective function from $\text{Fld}(X)$ into \bar{A} such that for all $x, y \in \text{Fld}(X)$

$$x <_X y \leftrightarrow f(x) < f(y)$$

We say that an embedding f of X into \bar{A} preserves all suprema and infima of X if

- I) whenever $x \cup y = z$ then $f(x) \cup f(y) = f(z)$,
- II) whenever $x \cap y = z$ then $f(x) \cap f(y) = f(z)$.

We say that an embedding f of X into \bar{A} destroys all suprema and infima of X if

- I) whenever $x \parallel_X y$ and $x \cup y = z$ then $f(x) \cup f(y) \neq f(z)$,
- II) whenever $x \parallel_X y$ and $x \cap y = z$ then $f(x) \cap f(y) \neq f(z)$.

Observe that if $x \leq_X y$ then $x \cup y = y$ and $x \cap y = x$, so for any embedding f of X into \bar{A} $f(x) \cup f(y) = f(x \cup y)$ and $f(x) \cap f(y) = f(x \cap y)$.

Thus an embedding of X into \bar{A} cannot destroy suprema and infima of X -comparable elements.

All the notions from recursion theory we use can be found in SHOENFIELD [2]. In particular $\text{Seq}(x)$ means that x codes a finite sequence of natural numbers, $\text{lh}(x)$ is the length of that sequence. If $\text{Seq}(x)$ then

$$x = \langle (x)_0, \dots, (x)_{\text{lh}(x)-1} \rangle. \text{ If } a = \langle a_1, \dots, a_n \rangle \text{ and } b = \langle b_1, \dots, b_n \rangle \text{ then } a * b = \langle a_1, \dots, a_n, b_1, \dots, b_n \rangle.$$

All the mentioned functions and relations are recursive.

If $A = \{a_1, \dots, a_k\}$ then x is called the code of A ($x = \langle A \rangle$) if x is the least number z such that $\text{seq}(z)$, $\text{lh}(z) = k$ and $\{(z)_i : i < \text{lh}(z)\} = A$. If $f(x_1, \dots, x_n)$ is a function then $\text{graph}(f) = \{(x_1, \dots, x_n, y) : f(x_1, \dots, x_n) = y\}$.

DEFINITION 2. Let $\bar{A} = \langle A, \cup, \cap, -, 0, 1 \rangle$ be a Boolean algebra on ω .

We say that f is recursive in \bar{A} if f is recursive in $\{A, \text{graph}(\cup), \text{graph}(\cap), \text{graph}(-)\}$.

Similarly we define that f is recursive in A, B where B is another Boolean algebra on ω or that f is recursive in X, A for a set X .

DEFINITION 3. Let A be a Boolean algebra. Suppose that A and B are sets of elements of A . Then

- I) if $a \leq b$ for all $a \in A, b \in B$ we write $A \leq B$,
- II) if $a < b$ for all $a \in A, b \in B$ we write $A < B$,
- III) if $\neg(a \leq b)$ for all $a \in A, b \in B$ we write $A \not\leq B$,
- IV) if $a \parallel b$ for all $a \in A, b \in B$ we write $A \parallel B$.

Instead of $\{a\} < A$ we write $a < A$. Similarly with other relations.

Observe that for every set A $\phi < A, A < \phi, \phi \not\leq A, A \not\leq \phi$ and $\phi \parallel A$.

If A is a finite set of elements of A then $\sup A$ denotes the least element a of A such that $A \leq a$, and $\inf A$ denotes the greatest element a of A such that $a \leq A$. Observe that $\sup \phi = 0$ and $\inf \phi = 1$. Recall that a Boolean algebra A is atomless if $0 < x$ implies for some y $0 < y < x$.

§2. EMBEDDINGS DESTROYING SUPREMA AND INFIMA

We prove in this section the following theorem:

Theorem 1. Let X be a partial ordering on ω and let A be an atomless Boolean algebra on ω . Then there exists an embedding f of X into A such that

- I) f destroys all suprema and infima of X
- II) f is recursive in X, A .

We present at first an informal idea of the proof. Let $\text{Fld}(X) = \{a_0, a_1, \dots\}$ be a recursive in X enumeration of $\text{Fld}(X)$. We want to build the required embedding by induction. Suppose that for $i \leq n$ we already defined some elements b_i of A such that


$$a_i <_X a_j \leftrightarrow b_i < b_j \quad \text{for } i, j \leq n.$$

We want to define an element b_{n+1} of A such that

$$a_i <_X^{a_j} \leftrightarrow b_i < b_j \quad \text{for } i, j \leq n+1. \quad (*)$$

If we do not impose any conditions on b_i -s we can be stuck. For example if $a_0 < a_2$, $a_1 < a_2$, $a_0 < a_3$, $a_1 < a_3$ and $a_3 < a_2$

(in the picture



and we chose b_0 , b_1 and b_2 in such a way that $b_0 \cup b_1 = b_2$ then there is no b_3 such that $b_3 < b_2$, $b_0 < b_3$ and $b_1 < b_3$.

In order to prevent such situations we choose b_i -s in a more careful way. For example the above difficulty would not occur if $b_0 \cup b_1 < b_2$. Thus we assume that the elements b_0, \dots, b_n satisfy certain additional property, namely that the set $\{b_0, \dots, b_n\}$ is normal (see Definition 4).

$$\begin{aligned} \text{Let } A &= \{b_i : a_i <_X^{a_{n+1}}, \quad i \leq n\} \\ B &= \{b_i : a_{n+1} <_X^{a_i}, \quad i \leq n\} \\ C &= \{b_i : a_{n+1} \parallel_X^{a_i}, \quad i \leq n\} \end{aligned}$$

Then $A \cup B \cup C = \{b_0, \dots, b_n\}$. Observe that $A < B$, $C \not\leq A$ and $B \not\leq C$. Since $A \cup B \cup C$ is a normal set we get from this that $\sup A < \inf B, C \not\leq \sup A$ and $\inf B \not\leq C$. We are looking for an element b_{n+1} such that $\sup A < b_{n+1} < \inf B$ and $b_{n+1} \parallel C$. Then $(*)$ holds. The existence of such a b_{n+1} is guaranteed by lemma 1.

But we want also to preserve our additional condition, so we claim also that the set $A \cup B \cup C \cup \{b_{n+1}\}$ is to be normal. The lemma 2 shows that the required b_{n+1} still can be found. Its proof uses the lemma 1, but in an approximately modified way.

Thus the induction step works. The obtained embedding destroys all suprema and infima of X which is an immediate consequence of the fact that for each n the set $\{b_0, \dots, b_n\}$ is normal.

Choosing at each time the smallest b_{n+1} satisfying the above conditions (look at the Definition of the function g in the proof of theorem 1) we ensure that the above embedding is recursive in X , A .

We present now the precise proof of the theorem :

We prove at first two lemmata

Lemma 1. *Let A be an atomless Boolean algebra. Suppose that $A \cup \{a, b\}$ is a finite set of elements of A , such that :*

1. $a < b$
2. $A \not\leq a$
3. $b \not\leq A$

Then there exists an element c of A , such that $a < c < b$ and $c \parallel A$.

Obviously the conditions 2. and 3. have to be satisfied if we want to prove the claim. The lemma shows that 2. and 3. are also sufficient conditions.

Proof. At first we "modify" A to a set A' such that $a < A' < b$. We find then an element c such that $a < c < b$ and $c \parallel A'$. It turns out that also $c \parallel A$.

Let

$A' = \{b \cap d : d \in A \text{ \& } a < b \cap d\} \cup \{a \cup d : d \in A \text{ \& } a \cup d < b\}$. Suppose that $x = b \cap d$ for some $d \in A$ such that $a < b \cap d$. Then $x \leq b$. If $x = b$ then $b \leq d$ which violates our assumptions. Thus $a < x < b$.

Suppose now that $x = a \cup d$ for some $d \in A$ such that $a \cup d < b$. Then $a \leq x$. If $a = x$ then $d \leq a$ which violates our assumptions. Thus $a < x < b$. So $a < A' < b$.

We can treat the set $B = \{x : a \leq x \leq b\}$ as a Boolean algebra with the operations induced by A .

$$\begin{aligned} x \dot{\cup} y &= x \cup y \\ x \dot{\cap} y &= x \cap y \\ \dot{0} &= a \\ \dot{1} &= b \\ \dot{-} x &= a \cup (b \cap \neg x) \end{aligned}$$

Let $A' = \{a_1, \dots, a_n\}$. We just proved that $\dot{0} < a_i$ and $\dot{0} < -a_i$ for all $i \leq n$.

Let $C = \{b_1 \cap \dots \cap b_n : \text{for all } i \leq n \ b_i = a_i \text{ or } b_i = -a_i\}$. Then each a_i or $-a_i$ is a sum of elements of C .

For each $j \leq 2n$ pick an element c_j from C such that

$$\begin{aligned} \dot{0} < c_j &\leq a_j \\ \dot{0} < c_{j+n} &\leq -a_j \end{aligned} \quad \text{for all } j \leq n.$$

A is atomless so there exist elements d_i such that for $i \leq 2n$

$$\dot{0} < d_i < c_i.$$

We can choose d_i -s in such a way that $d_i = d_j$ if $c_i = c_j$.

Finally let $c = d \cup \dots \cup d_{2n}$. We claim that c is the desired element.

We prove at first that $c \parallel A'$. Suppose that for some $1 \leq i \leq n$ $c \leq a_i$. Then

$$\dot{0} < d_{i+n} \leq a_i \text{ and } d_{i+n} < -a_i$$

which is clearly impossible. If for some $i \leq n$ $a_i \leq c$ then

$$c_i \cap -d_i \leq c_i \leq a_i \leq c.$$

Observe that for $x, y \in C$ either $x = y$ or $x \cap y = \dot{0}$. Hence for $k \leq n$ either $c_k = c_i$ or $c_k \cap c_i = \dot{0}$. In the first case $d_k = d_i$, in the second $d_k \cap (c_i \cap -d_i) = \dot{0}$. So in both cases we obtain $d_k \cap (c_i \cap -d_i) = \dot{0}$.

Finally we obtain :

$$c_i \cap -d_i = c_i \cap -d_i \cap c = \bigcup_{k=1}^{2n} d_k \cap (c_i \cap -d_i) = \dot{0}.$$

which contradicts the choice of d_i .

Observe that by construction $a < c < b$. We prove now that $c \parallel A$.

Suppose that $x \in A$. There are 3 possible cases

I) $x \parallel a$ and $x \parallel b$.

Then for every y such that $a \leq y \leq b$ $x \parallel y$, so in particular $x \parallel c$.

II) $x < b$

There are two possible cases

1. $a \cup x < b$

Then $a \cup x \in A'$, So $a \cup x \parallel c$.

If $x \leq c$ then $a \cup x \leq c$ which is impossible

If $c \leq x$ then $c \leq a \cup x$ which is impossible. Thus $c \parallel x$.

2. $a \cup x = b$

If $x \leq c$ then $a \cup x \leq c$, so $b \leq c$ which is impossible

If $c \leq x$ then $a \leq x$, so $a \cup x = x$ i.e. $b = x$ which contradicts our assumptions. Thus $c \parallel x$.

III) $a < x$

There two possible cases

1. $a < b \cap x$

Then $b \cap x \in A'$, so $b \cap x \parallel c$

If $x \leq c$ then $b \cap x \leq c$ which is impossible.

If $c \leq x$ then $c \leq b \cap x$ which is impossible.

Thus $x \parallel c$.

2. $a = b \cap x$

If $x \leq c$ then $x \leq b$, so $b \cap x = x$ i.e. $a = x$ which contradicts our assumptions.

If $c \leq x$ then $c \leq b \cap x$ i.e. $c \leq a$ which is impossible.

Thus $c \parallel x$.

This concludes the proof of the lemma. \square

DEFINITION 4. Let A be a Boolean algebra. A finite set T of elements of A is called normal if for all A and B , such that $A \cup B \subset T$.

$A < B$ implies $\sup A < \inf B$,

$A \not\leq B$ implies $\inf A \not\leq \sup B$.

Lemma 2. Let A be an atomless Boolean algebra. Suppose that for some finite sets A, B and C of elements of A

$A < B$

$C \not\leq A$

$B \not\leq C$

$A \cup B \cup C$ is normal

Then there exists an element a of A such that

$$\begin{aligned} \sup A &< a < \inf B \\ a \parallel C \\ A \cup B \cup C \cup \{a\} &\text{ is normal} \end{aligned}$$

Proof. Let S be a subalgebra of A generated by the set $A \cup B \cup C$. Let $T = \{x : x \in S \wedge \neg(x \leq \sup A) \wedge \neg(\inf B \leq x)\}$. The set T is of course finite.

Since $A \cup B \cup C$ is normal we get from our assumptions that $\sup A < a < \inf B$ and $a \parallel T$. We claim that a is the required element. If $c \in C$ then $c \not\leq A$ and $B \not\leq c$. Since $A \cup B \cup C$ is normal, $c \not\leq \sup A$ and $\inf B \not\leq c$. Thus $C \subseteq T$ i.e. $a \parallel C$.

It is left to prove that $A \cup B \cup C \cup \{a\}$ is normal.

Let $K \cup L \subseteq A \cup B \cup C$. We have to consider the following four possible cases

1. $K < L$ and $a < L$

We prove that then $\sup(K \cup \{a\}) < \inf L$

Always $\sup(K \cup \{a\}) \leq \inf L$. Suppose that $\sup(K \cup \{a\}) = \inf L$.

Then $\sup K \cup a = \inf L$, so

$$\inf L \cap -\sup K \leq a$$

which indicates that $\inf L \cap -\sup K \notin T$. There are two possibilities

- I) $\inf B \leq \inf L \cap -\sup K$. Then $\inf B \leq a$ which contradicts the choice of a
- II) $\inf L \cap -\sup K \leq \sup A$. Then

$$\inf L \leq \sup A \cup \sup K$$

$$\text{i.e. } \inf L \leq \sup(A \cup K).$$

The assumption $a < L$ implies by the choice of a that $L \subseteq B$.

Thus $A < L$ since $A < B$. So $A \cup K < L$. But $A \cup B \cup C$ is normal, so we get that $\sup(A \cup K) < \inf L$, which contradicts our previous statement.

2. $K < L$ and $K < a$

We prove that $\sup K < \inf(L \cup \{a\})$

Always $\sup K \leq \inf(L \cup \{a\})$. Suppose that $\sup K = \inf(L \cup \{a\})$. Then $\sup K = \inf L \cap a$, so $a \leq \sup K \cup -\inf L$.

This indicates that $\sup K \cup -\inf L \notin T$. There are two possibilities

I) $\sup K \cup -\inf L \leq \sup A$.

Then $a \leq \sup A$ which is impossible.

II) $\inf B \leq \sup K \cup -\inf L$.

Then $\inf B \cap \inf L \leq \sup K$

i.e. $\inf(B \cup L) \leq \sup K$.

But $K < a$, so $K \subset A$ i.e. $K < B$. Thus $K < L \cup B$. Since $A \cup B \cup C$ is normal we get

$\sup K < \inf(B \cup L)$,

which contradicts the former statement.

3. $K \not\leq L$ and $a \not\leq L$

We prove that $\inf(K \cup \{a\}) \not\leq \sup L$.

Suppose that $\inf(K \cup \{a\}) \leq \sup L$, i.e.

$\inf K \cap a \leq \sup L$.

Then $a \leq \sup L \cup -\inf K$, so $L \cup -\inf K \notin T$.

There are two possibilities

I) $\sup L \cup -\inf K \leq \sup A$.

Then $a \leq \sup A$ which contradicts the choice of a .

II) $\inf B \leq \sup L \cup -\inf K$.

Then $\inf B \cap \inf K \leq \sup L$, i.e.

$\inf(B \cup K) \leq \sup L$.

But $a \not\leq L$, so by the choice of a $B \not\leq L$ i.e. $B \cup K \not\leq L$. Since $A \cup B \cup C$ is normal we get that $\inf(B \cup K) \not\leq \sup L$, which contradicts the former statement.

4. $K \not\leq L$, $K \not\leq a$

We prove that $\inf K \not\leq \sup(L \cup \{a\})$. Suppose that $\inf K \leq \sup(L \cup \{a\})$

i.e. $\inf K \leq \sup L \cup a$. Then

$\inf K \cap -\sup L \leq a$, so $\inf K \cap -\sup L \notin T$.

There are two possibilities

I) $\inf K \cap - \sup L \leq \sup A$.

Then $\inf K \leq \sup A \cup \sup L$ i.e.

$\inf K \leq \sup(A \cup L)$

On the other hand $K \not\leq a$, so by the choice of a

$K \not\leq A$ i.e. $K \not\leq A \cup L$.

$A \cup B \cup C$ is normal, so $\inf K \leq \sup(A \cup L)$ which gives the contradiction.

II) $\inf B \leq \inf K \cap - \sup L$.

Then $\inf B \leq a$ which contradicts the choice of a .

This completes the proof that $A \cup B \cup C \cup \{a\}$ is normal, so the proof of the lemma is concluded. \square

Proof of the theorem 1.

Observe that the relation

$P(x) \leftrightarrow x$ is a code of a finite set

is recursive.

It is easy to see that the relation

$T(x) \leftrightarrow x$ is a code of a normal set of elements of A

is recursive to A .

Define a function g as follows:

$$g(x,y,z) = \begin{cases} \text{if } x,y \text{ and } z \text{ are respectively codes of the sets } A, B \\ \text{and } C \text{ satisfying the conditions of lemma 2} \\ \text{then } \mu a \text{ (} a \text{ satisfies the claim of lemma 2)} \\ \text{else } 0 \end{cases}$$

Then g is a total function recursive in A .

$\text{Fld}(X)$ is a recursive in X set, so for some total injective function $a(x)$, which is recursive in X .

$$\text{Fld}(X) = \{a(0), a(1), \dots\}$$

For any total function $h(x)$ and $n \geq 0$ let

$$A(h,n) = \{h(k) : a(k) <_X a(n+1), k \leq n\}$$

$$B(h,n) = \{h(k) : a(n+1) <_X a(k), k \leq n\}$$

$$C(h,n) = \{h(k) : a(k) \parallel_X a(n+1), k \leq n\}$$

Let b be an arbitrary element of A such that $0 < b < 1$.

Define a function h as follows:

$$h(0) = b$$

$$h(n+1) = g(\langle A(h,n) \rangle, \langle B(h,n) \rangle, \langle C(h,n) \rangle).$$

h is a well defined total function. It is easy to see that h is recursive in X, A .

Finally define

$$f(a(n)) = h(n) \quad \text{for } n \geq 0.$$

We claim that f is the required function.

Observe that

$$f(x) = y \leftrightarrow \exists n(x=a(n) \wedge y=h(n)).$$

so f is recursive in X, A .

By induction on k , we prove that for all k

- I) $a(i) <_X a(j)$ iff $f(a(i)) < f(a(j))$ for all $i, j \leq k$,
- II) the set $\{f(a(i)) : i \leq k\}$ is normal.

Observe that the set $\{f(a(0))\}$ is normal, so I) and II) is true for $k = 0$.

Suppose that I) and II) are true for k . Then I) implies that

$$A(h,k) < B(h,k)$$

$$C(h,k) \not\leq A(h,k)$$

$$B(h,k) \not\leq C(h,k)$$

Also $A(h,k) \cup B(h,k) \cup C(h,k) = \{f(a(i)) : i \leq k\}$ so it is a normal set.

Thus the sets $A = A(h,k)$, $B = B(h,k)$, $C = C(h,k)$ satisfy the claim of the lemma 2.

$$g(\langle A(h,k) \rangle, \langle B(h,k) \rangle, \langle C(h,k) \rangle) = f(a(k+1)),$$

so by the definition of the function g

$$\sup A(h,k) < f(a(k+1)) < \inf B(h,k),$$

$f(a(k+1)) \parallel C(h,k)$ and $A(h,k) \cup B(h,k) \cup C(h,k) \cup \{f(a(k+1))\}$ is a normal set.

Observe now that for $i < k + 1$

$$\begin{aligned} a(i) <_X a(k+1) &\leftrightarrow f(a(i)) \in A(h,k) \leftrightarrow f(a(i)) < f(a(k+1)) \\ a(k+1) <_X a(i) &\leftrightarrow f(a(i)) \in B(h,k) \leftrightarrow f(a(k+1)) < f(a(i)) \\ a(i) \parallel_X a(k+1) &\leftrightarrow f(a(i)) \in C(h,k) \leftrightarrow f(a(i)) \parallel f(a(k+1)) \end{aligned}$$

Thus I) and II) are true for $k + 1$.

Hence by induction for all i and j

$$a(i) <_X a(j) \leftrightarrow f(a(i)) < f(a(j))$$

Since f is also injective it is an embedding of X into A .

It is left to show that f destroys all suprema and infima.

Suppose that for some i, j, k $a(i) \parallel_X a(j)$ and $a(i) \cup a(j) = a(k)$.

Then $a(i) <_X a(k)$ and $a(j) <_X a(k)$, so $f(a(i)) < f(a(k))$ and $f(a(j)) < f(a(k))$.

The set $\{f(a(n)) : n \leq \max(i, j, k)\}$ is normal thus

$$f(a(i)) \cup f(a(j)) < f(a(k)).$$

i.e. f destroys the supremum $a(i) \cup a(j)$. The same argument applies in the case of infimum of X -incomparable elements.

This concludes the proof of the theorem. \square

§3 EMBEDDING PRESERVING SUPREMA AND INFIMA

Let $A = \{x : \text{Seq}(x) \wedge \forall i (i < \text{lh}(x) \rightarrow ((x)_i = 0 \vee (x)_i = 1))\}$

Thus A is the set of codes of all finite sequences of zeroes and ones.

Let \cup and \cap be some operations on A satisfying the following property;

if

$$\langle k_1, \dots, k_n \rangle \in A$$

then

$$\langle k_1, \dots, k_n, 0 \rangle \cup \langle k_1, \dots, k_n, 1 \rangle = \langle k_1, \dots, k_n \rangle$$

$$\langle k_1, \dots, k_n, 0 \rangle \cap \langle k_1, \dots, k_n, 1 \rangle = \langle 0 \rangle.$$

Let M be the Boolean algebra generated by A and by operations \cup and \cap satisfying the above property. It is well known that M is (isomorphic to) the Boolean algebra of all clopen subsets of the Cantor Space. The elements of M are just all the finite joins and meets of A .

It is easy to see that M is recursive, that is to say

$$M = \langle A_M, \cup, \cap, -, 0, 1 \rangle \quad \text{where}$$

A_M is a recursive set and the graphs of partial functions \cup, \cap and $-$ are recursive. M is an atomless Boolean algebra.

We prove the following theorem.

Theorem 2. There exists a recursive partial ordering X on ω , such that

- I) there is an embedding of X into M which preserves all suprema and infima of X .
- II) no such embeddings are recursive.

Proof. Let $P(x)$ be a $\Sigma_2^0 - \Pi_2^0$ relation. For some recursive R

$$P(x) \leftrightarrow \exists y \forall z R(x, y, z)$$

Define a partial function g as follows:

$$g(x, y) \simeq \langle x, y, \mu z \neg R(x, y, z) \rangle$$

Observe that graph (g) is recursive.

Define


$$h(x, y) \simeq \langle g(x, 0), \dots, g(x, y-1) \rangle \quad \text{where } y \div 1 = \max(y-1, 0)$$

Clearly h is a partial recursive function.

Observe that

1. $(h(x,y) \text{ is defined} \ \& \ z < y) \rightarrow (h(x,z) \text{ is defined})$
2. For all x $[\lambda y h(x,y) \text{ is total} \leftrightarrow \lambda y g(x,y) \text{ is total}]$
3. $\text{graph}(h)(x,y,z) \leftrightarrow \text{Seq}(z) \wedge 1h(z) = y \wedge \forall i (i < y \rightarrow \text{graph}(g)(x,i,(z)_i))$.

Our ordering X looks as follows



More formally

$$X = \{(\langle x \rangle, \langle x \rangle) : x \geq 0\} \cup \{(h(x,m), h(x,n)) : x \geq 0, n \geq m \geq 0\} \cup \\ \cup \{(h(x,m), \langle x+1 \rangle) : m \geq 0, x \geq 0\} \cup \{(h(x,m), \langle x+1 \rangle) : x \geq 0, m \geq 0\}.$$

X is clearly a recursive set.

Let now T be the following relation

$$T(x) \leftrightarrow \langle x \rangle \cap \langle x+1 \rangle \text{ exists}$$

Then

$$\begin{aligned} T(x) &\leftrightarrow \lambda y h(x,y) \text{ is not total,} \\ &\leftrightarrow \lambda y g(x,y) \text{ is not total,} \\ &\leftrightarrow \exists y (g(x,y) \text{ is not defined}), \\ &\leftrightarrow \exists y (\forall z R(x,y,z)), \\ &\leftrightarrow P(x). \end{aligned}$$

Hence T is a $\Sigma_2^0 - \Pi_2^0$ relation.

It is easy to see that there is an embedding of X into M which preserves

all suprema and infima of X .

Let f be such an embedding.

Then

$$T(x) \leftrightarrow \exists z(z \in \text{Fld}(X) \wedge (f(\langle\langle x \rangle\rangle) \cap f(\langle\langle x+1 \rangle\rangle) = f(z))).$$

Thus if f was recursive then T would be a Σ_1^0 set, which is not the case. Hence no such embeddings are recursive.

The above theorem shows that theorem 1 is not true when I) is changed for

I') f preserves all suprema and infima of X .

We pass now to the problem of recursive embeddings of Boolean algebras into Boolean algebras. Abian in ABIAN [1] proves the following lemma.

Lemma 3. (Abian). *Let A and B be countable Boolean algebras and let B be atomless. Let f be an isomorphism from a finite subalgebra A_1 of A onto a finite subalgebra B_1 of B . Then for every $a \in A - A_1$ there exists $b \in B - B_1$ such that the assignment $f(a) = b$ extends the isomorphism f from the subalgebra of A generated by $A_1 \cup \{a\}$ onto the subalgebra of B generated by $B_1 \cup \{b\}$.*

Using this lemma Abian gives an algebraic proof of the well known theorem that two countable atomless Boolean algebras are isomorphic. In fact this isomorphism is recursive in the considered algebras. More precisely we have the following theorem.

Theorem 3. Let A and B be countable Boolean algebras on ω and let B be atomless. Then

- I) There exists an embedding of A into B (as Boolean algebras) which is recursive in A, B .
- II) if A is atomless then there exists an isomorphism of A and B which is recursive in A, B .

Proof.

I) Due to the repeated use of the lemma 3.

II) Using repeatedly the lemma 3 in the back and forth way.

It is clear that in both cases the constructed embedding f is recursive in A, B .

REMARK. This paper is closely related with the VAN EMDE BOAS [2] paper. Van Emde Boas proves there that every recursive partial ordering can be recursively embedded into the Boolean algebra M defined on page 13. We obtained the theorem 1 independently of his paper.

REFERENCES

- [1] A. ABIAN, *Categoricity of denumerable atomless Boolean rings*, *Studia Logica*, vol. XXX (1972), 63-68.
- [2] P. VAN EMDE BOAS, *Mostowski's universal set-algebra*, Mathematisch Centrum report, ZW 14/73, 1973.
- [3] J.R. SHOENFIELD, *Mathematical logic*, Addison, Wesley, Reading, Massachusetts, 1967.
- [4] R. SIKORSKI, *Boolean algebras*, third edition, Berlin, 1969.