

# COMPOSITIO MATHEMATICA

J. VAN DER SLOT

## **An elementary proof of the Hewitt-Shirota theorem**

*Compositio Mathematica*, tome 21, n° 2 (1969), p. 182-184.

[http://www.numdam.org/item?id=CM\\_1969\\_\\_21\\_2\\_182\\_0](http://www.numdam.org/item?id=CM_1969__21_2_182_0)

© Foundation Compositio Mathematica, 1969, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

## An elementary proof of the Hewitt-Shirota theorem

by

J. van der Slot

The abbreviation "space" is used to denote "completely regular space".

A well-known theorem of Hewitt and Shirota [1] states that a realcompact completely regular space is homeomorphic with a closed subspace of a product of real lines. Many proofs of this fundamental theorem have appeared, among which are those applying general realcompactification methods and methods using embedding in complete uniform structures. The present note has the goal of giving an elementary self-contained proof of this theorem invoking only basic set theoretical properties of realcompactness.

Recall that a space is *realcompact* provided that each maximal centered family of zerosets with the countable intersection property has nonempty intersection.

If  $X$  is a space, then it is well-known (see [1] page 17) that each two disjoint zerosets of  $X$  have disjoint cozeroset neighborhoods. Moreover, each finite cover of  $X$  by cozerosets has a finite refinement consisting of zerosets. The proof of the last remark is similar to the proof of the statement that each finite open cover of a normal space has a finite closed refinement [2].

If  $X$  is a space and  $\alpha X$  an extension of  $X$  in which  $X$  is dense, then we shall say that  $\alpha X$  has property (Z) in case that for each countable collection of zerosets of  $X$  with empty intersection, the closures in  $\alpha X$  have empty intersection.

**LEMMA 1.** *Let  $\alpha X$  be an extension of  $X$  which contains  $X$  as a dense subspace and such that each continuous function on  $X$  has a continuous extension over  $\alpha X$ . Then  $\alpha X$  has property (Z).*

**PROOF.** If  $Z_1$  and  $Z_2$  are disjoint zerosets of  $X$ , then by the complete regularity of  $X$  there exists  $f \in C(X)$  satisfying  $f(Z_1) = 0$ ,  $f(Z_2) = 1$ . Let  $\bar{f}$  be the continuous extension of  $f$  over  $\alpha X$ . It follows that  $\bar{Z}_1^{\alpha X} \cap \bar{Z}_2^{\alpha X} \subset \bar{f}^{-1}(0) \cap \bar{f}^{-1}(1) = \emptyset$ . Moreover, if  $\{Z_1, \dots, Z_n\}$  is a finite collection of zerosets of  $X$  with empty

intersection, then according to the remark above there exists a finite collection of zerosets  $\{T_1, \dots, T_m\}$  which is a cover of  $X$  and which refines  $\{X \setminus Z_1, \dots, X \setminus Z_n\}$ . The fact that each two disjoint zerosets of  $X$  have disjoint closures in  $\alpha X$  implies that  $\{\overline{T_i^{\alpha X}} | i = 1, 2, \dots, m\}$  is a cover of  $\alpha X$  which refines

$$\{\alpha X \setminus \overline{Z_i^{\alpha X}} | i = 1, 2, \dots, n\}.$$

Hence

$$\cap \{\overline{Z_i^{\alpha X}} | i = 1, 2, \dots, n\} = \emptyset.$$

Now, let  $\{Z_i | i = 1, 2, \dots\}$  be a countable collection of zerosets of  $X$  with empty intersection. If there exists

$$p \in \cap \{\overline{Z_i^{\alpha X}} | i = 1, 2, \dots\},$$

then for  $i = 1, 2, \dots$  let  $f_i \in C(X)$  be such that  $0 \leq f_i \leq 1$  and  $Z_i = \{x \in X | f_i(x) = 0\}$ . The result proved in the last few lines above implies that an arbitrary (zeroset) neighborhood  $U$  of  $p$  in  $\alpha X$  intersects  $Z_1 \cap Z_2 \cap \dots \cap Z_k$  for each  $k$ , so the function  $f$  on  $X$  defined by

$$f(x) = \sum_{i=1}^{\infty} 2^{-i} f_i(x)$$

takes arbitrarily small values on  $U \cap X$ . It follows that the function  $1/f$  cannot be extended continuously over  $\alpha X$ , which contradicts our hypothesis.

**LEMMA 2.** *If  $X$  is a realcompact space and if  $\alpha X$  is an extension of  $X$  with property (Z), then  $\alpha X = X$ .*

**PROOF.** Denote the collection of zerosets of  $X$  by  $\mathcal{Z}$ . Assume that there exists  $p \in \alpha X \setminus X$ , and let  $\mathcal{Z}_1$  be the subcollection of  $\mathcal{Z}$  defined by  $\mathcal{Z}_1 = \{Z \in \mathcal{Z} | p \in \overline{Z^{\alpha X}}\}$ . Condition (Z) implies that  $\mathcal{Z}_1$  is a maximal centered family of zerosets of  $X$  with the countable intersection property; thus by realcompactness of  $X$  there exists  $q \in \cap \mathcal{Z}_1$ . Let  $G$  be a zeroset neighborhood of  $p$  in  $\alpha X$  which contains  $p$  and does not meet  $q$ . Then  $p \in \overline{G \cap X}^{\alpha X}$ , so  $G \cap X$  is a member of  $\mathcal{Z}_1$  which does not meet  $q$ . This is a contradiction.

We are now in a position to prove Hewitt-Shirota's theorem. We state it in the following way.

**THEOREM.** *Let  $X$  be a realcompact space. The mapping  $e : X \rightarrow \mathbf{R}^{C(X)}$  defined by  $e(x)_f = f(x)$  for  $f \in C(X)$  is a homeomorphism of  $X$  onto a closed subspace of  $\mathbf{R}^{C(X)}$ .*

PROOF. By the complete regularity of  $X$ ,  $e$  is a homeomorphism. By Lemma 1 the closure  $\overline{e(X)}$  of  $e(X)$  in  $\mathbb{R}^{C(X)}$  is an extension of  $X$  with property (Z) and by Lemma 2,  $\overline{e(X)} = e(X)$ . Thus  $e$  is a closed embedding.

#### REFERENCES

L. GILLMAN and M. JERISON

[1] Rings of continuous functions, van Nostrand, 1960.

P. ALEXANDROFF and H. HOPF

[2] Topologie, Berlin, 1935.

(Oblatum 18-9-68)