

stichting
mathematisch
centrum



AFDELING ZUIVERE WISKUNDE

ZW 19/74

JANUARY

A.E. BROUWER
TWO NUMBER THEORETIC SUMS

BIBLIOTHEEK MATHEMATISCH CENTRUM
AMSTERDAM

2e boerhaavestraat 49 amsterdam

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O), by the Municipality of Amsterdam, by the University of Amsterdam, by the Free University at Amsterdam, and by industries.

Let $g_k(s_k)$ denote the largest (smallest) prime divisor of the natural number $k \geq 2$ and let $g_1 = s_1 = 1$.

Recently J. van de Lune raised the problem of determining the asymptotic behaviour of the sums

$$\sum_{k \leq x} g_k \quad \text{and} \quad \sum_{k \leq x} s_k.$$

Van de Lune and E. Wattel showed that the upper and lower limits of

$$\frac{\log x}{x^2} \cdot \sum_{k \leq x} g_k$$

lie between $\frac{1}{2}$ and 1. In this report the following theorem will be proved.

THEOREM. (i)
$$\sum_{k \leq x} l_k = \frac{\pi^2}{12} \frac{x^2}{\log x} + O(x^2 \log^{-3/2} x \log \log x),$$

(ii)
$$\sum_{k \leq x} s_k = \frac{1}{2} \frac{x^2}{\log x} + O(x^2 \log^{-2} x).$$

PROOF.

(i) First $\sum_{k \leq x} l_k \leq \sum_{p \leq x} \left[\frac{x}{p} \right] p$, because the last sum contains for each k all prime factors of k instead of only the largest one. Also

$\sum_{k \leq x} l_k \geq \sum_{\sqrt{x} < p \leq x} \left[\frac{x}{p} \right] p$ since $p > \sqrt{x}$, $k \leq x$, $p | k$ imply $p = l_k$. Clearly

$\sum_{p \leq \sqrt{x}} \left[\frac{x}{p} \right] p \leq x \sqrt{x}$, and therefore

(1)
$$\sum_{k \leq x} l_k = \sum_{p \leq x} \left[\frac{x}{p} \right] p + O(x^{3/2}).$$

Let f be an increasing function such that:

(a)
$$\lim_{x \rightarrow \infty} f(x) = \infty$$

(b)
$$\exists \epsilon > 0: f(x) = O(x^{1-\epsilon}).$$

Then

$$\sum_{p \leq x/f(x)} \left[\frac{x}{p} \right] p \leq x \pi\left(\frac{x}{f(x)}\right),$$

and since $\pi(x) = O\left(\frac{x}{\log x}\right)$ it follows that

$$\begin{aligned}
\sum_{p \leq x} \left[\frac{x}{p} \right] p &= \sum_{x/f(x) \leq p \leq x} \left[\frac{x}{p} \right] p + o\left(\frac{x^2}{\log x \cdot f(x)}\right) = \\
&= \sum_{n=1}^{f(x)} \sum_{p \leq \frac{x}{n}} p + o\left(\frac{x^2}{\log x \cdot f(x)}\right) \quad \text{see (ii)} \\
&= \sum_{n=1}^{f(x)} \left(\frac{\left(\frac{x}{n}\right)^2}{2 \log \frac{x}{n}} + o\left(\frac{x^2}{\log^2 x}\right) \right) + o\left(\frac{x^2}{\log x \cdot f(x)}\right) = \\
&= \frac{x^2}{2 \log x} \sum_{n=1}^{f(x)} \frac{1}{n^2} \left(1 - \frac{\log n}{\log x}\right)^{-1} + o\left(\frac{x^2 f(x)}{\log^2 x}\right) + o\left(\frac{x^2}{\log x \cdot f(x)}\right) = \\
&= \frac{x^2}{2 \log x} \left(\frac{\pi^2}{6} + o\left(\frac{1}{f(x)}\right) \right) \left(1 + o\left(\frac{f(x) \log f(x)}{\log x}\right)\right) + \\
&+ o\left(\frac{x^2 f(x)}{\log^2 x}\right) + o\left(\frac{x^2}{\log x \cdot f(x)}\right) = \\
&= \frac{\pi^2}{6} \cdot \frac{x^2}{2 \log x} + o\left(\frac{x^2 f(x) \log f(x)}{\log^2 x}\right) + o\left(\frac{x^2}{\log x \cdot f(x)}\right).
\end{aligned}$$

Now take $f(x) = \log^{\frac{1}{2}} x$. Then

$$\frac{x^2 f(x) \log f(x)}{\log^2 x} = \frac{x^2 \log \log x}{2 \log^{3/2} x} \quad \text{and} \quad \frac{x^2}{\log x \cdot f(x)} = \frac{x^2}{\log^{3/2} x}$$

which proves (i).

(ii) First $\sum_{k \leq x} s_k \geq \sum_{p \leq x} p$; also

$$\sum_{k \leq x} s_k \leq \sum_{p \leq x} p + \sum_{k \leq x} \sqrt{k} \leq \sum_{p \leq x} p + x^{3/2}.$$

Thus

$$(2) \quad \sum_{k \leq x} s_k = \sum_{p \leq x} p + o(x^{3/2}).$$

Now

$$\begin{aligned}\sum_{p \leq x} p &= \int_2^x x \, d\pi(x) = \int_2^x \left(\frac{x}{\log x} + o\left(\frac{x}{\log^2 x}\right) \right) dx = \\ &= \frac{x^2}{2 \log x} + o\left(\frac{x^2}{\log^2 x}\right)\end{aligned}$$

which proves (ii).