

STICHTING  
MATHEMATISCH CENTRUM

2e BOERHAAVESTRAAT 49  
AMSTERDAM

ZW 1963-1512

Commutative polynomial semigroups on a segment

P.C. Baayen and Z. Hedrlín

Reprinted from

Commentationes Mathematicae

Universitatis Carolinae, 4(1963), p 173-179



1964

P.C. BAAYEN  
Z. HEDRLIN



4, 4 (1963)

REDACTORES

Z. FROLÍK

Z. HEDRLÍN

M. KATĚTOV

J. KOLOMÝ

V. KOŘÍNEK

F. NOŽIČKA

J. POTOČEK

A. ŠVEC

L. TRUKSA

COMMENTATIONES MATHEMATICAE  
UNIVERSITATIS CAROLINAE

Vol. 4 , fasc. 4

Fundavit

Academicus E. Č E C H

Redigit

A. Š V E C

litum cura Facultatis mathematico-physicalis  
Universitatis Carolinae Pragensis

Pragae mense decembri 1963

Matematický ústav university Karlovy,  
Sokolovská 83, Praha 8 Karlín,  
Československo (Czechoslovakia)

C O N S P E C T U S

I. MAREK, A note on K-positive operators	137
K. DRBOHLAV, Concerning representations of small categories	147
J. FIALÁ, Representation of generalized mea- sures by integrals	153
A. PULTR, A remark on common fixed sets of commuting mappings	157
Z. HEDRLÍN - A. PULTR, Remark on topological spaces with given semigroups	161
J. KOLOMÝ, Contribution to the solution of non-linear equations	165
P.C. BAAYEN - Z. HEDRLÍN, Commutative poly- nomial semigroups on a segment	173

AUTHORS' ADDRESSES :

P.C. Baayen, Math. Centrum, 2e Boerhaavestraat  
49, Amsterdam, Holland

I. Marek - K. Drbohlav - J. Fiala - A. Pultr -  
Z. Hedrlín and J. Kolomý:  
Matematicko-fyzikální fakulta  
Karlovy university, Praha Karlín  
Sokolovská 83

©

Matematický ústav KU, Sokolovská 83, Praha 8 Karlín

1. Introduction

A commutative semigroup of mappings of a set  $X$  is a family of mappings  $X \rightarrow X$  which is a commutative semigroup under composition of functions. A commutative polynomial semigroup of mappings of a subset  $X$  of the real line  $R$  (shortly: an  $X$ -cps) is a commutative semigroup of mappings  $X \rightarrow X$ , all elements of which are restrictions to  $X$  of (real) polynomials on  $R$ . Such a semigroup  $S$  is called maximal if every continuous map  $g : X \rightarrow X$  which commutes with all  $f \in S$  itself belongs to  $S$ , and entire if it contains (restrictions to  $X$  of) polynomials of every non-negative degree.

If  $S_1$  is a semigroup of continuous maps  $X_1 \rightarrow X_1$  ( $i = 1, 2$ ), and if  $\tau$  is a homeomorphism of  $X_1$  onto  $X_2$  such that  $S_2 = \{\tau \circ f \circ \tau^{-1} \mid f \in S_1\}$ , then  $S_1$  and  $S_2$  are called equivalent (by means of  $\tau$ ). In that case the transformation  $f \rightarrow \tau \circ f \circ \tau^{-1}$  is an isomorphism of the abstract semigroup  $S_1$  onto the abstract semigroup  $S_2$ .

In this note we determine, up to equivalence, all entire  $I$ -cps, where  $I$  is the closed unit segment  $[0, 1]$ . Moreover, we establish which of these  $I$ -cps are maximal and which not. We denote by  $J$  the segment  $[-1, 1]$ .

2. Commutative polynomial semigroups of mappings  $R \rightarrow R$  and  $J \rightarrow J$ .

It follows from results of J.F. Ritt [7, 8] and of H.D.

Block and H.P. Thielman [5] that every entire R-cps is equivalent by means of a linear transformation to one of the following three semigroups of polynomials:

(i) the semigroup  $P$ , consisting of the maps

$P_0, P_1, P_2, \dots$  with

$$P_n(x) = x^n;$$

(ii) the semigroup  $P^*$ , consisting of all  $P_n$ ,  $n \geq 1$

and the map  $P_0^*$  such that

$$P_0^*(x) = 0 \text{ for all } x;$$

(iii) the semigroup  $T$  of all Chebyshev polynomials

$T_0, T_1, T_2, \dots$ , where

$$T_n(x) = \cos(n \cdot \arccos x).$$

The first two semigroups are not maximal; e.g. consider  $x^{\frac{2}{3}}$ .

**Lemma 1.** There exists a unique maximal commutative semigroup  $\bar{P}$  ( $\bar{P}^*$ ) of continuous maps  $J \rightarrow J$  containing  $P|J$  ( $P^*|J$ , respectively). The semigroup  $\bar{P}$  ( $\bar{P}^*$ ) consists of the following maps: all maps  $x \rightarrow |x|^\epsilon$ ,  $\epsilon > 0$  a real number; all maps  $x \rightarrow |x|^\epsilon \cdot \text{sign } x$ ,  $\epsilon > 0$  a real number; and all maps in  $P$  (in  $P^*$ , respectively).

**Proof.** It is immediately verified that  $\bar{P}$  and  $P^*$  are commutative semigroups. In order to show their maximality, and the fact that they are the only maximal semigroups containing  $\bar{P}$  or  $\bar{P}^*$ , we proceed as follows.

Let  $f$  be any continuous map  $R \rightarrow R$  commuting with all maps in  $P$  or in  $P^*$ . Take any  $a$  with  $0 < a < 1$  and let  $f(a) = \alpha$ . As  $\alpha = P_2 f(\sqrt{a})$ ,  $\alpha \geq 0$  if  $\alpha = 0$ , it follows that  $f(a^r) = \alpha^r = 0$  for all rational  $r$ , because  $f \circ P_n = P_n \circ f$  for all natural  $n$ . Hence  $f(x) = 0$  for  $x \geq 0$ ; if  $x \leq 0$ ,  $P_2 f(x) = f(x^2) = 0$  implies again  $f(x) = 0$ . Thus  $f$  is identically zero.

Assume  $\alpha > 0$  and let  $\epsilon \in \mathbb{R}$  with  $a^\epsilon = \alpha$ . Then as  $f$  and  $P_n$  commute,  $f(a^r) = a^{r\epsilon}$  for all rational  $r$ ; hence  $f(x) = x^\epsilon$  for  $x \geq 0$ . If  $x < 0$ , then  $P_2 f(x) = f P_2(x) = (x^2)^\epsilon$ , hence  $f(x) = \pm |x|^\epsilon$ . As  $f$  is continuous, the lemma follows.

The situation is different for the semigroup  $T$ : this semigroup is maximal. In order to show this, we consider the following mappings of the unit interval  $I$  into itself, first introduced in [2]:

$$t_0(x) = 0 \text{ for all } x;$$

and, if  $n \geq 1$ :

$$\left\{ \begin{array}{l} t_n\left(\frac{2k}{n}\right) = 0, \quad t_n\left(\frac{2k+1}{n}\right) = 1 \quad (k = 0, 1, 2, \dots, \lfloor \frac{n}{2} \rfloor); \\ t_n \Big| \left[ \frac{k}{n}, \frac{k+1}{n} \right] \text{ is linear} \quad (k = 0, 1, 2, \dots, n-1). \end{array} \right.$$

These so-called multihats are easily seen to constitute a commutative semigroup  $M$ ; in fact,  $t_n \circ t_m = t_{n+m}$ . In [2] P.C. Baayen, W. Kuyk and M.A. Maurice proved much more: the semigroup of all  $t_n$ ,  $n = 0, 1, 2, \dots$ , is a maximal commutative semigroup of continuous maps  $I \rightarrow I$ .

Lemma 2. The semigroup  $M$  is equivalent to the semigroup  $T'$  of all Chebyshev polynomials  $T_n$ , restricted to the segment  $J$ , by means of the homeomorphism  $\tau: [0, 1] \rightarrow [-1, 1]$  such that

$$\tau x = \cos \pi x.$$

Proof: immediate.

Hence we have shown:

Lemma 3. The  $J$ -cps  $T$  is maximal.

This strengthens considerably a result of G. Baxter and J.T.

Joichi [3], who showed that  $T$  cannot be embedded in a 1-parameter semigroup of commuting functions.



We conclude this section with a triviality.

Lemma 4. Let  $Q_1, Q_2$  be polynomials commuting on some non-degenerate segment. Then  $Q_1$  and  $Q_2$  commute everywhere on  $R$ .

3. Commutative polynomial semigroups of mappings  $I \rightarrow I$

It follows from the results of section 2 that every entire  $I$ -cps is equivalent by means of a linear transformation to a semigroup  $S|A$ , where  $S$  is one of the  $R$ -cps  $T, P, P^*$  and  $A$  is a closed segment  $[a, b]$ ,  $a < b$ , that is invariant under  $S$ .

The only non-degenerate segment mapped into itself by  $T$  is  $[-1, +1]$ . The only non-trivial segments mapped into themselves by  $P$  are the segments  $[-a, 1]$ , with  $0 \leq a \leq 1$ ; we write  $P(a)$  for the  $[-a, 1]$ -cps of all  $P_n|[-a, 1]$ ,  $n = 0, 1, 2, \dots$ . The only non-trivial segments invariant under  $P^*$  are the segments  $[-a, b]$ , with  $0 \leq a \leq 1$ ,  $a^2 \leq b \leq 1$ ,  $b \neq 0$ ; we write  $P^*(a, b)$  for the  $[-a, b]$ -cps of all  $P_n|[-a, b]$ ,  $n \geq 1$  together with  $P_0|[-a, b]$ .

Lemma 5. Each of the semigroups  $P(a)$ ,  $0 \leq a \leq 1$ , is not maximal, and is contained in a unique maximal  $[-a, 1]$ -semigroup  $\overline{P(a)}$ . Similarly each  $P^*(a, b)$  is contained in a unique maximal  $[-a, b]$ -semigroup  $\overline{P^*(a, b)}$ .

Proof. In the same way as in the proof of Lemma 1 one shows that  $\overline{P(a)} = \overline{P} \parallel [-a, 1]$  is the unique maximal commutative semigroup of continuous maps  $[-a, 1] \rightarrow [-a, 1]$  containing  $P(a)$ . Similarly  $\overline{P^*(a, b)} = \overline{P^*} \parallel [-a, b]$ .

Remark: If  $S$  is a semigroup of mappings of a set  $X$  into itself, and if  $A \subset X$ , then  $S \parallel A$  denotes the semigroups of mappings of  $A$  into itself, consisting of all mappings  $f|A$  such that  $f \in S$  and  $f(A) \subset A$  (cf. [6]).

Theorem 1. There are two maximal entire I-cps; they are both equivalent to  $T'$  (or to  $M$ ).

Proof. Every maximal entire I-cps must be equivalent by means of a linear map to  $T' = T|[-1, +1]$ . There exist two linear maps of  $[-1, +1]$  onto  $I = [0, 1]$ .

Lemma 6. If  $0 < a, b < 1$ , then  $P(a)$  and  $P(b)$  are equivalent by means of the homeomorphism  $\tau$ ,

$$\tau(x) = \text{sign} x \cdot |x|^\varepsilon,$$

where  $\varepsilon = \frac{\log b}{\log a}$ .

Lemma 7. Let  $0 \leq a_i \leq 1$ ,  $a_i^2 \leq b_i \leq 1$ ,  $b_i \neq 0$  ( $i = 1, 2$ ). The semigroups  $P^*(a_1, b_1)$  and  $P^*(a_2, b_2)$  are equivalent if and only if there exists a real number  $\varepsilon \neq 0$  such that  $a_2 = a_1^\varepsilon$ ,  $b_2 = b_1^\varepsilon$ .

Proof. Suppose  $P^*(a_1, b_1)$  and  $P^*(a_2, b_2)$  are equivalent by means of  $\tau$ . Then we have, for arbitrary  $x \in [-a_1, b_1]$  and for arbitrary integers  $n \geq 1$ , that  $P_n(x) = (\tau^{-1} \circ P_n \circ \tau)(x)$ ; i.e.  $(\tau \circ P_n)(x) = (P_n \circ \tau)(x)$ . It follows (cf. lemma 1) that either  $\tau$  is of the form:  $\tau(x) = |x|^\varepsilon$ , for all  $x \in [-a_1, b_1]$ , where  $\varepsilon$  is some real number - as  $\tau$  is a homeomorphism this is only possible if  $a_1 = 0$  - or  $\tau$  is of the form:  $\tau(x) = |x|^\varepsilon \cdot \text{sign} x$ . As clearly we must have:  $\tau(a_1) = a_2$ ,  $\tau(b_1) = b_2$ , the assertion follows.

The next lemma is easily proved:

Lemma 8. No semigroup  $P(a)$  is equivalent to a semigroup  $P^*(b, c)$ .

Consequently we have:

Theorem 2. There are infinitely many non-equivalent non-maximal entire I-cps. Each of them is equivalent to one of the following semigroups, which are all mutually inequivalent:  $P(0)$ ,

$P(\frac{1}{2})$ ,  $P(1)$ ;  $P^*(a, 1)$ ,  $0 \leq a \leq 1$ ;  $P^*(a, \frac{1}{4})$ ,  $0 \leq a \leq \frac{1}{2}$ .

Theorem 3. Every entire I-cps is contained in a unique maximal commutative semigroup of continuous maps  $I \rightarrow I$ . Two entire I-cps are equivalent if and only if the maximal commutative semigroups in which they are contained are equivalent.

4. Remark on mappings commuting with  $T_n$  or  $P_n$ ,  $n \geq 2$ .

It was shown by P.C. Baayen and W. Kuyk in [1] that every open map of  $I$  into itself that commutes with  $t_2$  is itself a multihat  $t_n$ . From this it follows almost at once that every continuous map commuting with  $t_2$  is either a  $t_n$  or is everywhere oscillating (nowhere monotone).

This result has been improved very much by G. Baxter and J.T. Joichi [4], who showed the following theorem:

If a continuous map  $f: I \rightarrow I$  commutes with some multihat  $t_n$ ,  $n \geq 2$ , it is itself either a hat-function or a constant map.

Now we saw in section 2 that the semigroup  $M$  of all hats  $t_n$  is equivalent to the semigroup  $T'$  of all Chebyshev polynomials on  $[-1, +1]$ .

Hence we conclude:

Theorem 4. Every non-constant continuous map of  $[-1, +1]$  into itself that commutes with a Chebyshev polynomial  $T_n$  with  $n \geq 2$ , is itself a Chebyshev polynomial.

For the maps  $P_n$ ,  $n \geq 2$ , the situation is completely different. Consider e.g. continuous maps of  $[0, 1]$  into itself which commute with  $P_2$  on that interval.

There exist multitudes of such functions. For let  $0 < a < 1$ , and let  $f_0$  be any continuous function of  $[a^2, a]$  into

$(0, 1)$  such that  $[f_0(a)]^2 = f_0(a^2)$ . If we define:  
 $f(0) = 0$ ,  $f(1) = 1$ ,  $f(x) = [f_0(x^{2^{-n}})]^{2^n}$  if  $x \in [a^{2^{n+1}}, a^{2^n}]$

( $n$  integer),  $f$  will be a continuous map  $I \rightarrow I$  commuting with  $P_2$ .

R e f e r e n c e s

- [1] P.C. BAAAYEN and W.KUYK, Mappings commuting with the hat. Report ZW 1963-007, Mathematical Centre, Amsterdam, 1963.
- [2] P.C. BAAAYEN, W. KUYK, and M.A. MAURICE, On the orbits of the hat-function, and on countable maximal commutative semigroups of continuous mappings of the unit interval into itself, Report ZW 1962-018, Mathematical Centre, Amsterdam 1962.
- [3] G. BAXTER and J.T. JOICHI, On permutations induced by functions, and an embedding question, Submitted to Math. Scand.
- [4] G. BAXTER and J.T. JOICHI, On functions that commute with full functions, Mimeographed report (preprint). University of Minnesota, 1963.
- [5] H.D. BLOCK and H.P. THIELMAN, Commutative polynomials, Quart. J. Math. Oxford (2), 2 (1951), 241-243.
- [6] Z. HEDRLÍN, On commutativity of transformations, Report ZW 1962-015, Mathematical Centre, Amsterdam, 1962.
- [7] J.F. RITT, Prime and composite polynomials, Trans. Amer. Math. Soc. 23(1922), 51-66.
- [8] J.F. RITT, Permutable rational functions, Trans Amer. Math. Soc. 25(1923), 399-448.

CMUC, Commentationes Mathematicae Universitatis Carolinae is published by the Mathematical Institute of the Caroline University, Prague. It contains preliminary reports and scientific papers concerning all branches of mathematics. Reports from mathematical institutions in Czechoslovakia are also accepted for publication. Papers are published in either English, French, German or Russian.

Contributions are accepted from all authors. The Editorial Council decides on publication. Manuscripts must be submitted ready for publication also from the language point of view.

The author receives 50 copies of his paper.

CMUC is not for sale but intended to be exchanged for other publications. It is regularly sent to all institutions with which the Mathematical Institute and Departments of the Caroline University are in contact.

Журнал "Commentationes Mathematicae Universitatis Carolinae" (сокращено CMUC) издаётся Математическим институтом Карлова университета в Праге. В нём печатаются предварительные сообщения и научные работы по всем областям математики, а также информации из различных математических научных учреждений в ЧССР. Статьи принимаются как от чехословацких так и от заграничных авторов и печатаются на одном из следующих языков: английский, немецкий, русский и французский.

О публикации присланных статей решает редакционная коллегия. Рукописи должны прислаться вполне оформленными (также, что касается их языка) и подготовленными для передачи типографии.

Автор получает 50 оттисков своей работы.

CMUC не продается и служит для обмена за другие публикации; журнал посылается всем научным учреждениям, поддерживающим связь с математическим институтом и кафедрами Карлова университета.

- I. M A R E K, A note on K-positive operator
- K. D R B O H L A V, Concerning representations of small categories
- J. F I A L A, Representation of generalized measures by integrals
- A. P U L T R , A remark on common fixed sets of commuting mappings
- Z. H E D R L Í N - A. P U L T R , Remark on topological spaces with given semigroups
- J. K O L O M Ý, Contribution to the solution of non-linear equations
- P. C. B A A Y E N - Z. H E D R L Í N, Commutative polynomial semigroups on a segment