

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O), by the Municipality of Amsterdam, by the University of Amsterdam, by the Free University at Amsterdam, and by industries.

Some theorems concerning the number theoretical functions $\omega(n)$ and $\Omega(n)$ *)

J. van de Lune

Abstract

The functions ω and Ω are defined as follows: $\omega(1) = \Omega(1) = 0$ and if $n = p_1^{e_1} p_2^{e_2} \dots p_r^{e_r}$ is the canonical factorization of the natural number n , then $\omega(n) = r$ and $\Omega(n) = e_1 + e_2 + \dots + e_r$. It is known that $\sum_{n \leq x} (-1)^{\Omega(n)} = o(x)$, ($x \rightarrow \infty$). There seems to be no corresponding result in the literature for $\omega(n)$. In this report it is shown that $\sum_{n \leq x} (-1)^{\omega(n)} = o(x)$, ($x \rightarrow \infty$). Furthermore, it is shown that the series $\sum_{n=1}^{\infty} (-1)^{\omega(n)} / n$ converges to zero. Finally, the remarkable duality relation

$$1 = \sum_{d/n} z^{\omega(d)} (1-z)^{\Omega(\frac{n}{d})} = \sum_{d/r} z^{\Omega(d)} (1-z)^{\omega(\frac{n}{d})}$$

and some of its consequences are discussed.

*) This paper is not for review; it is meant for publication in a journal.

Introduction. As usual, let $\omega(n)$ denote the number of distinct prime divisors and $\Omega(n)$ the total number of prime divisors of the positive integer n . That is, $\omega(1) = \Omega(1) = 0$ and if

$$n = p_1^{e_1} p_2^{e_2} \cdots p_r^{e_r}$$

is the canonical factorization of n , then $\omega(n) = r$ and $\Omega(n) = e_1 + e_2 + \cdots + e_r$.

It is known [1, p. 123], [4, II, p. 617], [5, p. 74] that

$$\sum_{n \leq x} (-1)^{\Omega(n)} = o(x), \quad (x \rightarrow \infty).$$

There seems to be no corresponding result in the literature for $\omega(n)$. In this report we will prove that

$$S(x) \stackrel{\text{def}}{=} \sum_{n \leq x} (-1)^{\omega(n)} = o(x), \quad (x \rightarrow \infty).$$

We will also establish the convergence of

$$\sum_{n=1}^{\infty} \frac{(-1)^{\omega(n)}}{n}$$

and show that the sum of this series is zero. For the corresponding result for $\Omega(n)$ see [4, pp. 617-621]. The above results can be sharpened considerably, but we will not take the effort here to do so.

Finally we will prove a remarkable duality relation between ω and Ω and discuss some of its consequences.

1. *Proposition 1.1.* The function

$$\phi(s) \stackrel{\text{def}}{=} \sum_{n=1}^{\infty} \frac{(-1)^{\omega(n)}}{n^s},$$

2

which is obviously analytic for $\text{Re } s = \sigma > 1$, has an analytic continuation up to $\sigma \geq 1$.

Proof. From the definition of $\omega(n)$ it follows immediately that for $\sigma > 1$

$$\phi(s) = \sum_{n=1}^{\infty} \frac{(-1)^{\omega(n)}}{n^s} = \prod_p \left(1 - \frac{1}{p^s} - \frac{1}{p^{2s}} - \dots\right), \quad (p \text{ prime}).$$

Thus

$$\begin{aligned} \zeta(s)\phi(s) &= \prod_p \frac{1}{1 - \frac{1}{p^s}} \cdot \prod_p \left(1 - \frac{1}{p^{s-1}}\right) = \\ &= \prod_p \frac{p^{2s} - 2p^s}{(p^s - 1)^2} = \prod_p \left(1 - \frac{1}{(p^s - 1)^2}\right) \stackrel{\text{def}}{=} P(s). \end{aligned}$$

Since $p^s - 1$ has all its zeros on the imaginary axis, it follows that $P(s)$ is analytic for $\sigma > \frac{1}{2}$. Furthermore, it is well-known that $\frac{1}{\zeta(s)}$ is analytic for $\sigma \geq 1$ and it follows that

$$\phi(s) = \frac{1}{\zeta(s)} \cdot P(s)$$

is also analytic for $\sigma \geq 1$. This completes the proof. \square

THEOREM 1.1. $S(x) \stackrel{\text{def}}{=} \sum_{n \leq x} (-1)^{\omega(n)} = o(x), \quad (x \rightarrow \infty).$

Before proving this theorem we state the following special version of the well known [2, p. 124] WIENER-IKEHARA Tauberian theorem: Let $F(x)$ be non-negative and non-decreasing for $x \geq 0$.

Let

$$f(s) = \int_0^{\infty} e^{-sx} F(x) dx$$

converge for $\sigma > 1$. If $f(s)$ is analytic for $\sigma \geq 1$, except for a simple pole at $s = 1$ with residue A , then

$$\lim_{x \rightarrow \infty} \frac{F(x)}{e^x} = A.$$

Proof of Theorem 1.1. Note that

$$1 + (-1)^{\omega(n)} \geq 0 \text{ for } n = 1, 2, 3, \dots .$$

Hence $F(x) \stackrel{\text{def}}{=} [x] + S(x)$ is a non-negative, non-decreasing function not exceeding $2x$. It follows that for $\sigma > 1$,

$$\begin{aligned} \zeta(s) + \phi(s) &= \sum_{n=1}^{\infty} \frac{1 + (-1)^{\omega(n)}}{n^s} = \\ &= \sum_{n=1}^{\infty} \frac{F(n) - F(n-1)}{n^s} = \int_{1-0}^{\infty} x^{-s} dF(x) = \\ &= x^{-s} F(x) \Big|_{1-0}^{\infty} + s \int_1^{\infty} F(x) x^{-s-1} dx = \\ &= s \int_1^{\infty} F(x) x^{-s-1} dx = s \int_0^{\infty} e^{-sx} F(e^x) dx. \end{aligned}$$

Thus

$$\int_0^{\infty} e^{-sx} F(e^x) dx = \frac{\zeta(s) + \phi(s)}{s}, \quad (\sigma > 1).$$

Recall that $\zeta(s)$ is a meromorphic function with only one simple pole at $s = 1$ with residue $A = 1$. From Proposition 1.1 it now follows that the function

$$\frac{\zeta(s) + \phi(s)}{s} - \frac{1}{s-1}$$

is analytic for $\sigma \geq 1$. Now, applying the WIENER-IKEHARA theorem, we obtain

$$A = 1 = \lim_{x \rightarrow \infty} \frac{F(e^x)}{e^x} = \lim_{x \rightarrow \infty} \frac{F(x)}{x} =$$

$$= \lim_{x \rightarrow \infty} \frac{[x] + S(x)}{x} = 1 + \lim_{x \rightarrow \infty} \frac{S(x)}{x}.$$

Thus,

$$\lim_{x \rightarrow \infty} \frac{S(x)}{x} = 0 \text{ or } S(x) = o(x)$$

and this proves the theorem. \square

THEOREM 1.2. The series

$$\sum_{n=1}^{\infty} \frac{(-1)^{\omega(n)}}{n}$$

converges and its sum is zero.

Before proving this theorem we state the following well known result [1, p. 124].

Let

$$g(s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s}$$

be absolutely convergent for $\sigma > 0$ and suppose that $g(s)$ is analytic for $\sigma \geq 0$.

If in addition, $a(n) = o(1)$, then

$$\sum_{n=1}^{\infty} a(n)$$

converges and its sum is $g(0)$.

Proof of Theorem 1.2. Define

$$a(n) = \frac{(-1)^{\omega(n)}}{n}.$$

Then we have

$$g(s) = \sum_{n=1}^{\infty} \frac{(-1)^{\omega(n)}}{n^{s+1}} = \phi(s+1).$$

Since $\phi(s)$ is analytic for $\sigma \geq 1$, we see that $g(s)$ is analytic for $\sigma \geq 0$, and it follows that the series appearing in theorem 1.2 converges.

Moreover,

$$g(0) = \phi(1) = \lim_{s \rightarrow 1} \frac{P(s)}{\zeta(s)} = 0$$

and the theorem is proved. \square

Remark. Actually, $\phi(s)$ has a zero of order 2 at $s = 1$ because,

$$\begin{aligned} \lim_{s \rightarrow 1} \frac{\phi(s)}{(s-1)^2} &= \lim_{s \rightarrow 1} \left\{ \frac{1}{(s-1)\zeta(s)} \cdot \frac{1 - \frac{1}{(2^s-1)^2}}{(s-1)} \cdot \prod_{p>3} \left(1 - \frac{1}{(p^s-1)^2}\right) \right\} = \\ &= 4 \log 2 \cdot \prod_{p>3} \left(1 - \frac{1}{(p-1)^2}\right) \neq 0. \end{aligned}$$

Note by contrast that the corresponding function

$$\sum_{n=1}^{\infty} \frac{(-1)^{\Omega(n)}}{n^s} = \frac{\zeta(2s)}{\zeta(s)}$$

only has a simple zero at $s = 1$.

2. From their definitions it is hardly to be expected that there is much of a relation between $\omega(n)$ and $\Omega(n)$. However, we will exhibit below a remarkable duality relation between the two functions and discuss some of its consequences.

We first note that [3, p. 355]

$$\omega(n) = O\left(\frac{\log n}{\log \log n}\right),$$

from which it is easily seen that if $\operatorname{Re} s = \sigma > 1$ then

$$\sum_{n=1}^{\infty} \frac{z^{\omega(n)}}{n^s}$$

is an entire function of z . Also, for $\sigma > 1$ and $|z| < 2^\sigma$, it is clear that

$$\sum_{n=1}^{\infty} \frac{z^{\Omega(n)}}{n^s} = \prod_p \left(1 + \frac{z}{p^s} + \frac{z^2}{p^{2s}} + \dots\right) = \prod_p \frac{1}{1 - \frac{z}{p^s}}$$

is an analytic function of z .

THEOREM 2.1. If $|z| < 2^\sigma$ and $\sigma > 1$ then

$$(2.1) \quad \left(\sum_{n=1}^{\infty} \frac{z^{\omega(n)}}{n^s}\right) \cdot \left(\sum_{n=1}^{\infty} \frac{(1-z)^{\Omega(n)}}{n^s}\right) = \zeta(s).$$

Proof.

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{z^{\omega(n)}}{n^s} &= \prod_p \left(1 + \frac{z}{p^s} + \frac{z}{p^{2s}} + \frac{z}{p^{3s}} + \dots\right) = \\ &= \prod_p \left(1 + z \frac{1}{1 - \frac{1}{p^s}}\right) = \prod_p \frac{p^s + z - 1}{p^s - 1} \end{aligned}$$

and

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{(1-z)^{\Omega(n)}}{n^s} &= \prod_p \left(1 + \frac{1-z}{p^s} + \frac{(1-z)^2}{p^{2s}} + \frac{(1-z)^3}{p^{3s}} + \dots \right) = \\ &= \prod_p \left(1 + \frac{\frac{1-z}{p^s}}{1 - \frac{1-z}{p^s}} \right) = \prod_p \frac{p^s}{p^s + z - 1}. \end{aligned}$$

Hence

$$\begin{aligned} \left(\sum_{n=1}^{\infty} \frac{z^{\omega(n)}}{n^s} \right) \cdot \left(\sum_{n=1}^{\infty} \frac{(1-z)^{\Omega(n)}}{n^s} \right) &= \\ &= \prod_p \frac{p^s + z - 1}{p^s - 1} \prod_p \frac{p^s}{p^s + z - 1} = \prod_p \frac{1}{1 - \frac{1}{p^s}} = \zeta(s) \end{aligned}$$

and the theorem follows easily. \square

Performing Dirichlet multiplication in (2.1), equating coefficients and changing z into $1 - z$ we obtain:

$$\sum_{d|n} z^{\omega(d)} (1-z)^{\Omega\left(\frac{n}{d}\right)} = \sum_{d|n} z^{\Omega(d)} (1-z)^{\omega\left(\frac{n}{d}\right)} = 1,$$

a remarkable duality relation between ω and Ω .

We now study the analytic continuation of

$$g_z(s) \stackrel{\text{def}}{=} \sum_{n=1}^{\infty} \frac{z^{\omega(n)}}{n^s}, \quad (z \text{ fixed, } \sigma > 1)$$

as a function of s . We first have

Proposition 2.1. If $|1-z| \leq 2$ and $\sigma > 1$ then

$$(2.2) \quad \frac{g'_z(s)}{g_z(s)} = z \frac{\zeta'(s)}{\zeta(s)} + z(z-1) \sum_p \frac{\log p}{(p^s+z-1)(p^s-1)}$$

(where all derivatives are taken with respect to s).

Proof. From (2.1) we obtain

$$g_z(s) \cdot \prod_p \left(1 + \frac{1-z}{p^s} + \frac{(1-z)^2}{p^{2s}} + \dots\right) = \zeta(s),$$

which can be written as

$$g_z(s) = \zeta(s) \cdot \prod_p \frac{p^s + z - 1}{p^s}.$$

Taking logarithmic derivatives, we find

$$\begin{aligned} \frac{g'_z(s)}{g_z(s)} &= \frac{\zeta'(s)}{\zeta(s)} + \sum_p \left(\frac{p^s}{p^s + z - 1} - 1 \right) \log p = \\ &= \frac{\zeta'(s)}{\zeta(s)} + \sum_p \frac{1 - z}{p^s + z - 1} \log p = \\ &= \frac{\zeta'(s)}{\zeta(s)} + \sum_p \left(\frac{1 - z}{p^s + z - 1} - \frac{1-z}{p^s-1} \right) \log p + \sum_p \frac{1-z}{p^s-1} \log p = \\ &= z \cdot \frac{\zeta'(s)}{\zeta(s)} + z(z-1) \sum_p \frac{\log p}{(p^s+z-1)(p^s-1)}, \end{aligned}$$

which completes the proof. \square

If $|1-z| \leq \sqrt{2}$, then

$$R_z(s) \stackrel{\text{def}}{=} \sum_p \frac{\log p}{(p^{s+z}-1)(p^s-1)}$$

is regular for $\sigma > \frac{1}{2}$. It then follows that $\frac{g'_z(s)}{g_z(s)}$ is regular for $\sigma > \frac{1}{2}$ except at $s = 1$ and at the (possible) zeros of $\zeta(s)$ situated at the right of the line $\sigma = \frac{1}{2}$.

Integrating the formula


$$\frac{g'_z(s)}{g_z(s)} = z \frac{\zeta'(s)}{\zeta(s)} + z(z-1)R_z(s)$$

we get

$$g_z(s) = \zeta^z(s) \exp(P_z(s)),$$

where $P_z(s)$ is analytic for $\sigma > \frac{1}{2}$.

Thus $g_z(s)$ is analytic in the shaded region below, where the ρ 's stand for the (possible) zeros of $\zeta(s)$ which lie at the right of the line $\sigma = \frac{1}{2}$.



Hence, if (for example) z is irrational and $|1-z| \leq \sqrt{2}$, we find, surprisingly enough, regardless of which z is chosen subject to the above conditions, that the set of singularities of $g_z(s)$ in the halfplane $\sigma > \frac{1}{2}$ always consists of the same points, namely $s = 1$ and the zeros of $\zeta(s)$ lying in the halfplane $\sigma > \frac{1}{2}$. This seems to lend credence to the Riemann-hypothesis.

References

- [1] R. AYOUB, *An introduction to the analytic theory of numbers*, A.M.S., Providence, Rhode Island, 1963.
- [2] K. CHANDRASEKHARAN, *Introduction to analytic number theory*, Springer Verlag, 1968.
- [3] G.H. HARDY and E.M. WRIGHT, *Introduction to the theory of numbers*, 4th edition, Oxford, 1968.
- [4] E. LANDAU, *Handbuch der Lehre von der Verteilung der Primzahlen*, Chelsea, New York, 1953.
- [5] K. PRACHAR, *Primzahlverteilung*, Springer Verlag, 1957.

