

SMC WN30  
febr 70  
~~SECRET~~

Connected spaces in which all connected sets containing some fixed point are closed.

J.L. Hursch

A. Verbeek-Kroonenberg

Notations.

Let  $X$  be a connected  $T_1$ -space and  $x_0$  some fixed point of  $X$  such that any connected set containing  $x_0$  is closed.

The characters  $z, y, z, u, v, \dots$  denote points of  $X$ .

For a topological space  $Y$ , we write  $Y = A + B$  if  $Y$  is the topological sum of its subspaces  $A$  and  $B$ .

We will frequently apply the following two wellknown lemma's, most often with  $Y = \{z\}$  for some  $z \in Z$ :

Lemma 1. If  $Z$  and  $Y \subset Z$  are connected and  $Z \setminus Y = A + B$ , then  $Y \cup A$  (and  $Y \cup B$ ) is connected.

Lemma 2. If  $Z$  and  $Y \subset Z$  are connected and  $C$  is a component of  $Z \setminus Y$  then  $Z \setminus C$  is connected.

Let  $<$  be the relation (partial order) defined on  $X$  by:

$$\begin{aligned} x_0 < y & \text{ for all } y \in X \setminus \{x_0\} \\ x < y & \text{ if } x \text{ separates } x_0 \text{ and } y \end{aligned}$$

Then  $X$  and  $<$  have the following properties:

Proposition 1. The relation  $<$  is antisymmetric and transitive; i.e. is a partial order.

Proof. If  $x < y$  and  $y < x$  then there exist  $A, B, C, D \subset X$  such that  $X \setminus \{x\} = A + B$ ,  $X \setminus \{y\} = C + D$ ,  $x_0 \in A \cap B$ ,  $y \in B$  and  $x \in D$ . Now  $A \cup \{x\}$  is connected (lemma 1), contained in  $C + D$ , but meeting both  $C$  (in  $x_0$ ) and  $D$  (in  $x$ ). Contradiction.

If  $x < y$  and  $y < z$  then there exist  $A, B, C, D \subset X$  such that  $X \setminus \{x\} = A + B$ ,  $X \setminus \{y\} = C + D$ ,  $x_0 \in A \cap C$ ,  $y \in B$  and  $z \in D$ . Since  $D \cup \{y\}$  is connected (lemma 1) and intersects  $B$  (in  $y$ )  $D \cup \{y\} \subset B$ . Hence  $x < z$ . ■

Proposition 2. For each  $x \in X$   $\{y \mid y < x\}$  is linearly ordered (and wellordered by  $>$ , see 6)

Proof. Let  $y < x$ ,  $z < x$  but  $y \not\leq z$  and  $z \not\leq y$ . Then there exist  $A, B, C, D \subset X$  such that  $X \setminus \{y\} = A + B$ ,  $X \setminus \{z\} = C + D$ ,  $x_0 \in A \cap C$  and  $x \in B \cap D$ , but  $z \notin B$ ,  $y \notin D$  and so  $z \in A$  and  $y \in C$ . Since  $D \cup \{z\}$  is connected (lemma 1), and intersects  $A$  (in  $z$ ), but does not contain  $y$ ,  $D \cup \{z\} \subset A$ . This is contradictory to  $x \in D \setminus A$ . ■

Proposition 3. If  $x \in X$  and  $C$  is a component of  $X \setminus \{x\}$  which does not contain  $x_0$ , then  $C$  is open in  $X$ , and  $C^- = C \cup \{x\}$ . (If  $x_0 \in C$ , then  $C$  is closed in  $X$ ).

Proof.  $X \setminus C$  is connected (lemma 2), contains  $x_0$  and is hence closed. Now  $C$  cannot be closed in  $X$ , because  $X$  is connected. As  $C$  is closed in  $X \setminus \{x\}$ , only  $x$  can be another limitpoint of  $C$ . ■

Proposition 4. For any  $x \in X$   $\{y \mid x \leq y\}$  is the component of  $X \setminus \{x\}$  which contains  $x_0$ , and hence this set is closed. So its complement  $\{y \mid x \leq y\}$  is connected and open.

Proof. By definition of  $<$   $\{y \mid x \leq y\}$  is the quasicomponent of  $x_0$  in  $X \setminus \{x\}$ . If this set was not connected, then it would contain a component  $C$  of  $X \setminus \{x\}$  which does not contain  $x_0$ . But this  $C$  is open in  $X$  (by 3) and closed in  $X \setminus \{x\}$ . Thus  $\{y \mid x \leq y\}$  is not a quasicomponent.

The connectedness of  $\{y \mid x \leq y\}$  follows from lemma 2. ■

Proposition 5a. For each non empty linearly ordered  $A \subset X$  there is a (unique)  $x \in X$  such that  $x = \inf A$ .

5b. Each  $y \in X \setminus \{x_0\}$  has an immediate predecessor, which will be denoted by  $y'$ . For this point  $y'$ :

$$\{z \mid y \leq z\}^- = \{z \mid y \leq z\} \cup \{y'\}$$

Proof. (a) Let  $A^* = \{z \mid \exists a \in A \quad a < z\}$ . By 4 this set is open and hence not closed, as  $X$  is connected. Let  $x$  be a boundary point of  $A^*$ . At first we will show that  $x < a$  for all  $a \in A$  (or  $a \in A^*$ ). Since  $x \notin A^*$  we have  $a \not\leq x$  for all  $a \in A$ . Suppose  $x$  and some  $a \in A$  are not comparable. Then, by 2,  $x$  cannot be compared with any  $a \in A$ . But then again by 2,  $\{y \mid x \leq y\} \cap A^* = \emptyset$ . Since  $\{y \mid x \leq y\}$  is open (see 4),  $x \notin A^-$ .

Contradiction.

Secondly assume that for some  $y \quad x < y$  and  $y < a$  for all  $a \in A$ . For  $a \in A$  let  $C_a$  be the component of  $a$  in  $X \setminus \{y\}$ . Since, by 4,  $\{z \mid a \leq z\} \subset C_a$ , the family  $\{C_a \mid a \in A\}$  has no disjoint members. Hence it has a connected union. This means that for some component  $C$  of  $X \setminus \{y\}$   $A \subset C$ . By 3  $C^- = C \cup \{y\}$ , but  $C$  does not contain  $x$  since  $x < y$ . This contradicts  $x \in A^- \subset C^-$ .

So if  $A$  has no smallest element then  $x = \inf A$ .

(b) Let  $A = \{y\}$ ,  $y' = x =$  the boundary point of  $\{z \mid x \leq z\}$ . ■

For each ordertype  $\alpha$ , ordered by  $<$ , let  $\alpha^*$  denote the ordertype of  $\alpha$ , ordered by  $>$ . It follows immediately from 5a and 5b that for each  $x \in X$  the set  $\{y \mid y \leq x\}$  has ordertype  $\alpha^*$  for some ordinal  $\alpha$ . If  $A$  is a linearly ordered subset of  $X$ , and  $B$  is an infinite strictly increasing sequence, then by consequence  $B$  is cofinal with  $A$ . It follows from 4 and 5b that  $X$  cannot have maximal members.

Thus we proved:

Proposition 6. Let  $A$  be a linearly ordered subset of  $X$ , with ordertype  $\alpha$ . If  $A$  is bounded in  $X$  then  $\alpha = \beta^*$  for some ordinal  $\beta$ . If  $A$  is not bounded in  $X$  then  $\alpha = \sum_{n \in \mathbb{N}} \beta_n^*$ , for some suitable countable set of ordinals  $\{\beta_1, \beta_2, \dots\}$ .

We feel that the following facts deserve special attention

7. Any point of  $X \setminus \{x_0\}$  separates  $X$  in infinitely many components (as follows from 4).
8. Any connected space has a non-closed connected (proper) subset (else it were a space like  $X$ , but  $X \setminus \{x_0\}$  is non-closed and connected).
9. ZARANKIEWICZ [2]. If  $M$  is a connected separable metric space and  $D$  is the set of points  $x \in M$  for which  $M \setminus \{x\}$  has at least 3 components, then  $D$  is countable. On the other hand  $M$  has continuously many points.

Corollary.  $X$  is not separable metric.

Example of a Hausdorffspace  $X$ .

Let  $\mathbb{N}$  be the set of natural numbers, and  $P \subset \mathbb{N}$  the set of primenumbers.

Put  $X = \bigcup \{\mathbb{N}^n \mid n \in \mathbb{N}\} \cup \{0\}$ .

For  $x \in X$  we define  $\text{length } x = \begin{cases} 2 & \text{if } x = 0 \\ n+2 & \text{if } x \in \mathbb{N}^n. \end{cases}$

We define a partial order on  $X$  by taking  $0 \leq x$  for all  $x \in X$  and  $x \leq y$  if  $x$  is an initial sequent of  $y$ , i.e. if  $x \in \mathbb{N}^n$ ,  $y \in \mathbb{N}^m$ ,  $n \leq m$ , and there exist  $a_1, \dots, a_m \in \mathbb{N}$  such that  $x = (a_1, \dots, a_n)$ ,  $y = (a_1, \dots, a_m)$ . If  $x = (a_1, \dots, a_n)$  then let  $x' = (a_1, \dots, a_{n-1})$ .

As a subbase for the open sets we take all sets

- (i)  $\{z \mid x \leq z\}$  for each  $x \in X$
- (ii)  $\{z \mid x \leq z \wedge z \neq x'\}$  for each  $x \in X$
- (iii)  $\{z \mid \text{the only primes deviding length } y \text{ are } p_1, \dots, p_n\}$   
for each finite set of primes,  $p_1, \dots, p_n$ .

10.  $X$  is a Hausdorffspace

Let  $u, v \in X$ . We distinguish between

- (a)  $u < v$  and even  $u < v'$
- (b) neither  $u < v$  nor  $v < u$
- (c)  $u = v'$ .

- (a) In this case  $\{y \mid v \perp y \wedge y \neq v\}$  and  $\{z \mid v \leq z\}$  are disjoint neighbourhoods of  $u$  and  $v$ .
- (b) Now  $\{z \mid u \leq z\}$  and  $\{z \mid v \leq z\}$  are disjoint neighbourhoods of  $u$  and  $v$ , since  $u < z$  and  $v < z$  (for some  $z \in X$ ) would imply that  $u$  and  $v$  are comparable (by definition of  $X$ ).
- (c) Let  $p_1, \dots, p_n$  be the set of prime numbers which divide length  $u$ , and  $q_1, \dots, q_m$  idem for length  $v$ . Then  $\{p_1, \dots, p_n\} \cap \{q_1, \dots, q_m\} = \emptyset$  and so
- $$\{z \mid \forall p \in P \quad p \mid \text{length } z \Rightarrow p \in \{p_1, \dots, p_n\}\} \text{ and}$$
- $$\{z \mid \forall p \in P \quad p \mid \text{length } z \Rightarrow p \in \{q_1, \dots, q_m\}\}$$
- are disjoint neighbourhoods of  $u$  and  $v$ . ■

11. Any connected set  $C \subset X$  containing 0 is closed.

If  $u \in X \setminus C$  then we will show that  $C$  is disjoint from  $\{y \mid u \leq y\}$ ; hence  $C$  is closed. Suppose  $u \leq y$  for some  $y \in U$ , and  $u \in \mathbb{N}^n$ ,  $y = (a_1, \dots, a_m) \in \mathbb{N}^m$ . Now

$$C = (C \cap \{z \mid (a_1, \dots, a_{n+1}) \leq z\}) + (C \cap \{z \mid (a_1, \dots, a_{n+1}) \perp z \wedge z \neq u\}),$$

contradictory to the connectedness of  $C$ . ■

12.  $X$  is connected.

Lemma. For each  $u \in X \setminus \{0\}$  the points  $u$  and  $u'$  have no disjoint closed neighbourhoods.

Proof of the connectedness of  $X$ .

Suppose  $X = A + B$ ,  $0 \in A$ ,  $y \in B$  is such that length  $y$  is minimal. Then  $y' \in A$ , and  $A$  and  $B$  are disjoint closed neighbourhoods of  $y'$  and  $y$ . Contradiction. ■

Proof of the lemma. Let  $u = (a_1, \dots, a_1)$ .

For each point  $x \in X$  and each finite family  $\{x_1, \dots, x_n\}$  such that  $x_i \perp x$  and  $x' \neq x'_i$  we define the following neighbourhood of  $x$ :

$$U(x, \{x_1, \dots, x_n\}) = \{z \mid x \leq z\} \cap \bigcap_{p \in P} \{z \mid \forall p \in P \quad (p \mid \text{length } z) \Rightarrow (p \mid \text{length } x)\} \cap \bigcap_{i=1}^n \{z \mid x_i \perp z \wedge z \neq x'_i\}.$$

It should be clear that if the  $x_1, \dots, x_n$  vary we obtain a neighbourhoodbase of  $x$ . (We may also vary only over those  $x_i$  for which  $x < x_i'$ ).

For  $x = (a_1, \dots, a_n)$  we let  $\max x = \max\{a_1, \dots, a_n\}$ .

Now let  $U(u', \{x_1, \dots, x_n\})$  and  $U(u, \{x_{n+1}, \dots, x_m\})$  be two arbitrary basic neighbourhood of  $u'$  and  $u$ .

Put

$$\begin{aligned} N &= \max\{\max x_i \mid i=1, \dots, k\} + 1 \\ L &= (\text{length } x)(\text{length } x') - 2 \\ v &= (a_1, \dots, a_1, N, N, \dots, N) \in N^L. \end{aligned}$$

We will show that  $v \in U(u', \{x_1, \dots, x_n\}) \cap U(u, \{x_{n+1}, \dots, x_m\})$ .

Let  $U(v, \{x_{m+1}, \dots\})$  be an arbitrary neighbourhood of  $v$ . Put

$$N' = \max\{\max x_i \mid i=1, \dots, k, \dots, m, m+1, \dots\} + 1.$$

Let  $p, q \in \mathbb{P}$  be such that  $p \mid \text{length } u'$ ,  $q \mid \text{length } u$ , and choose  $r \in \mathbb{N}$  such that  $p^r > L$  and  $q^r > L$ . Then

$$\underbrace{(a_1, \dots, a_1, N, \dots, N, N', \dots, N')}_{\substack{L \text{ numbers} \\ p^r \text{ numbers}}} \in U(u', \{x_1, \dots, x_n\}) \cap U(v, \{x_{m+1}, \dots\})$$

$$\text{and } \underbrace{(a_1, \dots, a_1, N, \dots, N, N', \dots, N')}_{\substack{L \text{ numbers} \\ q^r \text{ numbers}}} \in U(u, \{x_{n+1}, \dots, x_m\}) \cap U(v, \{x_{m+1}, \dots\})$$

It is easily seen that if  $C \subset X$  is connected, then each  $x \in C$  disconnects  $C$ , except <sup>maybe</sup>  $\inf C$  (cf 11 and 4 and 5). In the terminology of [1]: each connected subset of  $X$  has at most one endpoint.

The points (1), (2), (3) are such that none of them separates the other 2. So this settles the problem mentioned in [1] p 24 remark 3.

## REFERENCES

- [1] H. Kok                    On conditions equivalent to the orderability  
of a connected space.  
Wiskundig Seminarium der Vrije Universiteit  
Rapport 6 November 1969.
- [2] C. Zarankiewicz  
and K. Kuratowski    Bull. Amer. Math. Soc. 33(1927) 571.

7601