

**stichting
mathematisch
centrum**



AFDELING TOEGEPASTE WISKUNDE
(DEPARTMENT OF APPLIED MATHEMATICS)

TW 174/78 JANUARI

O. DIERMANN, D. HILHORST & L.A. PELETIER

A SINGULAR BOUNDARY VALUE PROBLEM ARISING IN A
PRE-BREAKDOWN GAS DISCHARGE

Preprint

2e boerhaavestraat 49 amsterdam

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O).

A singular boundary value problem arising in a pre-breakdown gas discharge^{*)}

by

O. Diekmann, D. Hilhorst & L.A. Peletier

ABSTRACT

We consider the nonlinear two-point boundary value problem $\varepsilon xy'' + (g(x)-y)y' = 0$, $y(0) = 0$, $y(R) = k$, where g is a given concave function. We prove that the problem has a unique solution and we study the limiting behaviour of this solution as $R \rightarrow \infty$ and as $\varepsilon \downarrow 0$.

Furthermore, we show how a so-called pre-breakdown discharge in an ionized gas between two electrodes can be described by an equation of this form, and we interpret the results physically.

KEY WORDS & PHRASES: *singularly perturbed nonlinear two-point boundary value problem; pre-breakdown discharge in an ionized gas between two electrodes.*

^{*)} This report will be submitted for publication elsewhere.

1. INTRODUCTION

In this paper we study a model of the electric field which arises in the space between two electrodes. The equation in which we shall be engaged is

$$(1.1) \quad \varepsilon xy'' + (g(x)-y)y' = 0 \quad 0 < x < R,$$

where the given function g satisfies the following hypotheses

$$H_g: g \in C^2(\mathbb{R}_+; \mathbb{R}); \quad g(0) = 0; \quad g'(x) > 0 \quad \text{and} \quad g''(x) < 0 \quad \text{for all } x \geq 0.$$

We are interested in solutions of (1.1) which are subject to the boundary conditions

$$(1.2) \quad y(0) = 0,$$

$$(1.3) \quad y(R) = k.$$

We assume that the given numbers k and R satisfy $0 < k < g(\infty)$ and $R > x_0$, where x_0 is defined as the (unique) root of the equation $g(x) = k$.

In section 2 we shall derive some a priori estimates and we shall prove the existence of a solution of (1.1) - (1.3). In section 3 we prove that the solution is unique.

The main objective of this paper is the study of the dependence of the solution on the parameters ε and R . In section 3 we prove that the solution is a monotone function of ε and of R . From the physical point of view the interesting regions of the parameters are small ε and large R . In section 4 we analyse the limiting behaviour of the solution when R tends to infinity and ε is fixed. It turns out that the solution converges uniformly in x to a function \bar{y} which satisfies (1.1), (1.2) and the limiting form of (1.3), i.e. $\bar{y}(\infty) = k$, if and only if $\varepsilon \leq g(\infty) - k$. If, on the other hand, this inequality is violated, then the solution converges uniformly on compact subsets to a function \bar{y} which satisfies (1.1), (1.2) and $\bar{y}(\infty) = \max\{g(\infty) - \varepsilon, 0\}$. In particular this implies that \bar{y} is identically zero if $\varepsilon \geq g(\infty)$.

In section 5 we analyse the limiting behaviour of the solution when ε tends to zero and R is fixed. It turns out that the solution converges

uniformly in x to the function $\tilde{y}(x) = \min\{g(x), k\}$. Moreover we show that y' converges to \tilde{y}' uniformly for $x \in [0, x_0 - \delta] \cup [x_0 + \delta, R]$ for arbitrary $\delta > 0$. The behaviour of y' in the transition layer near x_0 will be discussed formally in section 6. Since the limits $\epsilon \downarrow 0$ and $R \rightarrow \infty$ (for $\epsilon \leq g(\infty) - k$) are interchangeable, the two separate limits give a complete picture of the limiting behaviour with respect to both parameters.

Finally, in section 7 we shall indicate how the problem arises in physics and we shall interpret the results physically.

Some partial results have been obtained for the case that g is neither increasing nor concave everywhere. This case will be studied in a forthcoming paper.

The problem (1.1) - (1.3) is, in many respects, similar to a singular boundary value problem of rotating fluids studied by HALLAM & LOPER [6]. We also draw attention to the recent paper of CLÉMENT & EMMERTH [2]. They study the limiting behaviour as $\epsilon \downarrow 0$ for more general problems by using completely different methods.

2. A PRIORI ESTIMATES AND THE EXISTENCE OF A SOLUTION

In this section we consider the problem (1.1) - (1.3) for fixed values of the parameters ϵ and R . By a solution we shall mean a function $y \in C^2([0, R]; \mathbb{R})$ which satisfies (1.1) - (1.3). We first derive some a priori estimates for a solution and its first two derivatives. Subsequently we prove that a solution actually exists by constructing an upper and lower solution and by verifying the appropriate Nagumo condition.

THEOREM 2.1. *Let y be a solution, then for all $x \in (0, R)$*

- (i) $0 < y(x) < \min\{g(x), k\}$;
- (ii) $0 < y'(x) < g'(0)$;
- (iii) $-\frac{(g'(0))^2}{\epsilon} < y''(x) < 0$.

PROOF. Let us first prove that $y'(x) > 0$ for all $x \in (0, R)$. Suppose that $y'(x_1) = 0$ for some $x_1 > 0$, then the standard uniqueness theorem for ordinary differential equations implies that $y(x) = y(x_1)$ for all x . Since this is not compatible with the two boundary conditions we conclude

that y' is sign-definite. Invoking the boundary conditions once more, we see that the sign has to be positive.

The positivity of y' implies that $0 < y(x) < k$ for $x \in (0, R)$. Next we shall prove that $y(x) < g(x)$. We begin by observing that this inequality holds for $x \geq x_0$. Suppose there is an interval $[x_1, x_2] \subset [0, x_0]$ such that $y - g$ is strictly positive in the interior of $[x_1, x_2]$ and $y(x_1) - g(x_1) = y(x_2) - g(x_2) = 0$. Then $y'(x_2) \leq g'(x_2) < g'(x_1) \leq y'(x_1)$. On the other hand the equation (1.1) implies that $y''(x) > 0$ for $x \in (x_1, x_2)$ and hence $y'(x_2) = y'(x_1) + \int_{x_1}^{x_2} y''(\xi) d\xi > y'(x_1)$. So our assumption must be false since it leads to a contradiction. Thus, $y(x) \leq g(x)$. Now, let us suppose that $y(x_1) = g(x_1)$ for some $x_1 > 0$, then necessarily $y'(x_1) = g'(x_1)$. However, because $y''(x_1) = 0$ (by (1.1)) and $g''(x_1) < 0$, this would imply that $y(x) > g(x)$ in a right-hand neighbourhood of x_1 , which is impossible. Hence the inequality is strict for $x \in (0, R]$, and this completes the proof of (i).

From (i), $y'(x) > 0$ and equation (1.1) we deduce that $y''(x) < 0$ for $x \in (0, R)$. Hence $y'(x) < y'(0) \leq g'(0)$ for $x \in (0, R)$ which completes the proof of (ii).

Finally, we note that H_g implies that $g(x) \leq g'(0)x$ and hence that $y''(x) = (\epsilon x)^{-1}(y(x) - g(x))y'(x) > -(\epsilon x)^{-1}g(x)g'(0) \geq -\epsilon^{-1}(g'(0))^2$. This proves property (iii). \square

THEOREM 2.2. *There exists a function $y \in C^2([0, R]; \mathbb{R})$ which satisfies (1.1) - (1.3).*

PROOF. We define two functions α and β by $\alpha(x) := 0$ and $\beta(x) := g(x)$ for $x \in [0, R]$. Moreover, we define a function f by $f(x, y, y') := (\epsilon x)^{-1}(y - g(x))y'$. Then $\alpha''(x) = 0 \geq 0 = f(x, \alpha(x), \alpha'(x))$ and $\beta''(x) = g''(x) < 0 = f(x, \beta(x), \beta'(x))$ for $x \in (0, R)$. Hence α and β are, respectively, a lower and an upper solution of (1.1). The existence of a solution now follows from [1, Theorem 1.5.1] if we can show that f satisfies a Nagumo condition with respect to the pair α, β . This amounts to finding a positive continuous function h on $[0, \infty)$ such that $|f(x, y, y')| \leq h(|y'|)$ for all $x \in [0, R]$, $\alpha(x) \leq y \leq \beta(x)$ and $y' \in \mathbb{R}$ and, furthermore, such that

$$\int_{R^{-1}\beta(R)}^{\infty} \frac{s}{h(s)} ds > \beta(R),$$

cf. [1, Definition 1.4.1]. The function h defined by $h(s) := \varepsilon^{-1}g'(0)(s+1)$ satisfies all these conditions. \square

3. A COMPARISON THEOREM

In order to emphasize that we are going to study the dependence of a solution on the parameters ε and R , we introduce the notation $P(\varepsilon, R)$ for the problem (1.1) - (1.3). The main result of this section is a comparison theorem which is proved by standard maximum principle arguments. As corollaries we obtain that the solution is unique and that it depends in a monotone fashion on both ε and R .

THEOREM 3.1. *Let y_i be a solution of $P(\varepsilon_i, R_i)$ for $i = 1, 2$ and suppose that $R_2 \geq R_1 > x_0$ and $\varepsilon_2 \geq \varepsilon_1$. Then $y_1(x) \geq y_2(x)$ for $0 < x < R_1$. Moreover, if one of the inequalities for the parameters is strict, then so is the inequality for the solutions.*

PROOF. Let the function m be defined by $m(x) := y_1(x) - y_2(x)$. Suppose that m achieves a nonpositive minimum on $(0, R_1)$, i.e. suppose that for some $x_1 \in (0, R_1)$, $m(x_1) \leq 0$, $m'(x_1) = 0$ and $m''(x_1) \geq 0$. By subtracting the equation for y_2 from the one for y_1 we obtain

$$\varepsilon_1 x_1 m''(x_1) - (\varepsilon_2 - \varepsilon_1) x_1 y_2''(x_1) - y_1'(x_1) m(x_1) = 0.$$

However, all the terms on the left-hand side of this equality are non-negative and if either $\varepsilon_2 > \varepsilon_1$ or $m(x_1) < 0$ at least one of them is positive. If $\varepsilon_1 = \varepsilon_2$ and $m(x_1) = 0$ then the uniqueness theorem for ordinary differential equations implies that $m(x) = 0$ for all $x \in [0, R_1]$, which cannot be true if $R_2 > R_1$. So we see that m cannot achieve a negative minimum and that m cannot become zero on $(0, R_1)$ if one of the inequalities for the parameters is strict. Since $m(0) = 0$ and $m(R_1) \geq 0$ this proves the theorem. \square

COROLLARY 3.2. *The problem $P(\epsilon, R)$ has one and only one solution.*

PROOF. We know that at least one solution exists (Theorem 2.2). Let both y_1 and y_2 satisfy $P(\epsilon, R)$, then Theorem 3.1 implies that $y_1(x) \geq y_2(x)$ but likewise that $y_2(x) \geq y_1(x)$. Hence, $y_1(x) = y_2(x)$ for $x \in [0, R]$. \square

COROLLARY 3.3. *Let $y = y(x; \epsilon, R)$ be the solution of $P(\epsilon, R)$. Then y is a monotone decreasing function of ϵ for each $R > x_0$ and each $x \in (0, R)$, and y is a monotone decreasing function of R for each $\epsilon > 0$ and each $x \in (0, R)$.*

4. THE LIMITING BEHAVIOUR AS $R \rightarrow \infty$

In this section we study the limiting behaviour as $R \rightarrow \infty$ of the solution $y = y(x; \epsilon, R)$ of the problem $P(\epsilon, R)$. Since y is a bounded and monotone function of R , the definition $\bar{y}(x; \epsilon) := \lim_{R \rightarrow \infty} y(x; \epsilon, R)$ makes sense for all $x, \epsilon > 0$. This definition implies at once that $\bar{y}(0; \epsilon) = 0$ and that \bar{y} is a nondecreasing function of x and a nonincreasing function of ϵ .

From the estimates in Theorem 2.1 we obtain, via the Arzela-Ascoli theorem, that both $y(\cdot; \epsilon, R)$ and $y'(\cdot; \epsilon, R)$ converge uniformly on compact subsets. Invoking equation (1.1) we see that the same must be true for $y''(\cdot; \epsilon, R)$. It follows that $\bar{y}(\cdot; \epsilon)$ belongs to $C^2(\mathbb{R}_+; \mathbb{R})$ and satisfies equation (1.1). Now it remains to determine $\bar{y}(\infty; \epsilon)$. We will estimate $\bar{y}(\infty; \epsilon)$ from below by constructing a more subtle lower solution for y . But first we prove a result which can be used to estimate $\bar{y}(\infty; \epsilon)$ from above.

LEMMA 4.1. *Let $z \in C^2(\mathbb{R}_+; \mathbb{R})$ satisfy equation (1.1) and $z(0) = 0$. Suppose that $z(\infty) := \lim_{x \rightarrow \infty} z(x)$ exists and satisfies $0 < z(\infty) < \infty$. Then $z(\infty) \leq g(\infty) - \epsilon$.*

PROOF. Both z and z' are positive on $(0, \infty)$ (cf. the proof of Theorem 2.1). For the purpose of contradiction, let us suppose that $z(\infty) > g(\infty) - \epsilon$. Let x_1 be such that $\beta := \epsilon^{-1}(z(x_1) - g(\infty)) > -1$. Then $z(x) - g(x) \geq z(x_1) - g(\infty) = \epsilon\beta$ for all $x \geq x_1$. Integrating equation (1.1) twice from x_1 to x we obtain

$$z(x) = z(x_1) + z'(x_1) \int_{x_1}^x \exp\left(\int_{x_1}^{\xi} \frac{z(\eta) - g(\eta)}{\epsilon\eta} d\eta\right) d\xi.$$

Thus, for $x \geq x_1$,

$$z(x) \geq z'(x_1) \int_{x_1}^x \exp\left(\beta \ln \frac{\xi}{x_1}\right) d\xi = \frac{x_1 z'(x_1)}{\beta + 1} \left(\left(\frac{x}{x_1}\right)^{\beta+1} - 1 \right).$$

Since $\beta+1 > 0$ this would imply that $z(x) \rightarrow \infty$ as $x \rightarrow \infty$. Hence the assumption that $z(\infty) > g(\infty) - \varepsilon$ must be false. \square

We define a function $s = s(x; \lambda, x_1, \nu)$ by

$$(4.1) \quad s(x; \lambda, x_1, \nu) := \lambda \left(1 - \left(\frac{x}{x_1} \right)^{-\nu} \right)$$

and we investigate which conditions for the parameters λ , x_1 and ν guarantee that $s'' \geq f(x, s, s')$ for $x \geq x_1$ (recall that $f(x, y, y') = (\varepsilon x)^{-1} (y - g(x)) y'$). A simple computation shows that this inequality holds indeed for all $x \geq x_1$ if and only if $g(x_1) - \lambda - \varepsilon \nu - \varepsilon \geq 0$, or equivalently, $\nu \leq \varepsilon^{-1} (g(x_1) - \lambda) - 1$. The latter inequality can be satisfied for some *positive* value of ν if and only if $\lambda < g(x_1) - \varepsilon$. In its turn this inequality can be satisfied for sufficiently large x_1 and some *positive* value of λ if and only if $g(\infty) - \varepsilon > 0$.

We now have all the ingredients at hand to prove the following theorem.

THEOREM 4.2.

- (i) If $\varepsilon \leq g(\infty) - k$ then $\bar{y}(\infty; \varepsilon) = k$ and $\lim_{R \rightarrow \infty} \sup_{0 \leq x \leq R} |y(x; \varepsilon, R) - \bar{y}(x; \varepsilon)| = 0$;
- (ii) if $g(\infty) - k < \varepsilon < g(\infty)$ then $\bar{y}(\infty; \varepsilon) = g(\infty) - \varepsilon$;
- (iii) if $\varepsilon \geq g(\infty)$ then $\bar{y}(x; \varepsilon) = 0$ for all $x \geq 0$.

PROOF. (i) For any $\lambda < k$ we can choose x_1 such that $\lambda < g(x_1) - \varepsilon$ and subsequently ν such that $0 < \nu \leq \varepsilon^{-1} (g(x_1) - \lambda) - 1$. For these values of the parameters, s is a lower solution on the interval $[x_1, R]$. The function t defined by $t(x) := k$ is an upper solution and f satisfies a Nagumo condition with respect to the pair s, t and the interval $[x_1, R]$. It follows that the inequality

$$s(x; \lambda, x_1, \nu) \leq y(x; \varepsilon, R) \leq k,$$

which holds for $x = x_1$ and for $x = R$, actually is satisfied for all

$x \in [x_1, R]$. By taking first the limit $R \rightarrow \infty$ and then the limit $x \rightarrow \infty$ we obtain

$$\lambda \leq \bar{y}(\infty; \varepsilon) \leq k.$$

Since this inequality holds for $\lambda < k$, necessarily $\bar{y}(\infty; \varepsilon) = k$. This result and the monotonicity of y with respect to x together imply that the convergence of y to \bar{y} is in fact uniform in x (we refer to [3, Lemma 2.4] for the proof of this statement).

(ii). If $g(\infty) - k \leq \varepsilon < g(\infty)$, we can make s into a lower solution by a suitable choice of x_1 and v if and only if $\lambda < g(\infty) - \varepsilon$. The argument we used in the proof of (i) now shows that $\bar{y}(\infty; \varepsilon) \geq g(\infty) - \varepsilon$. On the other hand, Lemma 4.1 implies that $\bar{y}(\infty; \varepsilon) \leq g(\infty) - \varepsilon$. So $\bar{y}(\infty; \varepsilon) = g(\infty) - \varepsilon$.

(iii) From Lemma 4.1 we deduce that no solution of (1.1) with a positive limit at infinity can exist if $\varepsilon \geq g(\infty)$. Hence $\bar{y}(\infty; \varepsilon) = 0$ and consequently $\bar{y}(x; \varepsilon) = 0$ for all $x \geq 0$. \square

The results of this section are at the same time results concerning the existence and non-existence of a solution of the problem $P(\varepsilon, \infty)$ defined by (1.1), (1.2) and $\lim_{x \rightarrow \infty} y(x) = k$. By exactly the same arguments which we used before one can derive the bounds of Theorem 2.1 and one can show that there exists at most one solution of $P(\varepsilon, \infty)$. For convenience we formulate this result in the following theorem.

THEOREM 4.3. *There exists a function $y \in C^2(\mathbb{R}_+; \mathbb{R})$ which satisfies (1.1), (1.2) and the condition $\lim_{x \rightarrow \infty} y(x) = k$ if and only if $\varepsilon \leq g(\infty) - k$. If it exists, it is unique and it satisfies the inequalities given in Theorem 2.1.*

5. THE LIMITING BEHAVIOUR AS $\varepsilon \downarrow 0$

Throughout this section $R > x_0$ will be fixed and we will suppress the dependence on R in the notation, because it is inessential. The solution y of (1.1) - (1.3) is a bounded and monotone function of ε and we define $\tilde{y}(x) := \lim_{\varepsilon \downarrow 0} y(x; \varepsilon)$. From Theorem 2.1(i) and (ii) and the Arzela-Ascoli theorem we deduce that \tilde{y} is continuous and that in fact $\lim_{\varepsilon \downarrow 0} \sup_{0 \leq x \leq R} |\tilde{y}(x) - y(x; \varepsilon)| = 0$.

THEOREM 5.1. $\tilde{y}(x) = \min\{g(x), k\}$.

PROOF. From Theorem 2.1(i) we know that $\tilde{y}(x) \leq \min\{g(x), k\}$. Take any $x < x_0$, then $\tilde{y}(x) < k$. We claim that this implies that $\liminf_{\varepsilon \downarrow 0} y'(x; \varepsilon) > 0$. Indeed, suppose that the sequence $\{\varepsilon_i\}$ is such that $\varepsilon_i \downarrow 0$ and $y'(x; \varepsilon_i) \downarrow 0$ as $i \rightarrow \infty$, then by taking the limit $i \rightarrow \infty$ in the relation

$$\begin{aligned} k &= y(R; \varepsilon_i) \\ &= y(x; \varepsilon_i) + \int_x^R y'(\xi; \varepsilon_i) d\xi \leq y(x; \varepsilon_i) + (R-x)y'(x; \varepsilon_i), \end{aligned}$$

we arrive at the conclusion that $\tilde{y}(x) \geq k$, which is impossible.

Integrating equation (1.1) from 0 to x we obtain

$$(5.1) \quad \varepsilon(y'(x; \varepsilon) - y'(0; \varepsilon)) = \int_0^x \frac{y(\xi; \varepsilon) - g(\xi)}{\xi} y'(\xi; \varepsilon) d\xi.$$

Suppose that $x < x_0$ and $\max_{0 \leq \xi \leq x} |\tilde{y}(\xi) - g(\xi)| > 0$ then, since $g'(0) > y'(\xi; \varepsilon) \geq y'(x; \varepsilon)$ for $0 < \xi \leq x$ and $\liminf_{\varepsilon \downarrow 0} y'(x; \varepsilon) > 0$, the right-hand side of (5.1) is bounded away from zero as $\varepsilon \downarrow 0$. However, this is impossible since the left-hand side tends to zero as $\varepsilon \downarrow 0$. So $\tilde{y}(x) = g(x)$ for all $x < x_0$, and by continuity $\tilde{y}(x_0) = k$. The function \tilde{y} , being the limit of monotone functions, is monotone nondecreasing. Hence $\tilde{y}(x) \geq k$ for $x > x_0$ and consequently $\tilde{y}(x) = k$ for $x > x_0$. \square

By taking $\varepsilon = 0$ in (1.1) we obtain the reduced equation

$$(5.2) \quad (g(x) - y)y' = 0.$$

The limiting function \tilde{y} satisfies the boundary conditions (1.2) and (1.3) and the equation (5.2) except at the point $x = x_0$, where \tilde{y}' is not defined. Motivated in part by the physical application (cf. section 7) we shall now investigate the limiting behaviour of $y'(x; \varepsilon)$ as $\varepsilon \downarrow 0$. It will then become even more apparent that $x = x_0$ is an exceptional point. The following lemma is needed in the proof of Theorem 5.3, but it is of some interest in itself.

LEMMA 5.2. *Let $\delta > 0$ be arbitrary. For any $\varepsilon_0 > 0$ there exists an $M > 0$ such that $0 < g(x) - y(x;\varepsilon) < M\varepsilon x$ for all $x \in [0, x_0 - \delta]$ and all $\varepsilon \in (0, \varepsilon_0)$.*

PROOF. Let $\delta > 0$ and $\varepsilon_0 > 0$ arbitrary. We define

$$m(\varepsilon) := \min_{x_0 - \delta \leq x \leq x_0 - \frac{1}{2}\delta} \{g(x) - y(x;\varepsilon)\}.$$

Then there exist positive constants C_i , $i = 1, 2, 3$, such that for $\varepsilon \in (0, \varepsilon_0)$

$$\begin{aligned} m(\varepsilon) &\leq C_1 \int_{x_0 - \delta}^{x_0 - \delta/2} (g(\xi) - y(\xi;\varepsilon)) d\xi \\ &\leq C_2 \int_{x_0 - \delta}^{x_0 - \delta/2} \frac{g(\xi) - y(\xi;\varepsilon)}{\xi} y'(\xi;\varepsilon) d\xi \leq C_3 \varepsilon \end{aligned}$$

(see the proof of Theorem 5.1 and in particular formula (5.1)). Let the function $v = v(x;\varepsilon)$ be defined by $v(x;\varepsilon) := g(x) - y(x;\varepsilon) - M\varepsilon x$, where the constant $M > 0$ is still at our disposal. Then v satisfies the equation

$$\varepsilon x v'' - y'(x;\varepsilon)v = \varepsilon x(g''(x) + M y'(x;\varepsilon))$$

and consequently $\varepsilon x v'' - \mu v > 0$ if $M > \gamma \mu^{-1}$, $\varepsilon \in (0, \varepsilon_0)$ and $x \in (0, x_0 - \frac{1}{2}\delta]$, where the positive numbers γ and μ are defined by

$$\gamma := - \inf_{0 < x \leq x_0 - \frac{1}{2}\delta} g''(x)$$

and

$$\mu := \inf_{0 < \varepsilon < \varepsilon_0} y'(x_0 - \frac{\delta}{2}; \varepsilon).$$

So if $M > \gamma \mu^{-1}$ and $\varepsilon \in (0, \varepsilon_0)$, then v cannot assume a nonnegative maximum on $(0, x_0 - \frac{1}{2}\delta)$. Let $x(\varepsilon)$ be such that $g(x) - y(x;\varepsilon)$ achieves its minimum on the set $[x_0 - \delta, x_0 - \frac{1}{2}\delta]$ in the point $x = x(\varepsilon)$. Then $v(x(\varepsilon);\varepsilon) = m(\varepsilon) - M\varepsilon x(\varepsilon) < 0$ if $M > (x_0 - \delta)^{-1} C_3$. Since $v(0;\varepsilon) = 0$, this implies that for $M > \max\{\gamma \mu^{-1}, (x_0 - \delta)^{-1} C_3\}$ $v(x;\varepsilon) < 0$ for $x \in (0, x(\varepsilon))$ and a fortiori for $x \in (0, x_0 - \delta)$. \square

THEOREM 5.3. *Let $\delta > 0$ be arbitrary. Then*

- (i) $\lim_{\varepsilon \downarrow 0} \sup_{0 \leq x \leq x_0 - \delta} |g'(x) - y'(x; \varepsilon)| = 0;$
(ii) $\lim_{\varepsilon \downarrow 0} \sup_{x_0 + \delta \leq x \leq R} |y'(x; \varepsilon)| = 0.$

PROOF. (i) From the equation (1.1), Theorem 2.1(ii) and Lemma 5.2 we deduce that $-g'(0)M < y''(x; \varepsilon) < 0$ for $x \in [0, x_0 - \delta]$ and $\varepsilon \in (0, \varepsilon_0)$. By the Arzela-Ascoli theorem this implies that the limit set of $\{y'(\cdot; \varepsilon) \mid \varepsilon > 0\}$ as $\varepsilon \downarrow 0$ is nonempty in $C([0, x_0 - \delta]; \mathbb{R})$. The result now follows from the fact that y tends to g on $[0, x_0 - \delta]$ as $\varepsilon \downarrow 0$.

(ii). Integrating equation (1.1) from $x_0 + \frac{1}{2}\delta$ to x we obtain

$$\varepsilon(y'(x; \varepsilon) - y'(x_0 + \frac{1}{2}\delta; \varepsilon)) = \int_{x_0 + \frac{1}{2}\delta}^x \frac{y(\xi; \varepsilon) - g(\xi)}{\xi} y'(\xi; \varepsilon) d\xi.$$

For $x \in [x_0 + \delta, R]$ the right-hand side is smaller than $\frac{1}{2}\delta R^{-1} (k - g(x_0 + \frac{\delta}{2})) y'(x; \varepsilon)$. Consequently $0 < y'(x; \varepsilon) < 2g'(0)\varepsilon R \delta^{-1} (g(x_0 + \frac{\delta}{2}) - k)^{-1}$. \square

In the next section we shall concentrate on a formal approximation for y' in the neighbourhood of $x = x_0$.

In section 4 it was shown that the problem $P(\varepsilon, \infty)$ has a unique solution for ε sufficiently small. The analysis of this section can be repeated, *mutatis mutandis*, to derive the analogous results concerning the limiting behaviour of this solution as $\varepsilon \downarrow 0$. In particular this implies that the limits $\varepsilon \downarrow 0$ and $R \rightarrow \infty$ are interchangeable.

6. THE TRANSITION LAYER

In Theorem 5.3 we have shown that y' converges nonuniformly on the interval $[0, R]$ as $\varepsilon \downarrow 0$. This feature is typical for a singular perturbation problem. In this section we use the standard method of the stretching of a variable to obtain more information about the behaviour of y' near the transition point $x = x_0$.

Our starting point is the equation

$$(6.1) \quad \epsilon x z'' + \epsilon z' - \epsilon x \frac{(z')^2}{z} + (g'(x) - z)z = 0,$$

which is obtained by differentiation of (1.1), followed by elimination of y and the substitution $y' = z$. By the stretching of x near x_0 we mean the introduction of a local coordinate ξ according to $x = x_0 + \xi \epsilon^\alpha$. If we make this substitution in (6.1) and if we subsequently retain only the terms of lowest order in ϵ , then it depends on the value of α what the resulting equation will be. One easily verifies that the choice $\alpha = \frac{1}{2}$ leads to a significant equation, namely to

$$(6.2) \quad x_0 z'' - x_0 \frac{(z')^2}{z} + (g'(x_0) - z)z = 0.$$

To this equation we add the condition that its solution should match the limits of y' to the left and to the right of x_0 , respectively. This amounts to the condition

$$(6.3) \quad \lim_{\xi \rightarrow -\infty} z(\xi) = g'(x_0), \quad \lim_{\xi \rightarrow +\infty} z(\xi) = 0.$$

We rewrite (6.2) as a two-dimensional autonomous first order system

$$(6.4) \quad \begin{aligned} z' &= u, \\ u' &= \frac{u^2}{z} + \frac{1}{x_0}(z - g'(x_0))z, \end{aligned}$$

and we analyse the trajectories in the (z, u) -phase plane. If we linearize the system about the equilibrium point $(g'(x_0), 0)$ we find the corresponding eigenvalues and eigenvectors to be

$$\lambda_{\pm} = \pm \sqrt{\frac{g'(x_0)}{x_0}}$$

and

$$(1, \lambda_{\pm})^T.$$


So $(g'(x_0), 0)$ is a saddle point and the unstable manifold is tangent to the vector $(1, \lambda_+)^T$.

Our next objective is to overcome the difficulty that the right-hand side of (6.4) is singular for $z = 0$. Consider the set Ω enclosed by the z -axis, the line $z = g'(x_0)$ and the arc θ which is defined by the parameterization $(z,u) = (\theta_1(t), \theta_2(t))$, $1 \leq t < \infty$, with

$$\theta_1(t) := g'(x_0) \exp\left\{\frac{g'(x_0)}{2x_0} (1-t^2)\right\}$$

and

$$\theta_2(t) := -\frac{(g'(x_0))^2}{x_0} t \exp\left\{\frac{g'(x_0)}{2x_0} (1-t^2)\right\}.$$



Along $\partial\Omega$ the vector field defined by the right-hand side of (6.4) is always pointing inward. For the part of $\partial\Omega$ where $u = 0$ or $z = g'(x_0)$ this is immediately clear from (6.4), and along θ it follows from the fact that the outcome of a calculation of the inner product of the vector field with $(\hat{\theta}_2, -\hat{\theta}_1)^T$, i.e. with the inward pointing normal to θ , is

$$\frac{(g'(x_0))^4}{x_0^2} t \exp\left\{\frac{3g'(x_0)}{2x_0} (1-t^2)\right\},$$

which is positive for $1 \leq t < \infty$.

In Ω the vector field is continuous if we define its value at $(0,0)$

to be zero. The unstable manifold of the saddle point which enters Ω can be continued and it cannot leave Ω through $\partial\Omega \setminus (0,0)$. Hence it enters the singular point $(0,0)$. So we have shown that the problem (6.2) - (6.3) has (modulo translation) a unique solution.

Equation (6.2) can be solved explicitly for ξ as a function of z . If we require that $\lim_{\xi \rightarrow -\infty} z(\xi) = g'(x_0)$ and $z(0) = z_0$ we find that

$$\xi = \sqrt{\frac{x_0}{2g'(x_0)}} \int_{\ln \frac{z_0}{g'(x_0)}}^{\ln \frac{z}{g'(x_0)}} \frac{du}{\sqrt{e^u - 1 - u}}.$$

From this expression we deduce that

$$z(\xi) \sim \exp\left(-\frac{g'(x_0)}{2x_0} \xi^2\right) \quad \text{for } \xi \rightarrow +\infty.$$

Let z be a solution of (6.2) - (6.3). The idea of singular perturbation theory is that z describes the shape of y' near $x = x_0$ for small values of ε , and that one can approximate y' uniformly on $[0,R]$ by using the building-stones z and \tilde{y}' . We shall not give a precise formulation of this idea. The interested reader is referred to a paper of FIFE [4] where similar problems are treated and rigorous proofs are given.

7. APPLICATION TO A THEORETICAL MODEL OF AN IONIZED GAS BETWEEN TWO ELECTRODES

7.1. The physical background

MARODE et al. [8] consider an ionized gas between two electrodes in which the ions can be represented by a density $n_i(\vec{r})$ and the electrons by a density $n_e(\vec{r})$. Due to the heavy weight of the ions, the density $n_i(\vec{r})$ may be considered as fixed, whereas the highly mobile electrons assume a spatial distribution in thermal equilibrium with the given $n_i(\vec{r})$. The problem is then to find $n_e(\vec{r})$ for given $n_i(\vec{r})$. A special situation of practical

interest is considered, namely a pre-breakdown discharge which spreads out in filamentary form, as discussed by GALLIMBERTI [5] and MARODE [7]. For this case, where there is cylindrical symmetry, MARODE et al. [8] have derived the following equation:

$$(7.1) \quad -\tilde{\epsilon} \frac{1}{r} \frac{d}{dr} \left(\frac{r}{n_e(r)} \frac{dn_e(r)}{dr} \right) + n_e(r) = n_i(r),$$

where $\tilde{\epsilon}$ is a small positive constant proportional to the temperature. In addition, $n_e(r)$ has to satisfy the boundary condition

$$(7.2) \quad \frac{dn_e}{dr}(0) = 0$$

and the condition

$$(7.3) \quad \int_0^{\infty} (n_i(r) - n_e(r)) r dr = N > 0,$$

where N represents the excess of ions.

The electron density $n_e(r)$ is to be determined from equations (7.1) - (7.3). We assume that $n_i(r)$ is a smooth decreasing function of r . An explicit choice of MARODE et al. [8] is $n_i(r) = e^{-r^2}$, because the ions are supposed to have a Gaussian distribution due to diffusion.

7.2. A transformation of the equation

Let us make the following change of function:

$$(7.4) \quad y(r^2) = \int_0^r n_e(\rho) \rho d\rho.$$

Obviously y represents the total quantity of electrons per unit length in a cylinder of radius r . We multiply (7.1) by r and integrate from 0 to r to find

$$-\tilde{\epsilon} \frac{r}{n_e(r)} \frac{dn_e(r)}{dr} + \int_0^r n_e(\rho) \rho d\rho = \int_0^r n_i(\rho) \rho d\rho,$$

whence upon putting $x = r^2$

$$-2\tilde{\epsilon}xy''(x) + y(x)y'(x) = \left(\int_0^{\sqrt{x}} n_e(\rho)\rho d\rho \right) y'(x).$$

We finally obtain the equation (1.1), the boundary condition (1.2) and the limiting form of the boundary condition (1.3), i.e. $y(\infty) = k$, if we put

$$\epsilon = 2\tilde{\epsilon}, \quad g(x) = \int_0^{\sqrt{x}} n_i(\rho)\rho d\rho, \quad \text{and } k = \int_0^{\infty} n_i(\rho)\rho d\rho - N.$$

Since there are fewer electrons than ions, we have $k < g(\infty)$, as assumed in the paper.

Problems (7.1) - (7.3) and $P(\epsilon, \infty)$ are equivalent under the change of function (7.4) and the change of variable $x = r^2$. (Since $y' > 0$, $n_e(r) = 2y'(r^2)$ is well defined.) In this paper, starting from the study of the corresponding problem on a finite interval, we have established the main characteristics of $P(\epsilon, \infty)$.

7.3. Physical interpretation of the results

Given the properties for y and y' which were proved in the previous sections, we deduce in particular the following properties for n_e :

(a) for $\tilde{\epsilon} \leq N/2$, there exists a unique solution of problem (7.1) - (7.3) and there does not exist any otherwise.

(b) Let $r_0 > 0$ be defined by $\int_{r_0}^{\infty} n_i(\rho)\rho d\rho = N$. Then for any $\delta \in (0, r_0)$

$$\lim_{\tilde{\epsilon} \rightarrow 0} \sup_{r \in [0, r_0 - \delta]} |n_e(r) - n_i(r)| = 0,$$

$$\lim_{\tilde{\epsilon} \rightarrow 0} \sup_{r \in [r_0 + \delta, \infty)} |n_e(r)| = 0.$$

(c) In the neighbourhood of r_0 , n_e exhibits a transition layer behaviour.

The most interesting physical result is that at temperatures such that $\tilde{\epsilon} \ll 1$, the density of electrons is not smoothly distributed in the ionized region. It is, on the contrary, almost confined to a cylindrical

neighbourhood of the discharge axis where, as a consequence, the plasma is quasi-neutral. At a certain distance from the axis a transition occurs to a region of positive space charge. The width of the transition layer is of order $\xi^{\frac{1}{2}}$, which MARODE et al. [8] show to have the form of a Debye shielding length, resulting from a balance between thermal and Coulomb energy.

ACKNOWLEDGEMENT. *The authors would like to thank E. Marode and I. Gallimberti for suggesting this problem to them and for helpful discussions, and M. Bakker for his numerical solution which inspired the treatment of this paper.*

REFERENCES

- [1] BERNFELD, S.R. & V. LAKSHMIKANTHAM, *An Introduction to Nonlinear Boundary Value Problems*, Academic Press, New York 1974.
- [2] CLÉMENT, Ph. & I.B. EMMERTH, *On the structure of continua of positive and concave solutions for twopoint nonlinear eigenvalue problems*, Univ. of Wisconsin MRC TSR 1766, 1977.
- [3] DIEKMANN, O., *Limiting behaviour in an epidemic model*, *Nonlinear Analysis, Theory, Methods & Applications* 1 (1977) 459-470.
- [4] FIFE, P.C., *Transition layers in singular perturbation problems*, *J. Diff. Eq.* 15 (1974) 77-105.
- [5] GALLIMBERTI, I., *A computer model for streamer propagation*, *J. Phys. D. Appl. Phys.* 5 (1972) 2179-2189.
- [6] HALLAM, T.G. & D.E. LOPER, *Singular boundary value problems arising in a rotating fluid flow*, *Arch. Rat. Mech. Anal.* 60 (1976) 355-369.
- [7] MARODE, E., *The mechanism of spark breakdown in air at atmospheric pressure between a positive point and a plane; I: Experimental: nature of the streamer track; II: Theoretical: Computer simulation of the streamer track*, *J. Appl. Phys.* 46 (1975) 2005-2015, 2016-2020.
- [8] MARODE, E. et al., in preparation.