

stichting
mathematisch
centrum



DEPARTMENT OF APPLIED MATHEMATICS

TW 156/76

MAY

T.H. KOORNWINDER & I.G. SPRINKHUIZEN-KUYPER

HYPERGEOMETRIC FUNCTIONS OF 2×2 MATRIX ARGUMENT ARE
SOLUTIONS OF THE PARTIAL DIFFERENTIAL EQUATIONS FOR
APPELL'S FUNCTION F_4

Prepublication

2e boerhaavestraat 49 amsterdam

BIBLIOTHEEK MATHEMATISCH CENTRUM
—AMSTERDAM—

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O), by the Municipality of Amsterdam, by the University of Amsterdam, by the Free University at Amsterdam, and by industries.

AMS(MOS) subject classification scheme (1970): 33A30

Hypergeometric functions of 2×2 matrix argument are solutions of the partial differential equations for Appell's function F_4 ^{*)}

by

T.H. Koornwinder & I.G. Sprinkhuizen-Kuyper.

ABSTRACT

The statement in the title is proved. As a result the hypergeometric function of 2×2 matrix argument is written as a linear combination of two hypergeometric Appell functions F_4 .

KEY WORDS & PHRASES: *hypergeometric functions of matrix argument; Appell's hypergeometric function F_4 .*

^{*)} This paper is not for review; it is meant for publication elsewhere.

1. INTRODUCTION

Hypergeometric functions of matrix argument, introduced by HERZ [6] are of considerable interest in multivariate statistics (cf. for instance CONSTANTINE [2] and JAMES [7]). Until now the hypergeometric functions of matrix argument on the one hand and of Appell and Lauricella type on the other hand seemed to be completely different generalizations of the one-variable case. In the present note it will be shown that at least in the two variable case some relationship exists between these generalized hypergeometric functions. It will be proved that both the hypergeometric function ${}_2F_1(a,b;c;Z)$ of 2×2 matrix argument Z and Appell's hypergeometric function in two variables $F_4(a,b;c-\frac{1}{2},1+a+b-c;x,y)$ are solutions of the same set of partial differential equations. Since a fundamental system of solutions around $(0,0)$ of the partial differential equations for F_4 is known, there follows an explicit expression of ${}_2F_1(a,b;c;Z)$ as a linear combination of two F_4 functions. For literature about F_4 functions see ERDÉLYI [4, Sect.5.7].

The idea that the results presented here would be valid, originated from our research on a class of orthogonal polynomials in two variables (cf. KOORNWINDER & SPRINKHUIZEN [9]). In a forthcoming paper we will extend the results presented here to the solutions of the partial differential equations of F_4 with arbitrary parameters, thus continuing the work initiated by ERDÉLYI [3].

2. PRELIMINARIES

CONSTANTINE [2] gives a series expansion for $F_{p,q}(Z)$ in terms of so-called zonal polynomials which were introduced by JAMES [7]. If Z is a 2×2 matrix then the zonal polynomials are explicitly known (cf. JAMES [8,(7.9)]). This results in the explicit expansion

$$(2.1) \quad {}_2F_1\left(a,b;c;\begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}\right) = \\ = \sum_{m=0}^{\infty} \sum_{\ell=0}^m \frac{(a)_m (a-\frac{1}{2})_{\ell} (b)_m (b-\frac{1}{2})_{\ell} (\frac{3}{2})_{m-\ell}}{(c)_m (c-\frac{1}{2})_{\ell} (\frac{3}{2})_m \ell! (\frac{1}{2})_{m-\ell}} (XY)^{\frac{1}{2}(m+\ell)} P_{m-\ell}\left(\frac{1}{2}\frac{X+Y}{\sqrt{XY}}\right),$$

$a, b, c \in \mathbb{C}$, $c \neq \frac{1}{2}, 0, -\frac{1}{2}, -1, \dots$, $X, Y \in \mathbb{C}$, $|X|, |Y| < 1$, where $P_n(x)$ denotes the Legendre polynomial of degree n (cf. ERDÉLYI [5]). Note that

$${}_2F_1\left(a, b; c; \begin{bmatrix} X & 0 \\ 0 & 0 \end{bmatrix}\right) = \sum_{m=0}^{\infty} \frac{(a)_m (b)_m}{(c)_m m!} X^m = {}_2F_1(a, b; c; X).$$

For ${}_2F_1\left(a, b; c; \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}\right)$ MUIRHEAD [10] gives the following set of partial differential equations

$$(2.2a) \quad X(1-X)z_{xx} + \left\{c - \frac{1}{2} - (a+b + \frac{1}{2})X + \frac{1}{2} \frac{X(1-X)}{X-Y}\right\} z_x - \frac{1}{2} \frac{Y(1-Y)}{X-Y} z_y = abz,$$

$$(2.2b) \quad Y(1-Y)z_{yy} + \frac{1}{2} \frac{X(1-X)}{X-Y} z_x + \left\{c - \frac{1}{2} - (a+b + \frac{1}{2})Y - \frac{1}{2} \frac{Y(1-Y)}{X-Y}\right\} z_y = abz,$$

and he proves that ${}_2F_1\left(a, b; c; \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}\right)$ is the unique solution of each of these differential equations that satisfies the additional properties

- (a) $z(X, Y)$ is a symmetrical function of X and Y , and
- (b) z is regular in $(0, 0)$ and $z(0, 0) = 1$.

The hypergeometric function $F_4(\alpha, \beta; \gamma, \gamma'; x, y)$ of two variables is defined by

$$(2.3) \quad F_4(\alpha, \beta; \gamma, \gamma'; x, y) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{(\alpha)_{m+n} (\beta)_{m+n}}{(\gamma)_m (\gamma')_n m! n!} x^m y^n,$$

$\alpha, \beta, \gamma, \gamma' \in \mathbb{C}$, $\gamma, \gamma' \neq 0, -1, -2, \dots$, $x, y \in \mathbb{C}$, $|x|^{\frac{1}{2}} + |y|^{\frac{1}{2}} < 1$, cf. APPELL & KAMPÉ DE FÉRIET [1] or SLATER [11]. Note that

$$F_4(\alpha, \beta; \gamma, \gamma'; 0, y) = \sum_{n=0}^{\infty} \frac{(\alpha)_n (\beta)_n}{(\gamma')_n n!} y^n = {}_2F_1(\alpha, \beta; \gamma'; y).$$

The function $F_4(\alpha, \beta; \gamma, \gamma'; x, y)$ satisfies the following set of partial differential equations

$$(2.4a) \quad x(1-x)z_{xx} - 2xyz_{xy} - y^2z_{yy} + (\gamma - (\alpha + \beta + 1)x)z_x - (\alpha + \beta + 1)yz_y = \alpha\beta z,$$

$$(2.4b) \quad -x^2 z_{xx} - 2xyz_{xy} + y(1-y)z_{yy} - (\alpha+\beta+1)xz_x + (\gamma' - (\alpha+\beta+1)y)z_y = \alpha\beta z.$$

In any simply connected region included in $\{(x,y) \in \mathbb{C}^2 \mid x \neq 0, y \neq 0, |x|^{\frac{1}{2}} + |y|^{\frac{1}{2}} < 1\}$ the general solution of (2.4a,b) reads

$$(2.5) \quad z(x,y) = AF_4(\alpha, \beta; \gamma, \gamma'; x, y) + \\ + By^{1-\gamma'} F_4(1+\alpha-\gamma', 1+\beta-\gamma'; \gamma, 2-\gamma'; x, y) + \\ + Cx^{1-\gamma} F_4(1+\alpha-\gamma, 1+\beta-\gamma; 2-\gamma, \gamma'; x, y) + \\ + Dx^{1-\gamma} y^{1-\gamma'} F_4(2+\alpha-\gamma-\gamma', 2+\beta-\gamma-\gamma'; 2-\gamma, 2-\gamma'; x, y),$$

provided that γ and γ' are noninteger (cf. APPELL & KAMPÉ DE FÉRIET [1]).

3. THE MAIN RESULT

Consider the transformation of variables

$$(3.1) \quad \begin{cases} x = XY, \\ y = (1-X)(1-Y). \end{cases}$$

THEOREM 3.1. *The function given by (2.1) is an analytic function of x and y in some neighbourhood of $(0,1)$ and it satisfies the system of partial differential equations (2.4a,b) with $\alpha: = a$, $\beta: = b$, $\gamma: = c - \frac{1}{2}$, $\gamma': = 1 + a + b - c$.*

PROOF. The first statement is clear from (2.1). For proving the second statement let S and T denote the equations (2.2a) and (2.2b), respectively. Then equations (2.4a) and (2.4b) are obtained by transforming the equations

$$\frac{1}{2}(S+T) - \frac{2-X-Y}{2(X-Y)}(S-T) \text{ and}$$

$$\frac{1}{2}(S+T) + \frac{X+Y}{2(X-Y)}(S-T)$$

in terms of x and y . \square

THEOREM 3.2.

$$\begin{aligned}
{}_2F_1\left(a, b; c; \begin{bmatrix} X & 0 \\ 0 & Y \end{bmatrix}\right) &= \\
&= \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} F_4(a, b; c-\frac{1}{2}, 1+a+b-c; XY, (1-X)(1-Y)) \\
&+ \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} \{(1-X)(1-Y)\}^{c-a-b} F_4(c-b, c-a; c-\frac{1}{2}, 1+c-a-b; \\
&\qquad\qquad\qquad XY, (1-X)(1-Y)),
\end{aligned}$$

with X and Y such that all terms are well-defined and provided that c is not an integer or half integer $\leq \frac{1}{2}$ and that $a + b - c$ is not an integer.

PROOF. Let $z(x, y)$ denote the function given by (2.1) considered as a function of x and y in a neighbourhood of $(0, 1)$. Then, because of (2.5) and since z is regular in $(0, 1)$, it follows that

$$\begin{aligned}
z &= A F_4(a, b; c-\frac{1}{2}, 1+a+b-c; x, y) + \\
&+ B y^{c-a-b} F_4(c-b, c-a; c-\frac{1}{2}, 1+c-a-b; x, y).
\end{aligned}$$

Restriction to $x = 0$ gives

$$\begin{aligned}
{}_2F_1(a, b; c; 1-y) &= A {}_2F_1(a, b; 1+a+b-c; y) + \\
&+ B y^{c-a-b} {}_2F_1(c-b, c-a; 1+c-a-b; y).
\end{aligned}$$

It follows from ERDÉLYI [4, §2.10] that

$$A = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}, \quad B = \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)}. \quad \square$$

LITERATURE

- [1] APPELL, P. & J. KAMPÉ DE FÉRIET, *Fonctions Hypergéométriques et Hypersphériques. Polynomes d' Hermite*, Gauthier-Villars et Cie, Paris, 1926.
- [2] CONSTANTINE, A.G., *Some non-central distribution problems in multivariate analysis*, Ann. Math. Statist. 34 (1963) 1270-1285.

- [3] ERDÉLYI, A., *Integration of the differential equations of Appell's function F_4* , Quart. J. Math., Oxford Ser. 11 (1941) 68-77.
- [4] ERDÉLYI, A., W. MAGNUS, F. OBERHETTINGER & F.G. TRICOMI, *Higher Transcendental Functions*, Vol. I, McGraw-Hill, New York, 1953.
- [5] ERDELYI, A., W. MAGNUS F. OBERHETTINGER & F.G. TRICOMI, *Higher Transcendental Functions*, Vol. II, McGraw-Hill, New York, 1953.
- [6] HERZ, C.S., *Bessel functions of matrix argument*, Ann. Math. 61 (1955) 474-523.
- [7] JAMES, A.T., *Distribution of matrix variates and latent roots derived from normal samples*, Ann. Math. Statist. 35 (1964) 475-501.
- [8] JAMES, A.T., *Calculation of zonal polynomial coefficients by use of the Laplace-Beltrami operator*, Ann. Math. Statist. 39 (1968) 1711-1718.
- [9] KOORNWINDER, T.H. & I.G. SPRINKHUIZEN-KUYPER, *Generalized power series expansions for a class of orthogonal polynomials in two variables*, Math. Centrum Amsterdam Report TW 155 (1976).
- [10] MUIRHEAD, R.J., *Systems of partial differential equations for hypergeometric functions of matrix argument*, Ann. Math. Statist. 41 (1970) 991-1001.
- [11] SLATER, L.J. *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.