

TW

**stichting
mathematisch
centrum**



AFDELING TOEGEPASTE WISKUNDE

TW 144/74

AUGUST

I.G. SPRINKHUIZEN-KUYPER
ORTHOGONAL POLYNOMIALS IN TWO VARIABLES.

A Further analysis of the polynomials orthogonal over a
region bounded by two lines and a parabola.

Prepublication.

2e boerhaavestraat 49 amsterdam

BIBLIOTHEEK MATHEMATISCH CENTRUM
AMSTERDAM

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O), by the Municipality of Amsterdam, by the University of Amsterdam, by the Free University at Amsterdam, and by industries.

Orthogonal polynomials in two variables.

A further analysis of the polynomial orthogonal over a region bounded by two lines and a parabola.*)

by

I.G. Sprinkhuizen-Kuyper

ABSTRACT

Some new results are obtained for the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$, introduced by KOORNWINDER, Proc. Kon. Ned. Akad. Wetensch., A 77, (1974), 48-66, which are orthogonal over a region bounded by two straight lines and a parabola. The most important results are a Rodrigues-type formula and the recurrence relations for $u_{n,k}^{\alpha,\beta,\gamma}(u,v)$. These recurrence relations contain 5 and 9 terms, respectively. Furthermore, the quadratic norm of $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ and the value of $p_{n,k}^{\alpha,\beta,\gamma}(2,1)$ are explicitly given.

*) This paper is not for review; it is meant for publication in a journal.

1. INTRODUCTION

In this paper the analysis of the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ introduced by T.H. KOORNWINDER [4] will be continued.

In many respects, this class of orthogonal polynomials in two variables can be compared with the important class of Jacobi polynomials. In this analysis some properties of the Jacobi polynomials are generalized to the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$.

The polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ form an orthogonal set over a region bounded by two perpendicular straight lines, $1 - u + v = 0$, $1 + u + v = 0$ and by the parabola $u^2 - 4v = 0$ touching these lines, with respect to the weight function $(1-u+v)^\alpha(1+u+v)^\beta(u^2-4v)^\gamma$ which is singular at the boundary of the orthogonality region. For reasons of convergence it is required that $\alpha, \beta, \gamma > -1$ and $\alpha + \gamma + 3/2, \beta + \gamma + 3/2 > 0$. The main results of KOORNWINDER's paper are summarized in section 2.

In the subsequent sections a further analysis is given, using as the main tools a number of partial differential operators. In [4] it is proved that the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ are eigenfunctions of a second order operator $D_1^{\alpha,\beta,\gamma}$ and a fourth-order operator $D_2^{\alpha,\beta,\gamma}$, which are algebraically independent. Furthermore two second-order operators D_-^γ and $D_+^{\alpha,\beta,\gamma}$ are derived with the property that $D_-^\gamma p_{n,k}^{\alpha,\beta,\gamma}(u,v) = \text{const. } p_{n-1,k}^{\alpha+1,\beta+1,\gamma}(u,v)$ and $D_+^{\alpha,\beta,\gamma} p_{n-1,k-1}^{\alpha,\beta,\gamma}(u,v) = \text{const. } p_{n,k}^{\alpha,\beta,\gamma}(u,v)$. Then $D_2^{\alpha,\beta,\gamma}$ is given by $D_2^{\alpha,\beta,\gamma} = D_+^{\alpha,\beta,\gamma} \circ D_-^\gamma$.

In section 4 of this paper another pair of differential operators is derived: these operators $E_-^{\alpha,\beta}$ and $E_+^{\alpha,\beta,\gamma}$ have the property that $E_-^{\alpha,\beta} p_{n,k}^{\alpha,\beta,\gamma}(u,v) = \text{const. } p_{n-1,k}^{\alpha,\beta,\gamma+1}(u,v)$ and $E_+^{\alpha,\beta,\gamma} p_{n-1,k}^{\alpha,\beta,\gamma+1}(u,v) = \text{const. } p_{n,k}^{\alpha,\beta,\gamma}(u,v)$. Then another fourth-order operator, which has the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ as eigenfunctions, can be defined by $D_3^{\alpha,\beta,\gamma} = E_+^{\alpha,\beta,\gamma} \circ E_-^{\alpha,\beta}$. This operator is explicitly expressed as a polynomial in $D_1^{\alpha,\beta,\gamma}$ and $D_2^{\alpha,\beta,\gamma}$. The operators D_-^γ and $E_-^{\alpha,\beta}$ together play a similar role for the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ to that played by the operator $\frac{d}{dx}$ for the Jacobi polynomials.

One of the first problems which arise is to find an explicit expression for $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$. We have succeeded in finding an explicit expression of the Rodrigues-type by using the second-order operators D_+ and E_+ . Expressing

D_+ and E_+ in $(D_-)^*$ and $(E_-)^*$ respectively we obtain a formula for $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ which is similar to the Rodrigues formula for the Jacobi polynomials, but with the two second-order operators $(D_-)^*$ and $(E_-)^*$ instead of $\frac{d}{dx}$ (section 5). However, the expression derived by us is rather complicated and so we have tried to find other expressions for $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$. If $\gamma = -\frac{1}{2}$ and $\gamma = +\frac{1}{2}$ the polynomials can be expressed as symmetric ([4], section 2) or antisymmetric (section 3) products of Jacobi polynomials. In section 10 the polynomials with $\alpha, \beta = \pm\frac{1}{2}$ are expressed in terms of Jacobi polynomials. The case $\gamma = +\frac{1}{2}$ is comparable with the determinants of orthogonal polynomials treated by KARLIN and MCGREGOR [3]. The orthogonal set of 2×2 determinants of Jacobi polynomials gives $p_{n,k}^{\alpha,\beta,+\frac{1}{2}}(x+y,xy)$ after dividing by $(x-y)$.

In section 6 the explicit value of the quadratic norm for the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ is given. The quadratic norm is important for finding coefficients in Fourier expansions with respect to the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ and will be used for the computation of some of the coefficients in the recurrence relations (section 9).

Without knowing an explicit expression for $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ it is possible to find the value of $p_{n,k}^{\alpha,\beta,\gamma}(2,1)$ by using the operators D_+ and E_+ (section 7). The point $(u,v) = (2,1)$ is a vertex of the orthogonality region, which probably plays a similar role for the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ to that played by the point $x=1$ for the Jacobi polynomials $P_n^{(\alpha,\beta)}(x)$. The (unproved) hypothesis is that $p_{n,k}^{\alpha,\beta,\gamma}(2,1)$ is the absolute maximum of $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ if $\alpha \geq \beta \geq -\frac{1}{2}$ and $\gamma \geq -\frac{1}{2}$. For $\gamma = -\frac{1}{2}$ this maximum property follows directly from the explicit expression of $p_{n,k}^{\alpha,\beta,-\frac{1}{2}}(u,v)$ and the maximum property of the Jacobi polynomials.

The analysis of these polynomials suggests that not all powers $\leq (n,k)$ of u and v appear in $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$. This is proved in section 8 and it has a number of consequences. An immediate consequence is that some theorems which give alternative definitions for $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ can be derived. Another is that the number of terms in the recurrence relations is uniformly bounded while for general polynomials in more than one variable this number depends on the degree of the polynomial. In section 9 the recurrence relations are explicitly given. For $u p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ and $v p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ we obtain a five-term and a nine-term recurrence relation, respectively. To build up $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$

using the recurrence relations we need the formula for $vp_{n,k}^{\alpha,\beta,\gamma}(u,v)$ only if $n = k$, and then six terms remain.

Finally, in section 10 two quadratic transformation formulas are given for the case $\alpha = \beta$. These formulas together with the explicit expressions for $\gamma = +\frac{1}{2}$, $\gamma = -\frac{1}{2}$ yield explicit expressions for the cases that α and β are $+\frac{1}{2}$ or $-\frac{1}{2}$.

2. PRELIMINARIES

In this section the main results obtained by KOORNWINDER [4] are summarized.

Let N be the set of pairs of integers (n,k) , $n \geq k \geq 0$, with a lexicographic ordering defined by


$$(2.1) \quad (m,\ell) \leq (n,k) \iff \{m < n \vee (m = n \wedge \ell \leq k)\}.$$

A polynomial $q(u,v)$ is said to have *degree* $(n,k) \in N$ if:

$$q(u,v) = \sum_{(m,\ell) \leq (n,k)} c_{m,\ell} u^{m-\ell} v^{\ell}, \text{ with } c_{n,k} \neq 0.$$

The region with the properties $1 - u + v > 0$, $1 + u + v > 0$ and $u^2 - 4v > 0$, is denoted by R (cf. figure 1). In the region R the weight function $\mu^{\alpha,\beta,\gamma}(u,v)$ is defined by:

$$(2.2) \quad \mu^{\alpha,\beta,\gamma}(u,v) = (1-u+v)^{\alpha} (1+u+v)^{\beta} (u^2-4v)^{\gamma}.$$



DEFINITION 2.1. For $(n,k) \in \mathbb{N}$ and $\alpha, \beta, \gamma > -1$, $\alpha + \gamma + \frac{3}{2}$, $\beta + \gamma + \frac{3}{2} > 0$ the polynomials $p_{n,k}^{\alpha, \beta, \gamma}(u, v)$ are given by

$$(i) \quad p_{n,k}^{\alpha, \beta, \gamma}(u, v) = u^{n-k} v^k + \text{a polynomial of degree lower than } (n, k),$$

$$(ii) \quad \iint_{\mathbb{R}} p_{n,k}^{\alpha, \beta, \gamma}(u, v) u^{m-\ell} v^\ell \mu^{\alpha, \beta, \gamma}(u, v) \, dudv = 0 \text{ if } (m, \ell) < (n, k).$$

Then $p_{n,k}^{\alpha, \beta, \gamma}(u, v)$ satisfies:

$$(2.3) \quad D_1^{\alpha, \beta, \gamma} p_{n,k}^{\alpha, \beta, \gamma}(u, v) = -[n(n+\alpha+\beta+2\gamma+2)+k(k+\alpha+\beta+1)] p_{n,k}^{\alpha, \beta, \gamma}(u, v),$$

$$(2.4) \quad D_2^{\alpha, \beta, \gamma} p_{n,k}^{\alpha, \beta, \gamma}(u, v) = k(k+\alpha+\beta+1)(n+\gamma+\frac{1}{2})(n+\alpha+\beta+\gamma+\frac{3}{2}) p_{n,k}^{\alpha, \beta, \gamma}(u, v),$$

$$(2.5) \quad D_-^{\gamma} p_{n,k}^{\alpha, \beta, \gamma}(u, v) = \begin{cases} k(n+\gamma+\frac{1}{2}) p_{n-1, k-1}^{\alpha+1, \beta+1, \gamma}(u, v) & \text{if } k > 0, \\ 0 & \text{if } k = 0, \end{cases}$$

$$(2.6) \quad D_+^{\alpha, \beta, \gamma} p_{n-1, k-1}^{\alpha+1, \beta+1, \gamma}(u, v) = (k+\alpha+\beta+1)(n+\alpha+\beta+\gamma+\frac{3}{2}) p_{n,k}^{\alpha, \beta, \gamma}(u, v) \\ \text{if } k > 0.$$

The operators are defined by:

$$(2.7) \quad D_1^{\alpha, \beta, \gamma} = (-u^2+2v+2) \frac{\partial^2}{\partial u^2} - 2u(v-1) \frac{\partial^2}{\partial u \partial v} + (u^2-2v^2-2v) \frac{\partial^2}{\partial v^2} + \\ + [-(\alpha+\beta+2\gamma+3)u+(2\beta-2\alpha)] \frac{\partial}{\partial u} + \\ + [(\beta-\alpha)u-(2\alpha+2\beta+2\gamma+5)v-(2\gamma+1)] \frac{\partial}{\partial v},$$

$$(2.8) \quad D_-^{\gamma} = \frac{\partial^2}{\partial u^2} + u \frac{\partial^2}{\partial u \partial v} + v \frac{\partial^2}{\partial v^2} + (\gamma + \frac{3}{2}) \frac{\partial}{\partial v},$$

$$(2.9) \quad D_+^{\alpha, \beta, \gamma} = (1-u+v)^{-\alpha} (1+u+v)^{-\beta} D_-^{\gamma} (1-u+v)^{\alpha+1} (1+u+v)^{\beta+1} = \\ = (1-u-v)(1+u+v) \left(\frac{\partial^2}{\partial u^2} + u \frac{\partial^2}{\partial u \partial v} + v \frac{\partial^2}{\partial v^2} \right) + \\ + [(\alpha-\beta)(u^2-2v-2) + (\alpha+\beta+2)u(v-1)] \frac{\partial}{\partial u} + \\ + [(\alpha+\beta+\gamma+7/2)(-u^2+2v) + (\alpha-\beta)u(v-1) + (2\alpha+2\beta+\gamma+\frac{11}{2})v^2 +$$

$$\begin{aligned}
& + (\gamma + \frac{3}{2}) \frac{\partial}{\partial v} + (\alpha - \beta)(\alpha + \beta + \gamma + \frac{5}{2})u + (\alpha + \beta + 2)(\alpha + \beta + \gamma + \frac{5}{2})v + \\
& + (\alpha - \beta)^2 + (\gamma + \frac{1}{2})(\alpha + \beta + 2),
\end{aligned}$$

$$(2.10) \quad D_2^{\alpha, \beta, \gamma} = D_+^{\alpha, \beta, \gamma} \circ D_-^{\gamma}.$$

Consideration of $(D_-^{\gamma})^*$, the adjoint operator to D_-^{γ} , yields:

$$(2.11) \quad (D_-^{\gamma})^* = D_-^{-\gamma} = (u^2 - 4v)^{\gamma} D_-^{\gamma} \circ (u^2 - 4v)^{-\gamma}.$$

Hence:

$$(2.12) \quad D_+^{\alpha, \beta, \gamma} = \{\mu^{\alpha, \beta, \gamma}(u, v)\}^{-1} (D_-^{\gamma})^* \circ \mu^{\alpha+1, \beta+1, \gamma}(u, v).$$

The operators $D_+^{\alpha, \beta, \gamma}$ and D_-^{γ} are related by:

$$\begin{aligned}
(2.13) \quad & \iint_R (D_+^{\alpha, \beta, \gamma} p(u, v)) q(u, v) \mu^{\alpha, \beta, \gamma}(u, v) \, dudv = \\
& = \iint_R p(u, v) (D_-^{\gamma} q(u, v)) \mu^{\alpha+1, \beta+1, \gamma}(u, v) \, dudv,
\end{aligned}$$

for any two polynomials $p(u, v)$ and $q(u, v)$.

Let

$$(2.14) \quad p_n^{\alpha, \beta}(x) = \frac{2^n n!}{(n + \alpha + \beta + 1)_n} P_n^{(\alpha, \beta)}(x),$$

where $P_n^{(\alpha, \beta)}(x)$ denotes the Jacobi polynomial of order (α, β) (for Jacobi polynomials see ERDÉLYI [2] or SZEGÖ [6]). Then

$$(2.15) \quad p_{n,k}^{\alpha, \beta, -\frac{1}{2}}(x+y, xy) = \begin{cases} p_n^{\alpha, \beta}(x) p_k^{\alpha, \beta}(y) + p_k^{\alpha, \beta}(x) p_n^{\alpha, \beta}(y) & \text{if } n > k, \\ p_n^{\alpha, \beta}(x) p_k^{\alpha, \beta}(y) & \text{if } n = k. \end{cases}$$

3. THE POLYNOMIALS $p_{n,k}^{\alpha,\beta,+\frac{1}{2}}(u,v)$ AS AN ANTISYMMETRIC PRODUCT OF JACOBI POLYNOMIALS

Consider the antisymmetric product of Jacobi polynomials:

$$(3.1) \quad f_{n,k}^{\alpha,\beta}(x,y) = p_{n+1}^{\alpha,\beta}(x) p_k^{\alpha,\beta}(y) - p_k^{\alpha,\beta}(x) p_{n+1}^{\alpha,\beta}(y),$$

where $p_n^{\alpha,\beta}(x)$ is defined by (2.14).

The polynomials $f_{n,k}^{\alpha,\beta}(x,y)$ form an orthogonal set of antisymmetric polynomials over the simplex $-1 \leq y \leq x \leq 1$ with respect to the weight function $((1-x)(1-y))^\alpha ((1+x)(1+y))^\beta$ (cf. [3]). Then $(x-y)^{-1} f_{n,k}^{\alpha,\beta}(x,y)$ is a symmetric polynomial in x and y which can be uniquely expressed as a polynomial in $x+y = u$ and $xy = v$ (see VAN DER WAERDEN [7, §33]).

LEMMA 3.1. $(x-y)^{-1} f_{n,k}^{\alpha,\beta}(x,y) = p_{n,k}^{\alpha,\beta,+\frac{1}{2}}(x+y, xy)$.

PROOF. Application of the definition of $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ (definition 2.1) yields:

$$(i) \quad (x-y)^{-1} f_{n,k}^{\alpha,\beta,\gamma}(x,y) = \sum_{(m,\ell) \leq (n,k)} c_{m,\ell} \frac{x^{m+1} y^\ell - x^\ell y^{m+1}}{x-y} \text{ with } c_{n,k} = 1$$

$$= (x+y)^{n-k} (xy)^k + \text{a polynomial in } (x+y) \text{ and } xy \text{ of degree lower than } (n,k).$$

$$(ii) \quad \{(x-y)^{-1} f_{n,k}^{\alpha,\beta}(x,y) \text{ is an orthogonal set with respect to the measure } (x-y)^2 ((1-x)(1-y))^\alpha ((1+x)(1+y))^\beta dx dy = \text{const. } (1-u+v)^\alpha (1+u+v)^\beta (u^2-4v)^{+\frac{1}{2}} dudv. \quad \square$$

Hence

$$(3.2) \quad p_{n,k}^{\alpha,\beta,+\frac{1}{2}}(x+y, xy) = (x-y)^{-1} \{ p_{n+1}^{\alpha,\beta}(x) p_k^{\alpha,\beta}(y) - p_k^{\alpha,\beta}(x) p_{n+1}^{\alpha,\beta}(y) \}.$$

4. A PAIR OF DIFFERENTIAL OPERATORS WHICH CHANGE n AND γ

A pair of differential operators which change n and γ can be found by using (2.15) and (3.2) and the differential operators for the Jacobi polynomials. Let us define:

$$(4.1) \quad D_{(x)}^{\alpha, \beta} = (1-x)^{-\alpha} (1+x)^{-\beta} \frac{\partial}{\partial x} (1-x)^{\alpha+1} (1+x)^{\beta+1} \frac{\partial}{\partial x} ;$$

then

$$D_{(x)}^{\alpha, \beta} p_n^{\alpha, \beta}(x) = c_n p_n^{\alpha, \beta}(x) \text{ where } c_n = -n(n+\alpha+\beta+1).$$

Hence

$$(4.2) \quad (x-y)^{-1} \{D_{(x)}^{\alpha, \beta} - D_{(y)}^{\alpha, \beta}\} p_{n,k}^{\alpha, \beta, -\frac{1}{2}}(x+y, xy) = \begin{cases} (c_n - c_k) p_{n-1,k}^{\alpha, \beta, +\frac{1}{2}}(x+y, xy) & \text{if } n > k, \\ 0 & \text{if } n = k, \end{cases}$$

and

$$(4.3) \quad \{D_{(x)}^{\alpha, \beta} - D_{(y)}^{\alpha, \beta}\} \circ (x-y) p_{n-1,k}^{\alpha, \beta, +\frac{1}{2}}(x+y, xy) = (c_n - c_k) p_{n,k}^{\alpha, \beta, -\frac{1}{2}}(x+y, xy) \quad \text{if } n > k.$$

Formulas (4.2) and (4.3) now suggest the following definition:

$$(4.4) \quad \begin{aligned} E_-^{\alpha, \beta} &= (x-y)^{-1} \{D_{(x)}^{\alpha, \beta} - D_{(y)}^{\alpha, \beta}\} = \\ &= u \frac{\partial^2}{\partial u^2} + 2(v+1) \frac{\partial^2}{\partial u \partial v} + u \frac{\partial^2}{\partial v^2} + (\beta-\alpha) \frac{\partial}{\partial v} + (\alpha+\beta+2) \frac{\partial}{\partial u} . \end{aligned}$$

If $(E_-^{\alpha, \beta})^*$ is the adjoint operator to $E_-^{\alpha, \beta}$ then

$$(4.5) \quad (E_-^{\alpha, \beta})^* = u \frac{\partial^2}{\partial u^2} + 2(v+1) \frac{\partial^2}{\partial u \partial v} + u \frac{\partial^2}{\partial v^2} - (\beta-\alpha) \frac{\partial}{\partial v} - (\alpha+\beta-2) \frac{\partial}{\partial u} .$$

Note that

$$(4.6) \quad (E_-^{\alpha, \beta})^* = E_-^{-\alpha, -\beta} = (1-u+v)^\alpha (1+u+v)^\beta E_-^{\alpha, \beta} \circ (1-u+v)^{-\alpha} (1+u+v)^{-\beta} .$$

Now we can define the operator $E_+^{\alpha, \beta, \gamma}$ as:

$$(4.7) \quad \begin{aligned} E_+^{\alpha, \beta, \gamma} &= \{\mu^{\alpha, \beta, \gamma}(u, v)\}^{-1} (E_-^{\alpha, \beta})^* \circ \mu^{\alpha, \beta, \gamma+1}(u, v) \\ &= (u^2 - 4v) \left(u \frac{\partial^2}{\partial u^2} + 2(v+1) \frac{\partial^2}{\partial u \partial v} + u \frac{\partial^2}{\partial v^2} \right) \end{aligned}$$

$$\begin{aligned}
& + [(\alpha+\beta+4\gamma+6)(u^2-4v) + 8(\gamma+1)(v-1)] \frac{\partial}{\partial u} \\
& + [(\beta-\alpha)(u^2-4v) + 4(\gamma+1)u(v-1)] \frac{\partial}{\partial v} \\
& + 2(\gamma+1)(\alpha+\beta+2\gamma+3)u - 4(\gamma+1)(\beta-\alpha).
\end{aligned}$$

LEMMA 4.1. *If*

$$q_{n,k}(u,v) = u^{n-k}v^k + \text{a polynomial of degree lower than } (n,k)$$

and

$$q_{n-1,k}(u,v) = u^{n-k-1}v^k + \text{a polynomial of degree lower than } (n-1,k)$$

then

$$E_{-}^{\alpha,\beta} q_{n,k}(u,v) = \begin{cases} (n-k)(n+k+\alpha+\beta+1)u^{n-k-1}v^k + \text{a polynomial of} \\ \text{degree lower than } (n-1,k) \text{ if } n > k, \\ \text{a polynomial of degree equal or lower than} \\ (n-1,n-1) \text{ if } n = k, \end{cases}$$

and

$$E_{+}^{\alpha,\beta,\gamma} q_{n-1,k}(u,v) = (n-k+2\gamma+1)(n+k+\alpha+\beta+2\gamma+2)u^{n-k}v^k + \text{a polynomial of degree lower than } (n,k) \text{ if } n > k.$$

PROOF. Lemma 4.1 follows immediately from (4.4) and (4.7) \square

KOORNWINDER proved:

LEMMA 4.2. *Let R be a bounded region in \mathbb{R}^2 such that certain polynomials*

$w_1(x,y), w_2(x,y), \dots, w_k(x,y)$ *are positive over R and the product*

$w_1 \cdot w_2 \cdot \dots \cdot w_k$ *is zero at the boundary ∂R .*

Let $X^{\alpha_1, \dots, \alpha_k}$ be a partial differential operator in x , and Y , its coefficients being polynomials in $x, y, \alpha_1, \dots, \alpha_k$.

Let the operator $Y^{\alpha_1, \dots, \alpha_k}$ be defined by:

$$Y^{\alpha_1, \dots, \alpha_k} = w_1^{-\alpha_1} \dots w_k^{-\alpha_k} \left(X^{\alpha_1, \dots, \alpha_k} \right)^* w_1^{\alpha_1+i_1} \dots w_k^{\alpha_k+i_k},$$

for certain non-negative integers i_1, \dots, i_k .

If this operator also has coefficients that are polynomials in $x, y, \alpha_1, \dots, \alpha_k$, then:

$$\begin{aligned} & \iint_R p\left(Y_1^{\alpha_1}, \dots, Y_k^{\alpha_k}\right) w_1^{\alpha_1} \dots w_k^{\alpha_k} dx dy \\ &= \iint_R \left(X_1^{\alpha_1}, \dots, X_k^{\alpha_k}\right) q w_1^{\alpha_1+i_1} \dots w_k^{\alpha_k+i_k} dx dy, \end{aligned}$$

for any two polynomials p and q , and for all real $\alpha_1, \dots, \alpha_k$ such that

$$\iint_R w_1^{\alpha_1} \dots w_k^{\alpha_k} dx dy < \infty.$$

PROOF. For sufficiently large $\alpha_1, \alpha_2, \dots, \alpha_k$ the equality follows from partial integration because the function $w_1^{\alpha_1} \dots w_k^{\alpha_k}$ and its partial derivatives up to a certain order are zero at the boundary ∂R . By analytic continuation the equality follows for all $\alpha_1, \dots, \alpha_k$ such that

$$\iint_R w_1^{\alpha_1} \dots w_k^{\alpha_k} dx dy < \infty. \quad \square$$

Rewriting this lemma for $E_-^{\alpha, \beta}$ and $E_+^{\alpha, \beta, \gamma}$ we obtain:

$$\iint_R p\left(E_+^{\alpha, \beta, \gamma} q\right) \mu^{\alpha, \beta, \gamma}(u, v) du dv$$

(4.8)

$$= \iint_R \left(E_-^{\alpha, \beta} p\right) q \mu^{\alpha, \beta, \gamma}(u, v) du dv,$$

for any two polynomials p and q .

From lemma 4.1, formula (4.8) and definition 2.1 it can be proved that:

COROLLARY.

$$(4.9) \quad E_-^{\alpha, \beta} P_{n, k}^{\alpha, \beta, \gamma}(u, v) = \begin{cases} (n-k)(n+k+\alpha+\beta+1) p_{n-1, k}^{\alpha, \beta, \gamma+1}(u, v) & \text{if } n > k \\ 0 & \text{if } n = k \end{cases}$$

and

$$(4.10) \quad E_+^{\alpha, \beta, \gamma} p_{n-1, k}^{\alpha, \beta, \gamma+1}(u, v) = (n-k+2\gamma+1)(n+k+\alpha+\beta+2\gamma+2) p_{n, k}^{\alpha, \beta, \gamma}(u, v)$$

if $n > k$

(cf. the proof of theorem 5.4 in [4]).

Let us define the fourth order operator:

$$(4.11) \quad D_3^{\alpha, \beta, \gamma} = E_+^{\alpha, \beta, \gamma} \circ E_-^{\alpha, \beta}.$$

The polynomials $p_{n, k}^{\alpha, \beta, \gamma}(u, v)$ are eigenfunctions of $D_3^{\alpha, \beta, \gamma}$:

$$(4.12) \quad D_3^{\alpha, \beta, \gamma} p_{n, k}^{\alpha, \beta, \gamma}(u, v) = (n-k)(n-k+2\gamma+1)(n+k+\alpha+\beta+1)(n+k+\alpha+\beta+2\gamma+2) p_{n, k}^{\alpha, \beta, \gamma}(u, v).$$

Hence $D_3^{\alpha, \beta, \gamma}$ can be uniquely expressed as a polynomial in $D_1^{\alpha, \beta, \gamma}$ and $D_2^{\alpha, \beta, \gamma}$ (cf. [4, theorem 6.5]). By considering the eigenvalues it is clear that

$$(4.13) \quad D_3^{\alpha, \beta, \gamma} = (D_1^{\alpha, \beta, \gamma})^2 - 4D_2^{\alpha, \beta, \gamma} - (2\gamma+1)(\alpha+\beta+1)D_1^{\alpha, \beta, \gamma}.$$

5. A RODRIGUES-TYPE FORMULA

Using (2.6) and (4.10), $p_{n, k}^{\alpha, \beta, \gamma}(u, v)$ can be expressed in terms of polynomials of lower degree.

In (2.6) and (4.10) we write $D_+^{\alpha, \beta, \gamma}$ and $E_+^{\alpha, \beta, \gamma}$ respectively as:

$$(2.11) \quad D_+^{\alpha, \beta, \gamma} = \{\mu^{\alpha, \beta, \gamma}(u, v)\}^{-1} (D_-^\gamma)^* \circ \mu^{\alpha+1, \beta+1, \gamma}(u, v)$$

and

$$(4.7) \quad E_+^{\alpha, \beta, \gamma} = \{\mu^{\alpha, \beta, \gamma}(u, v)\}^{-1} (E_-^{\alpha, \beta})^* \circ \mu^{\alpha, \beta, \gamma+1}(u, v).$$

An $(n-k)$ -fold application of (4.10) and a k -fold application of (2.6) to $p_{0, 0}^{\alpha+k, \beta+k, \gamma+n-k}(u, v) \equiv 1$ yields:

$$(5.1) \left\{ \begin{aligned} & (k+\alpha+\beta+1)_k (n+\alpha+\beta+\gamma+\frac{3}{2})_k (n-k+2\gamma+1)_{n-k} (n+k+\alpha+\beta+2\gamma+2)_{n-k} p_{n,k}^{\alpha,\beta,\gamma}(u,v) = \\ & = (1-u+v)^{-\alpha} (1+u+v)^{-\beta} (u^2-4v)^{-\gamma} \left\{ \frac{\partial^2}{\partial u^2} + u \frac{\partial^2}{\partial u \partial v} + v \frac{\partial^2}{\partial v^2} - (\gamma - \frac{3}{2}) \frac{\partial}{\partial v} \right\}^k \circ \\ & \circ \left\{ u \frac{\partial^2}{\partial u^2} + 2(v+1) \frac{\partial^2}{\partial u \partial v} + u \frac{\partial^2}{\partial v^2} - (\beta-\alpha) \frac{\partial}{\partial v} - (\alpha+\beta+2k-2) \frac{\partial}{\partial u} \right\}^{n-k} \\ & (1-u+v)^{\alpha+k} (1+u+v)^{\beta+k} (u^2-4v)^{\gamma+n-k}. \end{aligned} \right.$$

This is a Rodrigues-type formula for $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$. So far it is the only "explicit" expression for $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ in the case of general α, β, γ .

6. THE QUADRATIC NORM OF $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$

The quadratic norm $h_{n,k}^{\alpha,\beta,\gamma}$ of the polynomial $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ is defined by:

$$(6.1) \quad h_{n,k}^{\alpha,\beta,\gamma} = \iint_R \{p_{n,k}^{\alpha,\beta,\gamma}(u,v)\}^2 u^{\alpha,\beta,\gamma}(u,v) du dv.$$

The explicit value of $h_{n,k}^{\alpha,\beta,\gamma}$ is important for calculating the coefficients in Fourier expansions with respect to the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ (cf. section 9).

From (2.13) and (4.8) we obtain the following recurrence relations for $h_{n,k}^{\alpha,\beta,\gamma}$:

$$(6.2) \quad h_{n,k}^{\alpha,\beta,\gamma} = \frac{k(n+\gamma+\frac{1}{2})}{(k+\alpha+\beta+1)(n+\alpha+\beta+\gamma+\frac{3}{2})} h_{n-1,k-1}^{\alpha+1,\beta+1,\gamma}$$

and

$$(6.3) \quad h_{n,k}^{\alpha,\beta,\gamma} = \frac{(n-k)(n+k+\alpha+\beta+1)}{(n-k+2\gamma+1)(n+k+\alpha+\beta+2\gamma+2)} h_{n-1,k}^{\alpha,\beta,\gamma+1}.$$

By repeated application of (6.2) and (6.3) we find:

$$(6.4) \quad h_{n,k}^{\alpha,\beta,\gamma} = \frac{k!(n-k)!(n-k+\gamma+\frac{3}{2})_k (2k+\alpha+\beta+2)_{n-k}}{(k+\alpha+\beta+1)_k (n+\alpha+\beta+\gamma+\frac{3}{2})_k (n-k+2\gamma+1)_{n-k} (n+k+\alpha+\beta+2\gamma+2)_{n-k}} h_{0,0}^{\alpha+k,\beta+k,\gamma+n-k}.$$

LEMMA 6.1.

$$(6.5) \quad h_{0,0}^{\alpha,\beta,\gamma} = \frac{2^{2\alpha+2\beta+4\gamma+3}}{\sqrt{\pi}} \frac{\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(\gamma+1)}{\Gamma(\alpha+\beta+\gamma+\frac{5}{2})} \frac{\Gamma(\alpha+\gamma+\frac{3}{2})\Gamma(\beta+\gamma+\frac{3}{2})}{\Gamma(\alpha+\beta+2\gamma+3)}.$$

PROOF. $p_{0,0}^{\alpha,\beta,\gamma}(u,v) \equiv 1,$

thus:

$$h_{0,0}^{\alpha,\beta,\gamma} = \iint_R (1-u+v)^\alpha (1+u+v)^\beta (u^2-4v)^\gamma du dv.$$

This transforms under the substitution

$$u = x + y, \quad v = xy$$

into

$$h_{0,0}^{\alpha,\beta,\gamma} = \int_{x=-1}^1 \left\{ \int_{y=-1}^x (1-y)^\alpha (1+y)^\beta (x-y)^{2\gamma+1} dy \right\} (1-x)^\alpha (1+x)^\beta dx.$$

By making the substitution $t = (1+x)^{-1}(1+y)$ and using

$$\int_0^1 t^{b-1} (1-t)^{c-b-1} (1-tz)^{-a} dt = \frac{\Gamma(b)\Gamma(c-b)}{\Gamma(c)} {}_2F_1(a, b; c; z)$$

it follows that:

$$\begin{aligned} h_{0,0}^{\alpha,\beta,\gamma} &= \\ &= 2^\alpha \frac{\Gamma(\beta+1)\Gamma(2\gamma+2)}{\Gamma(\beta+2\gamma+3)} \int_{-1}^1 (1-x)^\alpha (1+x)^{2\beta+2\gamma+2} {}_2F_1(-\alpha, \beta+1; \beta+2\gamma+3; \frac{1+x}{2}) dx = \end{aligned}$$

$$\begin{aligned}
&= 2^{2\alpha+2\beta+2\gamma+3} \frac{\Gamma(\beta+1)\Gamma(2\gamma+2)}{\Gamma(\beta+2\gamma+3)} \int_0^1 (1-s)^\alpha s^{2\beta+2\gamma+2} {}_2F_1(-\alpha, \beta+1; \beta+2\gamma+3; s) ds = \\
&= 2^{2\alpha+2\beta+2\gamma+3} \frac{\Gamma(\alpha+1)\Gamma(\beta+1)\Gamma(2\gamma+2)\Gamma(2\beta+2\gamma+3)}{\Gamma(\beta+2\gamma+3)\Gamma(\alpha+\beta+2\gamma+4)} {}_3F_2(-\alpha, \beta+1, 2\beta+2\gamma+3; \\
&\quad ; \beta+2\gamma+3, \alpha+2\beta+2\gamma+4; 1).
\end{aligned}$$

This ${}_3F_2$ function is of type ${}_3F_2(a, b, c; l+a-b, l+a-c; 1)$ with $a = 2\beta+2\gamma+3$, $b = \beta+1$ and $c = -\alpha$ and so the theorem of Dixon can be applied (see BAILEY [1, Chap. 3.1] or SLATER [5, (2.3.3)]). This proves the lemma. \square

COROLLARY. The quadratic norm $h_{n,k}^{\alpha, \beta, \gamma}$ is equal to:

$$(6.6) \quad h_{n,k}^{\alpha, \beta, \gamma} = \frac{2^{4n+2\alpha+2\beta+4\gamma+3} k! (n-k)! (n-k+\frac{3}{2})_k (2k+\alpha+\beta+2)_{n-k}}{\sqrt{\pi} (k+\alpha+\beta+1)_k (n+\alpha+\beta+\gamma+\frac{3}{2})_k (n-k+2\gamma+1)_{n-k} (n+k+\alpha+\beta+2\gamma+2)_{n-k}} \cdot \frac{\Gamma(k+\alpha+1)\Gamma(k+\beta+1)\Gamma(n-k+\gamma+1)\Gamma(n+\alpha+\gamma+\frac{3}{2})\Gamma(n+\beta+\gamma+\frac{3}{2})}{\Gamma(n+k+\alpha+\beta+\gamma+\frac{5}{2})\Gamma(2n+\alpha+\beta+2\gamma+3)}.$$

7. THE VALUE OF $p_{n,k}^{\alpha, \beta, \gamma}(2, 1)$

It is possible to find the value of $p_{n,k}^{\alpha, \beta, \gamma}(2, 1)$ by using the operators $D_+^{\alpha, \beta, \gamma}$ and $E_+^{\alpha, \beta, \gamma}$. It is of interest to know this value because of the hypothesis that for $\alpha \geq \beta \geq -\frac{1}{2}$ and $\gamma \geq -\frac{1}{2}$ the inequality

$$|p_{n,k}^{\alpha, \beta, \gamma}(u, v)| \leq p_{n,k}^{\alpha, \beta, \gamma}(2, 1)$$

is valid. This hypothesis was proved for $\gamma = -\frac{1}{2}$. If $\gamma \geq -\frac{1}{2}$ then it is true if $\alpha = \beta = -\frac{1}{2}$. Further it holds for the polynomials $p_{n,n}^{\alpha, \beta+\frac{1}{2}}(u, v)$, $p_{n,n-1}^{\alpha, \alpha+\frac{1}{2}}(u, v)$ and $p_{n,0}^{+\frac{1}{2}, -\frac{1}{2}, \gamma}(u, v)$.

Considering (2.9) and (4.7) we obtain the following equalities:

$$(D_+^{\alpha, \beta, \gamma} p)(2, 1) = 4(\alpha+1)(\alpha+\gamma+\frac{3}{2})p(2, 1)$$

and

$$(E_+^{\alpha, \beta, \gamma} p)(2, 1) = 8(\gamma+1)(\alpha+\gamma+\frac{3}{2})p(2, 1) ,$$

for any polynomial $p(u, v)$.

Hence:

$$(7.1) \quad p_{n,k}^{\alpha, \beta, \gamma}(2, 1) = \frac{4(\alpha+1)(\alpha+\gamma+\frac{3}{2})}{(k+\alpha+\beta+1)(n+\alpha+\beta+\gamma+\frac{3}{2})} p_{n-1, k-1}^{\alpha+1, \beta+1, \gamma}(2, 1)$$

and

$$(7.2) \quad p_{n,k}^{\alpha, \beta, \gamma}(2, 1) = \frac{8(\gamma+1)(\alpha+\gamma+\frac{3}{2})}{(n-k+2\gamma+1)(n+k+\alpha+\beta+2\gamma+2)} p_{n-1, k}^{\alpha, \beta, \gamma+1}(2, 1)$$

From (7.1), (7.2) and $p_{0,0}^{\alpha, \beta, \gamma}(u, v) \equiv 1$ it follows that

$$(7.3) \quad p_{n,k}^{\alpha, \beta, \gamma}(2, 1) = \frac{2^{3n-k} (\alpha+1)_k (\gamma+1)_{n-k} (\alpha+\gamma+\frac{3}{2})_n}{(k+\alpha+\beta+1)_k (n+\alpha+\beta+\gamma+\frac{3}{2})_k (n-k+2\gamma+1)_{n-k} (n+k+\alpha+\beta+2\gamma+2)_{n-k}}$$

Remark. The relation

$$p_{n,k}^{\alpha, \beta, \gamma}(-2, 1) = (-1)^{n-k} p_{n,k}^{\alpha, \beta, \gamma}(2, 1) \quad (10.1)$$

immediately gives the value of $p_{n,k}^{\alpha, \beta, \gamma}(-2, 1)$.

8. THE COEFFICIENTS IN THE POWER SERIES OF $p_{n,k}^{\alpha, \beta, \gamma}(u, v)$

For the coefficients $a_{ij}(n, k, \alpha, \beta, \gamma)$ in the expansion

$$(8.1) \quad p_{n,k}^{\alpha, \beta, \gamma}(u, v) = \sum_{(i,j) \leq (n,k)} a_{i,j}(n, k, \alpha, \beta, \gamma) u^{i-j} v^j$$

the following theorem holds:

THEOREM 8.1. $a_{i,j}(n,k,\alpha,\beta,\gamma) = 0$ if $i+j > n+k$ or $i > n$.

At this point it is useful to define the following partial ordering for $N = \{(n,k) | n \geq k \geq 0, n,k \in \mathbb{N}\}$:

$$(8.2) \quad (i,j) \prec (n,k) \text{ iff } i \leq n \text{ and } i+j \leq n+k.$$

Thus

$$(i,j) \prec (n,k) \iff ((i,j) \leq (n,k) \wedge i+j \leq n+k).$$

Theorem 8.1 is equivalent to:

$$(8.3) \quad p_{n,k}^{\alpha,\beta,\gamma}(u,v) = \sum_{(i,j) \prec (n,k)} a_{i,j}(n,k,\alpha,\beta,\gamma) u^{i-j} v^j.$$

PROOF OF THEOREM 8.1. The second statement is a consequence of the definition of $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$, because if $i > n$ then $(i,j) > (n,k)$. The first statement is trivially true for the polynomials $p_{n,n}^{\alpha,\beta,\gamma}(u,v)$ because in that case $i+j > n+n$ implies $i > n$. It is clear from (4.7) that

$$E_+^{\alpha,\beta,\gamma} u^{m-l} v^l = \sum_{(i,j) \prec (m+1,l)} c_{i,j} u^{i-j} v^j,$$

for certain constants $c_{i,j}$. By repeated application of the operators E_+ to $p_{n,n}^{\alpha,\beta,\gamma}(u,v)$ and by using (4.10) the theorem follows. \square

Corrollaries of theorem 8.1 are the next two theorems:

THEOREM 8.2. *Let*

$$(i) \quad p(u,v) = \sum_{(m,l) \prec (n,k)} c_{m,l} u^{m-l} v^l,$$

for certain constants $c_{m,l}$, with $c_{n,k} = 1$, and

$$(ii) \quad \iint_{\mathbb{R}} p(u,v) u^{m-l} v^l \mu^{\alpha,\beta,\gamma}(u,v) du dv = 0 \text{ if } (m,l) \prec \neq (n,k),$$

then

$$p(u, v) = p_{n, k}^{\alpha, \beta, \gamma}(u, v).$$

THEOREM 8.3. *Let*

$$(i) \quad p(u, v) = \sum_{(m, \ell) \prec (n, k)} c_{m, \ell} u^{m-\ell} v^{\ell},$$

for certain constants $c_{m, \ell}$, with $c_{n, k} = 1$,

and

$$(ii) \quad D_1^{\alpha, \beta, \gamma} p(u, v) = \lambda p(u, v) \quad \text{for some } \lambda \in \mathbb{R},$$

then

$$p(u, v) = p_{n, k}^{\alpha, \beta, \gamma}(u, v)$$

and

$$\lambda = -[n(n+\alpha+\beta+2\gamma+2)+k(k+\alpha+\beta+1)].$$

PROOF OF THEOREM 8.2. From (i) it follows that $p(u, v)$ can be uniquely expressed as:

$$p(u, v) = \sum_{(m, \ell) \prec (n, k)} c'_{m, \ell} p_{m, \ell}^{\alpha, \beta, \gamma}(u, v) \quad \text{with } c'_{n, k} = 1.$$

Then (ii) yields:

$$c'_{m, \ell} = (h_{m, \ell}^{\alpha, \beta, \gamma})^{-1} \iint_{\mathbb{R}} p(u, v) p_{m, \ell}^{\alpha, \beta, \gamma}(u, v) \mu^{\alpha, \beta, \gamma}(u, v) du dv = 0$$

$$\text{if } (m, \ell) \prec_{\neq} (n, k).$$

This proves the theorem. \square

For the proof of theorem 8.3 we need the following lemma:

LEMMA 8.1. If $(m, \ell) \not\prec (n, k)$ then $\lambda_{m, \ell} \neq \lambda_{n, k}$, with

$$\lambda_{m, \ell} = -[m(m+\alpha+\beta+2\gamma+2)+\ell(\ell+\alpha+\beta+1)].$$

PROOF. The parameters α, β, γ satisfy $\alpha, \beta, \gamma > -1$, $\alpha + \gamma + \frac{3}{2}$, $\beta + \gamma + \frac{3}{2} > 0$.

Suppose that $(m, \ell) \not\prec (n, k)$ and $\lambda_{m, \ell} = \lambda_{n, k}$. Then

$$(n-m)(n+m+\alpha+\beta+2\gamma+2) = (\ell-k)(\ell+k+\alpha+\beta+1).$$

and the factors $n + m + \alpha + \beta + 2\gamma + 2$ and $\ell + k + \alpha + \beta + 1$ are positive. Hence $n - m > 0$ and $\ell - k > 0$. Observe that $n + m + \alpha + \beta + 2\gamma + 2 \geq 2\ell + 1 + \alpha + \beta + 2\gamma + 2 > 2\ell + \alpha + \beta + 1 \geq \ell + k + \alpha + \beta + 1$. Thus $n - m < \ell - k$ contradicting the hypothesis $(m, \ell) \not\prec (n, k)$. \square

PROOF OF THEOREM 8.3. From (i) it follows that $p(u, v)$ can be uniquely expressed as:

$$p(u, v) = \sum_{(m, \ell) \prec (n, k)} c'_{m, \ell} p_{m, \ell}^{\alpha, \beta, \gamma}(u, v) \text{ with } c'_{n, k} = 1.$$

Then (ii) yields:

$$\sum_{(m, \ell) \prec (n, k)} \lambda c'_{m, \ell} p_{m, \ell}^{\alpha, \beta, \gamma}(u, v) = \sum_{(m, \ell) \prec (n, k)} \lambda_{m, \ell} c'_{m, \ell} p_{m, \ell}^{\alpha, \beta, \gamma}(u, v).$$

From $c'_{n, k} = 1$ it follows that

$$\lambda = \lambda_{n, k} = -[n(n+\alpha+\beta+2\gamma+2)+(k+\alpha+\beta+1)]$$

and from $\lambda_{n, k} c'_{m, \ell} = \lambda_{m, \ell} c'_{m, \ell}$ and lemma 8.1 it follows that $c'_{m, \ell} = 0$ for $(m, \ell) \not\prec (n, k)$ \square

Application of $D_1^{\alpha, \beta, \gamma}$ to (8.3) and comparison of the coefficients of equal powers of u and v give the following explicit values for some of the coefficients $a_{i, j}(n, k, \alpha, \beta, \gamma)$ in (8.3), which will be used in section 9 for

the computation of the coefficients in the recurrence relations:

$$(8.4a) \quad a_{n,k}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = 1,$$

$$(8.4b) \quad a_{n,k-1}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = -(\beta-\alpha)k/(2k+\alpha+\beta),$$

$$(8.4c) \quad a_{n,k-2}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = -\frac{1}{2}k(k-1)\{1-(\beta-\alpha)^2/(2k+\alpha+\beta)\}/(2k+\alpha+\beta-1),$$

$$(8.4d) \quad a_{n-1,k+1}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = -(n-k)(n-k-1)/(n-k+\gamma-\frac{1}{2}),$$

$$(8.4e) \quad a_{n-1,k}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = \frac{(\beta-\alpha)(n-k)}{(2n+\alpha+\beta+2\gamma+1)} \left\{ \frac{2k(n-k+1)}{2k+\alpha+\beta} + \frac{(n-k-1)(k+1)}{n-k+\gamma-\frac{1}{2}} - 2 \right\},$$

$$(8.4f) \quad \left\{ \begin{array}{l} -2(n+k+\alpha+\beta+\gamma+\frac{1}{2})a_{n-1,k-1}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = \\ = (\beta-\alpha)ka_{n-1,k}^{\alpha,\beta,\gamma} + k(k+1)a_{n-1,k+1}^{\alpha,\beta,\gamma} + 2(n-k+2)(n-k+1)a_{n,k-2}^{\alpha,\beta,\gamma} \\ + 2(\beta-\alpha)(n-k+1)a_{n,k-1}^{\alpha,\beta,\gamma} + 2(n-2k-\gamma+\frac{1}{2}), \end{array} \right.$$

$a_{n,k-3}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma)$ and $a_{n,k-4}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma)$ do not depend on n and γ .

9. THE RECURRENCE RELATIONS

For a further analysis of the polynomials $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ it is useful to have formulas for the series expansions of $u p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ and $v p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ in terms of $p_{i,j}^{\alpha,\beta,\gamma}(u,v)$. These formulas give $p_{n+1,k}^{\alpha,\beta,\gamma}(u,v)$ and $p_{n+1,k+1}^{\alpha,\beta,\gamma}(u,v)$ as linear combinations of lower degree polynomials.

Case I: Expansion of $u p_{n,k}^{\alpha,\beta,\gamma}(u,v)$. Consider the following equality:

$$(9.1) \quad \begin{aligned} u p_{n,k}^{\alpha,\beta,\gamma}(u,v) &= \sum_{(m,\ell) \prec (n,k)} a_{m,\ell} u^{m-\ell+1} v^\ell = \\ &= \sum_{(m,\ell) \prec (n+1,k)} a'_{m,\ell} u^{m-\ell} v^\ell = \\ &= \sum_{(m,\ell) \prec (n+1,k)} b_{m,\ell}(n,k,\alpha,\beta,\gamma) p_{m,\ell}^{\alpha,\beta,\gamma}(u,v), \end{aligned}$$

with

$$(9.2) \quad b_{m,\ell}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = \{h_{m,\ell}^{\alpha,\beta,\gamma}\}^{-1} \iint_R u p_{n,k}^{\alpha,\beta,\gamma}(u,v) p^{\alpha,\beta,\gamma}(u,v) \mu^{\alpha,\beta,\gamma}(u,v) du dv.$$

From symmetry it follows that:

$$(9.3) \quad b_{m,\ell}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = h_{n,k}^{\alpha,\beta,\gamma} \{h_{m,\ell}^{\alpha,\beta,\gamma}\}^{-1} b_{n,k}^{\alpha,\beta,\gamma}(m,\ell,\alpha,\beta,\gamma).$$

Hence $b_{m,\ell}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) \neq 0$ only if $(m,\ell) \succ (n-1,k)$. And so the summation in (9.1) atmost runs through $(m,\ell) \in \{(n+1,k), (n+1,k-1), (n+1,k-2), (n,k+1), (n,k), (n,k-1), (n-1,k+2), (n-1,k+1), (n-1,k)\}$. The coefficients can be computed by means of (8.4), (9.1) and (9.3). The coefficients $b_{n+1,k-1}^{\alpha,\beta,\gamma}$, $b_{n-1,k+1}^{\alpha,\beta,\gamma}$, $b_{n+1,k-2}^{\alpha,\beta,\gamma}$ and $b_{n-1,k+2}^{\alpha,\beta,\gamma}$ turn out to be zero.

For the five remaining coefficients in (9.1) we obtain:

$$(9.4a) \quad b_{n+1,k}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = 1,$$

$$(9.4b) \quad b_{n-1,k}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = \frac{4(n+\gamma+\frac{1}{2})(n+\alpha+\gamma+\frac{1}{2})(n+\beta+\gamma+\frac{1}{2})(n+\alpha+\beta+\gamma+\frac{1}{2})}{(2n+\alpha+\beta+2\gamma)_3(2n+\alpha+\beta+2\gamma+1)}.$$

$$\cdot \frac{(n-k)(n-k+2\gamma)(n+k+\alpha+\beta+1)(n+k+\alpha+\beta+2\gamma+1)}{(n-k+\gamma-\frac{1}{2})(n-k+\gamma+\frac{1}{2})(n+k+\alpha+\beta+\gamma+\frac{1}{2})(n+k+\alpha+\beta+\gamma+\frac{3}{2})},$$

$$(9.4c) \quad b_{n,k+1}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = \frac{(n-k)(n-k+2\gamma)}{(n-k+\gamma-\frac{1}{2})(n-k+\gamma+\frac{1}{2})},$$

$$(9.4d) \quad b_{n,k-1}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = \frac{4k(k+\alpha)(k+\beta)(k+\alpha+\beta)(n+k+\alpha+\beta+1)(n+k+\alpha+\beta+2\gamma+1)}{(2k+\alpha+\beta-1)_3(2k+\alpha+\beta)(n+k+\alpha+\beta+\gamma+\frac{1}{2})(n+k+\alpha+\beta+\gamma+\frac{3}{2})},$$

$$(9.4e) \quad b_{n,k}^{\alpha,\beta,\gamma}(n,k,\alpha,\beta,\gamma) = (\beta-\alpha)(\alpha+\beta) \left\{ \frac{1}{(2n+\alpha+\beta+2\gamma+1)(2n+\alpha+\beta+2\gamma+3)} + \frac{(2n+\alpha+\beta+2)(2n+\alpha+\beta+4\gamma+2)}{(2n+\alpha+\beta+2\gamma+1)(2n+\alpha+\beta+2\gamma+3)(2k+\alpha+\beta)(2k+\alpha+\beta+2)} \right\}.$$

If we define

$$(9.5) \quad p_{n,k}^{\alpha,\beta,\gamma}(u,v) \equiv 0 \text{ if } n < k \text{ or if } k < 0$$

then the following five-term formula holds for $up_{n,k}^{\alpha,\beta,\gamma}(u,v)$ for all $n \geq k \geq 0$:

$$(9.6) \quad \left\{ \begin{array}{l} up_{n,k}^{\alpha,\beta,\gamma}(u,v) = p_{n+1,k}^{\alpha,\beta,\gamma}(u,v) + b_{n,k+1}(n,k,\alpha,\beta,\gamma)p_{n,k+1}^{\alpha,\beta,\gamma}(u,v) + \\ + b_{n,k}(n,k,\alpha,\beta,\gamma)p_{n,k}^{\alpha,\beta,\gamma}(u,v) + b_{n,k-1}(n,k,\alpha,\beta,\gamma)p_{n,k-1}^{\alpha,\beta,\gamma}(u,v) + \\ + b_{n-1,k}(n,k,\alpha,\beta,\gamma)p_{n-1,k}^{\alpha,\beta,\gamma}(u,v), \end{array} \right.$$

with $b_{m,\ell}(n,k,\alpha,\beta,\gamma)$ given by (9.4).

It follows that:

$$(9.7) \quad \begin{aligned} p_{n+1,k}^{\alpha,\beta,\gamma}(u,v) &= -b_{n,k+1}(n,k,\alpha,\beta,\gamma)p_{n,k+1}^{\alpha,\beta,\gamma}(u,v) + (u - b_{n,k}(n,k,\alpha,\beta,\gamma))p_{n,k}^{\alpha,\beta,\gamma}(u,v) \\ &\quad - b_{n,k-1}(n,k,\alpha,\beta,\gamma)p_{n,k-1}^{\alpha,\beta,\gamma}(u,v) - b_{n-1,k}(n,k,\alpha,\beta,\gamma)p_{n-1,k}^{\alpha,\beta,\gamma}(u,v) \end{aligned}$$

if $n \geq k \geq 0$.

Remark. By application of the quadratic transformation formulas (10.5) and (10.6) to (9.6), repeated application of D_-^Y and analytic continuation, it can be proved that

$$(9.8) \quad p_{n,k}^{\alpha,\beta,\gamma}(u,v) = p_{n,k}^{\alpha,\beta+1,\gamma}(u,v) + Ap_{n,k-1}^{\alpha,\beta+1,\gamma}(u,v) + Bp_{n-1,k}^{\alpha,\beta+1,\gamma}(u,v) + Cp_{n-1,k-1}^{\alpha,\beta+1,\gamma}(u,v)$$

if $n \geq k \geq 0$,

with A, B and C being functions of n, k, α, β and γ to be determined from the coefficients of $u^{m-\ell}v^\ell$.

Case II: Expansion of $vp_{n,k}^{\alpha,\beta,\gamma}(u,v)$. Consider the equality:

$$(9.9) \quad vp_{n,k}^{\alpha,\beta,\gamma}(u,v) = \sum_{(m,\ell) \prec (n+1,k+1)} c_{m,\ell}(n,k,\alpha,\beta,\gamma) p_{m,\ell}^{\alpha,\beta,\gamma}(u,v),$$

with

$$(9.10) \quad c_{m,\ell}(n,k,\alpha,\beta,\gamma) = \{h_{m,\ell}^{\alpha,\beta,\gamma}\}^{-1} \iint_R vp_{n,k}^{\alpha,\beta,\gamma}(u,v) p_{m,\ell}^{\alpha,\beta,\gamma}(u,v) u^{\alpha,\beta,\gamma}(u,v) dudv.$$

From symmetry it follows that:

$$(9.11) \quad c_{m,\ell}(n,k,\alpha,\beta,\gamma) = h_{n,k}^{\alpha,\beta,\gamma} \{h_{m,\ell}^{\alpha,\beta,\gamma}\}^{-1} c_{n,k}(m,\ell,\alpha,\beta,\gamma).$$

Hence $c_{m,\ell}(n,k,\alpha,\beta,\gamma) \neq 0$ only if $(m,\ell) \succ (n-1,k-1)$. And so the summation in (9.9) atmost runs through $(m,\ell) \in \{(n+1,k+1), (n+1,k), (n+1,k-1), (n+1,k-2), (n+1,k-3), (n,k+2), (n,k+1), (n,k), (n,k-1), (n,k-2), (n-1,k+3), (n-1,k+2), (n-1,k+1), (n-1,k), (n-1,k-1)\}$. The coefficients can be computed by means of (8.4), (9.9) and (9.10), and by comparison with the case $\gamma = -\frac{1}{2}$. The coefficients $c_{n+1,k-2}, c_{n-1,k+2}, c_{n+1,k-3}, c_{n-1,k+3}, c_{n,k+2}$ and $c_{n,k-2}$ turn out to be zero.

For the nine remaining coefficients in (9.9) we obtain:

$$(9.12a) \quad c_{n+1,k+1}(n,k,\alpha,\beta,\gamma) = 1,$$

$$(9.12b) \quad c_{n-1,k-1}(n,k,\alpha,\beta,\gamma) = \frac{2^4 (n+\gamma+\frac{1}{2})(n+\alpha+\gamma+\frac{1}{2})(n+\beta+\gamma+\frac{1}{2})(n+\alpha+\beta+\gamma+\frac{1}{2})}{(2n+\alpha+\beta+2\gamma)_3 (2n+\alpha+\beta+2\gamma+1)} \\ \cdot \frac{k(k+\alpha)(k+\beta)(k+\alpha+\beta)(n+k+\alpha+\beta)_2 (n+k+\alpha+\beta+2\gamma)_2}{(2k+\alpha+\beta-1)_3 (2k+\alpha+\beta)(n+k+\alpha+\beta+\gamma-\frac{1}{2})_3 (n+k+\alpha+\beta+\gamma+\frac{1}{2})_2},$$

$$(9.12c) \quad c_{n+1,k}(n,k,\alpha,\beta,\gamma) = \frac{(\beta-\alpha)(\alpha+\beta)}{(2k+\alpha+\beta)(2k+\alpha+\beta+2)},$$

$$(9.12d) \quad c_{n-1,k}(n,k,\alpha,\beta,\gamma) = \frac{4(\beta-\alpha)(\alpha+\beta)(n+\gamma+\frac{1}{2})(n+\alpha+\gamma+\frac{1}{2})(n+\beta+\gamma+\frac{1}{2})}{(2k+\alpha+\beta)(2k+\alpha+\beta+2)(2n+\alpha+\beta+2\gamma)_3} \\ \cdot \frac{(n+\alpha+\beta+\gamma+\frac{1}{2})(n-k)(n-k+2\gamma)(n+k+\alpha+\beta+1)(n+k+\alpha+\beta+2\gamma+1)}{(2n+\alpha+\beta+2\gamma+1)(n-k+\gamma-\frac{1}{2})(n-k+\gamma+\frac{1}{2})(n+k+\alpha+\beta+\gamma+\frac{1}{2})_2},$$

$$(9.12e) \quad c_{n+1,k-1}(n,k,\alpha,\beta,\gamma) = \frac{4k(k+\alpha)(k+\beta)(k+\alpha+\beta)}{(2k+\alpha+\beta-1)_3 (2k+\alpha+\beta)},$$

$$(9.12f) \quad c_{n-1,k+1}(n,k,\alpha,\beta,\gamma) = \frac{4(n+\gamma+\frac{1}{2})(n+\alpha+\gamma+\frac{1}{2})(n+\beta+\gamma+\frac{1}{2})(n+\alpha+\beta+\gamma+\frac{1}{2})}{(2n+\alpha+\beta+2\gamma)_3 (2n+\alpha+\beta+2\gamma+1)} \\ \cdot \frac{(n-k-1)_2 (n-k+2\gamma-1)_2}{(n-k+\gamma-\frac{3}{2})_3 (n-k+\gamma-\frac{1}{2})_2},$$

$$(9.12g) \quad c_{n,k+1}^{(n,k,\alpha,\beta,\gamma)} = \frac{(\beta-\alpha)(\alpha+\beta)(n-k)(n-k+2\gamma)}{(2n+\alpha+\beta+2\gamma+1)(2n+\alpha+\beta+2\gamma+3)(n-k+\gamma-\frac{1}{2})(n-k+\gamma+\frac{1}{2})},$$

$$(9.12h) \quad c_{n,k-1}^{(n,k,\alpha,\beta,\gamma)} = \frac{4(\beta-\alpha)(\alpha+\beta)k(k+\alpha)(k+\beta)(k+\alpha+\beta)}{(2n+\alpha+\beta+2\gamma+1)(2n+\alpha+\beta+2\gamma+3)(2k+\alpha+\beta-1)_3(2k+\alpha+\beta)} \cdot \frac{(n+k+\alpha+\beta+1)(n+k+\alpha+\beta+2\gamma+1)}{(n+k+\alpha+\beta+\gamma+\frac{1}{2})_2},$$

$$(9.12i) \quad c_{n,k}^{(n,k,\alpha,\beta,\gamma)} = a_{n-1,k-1}^{(n,k,\alpha,\beta,\gamma)} - a_{n,k}^{(n+1,k+1,\alpha,\beta,\gamma)}$$

$$-c_{n+1,k}^{(n,k,\alpha,\beta,\gamma)} a_{n,k}^{(n+1,k,\alpha,\beta,\gamma)} - c_{n+1,k-1}^{(n,k,\alpha,\beta,\gamma)} a_{n,k}^{(n+1,k-1,\alpha,\beta,\gamma)}$$

$$-c_{n,k+1}^{(n,k,\alpha,\beta,\gamma)} a_{n,k}^{(n,k+1,\alpha,\beta,\gamma)}.$$

If $\gamma = -\frac{1}{2}$ then $c_{n,k}^{(n,k,\alpha,\beta,-\frac{1}{2})}$ is given by:

$$(9.12i)' \quad c_{n,k}^{(n,k,\alpha,\beta,-\frac{1}{2})} = \frac{(\alpha+\beta)^2(\beta-\alpha)^2}{(2n+\alpha+\beta)(2n+\alpha+\beta+2)(2k+\alpha+\beta)(2k+\alpha+\beta+2)}.$$

Formula (9.9) holds, with the coefficients given by (9.12), for all $n \geq k \geq 0$, where the convention (9.5) is used again.

Formulas (9.6) and (9.9) together give an algorithm for calculating $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$. If $n \neq k$ then $p_{n,k}^{\alpha,\beta,\gamma}(u,v)$ can be expressed in terms of lower degree polynomials by the five-term relation (9.6). If $n = k$ then (9.9) provides a six-term relation which expresses $p_{n,n}^{\alpha,\beta,\gamma}(u,v)$ in terms of lower degree polynomials.

10. A QUADRATIC TRANSFORMATION

The reflection $u \rightarrow -u$ maps the region R onto itself and transforms the weightfunction $\mu^{\alpha,\beta,\gamma}(u,v)$ into $\mu^{\beta,\alpha,\gamma}(u,v)$. Hence, in view of definition 2.1 the following equality holds:

$$(10.1) \quad p_{n,k}^{\alpha,\beta,\gamma}(-u,v) = (-1)^{n-k} p_{n,k}^{\beta,\alpha,\gamma}(u,v).$$

If $\alpha = \beta$, then (10.1) becomes:

$$(10.2) \quad p_{n,k}^{\alpha,\alpha,\gamma}(-u,v) = (-1)^{n-k} p_{n,k}^{\alpha,\alpha,\gamma}(u,v).$$

Formula (10.2) means that if $(n-k)$ is even, then $p_{n,k}^{\alpha,\alpha,\gamma}(u,v)$ is a polynomial in u^2 and v , and if $(n-k)$ is odd, then $u^{-1} p_{n,k}^{\alpha,\alpha,\gamma}(u,v)$ is a polynomial in u^2 and v .

Consider now the new variables:

$$(10.3) \quad u' = 2v, \quad v' = u^2 - 2v - 1.$$

These variables satisfy the following properties:

- (i) Each polynomial in u^2 and v is a polynomial in u' and v' .
 - (ii) The half region R given by $R \cap \{(u,v) | u > 0\}$ is mapped onto $\tilde{R} = \{(u',v') | (1+u'+v') > 0 \wedge (1-u'+v') > 0 \wedge ((u')^2 - 4v') > 0\}$.
 - (iii) If $(u,v) = (2,1)$ then $(u',v') = (2,1)$.
- (The transformation of variables $u' = -2v$ and $v' = u^2 - 2v - 1$ also satisfies (i) and (ii)).

From (10.3) we obtain:

$$(10.4) \quad \begin{cases} u = \sqrt{1+u'+v'}, & v = \frac{1}{2}u', \\ (1+u+v)(1-u+v) = \frac{1}{4}((u')^2 - 4v'), & u^2 - 4v = 1 - u' + v', \\ dudv = \frac{1}{4}(1+u'+v')^{-\frac{1}{2}} du' dv'. \end{cases}$$

If $\alpha = \beta$ the following quadratic transformation formulas hold:

THEOREM 10.1.

$$(10.5) \quad p_{n+k,n-k}^{\alpha,\alpha,\gamma}(u,v) = 2^{-n+k} p_{n,k}^{\gamma,-\frac{1}{2},\alpha}(u',v'),$$

and

$$(10.6) \quad u^{-1} p_{n+k+1,n-k}^{\alpha,\alpha,\gamma}(u,v) = 2^{-n+k} p_{n,k}^{\gamma,+\frac{1}{2},\alpha}(u',v'),$$

with u' and v' given by (10.3).

PROOF.
$$p_{n+k, n-k}^{\alpha, \alpha, \gamma}(u, v) = \sum_{(i, j) \prec (n+k, n-k)} a_{ij} u^{i-j} v^j.$$

If $(i-j)$ is odd then $a_{i, j} = 0$, so we can substitute $i - j = 2\ell$ and $i + j = 2m$. By (8.2) $(i, j) \prec (n+k, n-k)$ iff $(m, \ell) \prec (n, k)$. Hence:

$$\begin{aligned} p_{n+k, n-k}^{\alpha, \alpha, \gamma}(u, v) &= \sum_{(m, \ell) \prec (n, k)} a_{m, \ell}' (u^2)^\ell v^{m-\ell} \\ &= \sum_{(m, \ell) \prec (n, k)} a_{m, \ell}'' (u^2 - 2v - 1)^\ell (2v)^{m-\ell} \\ &= \sum_{(m, \ell) \prec (n, k)} a_{m, \ell}'' (u')^{m-\ell} (v')^\ell, \quad \text{with } a_{n, k}'' = 2^{-n+k}. \end{aligned}$$

With respect to the orthogonality the following holds:

$$\begin{aligned} 0 &= \iint_{\mathbb{R}} p_{n+k, n-k}^{\alpha, \alpha, \gamma}(u, v) (2v)^{m-\ell} (u^2 - 2v - 1)^\ell \mu^{\alpha, \alpha, \gamma}(u, v) \, dudv = \\ &= \text{const.} \iint_{\mathbb{R}} p_{n+k, n-k}^{\alpha, \alpha, \gamma}(u, v) (u')^{m-\ell} (v')^\ell \mu^{\gamma, -\frac{1}{2}, \alpha}(u', v') \, du' dv', \\ &\hspace{20em} \text{if } (m, \ell) \prec (n, k). \\ &\hspace{20em} \neq \end{aligned}$$

Application of theorem 8.2 proves (10.5). A similar proof can be given for (10.6). \square

If $(u, v) = (2, 1)$, then $(u'v') = (2, 1)$; hence (10.5) and (10.6) can also be written as:

$$(10.5)' \quad \frac{p_{n+k, n-k}^{\alpha, \alpha, \gamma}(u, v)}{p_{n+k, n-k}^{\alpha, \alpha, \gamma}(2, 1)} = \frac{p_{n, k}^{\gamma, -\frac{1}{2}, \alpha}(2v, u^2 - 2v - 1)}{p_{n, k}^{\gamma, -\frac{1}{2}, \alpha}(2, 1)},$$

and

$$(10.6)' \quad \frac{2u^{-1} p_{n+k+1, n-k}^{\alpha, \alpha, \gamma}(u, v)}{p_{n+k+1, n-k}^{\alpha, \alpha, \gamma}(2, 1)} = \frac{p_{n, k}^{\gamma, +\frac{1}{2}, \alpha}(2v, u^2 - 2v - 1)}{p_{n, k}^{\gamma, +\frac{1}{2}, \alpha}(2, 1)}.$$

Formulas (10.5) and (10.6) in combination with (2.15) and (3.2) give an explicit expression for the polynomials $p_{n, k}^{\alpha, \beta, \gamma}(u, v)$ if α and β are $+\frac{1}{2}$ or $-\frac{1}{2}$:

$$(10.7) \quad P_{n,k}^{-\frac{1}{2}, -\frac{1}{2}, \gamma}(2xy, x^2+y^2-1) = \begin{cases} 2^{n-k} \{p_{n+k}^{\gamma, \gamma}(x)p_{n-k}^{\gamma, \gamma}(y) + p_{n-k}^{\gamma, \gamma}(x)p_{n+k}^{\gamma, \gamma}(y)\} & \text{if } k > 0 \\ 2^n p_n^{\gamma, \gamma}(x)p_n^{\gamma, \gamma}(y) & \text{if } k = 0, \end{cases}$$

$$(10.8) \quad P_{n,k}^{+\frac{1}{2}, -\frac{1}{2}, \gamma}(2xy, x^2+y^2-1) = 2^{n-k} (x-y)^{-1} \{p_{n+k+1}^{\gamma, \gamma}(x)p_{n-k}^{\gamma, \gamma}(y) - p_{n-k}^{\gamma, \gamma}(x)p_{n+k+1}^{\gamma, \gamma}(y)\},$$

$$(10.9) \quad P_{n,k}^{-\frac{1}{2}, +\frac{1}{2}, \gamma}(2xy, x^2+y^2-1) = 2^{n-k} (x+y)^{-1} \{p_{n+k+1}^{\gamma, \gamma}(x)p_{n-k}^{\gamma, \gamma}(y) + p_{n-k}^{\gamma, \gamma}(x)p_{n+k+1}^{\gamma, \gamma}(y)\},$$

$$(10.10) \quad P_{n,k}^{+\frac{1}{2}, +\frac{1}{2}, \gamma}(2xy, x^2+y^2-1) = 2^{n-k} (x^2-y^2)^{-1} \{p_{n+k+2}^{\gamma, \gamma}(x)p_{n-k}^{\gamma, \gamma}(y) - p_{n-k}^{\gamma, \gamma}(x)p_{n+k+2}^{\gamma, \gamma}(y)\}.$$

REFERENCES

- [1] BAILEY, W.N., *Generalized hypergeometric series*, Cambridge Tracts in Math. and Math. Phys. 32, Cambridge University Press, Cambridge, 1935.
- [2] ERDELYI, A., W. MAGNUS, F. OBERHETTINGER & F.G. TRICOMI, *Higher transcendental functions, vol. II*, McGraw-Hill, New York, 1953.
- [3] KARLIN, S. & J. MCGREGOR, *Determinants of orthogonal polynomials*, Bull. Amer. Math. Soc. 68 (1962), 204-209.
- [4] KOORNWINDER, T.H., *Orthogonal polynomials in two variables which are eigenfunctions of two algebraically independent partial differential operators I, II*, Proc. Kon. Ned. Akad. Wetensch. A 77 = Indag. Math. 36 (1974), 48-66.
- [5] SLATER, L.J., *Generalized hypergeometric functions*, Cambridge University Press, Cambridge, 1966.
- [6] SZEGÖ, G., *Orthogonal polynomials*, Amer. Math. Soc. Colloquium Publications, vol. 23, Providence, Rhode Island, Third Ed., 1967.
- [7] WAERDEN, B.L. van der, *Algebra, Erster Teil*, Springer-Verlag, Berlin, Siebenten Auflage, 1966.

