

TW

**stichting
mathematisch
centrum**



TW

AFDELING TOEGEPASTE WISKUNDE

TW 131/71

NOVEMBER

T.H. KOORNWINDER
THE ADDITION FORMULA FOR JACOBI POLYNOMIALS
I SUMMARY OF RESULTS

2e boerhaavestraat 49 amsterdam

BIBLIOTHEEK MATHEMATISCH CENTRUM
AMSTERDAM

Printed at the Mathematical Centre, 49, 2e Boerhaavestraat 49, Amsterdam.

The Mathematical Centre, founded the 11-th of February 1946, is a non-profit institution aiming at the promotion of pure mathematics and its applications. It is sponsored by the Netherlands Government through the Netherlands Organization for the Advancement of Pure Research (Z.W.O.), by the Municipality of Amsterdam, by the University of Amsterdam, by the Free University at Amsterdam, and by industries.

The addition formula for Jacobi polynomials.

I Summary of results (*).

Recently the author derived a Laplace integral representation, a product formula and an addition formula for Jacobi polynomials $P_n^{(\alpha, \beta)}$. The results are

$$(1) \frac{P_n^{(\alpha, \beta)}(\cos 2\theta)}{P_n^{(\alpha, \beta)}(1)} = \frac{2 \Gamma(\alpha+1)}{\sqrt{\pi} \Gamma(\alpha-\beta) \Gamma(\beta+\frac{1}{2})} \cdot$$

$$\cdot \int_{r=0}^1 \int_{\phi=0}^{\pi} (\cos^2 \theta - r^2 \sin^2 \theta + ir \cos \phi \sin 2\theta)^n \cdot (1-r^2)^{\alpha-\beta-1} r^{2\beta+1} (\sin \phi)^{2\beta} dr d\phi,$$

$$(2) \frac{P_n^{(\alpha, \beta)}(\cos 2\theta_1)}{P_n^{(\alpha, \beta)}(1)} \frac{P_n^{(\alpha, \beta)}(\cos 2\theta_2)}{P_n^{(\alpha, \beta)}(1)} = \frac{2 \Gamma(\alpha+1)}{\sqrt{\pi} \Gamma(\alpha-\beta) \Gamma(\beta+\frac{1}{2})} \cdot$$

$$\cdot \int_{r=0}^1 \int_{\phi=0}^{\pi} \frac{P_n^{(\alpha, \beta)}(2 |\cos \theta_1 \cos \theta_2 + r e^{i\phi} \sin \theta_1 \sin \theta_2|^2 - 1)}{P_n^{(\alpha, \beta)}(1)} \cdot (1-r^2)^{\alpha-\beta-1} r^{2\beta+1} (\sin \phi)^{2\beta} dr d\phi$$

and

$$(3) P_n^{(\alpha, \beta)}(2 |\cos \theta_1 \cos \theta_2 + r e^{i\phi} \sin \theta_1 \sin \theta_2|^2 - 1) =$$

$$= \sum_{k=0}^n \sum_{l=0}^k c_{n,k,l}^{(\alpha, \beta)} (\sin \theta_1 \sin \theta_2)^{k+1} (\cos \theta_1 \cos \theta_2)^{k-1} \cdot$$

(*) This report will be communicated by Professor A. van Wijngaarden to the Koninklijke Nederlandse Akademie van Wetenschappen.

$$P_{n-k}^{(\alpha+k+1, \beta+k-1)}(\cos 2\theta_1) P_{n-k}^{(\alpha+k+1, \beta+k-1)}(\cos 2\theta_2) .$$

$$P_1^{(\alpha-\beta-1, \beta+k-1)}(2r^2 - 1) r^{k-1} \frac{\beta+k-1}{\beta} C_{k-1}^\beta(\cos \phi) .$$

In formulas (1) and (2) it is supposed that

$$\alpha > \beta > -\frac{1}{2} .$$

In formula (3) the coefficient is

$$c_{n,k,l}^{(\alpha, \beta)} = \frac{(\alpha+\beta+n+1)_k (-1)^l (-n-\beta)_l}{(\beta+1)_k (\alpha+k)_l P_{n-k}^{(\alpha+k+1, \beta+k-1)}(1)} .$$

If $\beta = 0$ the factor

$$\frac{\beta+k-1}{\beta} C_{k-1}^\beta(\cos \phi)$$

in formula (3) is replaced by

$$2 \cos(k-1)\phi \quad \text{if } k-1 \neq 0 \quad \text{and by}$$

$$1 \quad \text{if } k-1 = 0 .$$

The analogous formulas in the ultraspherical case $\alpha = \beta$ are due to Gegenbauer [8] (cf. Erdélyi [3], §3.15, formulas (22), (20) and (19)). They can be obtained as degenerate cases of our formulas (1.1), (1.2) and (1.3) respectively.

It is easy to see that (2) follows from (3) by integration and that (1) follows from (2) by dividing both sides by $P_n^{(\alpha, \beta)}(\cos 2\theta_2)$ and then letting $|\cos 2\theta_2| \rightarrow \infty$.

Formula (3) was proved by the author by using group theoretical methods. For integer α the polynomials $P_n^{(\alpha, 0)}$ can be interpreted as spherical functions on the complex projective space $SU(\alpha+2)/U(\alpha+1)$ (cf. Cartan [1], [2]) or as spherical functions on the sphere $U(\alpha+2)/U(\alpha+1)$ (cf. Ikeda [9], Ikeda and Kayama [10], Ikeda and Seto [11]). In the last mentioned interpretation the classical notion

of spherical harmonics is refined for the case that the sphere is the unit sphere in a complex vector space and the symmetry group is the group of unitary transformations. The usual methods for classical spherical harmonics (cf. for instance Erdélyi [4], Ch.11, Müller [12] Vilenkin [14]) can be extended for this case. The author proved an abstract addition formula and constructed a canonical orthonormal system of generalized spherical harmonics expressed as products of certain Jacobi polynomials. Thus formula (3) was proved for $\alpha = 1, 2, \dots$ and $\beta = 0$.

By repeated differentiation with respect to $\cos \phi$ of both sides of (3) this formula can be proved for all integer α and β , such that $\alpha > \beta \geq 0$. Analytic continuation with respect to α and β finally gives the general case of formula (3). Detailed proofs will be published in one or more subsequent papers.

By the substitution

$$\cos \theta_3 e^{i\psi} = \cos \theta_1 \cos \theta_2 + \sin \theta_1 \sin \theta_2 r e^{i\phi}$$

the product formula (2) is transformed into another product formula which was obtained earlier in a different way by Gasper [7]. Gasper used his result to prove a convolution structure for Jacobi series.

Let for arbitrary complex ν and $1 \leq x < \infty$ Jacobi functions be defined as

$$R_{\nu}^{(\alpha, \beta)}(x) = {}_2F_1(-\nu, \nu + \alpha + \beta + 1; \alpha + 1; \frac{1-x}{2}) .$$

Then it follows from (1) and (2) that

$$(4) \quad R_{\nu}^{(\alpha, \beta)}(\operatorname{ch} 2t) = \frac{2 \Gamma(\alpha + 1)}{\sqrt{\pi} \Gamma(\alpha - \beta) \Gamma(\beta + \frac{1}{2})} \cdot$$

$$\cdot \int_{r=0}^1 \int_{\phi=0}^{\pi} (\operatorname{ch} t)^2 + r^2 (\operatorname{sh} t)^2 + r \cos \phi \operatorname{sh} 2t)^{\nu} \cdot$$

$$\cdot (1-r)^{\alpha-\beta-1} r^{2\beta+1} (\sin \phi)^{2\beta} dr d\phi$$

and

$$(5) \quad R_{\nu}^{(\alpha, \beta)}(\operatorname{ch} 2t_1) R_{\nu}^{(\alpha, \beta)}(\operatorname{ch} 2t_2) = \frac{2 \Gamma(\alpha+1)}{\sqrt{\pi} \Gamma(\alpha-\beta) \Gamma(\beta+\frac{1}{2})} \cdot$$

$$\cdot \int_{r=0}^1 \int_{\phi=0}^{\pi} R_{\nu}^{(\alpha, \beta)}(2 | \operatorname{ch} t_1 \operatorname{ch} t_2 + \operatorname{sh} t_1 \operatorname{sh} t_2 r e^{i\phi} |^2 - 1) \cdot$$

$$\cdot (1-r^2)^{\alpha-\beta-1} r^{2\beta+1} (\sin \phi)^{2\beta} dr d\phi$$

for $\alpha > \beta > -\frac{1}{2}$.

This is proved by analytic continuation with respect to ν where a theorem of Carlson is applied (cf. Titchmarsh [13], p.186). Formulas (4) and (5) generalize well-known results for Legendre and Gegenbauer functions (cf. [3], §3.15.2 (22) and §3.11.1 (2)).

Formulas (4) and (5) have applications for the harmonic analysis with respect to the functions $R_{\nu}^{(\alpha, \beta)}$. Flensted - Jensen who had already obtained some Paley - Wiener type theorems for this case in [5] could simplify his proofs in [6] by using formula (4). In joint work Flensted - Jensen and the author used formula (5) to develop a convolution structure for the harmonic analysis with respect to Jacobi functions. These results will also be published in a near future.

Acknowledgement

The author is due to Professor Richard Askey for suggesting him this problem. The research summarized here was done at the Institute Mittag - Leffler in Djursholm, Sweden. The author is also due to Professor Lennart Carleson for his kind hospitality.

References.

1. Cartan, E., Sur la détermination d'un système orthogonal complet dans un espace de Riemann symétrique clos, Rend. Circ. Mat. Palermo 53 (1929), 217-252.
2. Cartan, E., Leçons sur la géométrie projective complexe, Gauthier - Villars, Paris, 1931.
3. Erdélyi, A., et al., Higher Transcendental Functions, Vol.I, McGraw - Hill, New York, 1953.
4. Idem, Vol.II.
5. Flensted - Jensen, M., Paley - Wiener type theorems for a spectral decomposition of a differential operator with application to symmetric spaces. Report Institut Mittag - Leffler (Djursholm, Sweden), May 1971.
6. Idem, to appear.
7. Gasper, G., Positivity and the convolution structure for Jacobi series, Ann. of Math. 93 (1971), 112-118.
8. Gegenbauer, L., Über einige bestimmte Integrale, Wiener Sitzungsberichte 70 (1874), 433-443.
9. Ikeda, M., On spherical functions for the unitary group (I, II, III), Mem. Fac. Engineering Hiroshima Univ., 3 (1967), 17-75.
10. Ikeda, M. and T. Kayama, On spherical functions for the unitary group (IV), Mem. Fac. Engineering Hiroshima Univ. 3 (1967), 77-100.
11. Ikeda, M. and N. Seto, On expansion theorems in terms of spherical functions for the unitary group (I), Math. Jap. 13 (1968), 149-157.

12. Müller, C., Spherical harmonics, Lecture Notes in Math. 17 , Springer - Verlag, Berlin, 1966.
13. Titchmarsh, E.C., The theory of functions, Oxford University Press, second ed., 1939.
14. Vilenkin, N.J., Special functions and the theory of group representations, Am. Math. Soc. Transl. of Math. Monographs, vol.22, 1968.